



# Snow redistribution in an intermediate-complexity snow hydrology modelling framework

Louis Quéno<sup>1</sup>, Rebecca Mott<sup>1</sup>, Paul Morin<sup>1</sup>, Bertrand Cluzet<sup>1</sup>, Giulia Mazzotti<sup>1,2</sup>, and Tobias Jonas<sup>1</sup>

<sup>1</sup>WSL Institute for Snow and Avalanche Research (SLF), Davos, Switzerland

<sup>2</sup>Univ. Grenoble Alpes, Université de Toulouse, Météo-France, CNRS, CNRM, Centre d'Études de la Neige, Grenoble, France

**Correspondence:** Louis Quéno (queno@slf.ch)

Received: 8 September 2023 – Discussion started: 4 October 2023

Revised: 20 May 2024 – Accepted: 10 June 2024 – Published: 9 August 2024

**Abstract.** Snow hydrological regimes in mountainous catchments are strongly influenced by snowpack heterogeneity resulting from wind- and gravity-induced redistribution processes, requiring them to be modelled at hectometre and finer resolutions. This study presents a novel modelling approach to address this issue, aiming at an intermediate-complexity solution to best represent these processes while maintaining operationally viable computational times. To this end, the physics-based snowpack model FSM2oshd was complemented by integrating the modules SnowTran-3D and SnowSlide to represent wind- and gravity-driven redistribution, respectively. This new modelling framework was further enhanced by implementing a density-dependent layering to account for erodible snow without the need to resolve microstructural properties. Seasonal simulations were performed over a 1180 km<sup>2</sup> mountain range in the Swiss Alps at 25, 50 and 100 m resolution, using appropriate downscaling and snow data assimilation techniques to provide accurate meteorological forcing. In particular, wind fields were dynamically downscaled using WindNinja to better reflect topographically induced flow patterns. The model results were assessed using snow depths from airborne lidar measurements. We found a remarkable improvement in the representation of snow accumulation and erosion areas, with major contributions from saltation and suspension as well as avalanches and with modest contributions from snowdrift sublimation. The aggregated snow depth distribution curve, key to snowmelt dynamics, significantly and consistently matched the measured distribution better than reference simulations from the peak of winter to the end of the melt season, with improvements at all spatial resolutions. This outcome is

promising for a better representation of snow hydrological processes within an operational framework.

## 1 Introduction

Snow is a crucial water resource in mountainous areas, where snowmelt represents a significant part of the runoff (e.g. Li et al., 2017). In the context of fast and marked changes in the cryosphere and water resources in the European Alps (Beniston et al., 2018), monitoring the snow cover in mountainous countries like Switzerland is necessary to assess its contribution to the streamflow in watersheds (e.g. Griessinger et al., 2019), to estimate its response to climate change in terms of runoff (e.g. Bavay et al., 2013; Hanzer et al., 2018) or rain-on-snow events (e.g. Schirmer et al., 2022), or to better anticipate future consequences of water scarcity (e.g. Brunner et al., 2019).

When estimating the state of mountain snow cover, the main challenge is to capture its seasonal evolution and the strong spatial heterogeneity that occurs at different scales. Many studies have highlighted the benefits of using kilometre-resolution meteorological data from numerical weather prediction models as inputs for snowpack models to represent most of the sources of variability (e.g. orographic precipitation) at the mountain range scale (e.g. Vionnet et al., 2016; Quéno et al., 2016; Luijting et al., 2018; He et al., 2019; Raparelli et al., 2023). At such scales, snow redistribution can usually be considered part of the sub-grid processes.

At hectometre and finer resolutions, the fine-scale variability in snow distribution also has a significant impact on

catchment hydrology (e.g. Luce et al., 1998). Anderton et al. (2002) showed that the decametre to hectometre variability in snow cover is critical for larger-scale snowmelt runoff simulations. Several studies emphasized that the spatial distribution of snow cover prior to the melt season is more important than spatial differences in melt behaviour for estimating cumulative snowmelt dynamics in a catchment (e.g. Anderton et al., 2004; Egli et al., 2012). Brauchli et al. (2017) identified the effects of more heterogeneous snowpack on the melt season at the sub-basin scale, with an earlier onset of runoff and an extension of the melt season due to shallower and deeper snow-covered areas, respectively. Several redistribution processes contribute to the slope-scale variability: gravitational redistribution on steep slopes (e.g. Sommer et al., 2015; Mott et al., 2019), wind-driven snow transport (e.g. Pomeroy and Gray, 1995; Mott et al., 2018) and near-surface atmospheric effects on snowfall deposition patterns (e.g. Wang and Huang, 2017; Gerber et al., 2019). Sublimation of suspended snow can also have a significant local impact on the snowpack mass budget, although the overall contribution is usually small at the regional scale in alpine areas (Strasser et al., 2008; Bernhardt et al., 2012; Groot Zwaafink et al., 2013; Sexstone et al., 2018). Altogether, these redistribution processes drastically alter snow distribution, and their representation in snow cover models is crucial for snow hydrology beyond hectometre resolution (Clark et al., 2011).

Post-deposition snow redistribution processes, in particular wind-driven snow transport, have been studied for several decades (e.g. Dyunin and Kotlyakov, 1980; Föhn and Meister, 1983; Pomeroy and Gray, 1990), and many blowing snow models have been developed with a wide range of complexity depending on the study context and application. The complexity of blowing snow models can be broadly categorized according to the following three criteria.

- The three-dimensional turbulent diffusion equation can be resolved explicitly as in the snowdrift module of the Alpine3D model (Lehning et al., 2008), in the Snowdrift3D model (Schneiderbauer and Prokop, 2011), in the snow2blow model (Sauter et al., 2013), in the coupled MesoNH–Crocus model (Vionnet et al., 2014) or, with a steady-state assumption, in the PBSM-3D model (Marsh et al., 2020a). To mitigate associated high computational costs, some models alternatively use a parameterization by vertical integration, as the PBSM model (Pomeroy et al., 1993), the SnowTran-3D model (Liston et al., 2007) or, more recently, the SnowPappus model (Baron et al., 2024) have done.
- A snowpack model coupled to a snowdrift module can cover a wide range of complexity, from simple models that do not represent layer properties to detailed layered models that resolve snow microstructure. For example, studies based on SnowTran-3D (e.g. Bernhardt et al., 2009, 2010, 2012; Gascoïn et al., 2013; Sexstone et al., 2018; Reynolds et al., 2020) are embedded within the

SnowModel framework (one-layer snowpack; Liston and Elder, 2006). Recently, a Lagrangian multi-layer version of the latter model (SnowModel-LG) has been developed (Liston et al., 2020). Marsh et al. (2020a) associate PBSM-3D with the Snobal model (two-layer snowpack; Marks et al., 1999). Musselman et al. (2015) use PBSM within the Distributed Snow Model (three-layer snowpack). The aforementioned models solve the mass and energy budgets of the snowpack, hence providing snow layer properties such as density, temperature and liquid water content, but do not resolve the snow microstructure properties, unlike multi-layer models like SNOWPACK (Lehning et al., 2002), used in Alpine3D (e.g. Dadić et al., 2010; Mott et al., 2010) and CRYOWRF (Sharma et al., 2023) simulations, or Crocus (Vionnet et al., 2012) used by, e.g. Vionnet et al. (2014) or Baron et al. (2024). The latter snowpack models benefit from additional information on surface-snow properties, which can improve the determination of snow erodibility (Guyomarc'h and Mérindol, 1998; Lehning et al., 2000) compared to formulations based on air temperature (Li and Pomeroy, 1997a) or snow density (Liston et al., 2007) used with the first category of models.

- The meteorological data used to derive wind fields driving the models can vary from spatial and temporal interpolation of station measurements (e.g. Gascoïn et al., 2013), statistical or dynamical downscaling of wind fields (e.g. Reynolds et al., 2020), and deep learning methods of wind field downscaling (e.g. Le Toumelin et al., 2023) to high-resolution atmospheric models that either produce forcing fields (e.g. Bernhardt et al., 2009; Mott et al., 2010) or couple atmosphere and surface processes fully (e.g. Vionnet et al., 2014; Sharma et al., 2023).

The level of complexity adopted in studies depends on the size of the simulation area (from a few square kilometres to local mountain ranges) and study duration (from individual events to full seasons). These choices are guided by the necessity of managing computational constraints and achieving a suitable model–resource equilibrium. The present study derives its objectives and constraints from the context of the Swiss Operational Snow Hydrology Service (OSHD; Mott et al., 2023), performing physics-based snow cover simulations over a large alpine domain covering the whole of Switzerland at 250 m horizontal resolution. Snow redistribution is not currently incorporated into the model. However, users of the OSHD simulations, such as the Swiss Avalanche Warning Service, would benefit from simulations representing slope-scale variability. We investigate here the added value of modelling snow redistribution at hectometre or smaller scales in the particular framework of intermediate-complexity snowpack modelling, enabling calculations over large domains with hourly updates.


A few recent studies have explored different approaches to performing seasonal snowpack simulations, which encompass snow redistribution over large domains all while maintaining computationally viable costs. Mower et al. (2024) parallelized SnowModel, including the SnowTran-3D module, to enable distributed snow evolution simulations at 100 m horizontal resolution over the contiguous United States. Baron et al. (2024) have chosen to use a simplified one-dimensional advection–diffusion equation in their snowdrift module SnowPappus, which is coupled to the complex multi-layer snowpack model Crocus, with a target horizontal resolution of 250 m. Vionnet et al. (2021) performed distributed snowpack simulations, including parameterized gravitational redistribution (Bernhardt and Schulz, 2010) and snowdrift modelling with PBSM-3D, using a simplified three-dimensional advection–diffusion equation (Marsh et al., 2020a) with an adaptive mesh resolution (Marsh et al., 2020b). The present study introduces a different method to achieve an efficient solution: an enhanced snow cover modelling technique that comprehensively considers erodible snow layering and incorporates snow redistribution within an intermediate-complexity framework. This combination of methods offers a novel approach with the aim of facilitating operational applications over an entire mountain range throughout an entire winter season. After presenting modelling (Sect. 2) and evaluation methods (Sect. 3), the model will be assessed against spatially distributed snow depth measurements (Sect. 4.1), with quantification of the impact of redistribution on the modelled snow hydrological mass budget (Sect. 4.2). Results will be discussed in Sect. 5.

## 2 Modelling methods

### 2.1 Modelling domain

The domain used for this study covers an area of 31.6 km by 37.3 km (1178.7 km<sup>2</sup>) located in the eastern Swiss Alps around Davos (Fig. 1). This area covers a wide range of elevations (from 540 to 3417 m a.s.l.), mostly in open terrain (77 %), and includes valleys and ridges of different orientations. Forests and urbanized areas were excluded from the study to focus on redistribution processes in open terrain. The prevailing wind directions in the region range from northwest to southwest.

A digital elevation model (DEM) is generated for this domain at three different spatial resolutions (100, 50 and 25 m) over which the spatially distributed simulations are performed. The DEM is derived from the 25 m resolution DEM of the Federal Office of Topography, swisstopo. The 100, 50 and 25 m resolution domains contain 117 868, 471 472 and 1 885 888 grid points, respectively. Figure 1 also shows the evaluation sub-domains B0, D0, D1 and D2, where D1 and D2 are part of D0.



**Figure 1.** Overview of the study area: location of the domain in Switzerland (top with the domain in red, Swiss borders in orange. Source – swisstopo) and digital elevation model of the domain at 25 m resolution (bottom), with the borders of sub-domains B0, D0, D1 and D2. D1 and D2 are part of D0.

### 2.2 Snowpack modelling

FSM2 is an intermediate-complexity snowpack model (Esery, 2015; Mazzotti et al., 2020) that explicitly resolves the snowpack mass and energy balance, including fluxes between the snowpack and the atmosphere and fluxes between the snowpack and the underlying ground. However, contrary to detailed models like Crocus (Vionnet et al., 2012) or SNOWPACK (Lehning et al., 2002), it does not resolve snow microstructural properties. This model is therefore particularly suited for advanced snow hydrological simulations with a low computational cost (Magnusson et al., 2015). This is why a variant named FSM2oshd was developed and is currently used within the modelling framework of the OSHD (Mott et al., 2023). The differences between FSM2 and FSM2oshd are described in detail by Mott et al. (2023). Snow–canopy interaction processes are represented in both models (Mazzotti et al., 2020) but were not considered in the present study, which focuses only on open areas. A summary of all FSM2 variants mentioned in the present study is provided in Table 1.

**Table 1.** Summary of the different versions of FSM2 model mentioned in the study.

Model version	Description	References
FSM2	Original snowpack model	Essery (2015), Mazzotti et al. (2020)
FSM2oshd	Nationwide operational implementation of FSM2 including data assimilation	Mott et al. (2023)
FSM2ref	FSM2oshd + density-dependent layering	Present study
FSM2trans	FSM2ref + wind- and gravity-driven redistribution	Present study
FSM2trans aval.	FSM2ref + gravity-driven redistribution	Present study
FSM2trans wind	FSM2ref + wind-driven redistribution	Present study

In order to represent erosion and accumulation due to redistribution in this intermediate-complexity framework, a few modifications were implemented. Indeed, the default layering scheme of FSM2 and FSM2oshd is a fixed stratification with predefined thicknesses (Essery, 2015) independent of snow properties (a maximum of 3 layers with top layers of 10 and 20 cm in FSM2oshd), which is inherently limited in representing near-surface-snow evolution. Cristea et al. (2022) highlighted the significance of layering and the thickness of the upper layer in modelling the accumulation and melting processes. This becomes even more important for redistribution mechanisms, as snow erodibility can change significantly throughout the snowpack stratigraphy due to varying microstructural properties. Of specific importance is the erodibility of surface snow. In cases where these microstructural attributes are not directly resolved, snow density serves as the most suitable proxy for assessing erodibility. Hence we implemented a new density-dependent layering scheme, enabling finer layering near the surface (Fig. 2). Dynamic layering methods based on microstructural properties are common in complex snowpack models (e.g. Vionnet et al., 2012), but the novelty of the present method is its introduction of a simpler dynamic layering suitable for models like FSM2oshd.

The density-dependent layering is run at each time step and at each grid point. When snowdrift and avalanches are enabled, the re-layering scheme is also run after each redistribution process. It is based on the below steps and is constrained by three parameters:  $N_{\max}$ , the maximum number of layers allowed (here defined as 6);  $HS_{\min}$ , the minimum snow layer thickness allowed (here defined as 2 cm); and  $HS_{\text{fine}}$ , the maximum surface thickness where the snowpack will be finely layered (here defined as 50 cm). The layering is performed in the following sequence.


- Every time new snow accumulates, whether from snowfall, snowdrift or an avalanche, it adds a new layer to the top of the snowpack.
- All the snow deeper than  $HS_{\text{fine}}$  is moved to the basal snow layer.
- If one of the layers is thinner than  $HS_{\min}$ , it is merged with the adjacent layer that has the closest density.

- If there are more than  $N_{\max}$  layers, the two adjacent layers with the closest densities are merged.
- If the number of layers used is less than  $N_{\max}$  and sufficiently thick layers can be divided, a recursive process of splitting the thickest layers into two follows. This continues until the total layer count reaches  $N_{\max}$ , adhering to the established criteria of minimum layer thickness and maintaining a finely layered surface.

The aim of this routine is to provide fine layering near the snowpack surface to determine surface-snow conditions better, while using as many homogeneous density layers as possible to represent the snowpack historical stratigraphy without resolving the snow microstructure. Similar to the default layering routine in FSM2, mass and energy are conserved throughout the re-layering steps by tracking the thickness, ice and water content, and internal energy of all layers, with appropriate weighting when splitting or merging occurs.

Furthermore, wet or refrozen snow layers are identified as non-erodible (e.g. Li and Pomeroy, 1997b). To this end, a mechanism to keep track of past snow wetting events is implemented. The historical wetting variable  $\text{hist}_{\text{wet}}$  of a given layer is initialized as  $\text{hist}_{\text{wet}} = 0$  and set to  $\text{hist}_{\text{wet}} = 1$  when this layer reaches its maximum liquid capacity, leading to drainage. A weighted average is performed during re-layering. A layer is considered non-erodible by snowdrift when  $\text{hist}_{\text{wet}} > 0.5$  (independent of friction velocity erodibility calculations in the snowdrift module). In the example situation illustrated in Fig. 2, fresh snow accumulation creates a new top snow layer, snow deeper than  $HS_{\text{fine}}$  is added to the basal layer and as the fourth layer from the top gets too thin, it is merged with layer five, the layer with the closest density. Since layer two has been previously wetted, the erodible snow consists only of the top low-density layer.

The snowpack simulations with the new layering scheme were compared to the operational simulations with fixed layering and showed a very similar seasonal evolution, except for slightly increased settling and melting that was most likely due to the presence of finer layers. As the FSM2oshd parameters are tuned each year and the seasonal dynamics were close, this difference was not considered significant. For clarity, the reference version of FSM2oshd including the



**Figure 2.** The density-dependent layering scheme implemented in FSM2ref.

aforementioned developments specifically developed for the present study is called FSM2ref hereafter (Table 1).

### 2.3 Redistribution modelling

In the OSHD modelling framework, the spatially distributed FSM2ref model does not represent any lateral interaction between grid points in open terrain (Mott et al., 2023). Post-deposition redistribution processes were implemented within this OSHD framework. The process representation complexity needs to match the intermediate complexity of FSM2ref snowpack modelling, i.e. no snow microstructure resolution and an hourly time step to maintain an acceptable compromise for operational simulations. With that in mind, we chose to integrate and adapt SnowTran-3D (Liston and Sturm, 1998; Liston and Elder, 2006; Liston et al., 2007) for wind-driven redistribution modelling and SnowSlide (Bernhardt and Schulz, 2010) for gravity-driven redistribution modelling in a new version of FSM2ref hereafter called FSM2trans (Table 1).

#### 2.3.1 Wind-driven redistribution

SnowTran-3D is a model for wind-induced snow transport based on semi-empirical parameterizations (Liston and Sturm, 1998; Liston and Elder, 2006; Liston et al., 2007); i.e. it does not explicitly resolve the three-dimensional turbulent diffusion equation. Vertically integrated snow transport fluxes and sublimation rates are calculated in the saltation and suspension layer based on two-dimensional wind field inputs. This parameterization enables efficient computations that benefit large-scale or full-season simulations at spatial resolutions between 1 and 100 m with a large span of applica-

tions: from 2 m resolution simulations of local drift patterns around road beds in flat terrain (Liston et al., 2007) to 90 m resolution seasonal simulations in glaciated mountains (Gascoin et al., 2013). In particular, this scheme is well suited for intermediate-complexity snowpack models since the threshold friction velocity to initiate snowdrift is parameterized by a formulation that relates it to snow density through exponential laws (Liston et al., 2007). Thus, this scheme was implemented within FSM2trans, with default parameter values and without using Tabler (1975) equilibrium-snowdrift profiles, which are more suitable for higher resolutions than those used in our study (Liston et al., 2007).

We integrated a few adaptations and improvements, described below. In this new implementation, snow erosion is performed layer by layer, where the eroded snow depth is derived from the mass flux using the density of each layer rather than assuming a constant density. However, as in the original model, redeposited snow is assigned a constant density of  $300 \text{ kg m}^{-3}$ . Indeed, in the absence of snow microstructure modelling, laws parameterizing the compaction of redeposited snow during snowdrift, as proposed by Durand et al. (2001), are irrelevant. The compaction of the top snow layer under the influence of wind is taken into account, following the original SnowTran-3D parameterization. The identification of erodible snow is assessed for each layer based on existing liquid water content, past wetting history and the relevant threshold friction velocity. Consequently, the SnowTran-3D conventional two-layer concept (comprising soft surface snow and underlying hard snow) is enriched to incorporate a more sophisticated scheme, ensuring a more accurate representation of the layering within the snowpack.

#### 2.3.2 Gravity-driven redistribution

SnowSlide (Bernhardt and Schulz, 2010) is a model using a simple parameterization for gravitational snow transport. Avalanches are simulated when a slope threshold and a snow holding capacity are exceeded. The snow holding capacity is defined as a threshold in snow thickness (i.e. normal to the slope) dependent on the slope. The parameterization for snow holding capacity follows that used in the implementation of SnowSlide in the Canadian Hydrological Model (CHM; Marsh et al., 2020b). The process is solved sequentially from the highest elevation pixel to the lowest one in the domain. Snow exceeding the holding capacity is transported laterally to lower neighbouring pixels, proportional to their elevation difference. The physics of avalanches, from triggering to dynamics, is not explicitly solved. This scheme was implemented within FSM2trans with a few improvements to mitigate these limitations, in the form of simple hysteretic features described in the following. In this new implementation, the slope threshold of  $25^\circ$  is only used for avalanche triggering. However, no such threshold is applied to pixels that receive the snow released by avalanches, to enable a larger deposition area. Furthermore, the snow holding capac-

ity is decreased by 30 % for the time steps when a pixel receives an avalanche, to mimic avalanche dynamics.

In its original version, SnowSlide updates the DEM with the newly calculated snow depth at each time step, which allows us to update the slope and the order of the pixel calculations sorted by decreasing elevations. This version has been tested with no significant visible differences in avalanche deposition areas. However, the calculation of the new sorted elevation list at each time step has a high computational cost. Consequently, we decided to discard this step with a view to intermediate-complexity modelling applicable to operations. The implementation of the hysteretic features showed a more significant impact on the avalanche extent.

### 2.3.3 Integration of the redistribution sub-models in FSM2trans

The adapted SnowTran-3D and SnowSlide sub-models are integrated as subroutines within FSM2trans. After solving the one-dimensional snowpack processes (heat conduction, melting, sublimation, water percolation, compaction and fresh snow addition), the layering subroutine is called, followed by wind-driven redistribution, re-layering, gravity-driven redistribution and a final re-layering. SnowSlide follows SnowTran-3D, as snowdrift occurs at the exact time step (under the given snow and meteorological conditions), while avalanches may actually occur after a certain delay. It enables us to take into account potential avalanche triggering on slopes loaded by snowdrift and to avoid the immediate gravitational transport of fresh snow to lower elevations where snowdrift is less likely to happen.

Each sub-model can be activated independently via a name list. The FSM2trans model version with only gravity-driven redistribution is called FSM2trans aval., while the version with only wind-induced redistribution is called FSM2trans wind (Table 1).

## 2.4 Meteorological input

FSM2ref and FSM2trans require several meteorological inputs for the simulated grid points at an hourly time step: near-surface air temperature, relative humidity and wind speed (at a defined height above the ground); longwave irradiance; direct and diffuse shortwave irradiance; air pressure; rainfall; and snowfall. The snowdrift module of FSM2trans also requires wind direction. Similar to the operational version FSM2oshd (Mott et al., 2023), these fields are primarily derived from the hourly analysis fields of the regional weather forecast model COSMO with a spatial resolution of 1 km. To enhance the effect of fine-scale topographical influences, this information is subsequently downscaled to 100, 50 and 25 m spatial resolution. In particular, near-surface air temperature, relative humidity and air pressure are downscaled by linear interpolation with lapse rates. Direct and diffuse shortwave irradiances are dynamically downscaled follow-

ing the approach of Jonas et al. (2020). Longwave irradiance downscaling follows Helbig and Löwe (2014). Snow depth measurements at stations are assimilated to improve the solid precipitation estimate through a data assimilation scheme using optimal interpolation (Magnusson et al., 2014). Total precipitation is then linearly interpolated to the finer grid where the phase split between rain and snow is made according to the downscaled near-surface air temperature field, following the same formulation as FSM2oshd using a sigmoid function centred on a 10 m air temperature of 1.04 °C.

Snowdrift is strongly determined by local topographic effects on wind fields (e.g. Mott et al., 2018). Consequently, the downscaling of wind patterns must encompass the interplay between terrain and airflow dynamics, such as local acceleration and deceleration near ridges or channelling in valleys and gullies. For this purpose, we used the mass-conserving dynamical downscaling model WindNinja (Forthofer et al., 2014; Wagenbrenner et al., 2016) version 3.7.0, which was forced by COSMO 1 km resolution wind fields. WindNinja was run separately at 100, 50 and 25 m spatial resolution to produce downscaled wind fields for the snowpack simulations, providing horizontal wind speed and direction. Similar to Vionnet et al. (2021), the mass- and momentum-conservation option (Wagenbrenner et al., 2019) was not free of model instabilities in study areas with complex topography and was therefore not retained.

## 2.5 Simulation setup

Spatially distributed seasonal snowpack simulations were performed with FSM2ref and FSM2trans at hourly time steps over the entire study area from 1 September 2016 to 30 June 2017 and from 1 September 2019 to 30 June 2020, at a resolution of 25, 50 and 100 m. To further assess the distinct effects of avalanche and snowdrift modelling, complementary simulations were performed in a similar setup with FSM2trans aval. and FSM2trans wind.

## 3 Evaluation data and methods

Four distinct datasets of distributed snow depth measurements were used to evaluate the snowpack simulations. These measurements were acquired by airborne lidar technology. During the 2016–2017 hydrological year, three aerial surveys covered a region centred on the Dischma valley near Davos, with a spatial resolution of 1 m. To exclude forests and urbanized areas, we filtered out elevations under 2000 m a.s.l. (corresponding to the treeline), since the forest snow model instance of FSM2oshd was not used in the present study. The resulting sub-domain is named D0 (Fig. 1). The airborne surveys took place on three dates: 20 March 2017, which marked the transition from winter to the melt season; 31 March 2017; and 17 May 2017, the latter two dates covering the melting period. These datasets were

validated against more than 11 000 manual measurements, resulting in a bias of  $-4$  to  $0$  cm and a root mean square deviation (RMSD) of  $4$  to  $8$  cm (Mazzotti et al., 2019). During the 2019–2020 hydrological year, one airborne survey covered an area east of Lenzerheide with a spatial resolution of  $1$  m. Similarly, elevations under  $2000$  m a.s.l. were filtered out, resulting in sub-domain B0 (Fig. 1). The flight was conducted on 17 March 2020 (at the transition between winter and melt season). This dataset was validated against 79 manual measurements, resulting in a bias of  $-2$  cm and a RMSD of  $15$  cm.

The datasets were post-processed to mask out lakes and a few obvious outliers (e.g. those due to buildings or in very steep terrain). The resulting snow depth maps were then aggregated to a resolution of  $25$ ,  $50$  and  $100$  m. As the outliers were mostly below  $2000$  m a.s.l. and very isolated above  $2000$  m a.s.l., the masked data at high resolution were simply excluded from the mean over aggregated pixels. In a second step, glacier masks were applied to the averaged snow depth maps.

Simulation results were extracted from the whole domain and from the masked lidar sub-domains B0, D0, D1 and D2 (Fig. 1). Modelled snow depth maps were compared to lidar snow depth maps upscaled to the model grid at  $25$ ,  $50$  and  $100$  m resolution. Beyond visual comparison, results were aggregated by elevation bands and aspects to better identify distribution patterns. In addition, results were aggregated by topographic position index (TPI), a parameter that characterizes the relative height of a point in relation to its local surroundings. It was calculated using a  $25$  m resolution DEM with a  $2$  km radius neighbourhood. This terrain descriptor is particularly suited for a focused analysis of areas most susceptible to wind exposure; it is often used in wind down-scaling methods, e.g. by Winstral et al. (2017) and Dujardin and Lehning (2022). For this purpose, we defined a “ridges” category for pixels with a TPI exceeding  $200$  m. This classification facilitates specific analysis of these areas of interest.

To complement the visual comparison of simulated and measured snow depth maps, we introduced a quantitative approach based on the structural similarity index (SSIM; Wang et al., 2004). This metric was originally developed for image quality assessment to quantify the similarity between a distorted (e.g. compressed) image and a reference image. It is a combination of luminance, contrast and structure comparison. Pixels are compared with their neighbourhood using a Gaussian weighting function. We apply it to snow depth maps by considering them as greyscale images, where the snow depth is the intensity on a scale from  $0$  to  $5$  m, using a Gaussian radius of  $150$  m. The mean SSIM (MSSIM) indicates the overall similarity of the two snow depth maps. A random snow depth distribution in the  $0$  to  $5$  m range gives  $MSSIM = 0$ . Values closer to  $1$  indicate better similarity, and  $MSSIM = 1$  if and only if the snow depth maps are identical. The structural similarity index can be computed using the `ssim` function in the Matlab Im-

age Processing Toolbox (<https://www.mathworks.com/help/images/ref/ssim.html>, last access: 8 May 2024) or using the `scikit-image` library in Python ([https://scikit-image.org/docs/stable/auto\\_examples/transform/plot\\_ssim.html](https://scikit-image.org/docs/stable/auto_examples/transform/plot_ssim.html), last access: 8 May 2024).

## 4 Results

### 4.1 Comparison of simulated snow depth to lidar data

Initially, we present snow depth maps derived from FSM2trans simulations, offering a comparative analysis versus both lidar datasets and reference FSM2ref simulations that do not incorporate redistribution effects. This analysis illustrates the capabilities of FSM2trans in representing specific redistribution patterns at different stages of the snow season, from the peak of accumulation to the melt season. These maps are presented for sub-domains to assess local patterns with more clarity. Figure 3 shows the snow depth map for sub-domain B0 on 17 March 2020, while Fig. 4 shows sub-domain D2 on 31 March 2017 and Fig. 5 sub-domain D1 on 17 May 2017. We highlighted specific erosion and accumulation patterns on the maps with red arrows.

A necessary element for the interpretation of these maps is the elevation profile of snow depth at the peak of winter for sub-domain D0 (Fig. A1 in the Appendix). It shows a marked underestimation of snow depth by FSM2ref compared to the lidar at elevations above  $2600$  m a.s.l. despite the absence of redistribution and snowdrift sublimation that would further reduce mean snow depths at the highest elevations. This suggests a significant underestimation of solid precipitation at high elevations in the forcing of FSM2ref and FSM2trans, which is consistent with the findings of Mott et al. (2023).

Although FSM2trans produces too little snow at the highest elevations, the smaller-scale snow distribution patterns match observations well. In Fig. 3, arrow 1 indicates three avalanches following steep and narrow gullies. These avalanches have been simulated well by FSM2trans, and in particular, their deposition areas show a good match with observations. Arrow 2 shows another avalanche that was successfully simulated. However, the deposition area is underestimated by FSM2trans. This underestimation of avalanche area can be noted for several other cases. The location of simulated high snow accumulation behind ridges exhibits remarkable overall agreement with the observed data. This accumulation results from a combination of snowdrift deposition and subsequent avalanche initiation within these strong accumulation zones in steep terrain. Such cases are highlighted, for example, by arrows 3 and 4 (Fig. 3), arrows 6 and 7 (Fig. 4), and arrows 8 and 9 (Fig. 5). The particular example of arrow 6 indicates an accurate location of a large accumulated snow mass transported by wind to the north-eastern lee side of the ridge and carried by avalanches to lower elevations extending further than the immediate foot

of the steep slopes. However, the extension of the deposition areas in the simulations sometimes remains too limited, with uncertainty in attributing this shortcoming to the snowdrift model or the avalanche model (e.g. arrows 4 and 7). These results are consistent throughout the season, with deposition patterns being particularly visible in spring (Fig. 5). High-elevation ridges show strong erosion patterns in the simulations, which are consistently overestimated compared to the observations. Strong variability in intermediate slopes is sometimes underestimated by the model (e.g. arrow 5 in Fig. 3). The high local variability is still partially represented on a high elevation pass particularly exposed to wind (arrow 10 in Fig. 5).

The added value of the FSM2trans representation of redistribution processes is particularly noteworthy when compared to the maps resulting from FSM2ref simulations (inset panels in Figs. 3, 4 and 5). The snow depth simulated by FSM2ref above 2000 m a.s.l. is notably homogeneous at the end of winter (Fig. 3), with variability introduced mainly throughout the melt season due to differences in melt energy between slopes (Fig. 5). These maps confirm that simulations that do not include redistribution processes cannot represent a significant part of the snowpack spatial variability at 25 m resolution, even if Mott et al. (2023) showed that such simulations (performed with the FSM2oshd variant) captured the average state of the snowpack well when compared to station measurements at all elevation bands, except for the highest elevations where snow measurements and data assimilation were lacking.

Figure 6 shows MSSIM values of snow depth maps simulated by FSM2trans, FSM2trans aval., FSM2trans wind and FSM2ref compared to the four lidar datasets. Simulations at a resolution of 25, 50 and 100 m are represented by increasing transparency (darkest shading for 25 m). At 25 m resolution, the highest similarity is obtained for FSM2trans simulations (0.39 to 0.45) and the lowest similarity for FSM2ref simulations (0.14 to 0.18). FSM2trans always has a higher MSSIM than FSM2trans aval. (0.35 to 0.38) and FSM2trans wind (0.19 to 0.24), highlighting the importance of modelling the interplay of avalanches and snowdrift. The MSSIM is higher when only avalanches are represented than when only snowdrift is represented; in a domain with a lot of steep terrain, simulating avalanches is easier because they are confined to steep slopes, whereas snowdrift is more widespread and hence has a less obvious spatial structure. At coarser resolutions, the MSSIM increases for FSM2ref simulations: the reference lidar map is smoother, so the similarity to simulations with less spatial variability increases. The MSSIM of FSM2ref remains clearly inferior to all other simulations representing one or both redistribution processes at all resolutions, proving that there are still clear benefits to modelling redistribution even at 100 m resolution. Following the same logic as FSM2ref, the MSSIM of FSM2trans wind increases with coarser resolutions. However, the MSSIM of FSM2trans aval. decreases from 50 to 100 m resolution, reaching values

similar to those of FSM2trans wind: with smoother terrain, some avalanche couloirs are not represented anymore and slopes are lower, hence less avalanche triggering occurs.

Figure 7 shows the snow depth distribution on sub-domains B0 and D0 for the four lidar datasets and the snow depth distribution simulated by FSM2ref and FSM2trans. These plots quantitatively confirm the visual assessment of redistribution patterns. In FSM2ref (blue curve), a frequency peak is observed in all plots throughout the season, even if the distribution gets flatter with differential melting during spring. In particular, compared to lidar observations (black curve), low snow depths are underrepresented and high snow depths (typically more than 2 m) are absent. FSM2trans (red curve) clearly improves the snow depth distribution, with a flatter curve matching the lidar curve better. Low and high snow depths are better represented, even though the spread remains lower than in observations. Discrepancies compared to observations are further influenced by uncertainties in precipitation input and modelling of compaction and melting processes.


In order to focus on the most wind-exposed areas, Fig. 8 represents the same frequency plots restricted to the areas where  $TPI > 200$  m, i.e. ridges and their surroundings. The match of FSM2trans to the lidar is even better than when all TPIs are considered, with a clear improvement compared to FSM2ref. In the four cases, shallow snowpacks are slightly overrepresented by FSM2trans, which confirms the visual observation of excessive ridge erosion.

FSM2trans simulations at 50 and 100 m grid resolution have also been assessed. Corresponding maps and plots are included in Appendix B and C. Despite the lower spatial resolution, redistribution patterns remain consistent between resolutions and compared to the lidar data. Fine-scale redistribution patterns are logically less present due to the absence of fine-scale terrain features such as narrow gullies channelling avalanches. However, even at 100 m resolution, the snow depth frequency curve shows a similar improvement to the higher resolution simulations compared to FSM2ref (Fig. 9), which confirms the outcomes of the structural similarity analysis.


## 4.2 Cumulated effect of snow redistribution

In this analysis, we examine the cumulative effects of each redistribution process over the course of a winter season, aiming to identify their individual contributions to the snowpack dynamics while also exploring their interrelationships.


Figure 10 represents the net seasonal effect of four distinct processes on snow water equivalent (SWE) within the designated sub-domain D1, spanning the period from 1 September 2016 to 30 June 2017. These processes encompass (a) saltation and suspension, (b) avalanches, (c) snowdrift sublimation, and (d) surface sublimation due to turbulent fluxes. Avalanches locally contribute very strongly to SWE change (up to 1000 mm gain or loss). The extension of avalanche




**Figure 3.** Maps of snow depth on 17 March 2020 for sub-domain B0 (a) as measured by lidar and aggregated to 25 m resolution, (b) as simulated at 25 m resolution by FSM2trans, and (c) as simulated at 25 m resolution by FSM2ref. (d) Indicative digital elevation model (DEM) of sub-domain B0 with higher elevations in lighter grey.



**Figure 4.** Maps of snow depth on 31 March 2017 for sub-domain D2 (a) as measured by lidar and aggregated to 25 m resolution, (b) as simulated at 25 m resolution by FSM2trans, and (c) as simulated at 25 m resolution by FSM2ref. (d) Indicative DEM of sub-domain D2 with higher elevations in lighter grey.




**Figure 5.** Maps of snow depth on 17 May 2017 for sub-domain D1 (a) as measured by lidar and aggregated to 25 m resolution, (b) as simulated at 25 m resolution by FSM2trans, and (c) as simulated at 25 m resolution by FSM2ref. (d) Indicative DEM of sub-domain D1 with higher elevations in lighter grey.



**Figure 6.** MSSIM for simulation results from FSM2trans (red), FSM2trans aval. (yellow), FSM2trans wind (grey) and FSM2ref (blue) versus the four lidar datasets. Lighter shades of colour indicate lower resolutions (25 m for the darkest shading, 50 m for the intermediate shading and 100 m for the lightest shading).

effects is, however, spatially restricted to steep slopes and their immediate surroundings. The cumulative impact resulting from snowdrift-induced saltation and suspension can lead to gains or losses exceeding 500 mm. Notably, these substantial changes in SWE are most pronounced upon or in close proximity to ridgelines. The extension of saltation and suspension effects is still more widespread than avalanches, with changes in SWE also occurring on intermediate slopes, but very limited down the valleys (typically less than 10 mm).

The location of snow accumulation due to saltation and suspension is consistent with the prevailing northwest to southwest wind directions, with more frequent net SWE gain on the eastern side of the ridges. Strong southwesterly Foehn wind events can also explain the erosion on the southern slopes. SWE losses due to snowdrift sublimation cover a similar spatial extent, with a lower quantitative impact (up to 100 mm on the ridges). When compared to surface sublimation due to turbulent fluxes, snowdrift sublimation reaches




**Figure 7.** Snow depth frequency plots in intervals of 10 cm at 25 m resolution for lidar measurements (black), FSM2trans simulation (red) and reference FSM2ref simulation without redistribution (blue) for sub-domains (a) B0 on 17 March 2020, (b) D0 on 20 March 2017, (c) D0 on 31 March 2017 and (d) D0 on 17 May 2017.


higher extreme values locally, while surface sublimation is more intense in valleys and at low elevations. Surface deposition dominates at high elevations (apart from ridges), which tends to partly compensate for snowdrift sublimation losses.

Figure 11 illustrates the proportional influence of each individual process in relation to the total snowfall occurring between 1 September 2016 and 30 June 2017. Values are aggregated over the whole domain, encompassing 8 aspects and 12 elevation bands, each spanning a 100 m range above 2000 m a.s.l. In the most wind-exposed regions, specifically the northwestern to southwestern aspects at higher elevations, a combination of saltation and suspension processes contribute on average to a loss of approximately 50 % of the total snowfall. On wind-sheltered aspects, accumulation due to saltation and suspension represent on average up to approximately 25 % of the total snowfall. The areas with the strongest avalanche erosion are located preferentially in areas where snowdrift accumulation is prominent: it reaches on average approximately 50 % of the total snowfall there. Avalanche deposits are more widespread across elevations, so their average represents approximately 5 % of the total snowfall. Snowdrift sublimation shows an increasing trend with elevation, independently of the aspect, and reaches on average up to 5 % of the total snowfall at the highest elevations. Ridges show extreme snowdrift sublimation values

locally of 20 % of the total snowfall. The areas of strongest snowdrift sublimation correspond to areas with the lowest intensity of surface sublimation or deposition due to turbulent fluxes (deposition on high-elevation northern slopes is up to 2 % of the total snowfall), while surface sublimation can reach up to 8 % of the total snowfall at lower elevations. When considering all elevations in the whole domain, snowdrift sublimation represents a 1.0 % loss of the total snowfall in 2016–2017 (1.4 % in 2019–2020), which is a smaller contribution than surface sublimation loss (4.3 % in 2016–2017 and 3.5 % in 2019–2020). These contributions become similar when considering only elevations above 2000 m a.s.l.: 1.8 % in 2016–2017 and 2.3 % in 2019–2020 for snowdrift sublimation and 2.9 % in 2016–2017 and 2.1 % in 2019–2020 for surface sublimation. Snowdrift sublimation significantly dominates at elevations above 3000 m a.s.l.: 4.4 % in 2016–2017 and 3.9 % in 2019–2020 versus 0.2 % in 2016–2017 and 0.5 % in 2019–2020 for surface sublimation.



**Figure 8.** Snow depth frequency plots in intervals of 10 cm at 25 m resolution for lidar measurements (black), FSM2trans simulation (red) and reference FSM2ref simulation without redistribution (blue) for areas where TPI > 200 m for sub-domains (a) B0 on 17 March 2020, (b) D0 on 20 March 2017, (c) D0 on 31 March 2017 and (d) D0 on 17 May 2017.




**Figure 9.** Snow depth frequency plots in intervals of 10 cm for sub-domain B0 on 17 March 2020 for lidar measurements (black), FSM2trans simulation (red) and reference FSM2ref simulation without redistribution (blue) at (a) 25 m resolution, (b) 50 m resolution and (c) 100 m resolution.

## 5 Discussion


### 5.1 Added value of snow redistribution modelling

We have implemented a modelling approach to capture wind- and gravity-driven snow transport within a spatially distributed snow cover framework used for operational snow hydrological modelling. Given that the primary objective of this implementation was to improve snow distribution patterns, we conducted a comprehensive evaluation by comparing it with lidar snow depth maps. The outcomes of this evalua-

tion reveal notable enhancement in the ability of FSM2trans to accurately depict the formation of strong snow accumulation positioned on the correct slopes of ridges. In contrast, FSM2ref represents a very homogeneous snowpack across elevations exceeding 2000 m a.s.l. with no aspect differences apart from differentiated melting (Figs. 3, 4 and 5). Certain accumulation and erosion patterns can be clearly attributed to either snowdrift or avalanches. Examples include the marked erosion of ridges due to strong winds and the concentration of snow slides within steep couloirs. However, it is important to acknowledge the interplay between snowdrift and



**Figure 10.** Map of net SWE change cumulated from 1 September 2016 to 30 June 2017 for sub-domain D1 as simulated by FSM2trans due to (a) saltation and suspension, (b) avalanches, (c) snowdrift sublimation, and (d) surface sublimation. Note that the colour axes differ between panels.



**Figure 11.** Net SWE change relative to total snowfall cumulated from 1 September 2016 to 30 June 2017 as simulated by FSM2trans, aggregated for the whole domain by aspect and 100 m elevation band above 2000 m a.s.l. due to (a) saltation and suspension, (b) avalanches, (c) snowdrift sublimation, and (d) surface sublimation. Note that the colour axes differ between panels.

avalanches; for example, strong accumulation due to wind-induced transport can create an overload that triggers or enhances an avalanche, as can be seen on many of the upper slopes of the maps. This interdependence justifies the need to combine snowdrift and avalanche modelling and a global assessment of all redistribution processes when compared to spatially distributed snow depth measurements. The structural similarity analysis of snow depth maps confirmed and quantified increased spatial similarity to lidar data for both avalanche and snowdrift modelling, with the best improvement when both processes are combined.

Snow depth frequency plots highlight the strong added value of FSM2trans, becoming particularly evident from the onset of the melt season onward. This added value is most prominently observed on the upper slopes. The better match of the FSM2trans snow depth distribution curve with the lidar data at the peak of winter (Fig. 7a and b) is significant progress towards a better determination of catchment snow hydrological regimes, as inferred by e.g. Anderton et al. (2004), Egli et al. (2012) and Brauchli et al. (2017). It is confirmed by the persistence of this improvement in late melt season curves (Fig. 7d), when the spatial fluctuations in melt energy become a critical element for progressive snow cover depletion (DeBeer and Pomeroy, 2017). The spatial hetero-

geneity of melt energy is represented in our model through a fine downscaling of the meteorological input (in particular incoming radiation) and a representation of the effects of terrain features (e.g. slope, aspect and surface roughness) on the snowpack energy budget. The addition of snow redistribution allows for the presence of locally strong snow accumulation and eroded areas that further influence the timing of snowmelt.

The most realistic redistribution patterns were obtained with 25 m resolution simulations. It is important to note, however, that a significant positive influence on snow distribution remained even at 50 m and 100 m resolution, as quantified by the structural similarity analysis. This result is encouraging for the potential application of such modelling approaches in an operational framework as it allows the inclusion of redistribution effects in a computationally efficient manner, enabling large-domain or ensemble simulations that are often required in the context of data assimilation. For studies focusing on specific limited areas of interest, conducting simulations at 25 m resolution could still be viably executed by employing nested runs within larger-scale simulations. In that regard, the recent development of an unstructured triangular mesh (Marsh et al., 2018) within a snowdrift-

resolving snowpack model (Marsh et al., 2020a, b; Vionnet et al., 2021) offers a promising alternative solution.

Ultimately, the FSM2trans modelling framework retains its efficiency in operational contexts, requiring only a marginal increase in computational time compared to FSM2ref. For reference, performing a full seasonal simulation (i.e. 10 months from 1 September to 30 June) for the present study domain (Fig. 1) at hourly time steps on a personal computer without code parallelization yields the following approximate time frames.

- 100 m resolution: 1 h 40 min for FSM2trans (+10 % compared to FSM2ref).
- 50 m resolution: 6 h 20 min for FSM2trans (+20 % compared to FSM2ref).
- 25 m resolution: 25 h 30 min for FSM2trans (+30 % compared to FSM2ref).

The relatively modest increase in computational time can be explained by the fact that the modelling chain is highly computationally bound by input and output processing steps. The computation times of FSM2trans are less than the computation times required to generate downscaled wind fields using WindNinja, parallelized over 8 cores (a total of 91 h yr<sup>-1</sup> including all three resolutions).

## 5.2 Limitations

Model assessment has pointed to a number of limitations. These limitations mostly stem from the necessary trade-offs and adjustments inherent to accommodating an intermediate-complexity framework for snow cover modelling.

First, the redistribution patterns show some shortcomings that persist across all dataset comparisons. Both maps and distribution curves consistently reveal excessive snow erosion along ridges. A first explanation is the effect of grid resolution on snowdrift modelling. Small-scale topographic features locally enable the retention of snow in saltation on windward slopes, which cannot be represented at resolutions of 25 m or coarser (Mott and Lehning, 2010). Moreover, in the absence of sub-grid parameterization, the extent of strong ridgeline erosion is likely overestimated since the modelled ridgeline pixel is 25, 50 or 100 m wide. Previous studies have also demonstrated that blowing snow models reliant on two-dimensional wind inputs, such as SnowTran-3D, are notably influenced by the specifics of the input wind fields (e.g. Musselman et al., 2015). Consequently, the observed excessive snow erosion on the ridges could be attributed to WindNinja's simulation of strong wind speeds. A preliminary evaluation of WindNinja wind speeds for the month of March 2017 against measurements from 13 automated weather stations positioned across the study domain, mostly at high elevations, showed a positive bias of 1.1 ms<sup>-1</sup>. Given WindNinja's established proficiency at accurately capturing ridge accelerations (Forthofer et al., 2014), it is plausible that

this overestimation is even more pronounced at ridge tops. Moreover, the absence of momentum conservation in the version of WindNinja used in this study (Sect. 2.4) prohibits the modelling of lee-side recirculation (Wagenbrenner et al., 2016), with consequences for blowing snow redistribution patterns. The observed deficiency in the extent of snowdrift deposition behind ridges could be attributed to the omission of these crucial terrain-induced influences on wind fields. To tackle that issue, Vionnet et al. (2021) used wind libraries in conjunction with WindNinja (Marsh et al., 2023), introducing a modification to mitigate wind speeds on lee sides. Upcoming research with FSM2trans will address the sensitivity of redistribution modelling to wind downscaling techniques.

Secondly, errors in meteorological inputs propagate to the ultimate snow distribution. For example, the lack of snow on ridges may also partly arise from the lack of simulated precipitation at high elevations. This can be attributed to the fact that the precipitation input derived by optimal interpolation has been effective in mitigating the COSMO precipitation forecast bias; however, it still retains a negative bias above about 2500 m a.s.l. (Mott et al., 2023) where the assimilated station data become sparse. Moreover, near-surface winter precipitation processes such as preferential deposition of snowfall (Lehning et al., 2008) were not accounted for in the present study despite their significant impact near the ridges (Gerber et al., 2019) and subsequent effect on post-deposition processes. All in all, the use of a high-resolution atmospheric model may be necessary to further improve the precipitation input at such resolutions in complex terrain. The atmospheric downscaling model HICAR (Reynolds et al., 2023), recently developed with a focus on computational efficiency in complex terrain, is a promising solution to provide precipitation and wind field inputs for FSM2trans that better account for complex atmosphere–topography interactions.

Finally, the coordinated use of modified versions of SnowTran-3D and SnowSlide within our modelling framework has shown satisfactory outcomes with regard to the initial objective of improving the heterogeneity of the snow cover distribution, outweighing the shortcomings resulting from the simplified parameterizations in each model. If spatially distributed snow depth measurements (e.g. acquired by airborne lidar surveys) enable us to evaluate the position and extent of erosion and deposition zones, more validation data are needed to assess the amount of snow lost by sublimation in the suspension layer. Our results (an average loss of approximately 1 %, reaching on average 4 % at high elevations) are consistent with previous studies based on SnowTran-3D in alpine terrain, e.g. 4.1 % for Strasser et al. (2008), 1.6 % for Bernhardt et al. (2012) and 3.4 % for Sexstone et al. (2018). However, the parameterization of SnowTran-3D does not model the atmospheric feedback due to the latent-heat exchange of snowdrift sublimation. Groot Zwaafink et al. (2013) showed that the feedback processes largely reduce the snowdrift sublimation to 0.1 %. Thus, the lack of atmo-

spheric feedback could explain extreme sublimation values on ridges (Sect. 4.2), contributing to their over-erosion. A partial coupling of FSM2trans with an atmospheric model such as HICAR (Reynolds et al., 2023) can be considered to further investigate this effect.

The most obvious limitation of SnowSlide is its non-dynamic representation of avalanche processes, which makes it challenging to model large deposit areas due to big avalanches, although the new hysteretic features we have introduced partly mitigate that issue. Figure D1 in the Appendix shows the snow depth distribution on slopes steeper than 40° in the lidar data, in FSM2ref and in FSM2trans. The average snow depth on steep slopes is improved (roughly divided by 2, matching the lidar average), but the variability is degraded: strong accumulation is no longer possible on steep slopes, while the roughness of steep rocky faces sometimes allows snow to be retained (Sommer et al., 2015).

## 6 Conclusions


The modelling of snow redistribution induced by wind or gravity becomes necessary at hectometre and finer resolutions in order to better represent the resulting snowpack heterogeneity, which strongly influences the snow hydrological regimes in mountainous catchments. This study presents a new strategy developed in the Swiss operational snow hydrology modelling framework to address this issue, aiming at an intermediate-complexity solution to best represent the processes while maintaining operationally viable computational times.

The present work offers a novel combination of approaches compared to existing models. It builds on the existing physics-based snowpack model FSM2oshd (Mott et al., 2023). A new density-dependent layering was included to represent the snowpack stratigraphy more realistically without resolving its microstructure, providing in particular a finer layering at the surface to determine erodible snow. These developments allowed for the inclusion of redistribution modules adapted to the new layering features. Wind-induced snow transport and sublimation were modelled by the snowdrift module SnowTran-3D (Liston et al., 2007). The avalanche module SnowSlide (Bernhardt and Schulz, 2010) was also included to represent gravity-induced snow transport, with the addition of simple hysteretic features to enable more realistic runout distances. Meteorological input fields were downscaled from a weather forecast model using the dynamical downscaling model WindNinja (Forthofer et al., 2014) for wind fields. The simulations of the new FSM2trans model at 25, 50 and 100 m resolution were compared to four spatially distributed snow depth datasets acquired by airborne lidar surveys in order to assess the added value of redistribution modelling for capturing catchment snowpack heterogeneity.

The FSM2trans snow depth maps showed a remarkable improvement in the representation of strong snow accumulation resulting from the interplay of snowdrift and avalanche processes, in terms of deposit positions and amounts. This improvement was quantified by an original structural similarity analysis of snow depth maps. The erosion and deposition areas were generally captured well in terms of aspect and slope. The main shortcomings were identified as an overestimation of ridgetop erosion and an underestimation of the extent of depositional areas. Saltation and suspension transport as well as avalanches were shown to be major contributors to the mass budget on the most wind-exposed slopes and at high elevations. Snowdrift sublimation had a much smaller overall effect, except for a locally significant contribution to ridgetop erosion. The snow depth distribution plots confirmed a significant enhancement in the variability compared to the reference simulations, with the FSM2trans distribution curve consistently matching the measured distribution curve better from the peak of winter to the end of the melt season. As the snow depth distribution curve is a key control of snowmelt dynamics, this is a promising outcome to better represent snow hydrological processes at the catchment scale. Further research should quantify the actual impact of redistribution on modelled catchment snowmelt runoff.

The most realistic snow distribution patterns were obtained at 25 m resolution, but redistribution at 50 and 100 m resolution also had a positive effect on the snow distribution, making our approach viable for operational applications over large extents that cannot afford resolutions as high as 25 m. These model developments have limited computational impact and remain feasible within an operational framework. The possibility of practical application at the national scale still needs to be clarified. Finally, further enhancements of the representation of physical processes to mitigate current modelling limitations can still be achieved within the current intermediate-complexity framework. An atmospheric downscaling model like HICAR (Reynolds et al., 2023) could provide precipitation fields accounting for local terrain effects and wind fields accounting for lee-side recirculation, as well as potential atmospheric feedback on snowdrift sublimation, at reasonable computational costs. Associated with the representation of forest snow processes (Mazzotti et al., 2020), such studies show that the representation of physical processes can be implemented in operational setups with significant benefits.


## Appendix A: Elevation profile of snow depth




**Figure A1.** Elevation profile of mean snow depth on 20 March 2017 in 50 m elevation bands for sub-domain D0 for lidar measurements (black) and FSM2ref simulations (blue).

## Appendix B: Simulated snow depth maps at different spatial resolutions


## B1 Maps at 50 m resolution



**Figure B1.** Maps of snow depth on 17 March 2020 for sub-domain B0 (a) as measured by lidar and aggregated to 50 m resolution, (b) as simulated at 50 m resolution by FSM2trans, and (c) as simulated at 50 m resolution by FSM2ref. (d) Indicative DEM of sub-domain B0 with higher elevations in lighter grey.




**Figure B2.** Maps of snow depth on 31 March 2017 for sub-domain D2 (a) as measured by lidar and aggregated to 50 m resolution, (b) as simulated at 50 m resolution by FSM2trans, and (c) as simulated at 50 m resolution by FSM2ref. (d) Indicative DEM of sub-domain D2 with higher elevations in lighter grey.




**Figure B3.** Maps of snow depth on 17 May 2017 for sub-domain D1 (a) as measured by lidar and aggregated to 50 m resolution, (b) as simulated at 50 m resolution by FSM2trans, and (c) as simulated at 50 m resolution by FSM2ref. (d) Indicative DEM of sub-domain D1 with higher elevations in lighter grey.


## B2 Maps at 100 m resolution



**Figure B4.** Maps of snow depth on 17 March 2020 for sub-domain B0 (a) as measured by lidar and aggregated to 100 m resolution, (b) as simulated at 100 m resolution by FSM2trans, and (c) as simulated at 100 m resolution by FSM2ref. (d) Indicative DEM of sub-domain B0 with higher elevations in lighter grey.




**Figure B5.** Maps of snow depth on 31 March 2017 for sub-domain D2 (a) as measured by lidar and aggregated to 100 m resolution, (b) as simulated at 100 m resolution by FSM2trans, and (c) as simulated at 100 m resolution by FSM2ref. (d) Indicative DEM of sub-domain D2 with higher elevations in lighter grey.




**Figure B6.** Maps of snow depth on 17 May 2017 for sub-domain D1 (a) as measured by lidar and aggregated to 100 m resolution, (b) as simulated at 100 m resolution by FSM2trans, and (c) as simulated at 100 m resolution by FSM2ref. (d) Indicative DEM of sub-domain D1 with higher elevations in lighter grey.

### Appendix C: Snow depth frequency plots at different spatial resolutions




**Figure C1.** Snow depth frequency plots at intervals of 10 cm at 50 m resolution for lidar measurements (black), FSM2trans simulation (red) and reference FSM2ref simulation without redistribution (blue) for sub-domains (a) B0 on 17 March 2020, (b) D0 on 20 March 2017, (c) D0 on 31 March 2017 and (d) D0 on 17 May 2017.



**Figure C2.** Snow depth frequency plots at intervals of 10 cm at 100 m resolution for lidar measurements (black), FSM2trans simulation (red) and reference FSM2ref simulation without redistribution (blue) for sub-domains (a) B0 on 17 March 2020, (b) D0 on 20 March 2017, (c) D0 on 31 March 2017 and (d) D0 on 17 May 2017.

**Appendix D: Snow depth distribution on steep slopes**



**Figure D1.** Box plot of snow depth distribution on 31 March 2017 aggregated over slopes steeper than 40° for sub-domain D0 for lidar measurements, FSM2ref and FSM2trans simulations.

*Code availability.* FSM2trans code is available at <https://doi.org/10.16904/envidat.509> (Quéno, 2024). This code builds on the existing models FSM2 (Essery, 2015), FSM2oshd (Mott et al., 2023), SnowTran-3D (Liston et al., 2007) and SnowSlide (Bernhardt and Schulz, 2010).

*Data availability.* The lidar data used in this study will be published in a forthcoming data paper. It is available on request from the authors.

*Author contributions.* LQ designed the study and made the modelling developments for FSM2trans. PM, TJ and LQ set up the WindNinja model and provided downscaled input fields. LQ post-processed the lidar data with help from RM. LQ performed the simulations and analysed the results with input from TJ and RM. BC, GM, LQ, RM and TJ contributed to the development of the OSHD modelling framework. LQ wrote the paper with feedback from all authors.

*Competing interests.* The contact author has declared that none of the authors has any competing interests

*Disclaimer.* Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes every effort to include appropriate place names, the final responsibility lies with the authors.

*Acknowledgements.* Grammar and vocabulary of some sentences in the original manuscript were corrected using DeepL (<https://www.deepl.com>, last access: 6 August 2024).

The authors thank Eliane Brändle for the first tests and assessments of the WindNinja simulations, as well as the two anonymous reviewers for their detailed comments that helped to improve the paper. The OSHD model (FSM2oshd) development and implementation were largely funded by the Swiss Federal Office for the Environment (FOEN).

*Review statement.* This paper was edited by Carrie Vuyovich and reviewed by two anonymous referees.

## References

- Anderton, S. P., White, S. M., and Alvera, B.: Micro-scale spatial variability and the timing of snow melt runoff in a high mountain catchment, *J. Hydrol.*, 268, 158–176, [https://doi.org/10.1016/S0022-1694\(02\)00179-8](https://doi.org/10.1016/S0022-1694(02)00179-8), 2002.
- Anderton, S. P., White, S. M., and Alvera, B.: Evaluation of spatial variability in snow water equivalent for a

high mountain catchment, *Hydrol. Process.*, 18, 435–453, <https://doi.org/10.1002/hyp.1319>, 2004.

- Baron, M., Haddjeri, A., Lafaysse, M., Le Toumelin, L., Vionnet, V., and Fructus, M.: SnowPappus v1.0, a blowing-snow model for large-scale applications of the Crocus snow scheme, *Geosci. Model Dev.*, 17, 1297–1326, <https://doi.org/10.5194/gmd-17-1297-2024>, 2024.
- Bavay, M., Grünewald, T., and Lehning, M.: Response of snow cover and runoff to climate change in high Alpine catchments of Eastern Switzerland, *Adv. Water Resour.*, 55, 4–16, <https://doi.org/10.1016/j.advwatres.2012.12.009>, 2013.
- Beniston, M., Farinotti, D., Stoffel, M., Andreassen, L. M., Coppola, E., Eckert, N., Fantini, A., Giacona, F., Hauck, C., Huss, M., Huwald, H., Lehning, M., López-Moreno, J.-I., Magnusson, J., Marty, C., Morán-Tejeda, E., Morin, S., Naaim, M., Provenzale, A., Rabatel, A., Six, D., Stötter, J., Strasser, U., Terzago, S., and Vincent, C.: The European mountain cryosphere: a review of its current state, trends, and future challenges, *The Cryosphere*, 12, 759–794, <https://doi.org/10.5194/tc-12-759-2018>, 2018.
- Bernhardt, M. and Schulz, K.: SnowSlide: A simple routine for calculating gravitational snow transport, *Geophys. Res. Lett.*, 37, L11502, <https://doi.org/10.1029/2010GL043086>, 2010.
- Bernhardt, M., Zängl, G., Liston, G. E., Strasser, U., and Mauser, W.: Using wind fields from a high-resolution atmospheric model for simulating snow dynamics in mountainous terrain, *Hydrol. Process.*, 23, 1064–1075, <https://doi.org/10.1002/hyp.7208>, 2009.
- Bernhardt, M., Liston, G. E., Strasser, U., Zängl, G., and Schulz, K.: High resolution modelling of snow transport in complex terrain using downscaled MM5 wind fields, *The Cryosphere*, 4, 99–113, <https://doi.org/10.5194/tc-4-99-2010>, 2010.
- Bernhardt, M., Schulz, K., Liston, G. E., and Zängl, G.: The influence of lateral snow redistribution processes on snow melt and sublimation in alpine regions, *J. Hydrol.*, 424–425, 196–206, <https://doi.org/10.1016/j.jhydrol.2012.01.001>, 2012.
- Brauchli, T., Trujillo, E., Huwald, H., and Lehning, M.: Influence of Slope-Scale Snowmelt on Catchment Response Simulated With the Alpine3D Model, *Water Resour. Res.*, 53, 10723–10739, <https://doi.org/10.1002/2017WR021278>, 2017.
- Brunner, M. I., Björnsen Gurung, A., Zappa, M., Zekollari, H., Farinotti, D., and Stähli, M.: Present and future water scarcity in Switzerland: Potential for alleviation through reservoirs and lakes, *Sci. Total Environ.*, 666, 1033–1047, <https://doi.org/10.1016/j.scitotenv.2019.02.169>, 2019.
- Clark, M. P., Hendrikx, J., Slater, A. G., Kavetski, D., Anderson, B., Cullen, N. J., Kerr, T., Örn Hreinsson, E., and Woods, R. A.: Representing spatial variability of snow water equivalent in hydrologic and land-surface models: A review, *Water Resour. Res.*, 47, W07539, <https://doi.org/10.1029/2011WR010745>, 2011.
- Cristea, N. C., Bennett, A., Nijssen, B., and Lundquist, J. D.: When and Where Are Multiple Snow Layers Important for Simulations of Snow Accumulation and Melt?, *Water Resour. Res.*, 58, e2020WR028993, <https://doi.org/10.1029/2020WR028993>, 2022.
- Dadic, R., Mott, R., Lehning, M., and Burlando, P.: Parameterization for wind-induced preferential deposition of snow, *Hydrol. Process.*, 24, 1994–2006, <https://doi.org/10.1002/hyp.7776>, 2010.

- DeBeer, C. M. and Pomeroy, J. W.: Influence of snowpack and melt energy heterogeneity on snow cover depletion and snowmelt runoff simulation in a cold mountain environment, *J. Hydrol.*, 553, 199–213, <https://doi.org/10.1016/j.jhydrol.2017.07.051>, 2017.
- Dujardin, J. and Lehning, M.: Wind-Topo: Downscaling near-surface wind fields to high-resolution topography in highly complex terrain with deep learning, *Q. J. Roy. Meteorol. Soc.*, 148, 1368–1388, <https://doi.org/10.1002/qj.4265>, 2022.
- Durand, Y., Guyomarc'h, G., and Mérindol, L.: Numerical experiments of wind transport over a mountainous instrumented site: I. Regional scale, *Ann. Glaciol.*, 32, 187–194, <https://doi.org/10.3189/172756401781819445>, 2001.
- Dyunin, A. K. and Kotlyakov, V. M.: Redistribution of snow in the mountains under the effect of heavy snow-storms, *Cold Reg. Sci. Technol.*, 3, 287–294, [https://doi.org/10.1016/0165-232X\(80\)90035-X](https://doi.org/10.1016/0165-232X(80)90035-X), 1980.
- Egli, L., Jonas, T., Grünwald, T., Schirmer, M., and Burlando, P.: Dynamics of snow ablation in a small Alpine catchment observed by repeated terrestrial laser scans, *Hydrol. Process.*, 26, 1574–1585, <https://doi.org/10.1002/hyp.8244>, 2012.
- Essery, R.: A factorial snowpack model (FSM 1.0), *Geosci. Model Dev.*, 8, 3867–3876, <https://doi.org/10.5194/gmd-8-3867-2015>, 2015.
- Föhn, P. M. B. and Meister, R.: Distribution of Snow Drifts on Ridge Slopes: Measurements and Theoretical Approximations, *Ann. Glaciol.*, 4, 52–57, <https://doi.org/10.3189/S0260305500005231>, 1983.
- Forthofer, J. M., Butler, B. W., and Wagenbrenner, N. S.: A comparison of three approaches for simulating fine-scale surface winds in support of wildland fire management. Part I. Model formulation and comparison against measurements, *Int. J. Wildland Fire*, 23, 969–981, <https://doi.org/10.1071/WF12089>, 2014.
- Gascoïn, S., Lhermitte, S., Kinnard, C., Bortels, K., and Liston, G. E.: Wind effects on snow cover in Pascua-Lama, Dry Andes of Chile, *Adv. Water Resour.*, 55, 25–39, <https://doi.org/10.1016/j.advwatres.2012.11.013>, 2013.
- Gerber, F., Mott, R., and Lehning, M.: The Importance of Near-Surface Winter Precipitation Processes in Complex Alpine Terrain, *J. Hydrometeorol.*, 20, 177–196, <https://doi.org/10.1175/JHM-D-18-0055.1>, 2019.
- Griessinger, N., Schirmer, M., Helbig, N., Winstral, A., Michel, A., and Jonas, T.: Implications of observation-enhanced energy-balance snowmelt simulations for runoff modeling of Alpine catchments, *Adv. Water Resour.*, 133, 103410, <https://doi.org/10.1016/j.advwatres.2019.103410>, 2019.
- Groot Zwaafink, C. D., Mott, R., and Lehning, M.: Seasonal simulation of drifting snow sublimation in Alpine terrain, *Water Resour. Res.*, 49, 1581–1590, <https://doi.org/10.1002/wrcr.20137>, 2013.
- Guyomarc'h, G. and Mérindol, L.: Validation of an application for forecasting blowing snow, *Ann. Glaciol.*, 26, 138–143, <https://doi.org/10.3189/1998AoG26-1-138-143>, 1998.
- Hanzer, F., Förster, K., Nemeč, J., and Strasser, U.: Projected cryospheric and hydrological impacts of 21st century climate change in the Ötztal Alps (Austria) simulated using a physically based approach, *Hydrol. Earth Syst. Sci.*, 22, 1593–1614, <https://doi.org/10.5194/hess-22-1593-2018>, 2018.
- He, C., Chen, F., Barlage, M., Liu, C., Newman, A., Tang, W., Ikeda, K., and Rasmussen, R.: Can Convection-Permitting Modeling Provide Decent Precipitation for Offline High-Resolution Snowpack Simulations Over Mountains?, *J. Geophys. Res.-Atmos.*, 124, 12631–12654, <https://doi.org/10.1029/2019JD030823>, 2019.
- Helbig, N. and Löwe, H.: Parameterization of the spatially averaged sky view factor in complex topography, *J. Geophys. Res.-Atmos.*, 119, 4616–4625, <https://doi.org/10.1002/2013JD020892>, 2014.
- Jonas, T., Webster, C., Mazzotti, G., and Malle, J.: HPEval: A canopy shortwave radiation transmission model using high-resolution hemispherical images, *Agric. For. Meteorol.*, 284, 107903, <https://doi.org/10.1016/j.agrformet.2020.107903>, 2020.
- Le Toumelin, L., Gouttevin, I., Helbig, N., Galiez, C., Roux, M., and Karbou, F.: Emulating the Adaptation of Wind Fields to Complex Terrain with Deep Learning, *Artif. Intell. Earth Syst.*, 2, e220034, <https://doi.org/10.1175/AIES-D-22-0034.1>, 2023.
- Lehning, M., Doorschot, J., and Bartelt, P.: A snowdrift index based on SNOWPACK model calculations, *Ann. Glaciol.*, 31, 382–386, <https://doi.org/10.3189/172756400781819770>, 2000.
- Lehning, M., Bartelt, P., Brown, B., Fierz, C., and Satyawali, P.: A physical SNOWPACK model for the Swiss avalanche warning: Part II. Snow microstructure, *Cold Reg. Sci. Technol.*, 35, 147–167, [https://doi.org/10.1016/S0165-232X\(02\)00073-3](https://doi.org/10.1016/S0165-232X(02)00073-3), 2002.
- Lehning, M., Löwe, H., Ryser, M., and Raderschall, N.: Inhomogeneous precipitation distribution and snow transport in steep terrain, *Water Resour. Res.*, 44, W07404, <https://doi.org/10.1029/2007WR006545>, 2008.
- Li, D., Wrzesien, M. L., Durand, M., Adam, J., and Lettenmaier, D. P.: How much runoff originates as snow in the western United States, and how will that change in the future?, *Geophys. Res. Lett.*, 44, 6163–6172, <https://doi.org/10.1002/2017GL073551>, 2017.
- Li, L. and Pomeroy, J. W.: Estimates of Threshold Wind Speeds for Snow Transport Using Meteorological Data, *J. Appl. Meteor. Climatol.*, 36, 205–213, [https://doi.org/10.1175/1520-0450\(1997\)036<0205:EOTWSF>2.0.CO;2](https://doi.org/10.1175/1520-0450(1997)036<0205:EOTWSF>2.0.CO;2), 1997a.
- Li, L. and Pomeroy, J. W.: Probability of occurrence of blowing snow, *J. Geophys. Res.-Atmos.*, 102, 21955–21964, <https://doi.org/10.1029/97JD01522>, 1997b.
- Liston, G. E. and Elder, K.: Distributed Snow-Evolution Modeling System (SnowModel), *J. Hydrometeorol.*, 7, 1259–1276, <https://doi.org/10.1175/JHM548.1>, 2006.
- Liston, G. E. and Sturm, M.: A snow-transport model for complex terrain, *J. Glaciol.*, 44, 498–516, <https://doi.org/10.3189/S0022143000002021>, 1998.
- Liston, G. E., Haehnel, R. B., Sturm, M., Hiemstra, C. A., Berzovskaya, S., and Tabler, R. D.: Simulating complex snow distributions in windy environments using SnowTran-3D, *J. Glaciol.*, 53, 241–256, <https://doi.org/10.3189/172756507782202865>, 2007.
- Liston, G. E., Itkin, P., Stroeve, J., Tschudi, M., Stewart, J. S., Pedersen, S. H., Reinking, A. K., and Elder, K.: A Lagrangian Snow-Evolution System for Sea-Ice Applications (SnowModel-LG): Part I – Model Description, *J. Geophys. Res.-Oceans*, 125, e2019JC015913, <https://doi.org/10.1029/2019JC015913>, 2020.
- Luce, C. H., Tarboton, D. G., and Cooley, K. R.: The influence of the spatial distribution of snow on basin-averaged snowmelt, *Hydrol.*

- Process., 12, 1671–1683, [https://doi.org/10.1002/\(SICI\)1099-1085\(199808/09\)12:10/11<1671::AID-HYP688>3.0.CO;2-N](https://doi.org/10.1002/(SICI)1099-1085(199808/09)12:10/11<1671::AID-HYP688>3.0.CO;2-N), 1998.
- Luijting, H., Vikhamar-Schuler, D., Aspeli, T., Bakketun, Å., and Homleid, M.: Forcing the SURFEX/Crocus snow model with combined hourly meteorological forecasts and gridded observations in southern Norway, *The Cryosphere*, 12, 2123–2145, <https://doi.org/10.5194/tc-12-2123-2018>, 2018.
- Magnusson, J., Gustafsson, D., Hüsler, F., and Jonas, T.: Assimilation of point SWE data into a distributed snow cover model comparing two contrasting methods, *Water Resour. Res.*, 50, 7816–7835, <https://doi.org/10.1002/2014WR015302>, 2014.
- Magnusson, J., Wever, N., Essery, R., Helbig, N., Winstral, A., and Jonas, T.: Evaluating snow models with varying process representations for hydrological applications, *Water Resour. Res.*, 51, 2707–2723, <https://doi.org/10.1002/2014WR016498>, 2015.
- Marks, D., Domingo, J., Susong, D., Link, T., and Garen, D.: A spatially distributed energy balance snowmelt model for application in mountain basins, *Hydrol. Process.*, 13, 1935–1959, [https://doi.org/10.1002/\(SICI\)1099-1085\(199909\)13:12/13<1935::AID-HYP868>3.0.CO;2-C](https://doi.org/10.1002/(SICI)1099-1085(199909)13:12/13<1935::AID-HYP868>3.0.CO;2-C), 1999.
- Marsh, C. B., Spiteri, R. J., Pomeroy, J. W., and Wheeler, H. S.: Multi-objective unstructured triangular mesh generation for use in hydrological and land surface models, *Comput. Geosci.*, 119, 49–67, <https://doi.org/10.1016/j.cageo.2018.06.009>, 2018.
- Marsh, C. B., Pomeroy, J. W., Spiteri, R. J., and Wheeler, H. S.: A Finite Volume Blowing Snow Model for Use With Variable Resolution Meshes, *Water Resour. Res.*, 56, e2019WR025307, <https://doi.org/10.1029/2019WR025307>, 2020a.
- Marsh, C. B., Pomeroy, J. W., and Wheeler, H. S.: The Canadian Hydrological Model (CHM) v1.0: a multi-scale, multi-extent, variable-complexity hydrological model – design and overview, *Geosci. Model Dev.*, 13, 225–247, <https://doi.org/10.5194/gmd-13-225-2020>, 2020b.
- Marsh, C. B., Vionnet, V., and Pomeroy, J. W.: Windmapper: An efficient wind downscaling method for hydrological models, *Water Resour. Res.*, 59, e2022WR032683, <https://doi.org/10.1029/2022WR032683>, 2023.
- Mazzotti, G., Currier, W. R., Deems, J. S., Pflug, J. M., Lundquist, J. D., and Jonas, T.: Revisiting Snow Cover Variability and Canopy Structure Within Forest Stands: Insights From Airborne Lidar Data, *Water Resour. Res.*, 55, 6198–6216, <https://doi.org/10.1029/2019WR024898>, 2019.
- Mazzotti, G., Essery, R., Moeser, C. D., and Jonas, T.: Resolving Small-Scale Forest Snow Patterns Using an Energy Balance Snow Model With a One-Layer Canopy, *Water Resour. Res.*, 56, e2019WR026129, <https://doi.org/10.1029/2019WR026129>, 2020.
- Mott, R. and Lehning, M.: Meteorological Modeling of Very High-Resolution Wind Fields and Snow Deposition for Mountains, *J. Hydrometeorol.*, 11, 934–949, <https://doi.org/10.1175/2010JHM1216.1>, 2010.
- Mott, R., Schirmer, M., Bavay, M., Grünewald, T., and Lehning, M.: Understanding snow-transport processes shaping the mountain snow-cover, *The Cryosphere*, 4, 545–559, <https://doi.org/10.5194/tc-4-545-2010>, 2010.
- Mott, R., Vionnet, V., and Grünewald, T.: The Seasonal Snow Cover Dynamics: Review on Wind-Driven Coupling Processes, *Front. Earth Sci.*, 6, 197, <https://doi.org/10.3389/feart.2018.00197>, 2018.
- Mott, R., Wolf, A., Kehl, M., Kunstmann, H., Warscher, M., and Grünewald, T.: Avalanches and micrometeorology driving mass and energy balance of the lowest perennial ice field of the Alps: a case study, *The Cryosphere*, 13, 1247–1265, <https://doi.org/10.5194/tc-13-1247-2019>, 2019.
- Mott, R., Winstral, A., Cluzet, B., Helbig, N., Magnusson, J., Mazzotti, G., Quéno, L., Schirmer, M., Webster, C., and Jonas, T.: Operational snow-hydrological modeling for Switzerland, *Front. Earth Sci.*, 11, 1228158, <https://doi.org/10.3389/feart.2023.1228158>, 2023.
- Mower, R., Gutmann, E. D., Liston, G. E., Lundquist, J., and Rasmussen, S.: Parallel SnowModel (v1.0): a parallel implementation of a distributed snow-evolution modeling system (SnowModel), *Geosci. Model Dev.*, 17, 4135–4154, <https://doi.org/10.5194/gmd-17-4135-2024>, 2024.
- Musselman, K. N., Pomeroy, J. W., Essery, R. L. H., and Leroux, N.: Impact of windflow calculations on simulations of alpine snow accumulation, redistribution and ablation, *Hydrol. Process.*, 29, 3983–3999, <https://doi.org/10.1002/hyp.10595>, 2015.
- Pomeroy, J. W. and Gray, D. M.: Saltation of snow, *Water Resour. Res.*, 26, 1583–1594, <https://doi.org/10.1029/WR026i007p01583>, 1990.
- Pomeroy, J. W. and Gray, D. M.: Snowcover Accumulation, Relocation and Management, NHRI Science Report No. 7, National Hydrology Research Institute, Saskatoon, Canada, <https://publications.gc.ca/site/eng/9.892773/publication.html> (last access: 6 August 2024), 1995.
- Pomeroy, J. W., Gray, D. M., and Landine, P. G.: The Prairie Blowing Snow Model: characteristics, validation, operation, *J. Hydrol.*, 144, 165–192, [https://doi.org/10.1016/0022-1694\(93\)90171-5](https://doi.org/10.1016/0022-1694(93)90171-5), 1993.
- Quéno, L.: FSM2trans code, EnviDat [code], <https://doi.org/10.16904/envidat.509>, 2024.
- Quéno, L., Vionnet, V., Dombrowski-Etchevers, I., Lafaysse, M., Dumont, M., and Karbou, F.: Snowpack modelling in the Pyrenees driven by kilometric-resolution meteorological forecasts, *The Cryosphere*, 10, 1571–1589, <https://doi.org/10.5194/tc-10-1571-2016>, 2016.
- Raparelli, E., Tuccella, P., Colaiuda, V., and Marzano, F. S.: Snow cover prediction in the Italian central Apennines using weather forecast and land surface numerical models, *The Cryosphere*, 17, 519–538, <https://doi.org/10.5194/tc-17-519-2023>, 2023.
- Reynolds, D., Gutmann, E., Kruyt, B., Haugeneder, M., Jonas, T., Gerber, F., Lehning, M., and Mott, R.: The High-resolution Intermediate Complexity Atmospheric Research (HICAR v1.1) model enables fast dynamic downscaling to the hectometer scale, *Geosci. Model Dev.*, 16, 5049–5068, <https://doi.org/10.5194/gmd-16-5049-2023>, 2023.
- Reynolds, D. S., Pflug, J. M., and Lundquist, J. D.: Evaluating Wind Fields for Use in Basin-Scale Distributed Snow Models, *Water Resour. Res.*, 57, e2020WR028536, <https://doi.org/10.1029/2020WR028536>, 2020.
- Sauter, T., Möller, M., Finkelnburg, R., Grabiec, M., Scherer, D., and Schneider, C.: Snowdrift modelling for the Vestfonna ice cap, north-eastern Svalbard, *The Cryosphere*, 7, 1287–1301, <https://doi.org/10.5194/tc-7-1287-2013>, 2013.

- Schirmer, M., Winstral, A., Jonas, T., Burlando, P., and Peleg, N.: Natural climate variability is an important aspect of future projections of snow water resources and rain-on-snow events, *The Cryosphere*, 16, 3469–3488, <https://doi.org/10.5194/tc-16-3469-2022>, 2022.
- Schneiderbauer, S. and Prokop, A.: The atmospheric snow-transport model: SnowDrift3D, *J. Glaciol.*, 57, 526–542, <https://doi.org/10.3189/002214311796905677>, 2011.
- Sexstone, G. A., Clow, D. W., Fassnacht, S. R., Liston, G. E., Hiemstra, C. A., Knowles, J. F., and Penn, C. A.: Snow Sublimation in Mountain Environments and Its Sensitivity to Forest Disturbance and Climate Warming, *Water Resour. Res.*, 54, 1191–1211, <https://doi.org/10.1002/2017WR021172>, 2018.
- Sharma, V., Gerber, F., and Lehning, M.: Introducing CRYOWRF v1.0: multiscale atmospheric flow simulations with advanced snow cover modelling, *Geosci. Model Dev.*, 16, 719–749, <https://doi.org/10.5194/gmd-16-719-2023>, 2023.
- Sommer, C. G., Lehning, M., and Mott, R.: Snow in a Very Steep Rock Face: Accumulation and Redistribution During and After a Snowfall Event, *Front. Earth Sci.*, 3, 73, <https://doi.org/10.3389/feart.2015.00073>, 2015.
- Strasser, U., Bernhardt, M., Weber, M., Liston, G. E., and Mauser, W.: Is snow sublimation important in the alpine water balance?, *The Cryosphere*, 2, 53–66, <https://doi.org/10.5194/tc-2-53-2008>, 2008.
- Tabler, R. D.: Predicting profiles of snowdrifts in topographic catchments, in: Proceedings of the 43rd Annual Western Snow Conference, Coronado, California, 1975, 87–97, <https://westernsnowconference.org/bibliography/1975Tabler.pdf> (last access: 6 August 2024), 1975.
- Vionnet, V., Brun, E., Morin, S., Boone, A., Faroux, S., Le Moigne, P., Martin, E., and Willemet, J.-M.: The detailed snowpack scheme Crocus and its implementation in SURFEX v7.2, *Geosci. Model Dev.*, 5, 773–791, <https://doi.org/10.5194/gmd-5-773-2012>, 2012.
- Vionnet, V., Martin, E., Masson, V., Guyomarc'h, G., Naaim-Bouvet, F., Prokop, A., Durand, Y., and Lac, C.: Simulation of wind-induced snow transport and sublimation in alpine terrain using a fully coupled snowpack/atmosphere model, *The Cryosphere*, 8, 395–415, <https://doi.org/10.5194/tc-8-395-2014>, 2014.
- Vionnet, V., Dombrowski-Etchevers, I., Lafaysse, M., Quéno, L., Seity, Y., and Bazile, E.: Numerical Weather Forecasts at Kilometer Scale in the French Alps: Evaluation and Application for Snowpack Modeling, *J. Hydrometeorol.*, 17, 2591–2614, <https://doi.org/10.1175/JHM-D-15-0241.1>, 2016.
- Vionnet, V., Marsh, C. B., Menounos, B., Gascoin, S., Wayand, N. E., Shea, J., Mukherjee, K., and Pomeroy, J. W.: Multi-scale snowdrift-permitting modelling of mountain snowpack, *The Cryosphere*, 15, 743–769, <https://doi.org/10.5194/tc-15-743-2021>, 2021.
- Wagenbrenner, N. S., Forthofer, J. M., Lamb, B. K., Shannon, K. S., and Butler, B. W.: Downscaling surface wind predictions from numerical weather prediction models in complex terrain with WindNinja, *Atmos. Chem. Phys.*, 16, 5229–5241, <https://doi.org/10.5194/acp-16-5229-2016>, 2016.
- Wagenbrenner, N. S., Forthofer, J. M., Page, W. G., and Butler, B. W.: Development and Evaluation of a Reynolds-Averaged Navier–Stokes Solver in WindNinja for Operational Wildland Fire Applications, *Atmosphere*, 10, 672, <https://doi.org/10.3390/atmos10110672>, 2019.
- Wang, Z. and Huang, N.: Numerical simulation of the falling snow deposition over complex terrain, *J. Geophys. Res.-Atmos.*, 122, 980–1000, <https://doi.org/10.1002/2016JD025316>, 2017.
- Wang, Z., Bovik, A. C., Sheikh, H. R., and Simoncelli, E. P.: Image quality assessment: from error visibility to structural similarity, *IEEE Trans. Image Process.*, 13, 600–612, <https://doi.org/10.1109/TIP.2003.819861>, 2004.
- Winstral, A., Jonas, T., and Helbig, N.: Statistical Downscaling of Gridded Wind Speed Data Using Local Topography, *J. Hydrometeorol.*, 18, 335–348, <https://doi.org/10.1175/JHM-D-16-0054.1>, 2017.