



Bias in modeled Greenland Ice Sheet melt revealed by ASCAT

Anna Puggaard^{1,2}, Nicolaj Hansen², Ruth Mottram², Thomas Nagler³, Stefan Scheiblauer³, Sebastian B. Simonsen¹, Louise S. Sørensen¹, Jan Wuite³, and Anne M. Solgaard⁴

¹Geodesy and Earth Observation, DTU-Space, Technical University of Denmark, Lyngby, Denmark

²National Centre for Climate Research, Danish Meteorological Institute (DMI), Copenhagen, Denmark

³Environmental Earth Observation Information Technology (ENVEO) IT GmbH, Innsbruck, Austria

⁴Glaciology and Climate, Geological Survey of Denmark and Greenland (GEUS), Copenhagen, Denmark

Correspondence: Anna Puggaard (annpu@space.dtu.dk)

Received: 12 April 2024 – Discussion started: 6 June 2024

Revised: 17 May 2025 – Accepted: 20 May 2025 – Published: 12 August 2025

Abstract. The runoff of surface melt is the primary driver of mass loss over the Greenland Ice Sheet. An accurate representation of surface melt is crucial for understanding the surface mass balance and, ultimately, the ice sheet's total contribution to sea level rise. Regional climate models (RCMs) model ice-sheet-wide melt volume but exhibit large variability in estimates among models, requiring validation with observed melt. Here, we explore a novel data processing method from the Advanced SCATterometer (ASCAT) instrument on board the EUMETSAT MetOp satellites, which provides estimates of the spatiotemporal variability of melt extent over the Greenland Ice Sheet between 2007 and 2020. We apply ASCAT wet-snow maps to pinpoint differences in the melt products from three distinct RCMs, where one is forced at the boundary with two different reanalyses. Using automatic weather station (AWS) air temperature observations, we assess how well RCM-modeled melt volume aligns with in situ temperatures. With this assessment, we establish a threshold for the RCMs to identify how much meltwater is in the models before it is observed at the AWSs and ultimately infer the melt extent in the RCMs. We show that applying thresholds, informed by in situ measurements, reduces the differences between ASCAT and RCMs and minimizes the discrepancies between different RCMs. Differences between modeled melt extent and melt extent observed by ASCAT are used to pinpoint (i) biases in the RCMs, which include variability in their albedo schemes, snowfall, turbulent heat fluxes, and temperature, as well as differences in radiation schemes, and (ii) limitations of the liquid water detection by ASCAT, including misclassification in the ablation zone and a temporal melt onset bias. Overall, we find that the

RCMs tend to have a later melt onset than ASCAT and an earlier end to the melt season, with a similar but slightly smaller melt area compared to that identified in ASCAT. Biases, however, vary spatially between models and with compensating errors in different regions, suggesting that one RCM can sometimes represent the present-day surface across the entire ice sheet more effectively than the ensemble mean.

1 Introduction

The Greenland Ice Sheet significantly contributes to the rise in global sea levels, contributing approximately 15 % since the early 1990s (WCRP Global Sea Level Budget Group, 2018). Between 1992 and 2020, satellite observations have shown that the Greenland Ice Sheet has lost 4892 ± 457 Gt of ice or 13.6 ± 1.3 mm sea level equivalent (Otosaka et al., 2023), with half of the mass loss attributed to a decrease in the surface mass balance (SMB; van den Broeke et al., 2016). However, the rate of mass loss has exhibited considerable annual variability in recent years, ranging from 86 ± 75 Gt in 2017 to 444 ± 93 Gt in 2019, with the latter being driven by exceptional surface melting during the summer (Tedesco and Fettweis, 2020). Modeling studies have shown that surface melt on the Greenland Ice Sheet has generally doubled since the 1990s due to a rise in temperature (Tedesco and Fettweis, 2020; van den Broeke et al., 2016). Meanwhile, snow accumulation has remained nearly constant (van den Broeke et al., 2016). In the warm summer months, the surface temperature rises above the melting point of ice, and surface melt

occurs. Depending on snow and firn characteristics, meltwater generated at the surface can collect at the surface and form supraglacial meltwater lakes (Koenig et al., 2015); run off as surface meltwater (Smith et al., 2015); or percolate into the snowpack, where it either refreezes (Forster et al., 2014; Harper et al., 2012) or runs off englacially (Chandler et al., 2013).

At present, regional climate models (RCMs) provide the most comprehensive approach for obtaining ice-sheet-wide estimates of meltwater volumes and runoff, with simulations showing the best agreement with observations (Fettweis et al., 2020). However, these models are influenced by the chosen modeling approach, and substantial disparities persist among models (Rae et al., 2012; Vernon et al., 2013; Fettweis et al., 2020; Glaude et al., 2024). In particular, recent studies suggest that small differences between models in the present day, representing differences in parameterizations, have large effects on projections of melt, runoff, and surface mass balance when run into the future, causing greater uncertainty in sea level rise estimates than is desirable for climate adaptation purposes (Goelzer et al., 2020). Thus, it is crucial to develop methods to evaluate melt estimates from RCMs against observations of melt to understand these discrepancies and ultimately evaluate which RCMs most realistically model melt (Langen et al., 2017). While melt intensity can be derived from in situ observations at automatic weather stations (AWSs), the sparse distribution of these stations across the ice sheet limits the evaluation of melt estimates beyond local scales (Fausto et al., 2018). On the other hand, satellite remote sensing can observe the presence of meltwater in the upper part of the firn pack, while melt intensity has not yet been measured successfully using remote sensing. Recent approaches, such as those by Dethinne et al. (2023) and Picard et al. (2022), have assimilated remote sensing datasets into more detailed modeling of surface melt processes to estimate meltwater volume. Although including more observational data generally improves the representation of current surface conditions in the RCMs (Langen et al., 2017; Dethinne et al., 2023), there remains a strong need for independent observational datasets to assess model outputs. Remote sensing satellites provide information on Greenland Ice Sheet surface melt through observations from the visible to the microwave spectrum, where some of the widely used sensors are the Advanced SCATterometer (ASCAT), Sentinel-1, the Moderate Resolution Imaging Spectroradiometer (MODIS), and the Special Sensor Microwave Imager/Sounder (SSMIS) (Husman et al., 2023). Microwave sensors have the advantage of offering measurements independently of cloud cover, weather conditions, and polar darkness. Over the ice sheets, the backscattering of the microwave signals from snow and ice depends on roughness geometry and electrical properties, which in turn depend on the physical characteristics of the snow and ice (Wismann, 2000; Long, 2017). During the winter, the backscatter signal can exhibit a gradual decrease due

to snow accumulation attenuating the volume scattering in the snowpack. As the temperature increases, meltwater at the surface is introduced, and the backscatter signal experiences a substantial drop. This sensitivity to meltwater has enabled several studies to estimate melt over both ice sheets using passive and active microwave measurements with a threshold method to detect the onset of melt and its extent (Long and Drinkwater, 1994; Wismann, 2000; Ashcraft and Long, 2006; Fettweis et al., 2011; Colosio et al., 2021; Husman et al., 2023). The magnitude of the decrease in backscatter varies due to factors such as the snow water content and the specific properties of the snowpack, such as grain size and the presence of ice layers and lenses, which influence dielectric properties and roughness geometries (Wismann, 2000; Long, 2017). Refrozen meltwater from the previous melt season can percolate into the firn, leading to the formation of subsurface features such as firn aquifers and ice lenses, which can potentially amplify the backscatter signal prior to the current melting season (Brangers et al., 2020). Further, meltwater in the subsurface can still be detected after refreezing of the surface layer as the low-frequency signals can still penetrate into the refrozen surface layer. Using a threshold method proposed by Ashcraft and Long (2006), Husman et al. (2023) showed that C-band (4–8 GHz) active microwave sensors detected more melt days than K-band (18–27 GHz) passive microwave sensors in areas with meltwater in the subsurface in Antarctica due to differences in penetration depth. Thus, for the correct identification of surface melt using microwave satellite observations, it is important to account for penetration depth, changes in dielectric properties, and roughness geometries of the snowpack. The formation of subsurface ice features and subsurface penetration can lead to significant misclassification of surface melt if a simple threshold method is used (Ashcraft and Long, 2006; Long, 2017). Instead of using a simple threshold method, ASCAT wet-snow maps utilize an algorithm that incorporates the temporal behavior of the backscattered signal. With this method, the ASCAT wet-snow maps can not only detect the presence of liquid water on the surface but also distinguish between melting and subsequent refreezing of the surface meltwater. This makes ASCAT wet-snow maps a unique product as they allow for a fairer comparison between the observed liquid meltwater extent and the surface melt extent simulated by RCMs. Furthermore, by applying an annual recalibration of the winter signal, the product accounts for the formation of subsurface features from the previous melt season (Nagler et al., 2024). Again, this ensures better classification of melt signal compared to previous melt extent products from both active and passive microwave measurements over the Greenland Ice Sheet, such as those by Abdalati and Steffen (1995), Wismann (2000), Nghiem et al. (2001), Tedesco (2007), Fettweis et al. (2011), and Colosio et al. (2021).

By using ASCAT wet-snow maps, we aim to establish a framework for evaluating the performance of RCMs in simulating the temporal variability of present-day melt extent.

As RCMs are often calibrated with respect to basin-wide surface mass balance, incorporating an independent satellite dataset like ASCAT wet-snow maps enables a more comprehensive assessment of model performance. By including HIRHAM5, the Regional Atmospheric Climate Model (RACMO2.3p2), and the Modèle Atmosphérique Régional (MARv3.12), we assess how well each model captures surface melt patterns, focusing not on internal model parameterizations, e.g., albedo and near-surface temperature, but on the representation of melt extent. Here, we employ a liquid water detection algorithm that integrates the temporal behavior of the backscatter signal by classifying the first- and second-time derivatives. Rather than solely identifying the period of meltwater presence, the method distinguishes between surface melting and subsequent refreezing of the surface meltwater. However, this method does not estimate the meltwater volume, meaning we can only use the satellite-observed melt to assess the RCMs' ability to represent the extent of surface melt realistically. Hence, we compare the melt extent observed by ASCAT to the melt extent modeled by RCMs. To ensure that the RCM-modeled melt aligns with the in situ observations, we compare modeled melt volumes to observed 2 m temperatures at automatic weather stations (AWSs) to establish the melting threshold (millimeters of water equivalent per day, mm w.e. d^{-1}) to in turn identify how much meltwater is in the models before it is observed at the AWSs. Once the melt extent is inferred from the RCMs, it is possible to compare it with the melt extent observed by ASCAT to identify biases within the RCMs and limitations in the ASCAT melt observations.

2 Data

2.1 PROMICE AWS

The Programme for Monitoring of the Greenland Ice Sheet and Greenland Climate Network (PROMICE GC-net AWS) offers hourly and daily meteorological and glaciological in situ measurements for 54 weather stations on the Greenland Ice Sheet, tundra, and peripheral glaciers (Fausto et al., 2021; How et al., 2022). Here, we include 34 stations on the Greenland Ice Sheet and measurements of air temperature between 2007 and 2020. PROMICE GC-net only includes active weather stations, but the historical GC-net data include several discontinued stations. Here data from the Summit, GITS, and PertermannELA stations are included, as they overlap in time with ASCAT data (Steffen et al., 2022; Vandecrux et al., 2023). See Fig. 1 for the locations of the AWSs.

2.2 Regional climate models

The melt volume from RCMs is derived by closing the surface energy budget. When skin temperature exceeds 0°C , additional energy contributes towards melting, resetting the skin temperature to 0°C (Langen et al., 2015; Noël et al.,




Figure 1. Locations of AWSs included in the study. Only AWSs on the ice sheet (ASCAT liquid water detection domain) are included in this study. Further, the Rignot and Mouginot (2012) drainage basins that are utilized in the subsequent evaluation of RCM-modeled melt extent against ASCAT-observed melt extent are also included.

2018). However, different model setups, such as horizontal and vertical resolutions, and choices of parameters, like surface albedo and subsurface schemes, impact the surface energy balance simulated within these models and thereby result in different melt volumes. We compare the melt extent observed by ASCAT with the modeled melt extent from three RCMs: HIRHAM5, RACMO2.3p2, and MARv3.12; see Figs. 2 and 3. A detailed description of model differences is provided in the following.

HIRHAM5

HIRHAM5 (Lucas Picher et al., 2012) utilizes a rotated polar grid at $0.05^{\circ} \times 0.05^{\circ}$ horizontal resolution, which for the Greenland Ice Sheet corresponds to approximately 5.5 km. HIRHAM5 was developed from the dynamics of the numerical weather forecast High-Resolution Limited Area Model (HIRLAM; Undén et al., 2002) combined with the physics from the ECHAM5 general circulation model (Roeckner et al., 2003) to ensure accurate simulation of the surface




Figure 2. Modeled mean annual melt by different RCMs inside the ASCAT data mask (2007–2020).

energy balance. HIRHAM5 is forced on the lateral boundary with 6 h global reanalysis temperature, relative humidity, wind vectors, and pressure fields. Further, daily sea ice concentration and sea surface temperature fields are also used to force the model (Langen et al., 2017). Here, we include HIRHAM5 run with both ERA-Interim for the period of 1979–2019 (Dee et al., 2011) and ERA5 for the period of 1960–2020 (Hersbach et al., 2020). HIRHAM5 forced with ERA-Interim is referred to as HIRHAM5-ERA1, and HIRHAM5 forced with ERA5 is referred to as HIRHAM5-ERA5. Further, the surface albedo in HIRHAM5-ERA1 is derived from MODIS gridded surface albedo (Box et al., 2012), described in Langen et al. (2017), while the surface albedo in HIRHAM5-ERA5 is computed internally as a linear function of temperature, described in Langen et al. (2015). The outputs from both HIRHAM5 runs are used to force the offline subsurface model described in Langen et al. (2017). The offline subsurface model includes a multilayer surface snow and mass balance scheme that simulates melt percolation, retention, and refreezing with a vertical resolution down to 60 m w.e. (Langen et al., 2017).

RACMO2.3p2

The polar Regional Atmospheric Climate Model (RACMO2.3p2) is run on an approximately 5.5 km ($0.05^\circ \times 0.05^\circ$) horizontal resolution (Noël et al., 2018, 2019). RACMO2.3p2 also integrates the numerical weather forecasting dynamics from HIRLAM (Undén et al., 2002) with the physics from the European Centre for Medium-Range Weather Forecasts–Integrated Forecast System cycle CY33r1 (ECMWF, 2009) to accurately simulate the surface energy balance. On the lateral boundary, RACMO2.3p2 uses the same 6 h and daily ERA5 fields as HIRHAM5-ERA5 (Hersbach et al., 2020). The snow

albedo is derived from the snow grain size, cloud optical thickness, solar zenith angle, and impurity concentration in snow. However, to account for temporal variability in the ablation zone, gridded MODIS observations of bare ice (Box et al., 2012) are included in these areas (Van Angelen et al., 2012). The model includes a multilayer snow module for simulating surface melt partitioning into percolation, retention, refreezing, and – ultimately – runoff (Ettema et al., 2010). The 5.5 km product is statistically downscaled to a horizontal resolution of 1 km to represent the steep SMB gradients found over narrow glaciers and confined ablation zones at the rugged margins of the ice sheet (Noël et al., 2016).

MARv3.12

The Modèle Atmosphérique Régional, MARv3.12, is run at a 10 km horizontal grid resolution (Fettweis et al., 2013, 2017; Tedesco and Fettweis, 2020). MARv3.12 combines atmospheric modeling with a soil–ice–snow–vegetation–atmosphere transfer scheme (SISVATS) (Gallée and Schayes, 1994) to simulate the surface energy balance and mass balance processes over the ice sheet. MARv3.12 is forced at the lateral boundaries with ERA5 (Hersbach et al., 2020) at a 6 h temporal resolution. Output values are averaged to obtain daily values (Tedesco and Fettweis, 2020). MARv3.12 includes the snow model Crocus (Brun et al., 1992), which simulates a number of layers of snow, ice, and firn of variable thickness and energy and mass transport between each layer. The snow model also provides snow grain properties, which are used in combination with density, age, and type to simulate snow albedo (Brun et al., 1992; Fettweis et al., 2017; Antwerpen et al., 2022). MARv3.12 also has an albedo range for bare ice between 0.4 and 0.55, depending on the cleanliness of the ice (Fettweis et al., 2017).




Figure 3. RCM-simulated variables from 2007–2020 for HIRHAM-ERA5, HIRHAM-ERA1, MARv3.12, and RACMO2.3p2. (a–d) Mean JJA temperatures in $^{\circ}\text{C}$, (e–h) the JJA surface albedo, (i–l) the JJA downward shortwave radiation (SWD) in W m^{-2} , and (m–p) the mean annual snowfall in mm w.e. yr^{-1} .

While both RACMO2.3p2 and HIRHAM5-ERA5 incorporate MODIS observations into the albedo computation, the surface albedo in MARv3.12 is only based on the internally computed broadband albedo (Brun et al., 1992).

2.3 ASCAT wet-snow maps

The ASCAT instrument on board the EUMETSAT MetOp polar-orbiting satellites has provided measurements since 2007. ASCAT is a real aperture radar (RAR) instrument and measures radar backscatter in vertical polarization at a range of incidence angles from 25 to 65° at a frequency of 5.255 GHz (C band) (Figa-Saldaña et al., 2002; Long, 2017). With a spatial resolution of 25 km, ASCAT has limited land applications, but with the introduction of the scatterometer image reconstruction (SIR) algorithm, the utility of ASCAT data has improved significantly (Lindsley and Long, 2010). The SIR algorithm takes advantage of image reconstruction techniques and the spatial overlap of measurements over a region to enhance the effective spatial resolution over a time interval. SIR depends on the number of measurements, where increasing the number of measurements leads to a reduction in noise and an improvement in the spatial resolution. However, the radar characteristics must remain constant between passes during the interval time. We refer to Long et al. (1993) and Long and Drinkwater (1994) for a more detailed description of the SIR algorithm. For Greenland, a 4 d time interval is appropriate as it ensures sufficiently dense sampling while justifying the assumption of constant radar characteristics (Lindsley and Long, 2010). With the 4 d time interval, diurnal variations in backscatter signal over ice, such as melt during the day and refreezing during the night, are averaged out, which introduces additional uncertainty. Additionally, the resolution-enhanced ASCAT product may not capture short melt events in the spring and intense precipitation events, as these signals are averaged. Furthermore, potential azimuth angle dependencies are not considered in the construction of the SIR products (Long and Drinkwater, 1994; Lindsley and Long, 2010).

The ASCAT SIR product is available from 2007 to 2020 and is used to identify four different melt stages by applying a hierarchical decision tree using dynamic thresholds based on the previous winter reference month (JFM) and the first- and second-time derivatives of the backscattered signal, σ^0 (Nagler et al., 2024). During the winter months, the backscatter signal is relatively stable, and only minor variations may occur due to changes in snowpack properties (labeled Stage 1 (ST-1), no melt). To account for the possibility of remnant changes in the snowpack from the previous melt season, the winter signal is recalibrated annually pixel by pixel. As the snow surface starts to melt in spring, rising above a liquid water content of $\sim 1\%$ volume (Mätzler, 1987), the backscatter signal decreases significantly. This rapid drop in the backscatter coefficient is utilized to identify the onset and occurrence of surface melt (labeled ST-2A, surface melt).

During intense melt events, the backscatter signal becomes fully saturated, which means that an increase in the melt intensity does not lead to further lowering of the backscatter signal (labeled ST-2B, wet-snow layer). As the meltwater starts to refreeze, the backscatter signal gradually increases again, returning to stable winter conditions once the meltwater down to the radar penetration depth is refrozen (labeled ST-3, an increase in the refrozen layer). For a more detailed description of the melt stage classification, we refer to Nagler et al. (2024). When comparing ASCAT wet-snow maps and the melt extent modeled by the RCMs, we excluded ST-3 (an increase in the refrozen layer) in order to make the fairest comparison with RCM surface melt. Throughout this paper, we use the phrase melt days when referring to days labeled ST-2A (surface melt) and ST-2B (wet-snow layer) in the ASCAT wet-snow maps. Similarly, melt extent refers to pixels labeled ST-2A and ST-2B. It should be noted that active surface melting may not necessarily occur on the ASCAT-derived melt days, but meltwater may simply be present without active melting. However, we mitigate the potential differences by excluding days classified as refreezing and recalibrating the winter baseline to account for liquid water that persisted through the winter.

As an example, Fig. 4a shows the backscatter measurements, and the liquid water classification is shown for the KAN_U AWS. The winter dry signal changes after at least one melt season, highlighting the melt classification method's ability to correctly detect liquid water despite a difference in the winter signal before and after the melt season. When comparing the liquid water extent observed by ASCAT and the melt extent modeled by the RCMs, we excluded ST-3 (an increase in the refrozen layer) since we are only interested in comparing surface melt. Figure 4b illustrates a snapshot of the detection and classification from 7 August 2017. Refreezing or no melting is observed in pixels close to the ice sheet margin when melt occurs at higher elevations on the ice sheet. Further, when looking at the mean annual number of melt days, the lower ablation zone exhibits fewer melt days than at higher elevations. This seemingly incorrect classification of liquid water near the ice margin is an effect of changes in the surface roughness associated with melt. Changes in surface roughness can counteract the decrease in backscatter associated with melt, effectively concealing liquid water in the backscatter signal. Therefore, when comparing ASCAT and RCMs, we mask out these areas by applying the maximum elevation of the snow line between 2007–2020 (Box et al., 2012), where pixels with an elevation below this snow line are excluded in the comparison (Fig. 4d).

3 Methods

Modeled surface melt in the RCMs is subject to large variability among models, as seen in Fig. 2 and discussed in, e.g., Fettweis et al. (2020). The Greenland Ice Sheet SMB




Figure 4. (a) The 2016–2018 melt detection and the backscatter signal at KAN_U. (b–c) Two examples of the spatial distribution of liquid water detection on 8 June 2016 and 7 August 2017. (d) Mean annual number of days with detected liquid water but without refreezing. In (d) we apply the maximum elevation of the snow line between 2007–2020 (black line) to mask out areas where ASCAT persistently cannot correctly detect the presence of liquid water.

model intercomparison project (GrSMBMIP) suggested that discrepancies between RCMs are not systematic (Fettweis et al., 2020); thus there is a need for individual evaluation of each modeled melt volume product before we can compare the extent observed by ASCAT. We refer to Fig. 5 for an overview of the evaluation taken prior to comparing RCMs with ASCAT. To establish a threshold (in mm w.e. d^{-1}) to infer the melt extent from the simulated RCM melt volume, we compare it to the PROMICE GC-net AWS. With this comparison we identify how much meltwater must be in the models before we can also observe it at the AWSs. We compare each AWS to the RCM grid cell, which has the closest center point to the AWS location; see Fig. 1 for AWS locations. It is important to note that the AWS measurements are significantly affected by local-scale weather conditions, so this approach only ensures that the melt modeled by the RCMs aligns primarily at these specific locations. Therefore, we include the maximum number of AWSs to align with, rather than being limited to those above the maximum snow line extent, which is where ASCAT and the inferred RCM melt extents are compared. Since melt is not directly measured at

the AWSs, we use 2 m air temperature as a proxy for melt conditions, as near-surface air temperature is closely linked to melt processes. This approach allows us to identify and quantify temperature biases in each of the RCMs and assess how well the models simulate melt compared to in situ observations.

Air temperature is strongly correlated with melt since melt is a response to a positive surface energy balance, which occurs when the surface temperature reaches 0°C (Cuffey and Paterson, 2010). However, it is important to note that air temperature and surface temperature are not the same; while air temperature influences surface conditions, surface temperature depends on a combination of energy exchanges at the surface. Additionally, the local properties of the snowpack can also affect when melt occurs, and melt can occur in the snowpack when air temperatures are below 0°C . Thus, it is crucial to consider that the heat content of the overlying atmosphere is not the sole driver of melt when we compare the modeled melt volume against the days with observed melt, as indicated by AWS temperature measurements. With this approach, we apply a threshold to identify days with significant




Figure 5. Diagram describing the evaluation process of RCM melt volume and processing of ASCAT wet-snow maps prior to the comparison between ASCAT liquid water extent and RCM-inferred melt extent.

Table 1. Melting thresholds for the different RCMs based on in situ PROMICE AWS observations of air temperature. The table also gives examples of the mean July to August air temperatures at six selected AWSs and the mean across all stations. The corresponding mean July to August air temperatures are showcased for each RCM. The locations of all stations included in the study are shown in Fig. 1.

	Thresholds [mm w.e. d ⁻¹]	Mean air temperature [°C]						
		CP1	DY2	KAN_U	KPC_U	NUK_U	SDL	All AWSs
HIRHAM5-ERA5	4.1	−6.36	−6.78	−5.08	−1.13	0.74	−9.62	−4.02
HIRHAM5-ERA1	0.4	−3.92	−4.04	−2.53	0.29	1.12	−6.50	−2.35
RACMO2.3p2	0.7	−4.66	−4.83	−2.95	−0.80	2.03	−7.14	−1.87
MARv3.12	1.0	−5.16	−5.14	−3.31	−0.93	1.24	−7.71	−2.64
PROMICE GC-net	–	−4.96	−4.70	−3.31	−0.85	−0.84	−6.80	−3.39

melt in the RCMs, but we also implement a threshold for the temperature observations to identify when melt occurs in the snowpack at the AWSs. We explore various thresholds for temperature observations to account for other factors in the snowpack that influence when surface melt occurs.

A logical measure to identify how well the models align with the in situ observations is to maximize the number of days on which both the models and the in situ observations agree on either melting or no melting. However, since the number of days with melt is fewer than the number without melt, the dataset exhibits an imbalance. In the ablation zone, the imbalance is less pronounced, with prolonged periods of melt in the summer, whereas in the accumulation zone, melt occurs for shorter periods. This data imbalance means that many days with agreement between RCMs and AWSs can

be attributed to seasonal patterns, concealing disagreement in the melt season. Instead, we utilize the receiver operating characteristic (ROC) curve (Peterson et al., 1954) and the precision-recall (PR) curve (Davis and Goadrich, 2006) to provide a more nuanced understanding of the alignment between RCMs and AWSs. Given that TP is the number of true positives, TN is the number of true negatives, FP is the number of false positives, and FN is the number of false negatives, the ROC curve is a measure of the ability to distinguish between two classes across all thresholds and consists of a graph that shows the true positive rate ($TPR = \frac{TP}{TP+FN}$) vs. the false positive rate ($FPR = 1 - \frac{TN}{TN+FP}$). Here, we define the AWSs as true. Thus, we define a TP as a case where the RCM and AWSs agree that melt is present and a TN as




Figure 6. ROC and PR curves for possible RCM melting thresholds using varying AWS temperature thresholds. The black dot indicates the optimal melting threshold for each RCM given in Table 1.

a case where both the AWSs and the RCM observe no melt. When melt is observed at the AWSs but not by the RCM, it is defined as an FN and vice versa for FP. The ROC curve provides the total performance measure across all potential classification thresholds where a random model will produce a diagonal line; see Fig. 5 as an example. In contrast, a perfect model will have a ROC curve that is composed of the left and upper boundary lines. The goal is to choose a melting threshold that maximizes the TPR while minimizing the FPR. However, in an imbalanced dataset, it is possible to produce a good ROC curve by making a large number of FP predictions, especially when the positive class is rare. Thus, for an imbalanced dataset, it is important to also consider the PR curve, which is the fraction of TP among the positive predictions ($\text{Precision} = \frac{TP}{TP+FP}$) vs. the TP among the actual positives ($\text{Recall} = \frac{TP}{TP+FN}$). When evaluating what melting threshold to apply to each RCM, we seek the RCM melting thresholds and AWS temperature thresholds that maximize the TPR while minimizing the FPR and maximizing precision and recall.

Figure 6 shows the ROC curves and PR curves for possible RCM melting thresholds using different AWS temperature thresholds. The black dot indicates the optimal melt-

ing threshold for each RCM (Table 1). For HIRHAM5-ERA5 and HIRHAM5-ERA5, we see that both the best ROC curve and the best PR curve match, although they suggest very different melting thresholds. For MARv3.12 and RACMO2.3p2, the ROC curve suggests using temperature thresholds between -0.5 and 1 °C to find the best RCM melting threshold. However, the PR curve suggests a lower temperature threshold between -1.0 and -2.0 °C, which yields better results. Table 1 showcases the applied thresholds informed by PROMICE temperature observations.

To ensure consistency across datasets, all RCMs are re-gridded to a common grid – in this case, the ASCAT grid of 5.5 km resolution. For HIRHAM5 and RACMO2.3p2, we upscale data and apply the nearest neighbor interpolation method. For MARv3.12, we downscale data and use a cubic interpolation method. It is important to note that re-gridding can potentially introduce a bias into the RCM output, meaning systematic errors not associated with internal parameterization choices within the RCMs. The potential implication of the re-gridding biases is considered when choosing a baseline threshold. Here, the aim is to apply a baseline threshold to all RCMs independent of warm/cold bias within the RCMs. Therefore, the baseline threshold was set to the smallest value possible ($0.1 \text{ mm w.e. d}^{-1}$) without allowing

regridding biases to impact the number of melt days. When comparing the RCM melt days and ASCAT melt days, we apply a snow line mask to the RCMs similar to that used for ASCAT.

4 Results

Table 1 showcases the mean air temperature for July and August observed by the AWSs and modeled by the RCMs at the AWSs, suggesting a cold bias in HIRHAM5-ERA5, whereas MARv3.12 and HIRHAM5-ERA1 have a warm bias. However, the mean air temperature at selected stations highlights that the temperature bias is not systematic across the ice sheet. To evaluate the difference between the mean annual number of melt days in ASCAT and each RCM, we compute the RMSE for the whole ice sheet and each drainage basin (Rignot and Mouginot, 2012). Besides applying the in-situ-informed thresholds found in Table 1, we also apply the baseline threshold of $0.1 \text{ mm w.e. d}^{-1}$. Table 2 shows the mean annual number of melt days and the mean duration of the melt season for both the RCMs with the two thresholds applied and ASCAT. We define the start of the melt season as the point when at least one grid point experiences melting. The melt extent using the in-situ-informed thresholds tends to align better with the ASCAT-observed mean number of melt days and the duration of the melt season. Furthermore, when the baseline threshold of $0.1 \text{ mm w.e. d}^{-1}$ is applied, melting occurs in parts of the SW basin all year. We apply a Mann–Whitney U test (Mann and Whitney, 1947) to test whether using an in-situ-informed threshold has no effect. The p value in Table 2 suggests that it is very unlikely that there is no effect from using an in-situ-informed threshold.

The RMSEs between ASCAT and RCMs are showcased in Table 3, where we see that applying the in-situ-informed threshold improves the discrepancies between modeled melt and ASCAT, except for HIRHAM5-ERA1. RACMO2.3p2 and MARv3.12 show the biggest improvement, although the improvement is not evenly distributed between drainage basins. Since the in-situ-informed thresholds generally reduce the differences between RCMs and ASCAT compared to the baseline threshold of $0.1 \text{ mm w.e. d}^{-1}$ (Table 3), we apply only the in-situ-informed thresholds in the following. A detailed comparison of the number of melt days using the baseline threshold is provided in Appendix A, showing larger discrepancies between ASCAT and RCMs, corresponding with the findings in Table 3.

Figure 7 shows the spatial variability of the number of melt days both within each RCM (upper panel) and in comparison with ASCAT (lower panel). Looking at the spatial variability in the annual number of melt days for the RCMs and ASCAT (Fig. 4d), areas with > 100 d of melt lie near the lower ablation zone, but the extent of areas with > 100 d of melt varies between ASCAT and among the models. Most notable is HIRHAM5-ERA5, which has substantially larger

areas with > 120 d of melt. Areas with < 1 d of melt on average are shown in white in Fig. 7, illustrating areas where melt very rarely occurs. Again, there is large variability among the models in terms of modeling areas with almost no melt days.

While Table 3 illustrates the variability in error between ASCAT and RCMs across drainage basins, the lower panels in Fig. 7 reveal that the largest differences are concentrated near the ice margin across all basins. Generally, the SW and SE basins have the highest RMSEs, explained by relatively large areas with 20 or more days of difference in these basins (Fig. 7). Similarly, HIRHAM5-ERA5 exhibits a high RMSE in the western basins, corresponding to large areas where HIRHAM-ERA5 models 20 or more melt days compared to what ASCAT observes. Although the in-situ-informed threshold reduces the differences, HIRHAM5-ERA5 continues to produce more melt days than ASCAT or any other model, even when the melting threshold is vastly greater than that of the other RCMs. On the other hand, HIRHAM5-ERA1 consistently underestimates the number of melt days across all drainage basins. MARv3.12 and RACMO2p2.3 show similar patterns of variability, where the only large differences to ASCAT occur close to the maximum snow line elevation.

Results from all RCMs and ASCAT (Fig. 8a) indicate that, on average, the melting season starts at the beginning of May and culminates around July when the greatest melt extent occurs. While RCMs suggest that the melt season, on average, ends around mid-September, small melt areas are still observed in ASCAT well into October. The maximum melt extent is, on average, approximately 30 % of the ice sheet, except for HIRHAM5-ERA5 with > 35 %. At the beginning of the melt season, ASCAT detects the increase in melt extent 10–15 d earlier compared to when the RCMs simulate an increase in the melt extent. However, the decrease in ASCAT melt extent at the end of the melt season corresponds well to the modeled melt extent. In August and at the beginning of September, the melt extent decreases but with small periodical increases, which ASCAT also detects prematurely, although not as prominently as at the beginning of the melt season.

On 12 July 2012, an extreme melting event was observed across almost the entire ice sheet (Nghiem et al., 2012). We can use this year to showcase how well an extreme but relatively short melt event is captured in the RCMs and by ASCAT. Results show that when using the in-situ-informed thresholds, only RACMO2p2.3 simulates melting of > 90 % of the ice sheet, while the rest only predict approximately 80 % (Fig. 8b). Figure 8b also shows that liquid water is detected earlier by ASCAT at the start of the season. During the increase in melt extent from July to mid-August, ASCAT was able to capture an increase in melt extent, but the magnitude of the increase was smaller compared to that of the models.

Table 2. Mean annual melt days and mean duration of the melt season for each model using two different thresholds and for ASCAT. The melt season duration is defined as starting when at least one grid point experiences melting. A Mann–Whitney *U* test (Mann and Whitney, 1947) was applied to assess whether using an in-situ-informed threshold has no effect. Additionally, the rank-biserial correlation (*r* value, Cureton, 1956) was computed to indicate the magnitude of the difference between thresholds.

Threshold method	Mean melt days		Duration of melt season		<i>p</i> value	<i>r</i> value
	Baseline	In situ	Baseline	In situ		
HIRHAM5-ERA5	25	21	225	204	> 0.001	0.48
HIRHAM5-ERA1	16	16	205	201	> 0.001	0.49
RACMO2.3p2	24	18	365	344	> 0.001	0.47
MARv3.12	25	18	274	166	> 0.001	0.45
ASCAT	18		154		–	–

Table 3. RMSE of the mean annual number of melt days modeled by RCMs using a baseline threshold and in-situ-informed thresholds and ASCAT across the whole ice sheet and for each Rignot and Mouginot (2012) drainage basin. RMSE is only computed inside the snow line data mask to mask out areas where ASCAT cannot correctly detect wet snow; see Fig. 1.

Threshold	HIRHAM5-ERA5		HIRHAM5-ERA1		RACMO2.3p2		MARv3.12	
	0.1	4.1	0.1	0.4	0.1	0.7	0.1	1.0
Full ice sheet	10.70	7.67	7.39	7.76	28.77	6.27	17.78	4.93
NW	8.23	5.13	3.17	3.44	1.97	1.77	3.12	2.00
CW	9.01	5.80	3.30	3.74	2.10	2.10	3.11	2.65
SW	6.88	5.40	6.10	6.69	12.14	4.87	9.68	3.73
SE	5.65	4.89	5.91	6.07	31.56	5.78	18.75	4.08
NE	4.53	3.17	3.13	3.08	2.94	2.15	3.21	1.93
NO	5.11	3.35	2.49	2.68	1.56	1.38	1.66	1.76

5 Discussion

The goal of this study is to use ASCAT to understand how RCMs model ice-sheet-wide melt extent compared to observations and to further pinpoint biases within models that lead to the observed discrepancies. By comparing RCMs with PROMICE GC-net temperature observations, we are able to determine how well the RCMs align with in situ observations. While the uneven distribution of AWSs, concentrated in the ablation zone, may result in certain areas being better represented in the assessment than regions with fewer AWSs, Table 3 shows that by ensuring that the RCMs align with in situ measurements at specific locations, the modeled melt extent and satellite-observed melt extent show better agreement. Further, with this approach, we identify that HIRHAM5-ERA5 shows large deviations from PROMICE GC-net air temperature observations. The melt threshold in HIRHAM5-ERA5 is considerably higher than the remaining melt estimates, indicating that HIRHAM-ERA5 overestimates melt. The identified melting threshold of 4.1 mm w.e. d^{−1} suggests potential issues with the representation of melt in HIRHAM5 when forced with ERA5. Consequently, this highlights the importance of carefully considering model performance. When we compare the number of melt days in the RCMs with that observed by AS-

CAT, MARv3.12 has the lowest RMSE for the whole ice sheet. However, MARv3.12 performs predominantly better on the eastern side of the ice sheet, whereas RACMO2p2.3 performs better on the west coast and in the northernmost areas. While none of the RCMs exhibit perfect agreement with either PROMICE GC-net or ASCAT observations when looking at the RMSEs and the spatial differences between the annual number of melt days, neither MARv3.12 nor RACMO2p2.3 has an advantage over the other. On the other hand, HIRHAM5-ERA5 and HIRHAM5-ERA1 display considerable discrepancies in terms of the number of melt days when compared to ASCAT, each displaying unique patterns of deviation. Our analysis shows the value of using common, independent spatially and temporally varying datasets to evaluate and improve models, and we suggest that this ASCAT dataset is a useful addition to better understand biases in models, especially as it is independent of other datasets, such as MODIS, which has been used to develop models.

5.1 Biases in RCMs

To obtain the most valid comparison between each RCM and ASCAT, we utilize in situ observations to assess biases and to determine an appropriate threshold for the melt extent in RCMs. By fitting each RCM to in situ observa-




Figure 7. (a–d) The mean annual number of melt days modeled by the RCMs using an in-situ-informed melt threshold to defined days with significant melt. Pixels with < 1 d of melt on average are marked as white, showcasing areas where melt rarely occurs. (e–h) The mean annual difference between the number of melt days in ASCAT and RCMs areas above the 2007–2020 maximum snow line elevation (Fig. 4d). Red areas correspond to more melt days in ASCAT on average and blue areas correspond to more melt days in the RCM on average. Melt in ASCAT is defined as Label ST-2A and ST-2B.

tions, we minimize the differences that are introduced due to model setups, such as resolution and parameterization. Thus, we reduce overall inter-model discrepancies and differences in melt extent compared to those observed by ASCAT. Despite applying the in-situ-informed thresholds, persistent patterns between RCMs and ASCAT remain. HIRHAM5-ERA5 is the only model that, on average, predicts more melt days than ASCAT while also having a substantially higher melting threshold compared to the other RCM melt outputs; see Tables 1 and 2. Using the in-situ-informed thresholds, RACMO2.3p2 and MARv3.12 model the same mean number of melt days as ASCAT, while HIRHAM5-ERA1 mod-

els slightly fewer melt days. Looking at Fig. 7, HIRHAM5-ERA1, RACMO2.3p2, and MARv3.12 all model fewer melt days than ASCAT on average in the lower accumulation zone, indicating either a limitation in the observation of melt by ASCAT or an underrepresentation of melt by the RCMs in this area. However, the magnitude of disagreement between the RCMs and ASCAT varies from RCM to RCM and region to region (Table 3), again suggesting that the melt and melt extent in these areas are not well represented by the RCMs or that the melt threshold should be considered non-stationary according to local conditions. While HIRHAM-ERA5 exhibits the largest meltwater pro-




Figure 8. (a) Mean seasonal melt extent (%) modeled by RCMs using in-situ-informed thresholds and retrieved from ASCAT (2007–2020). In-situ-informed thresholds are given in Table 3. (b) Daily melt extent (%) modeled by RCMs using in-situ-informed thresholds and retrieved from ASCAT in 2012. In-situ-informed thresholds are given in Table 3. The melt extent in both (a) and (b) is constrained to areas above the maximum elevation of the snow line.

duction (Fig. 2), HIRHAM-ERA5 is also characterized by the lowest mean JJA air temperatures (Fig. 3a–d). In general, HIRHAM-ERA5, MARv3.12, and RACMO2.3p2, all forced with ERA5, tend to report lower air temperatures compared to HIRHAM5-ERA1. This trend likely originates from ERA5 exhibiting lower air temperatures than its predecessor, ERA1, as highlighted by Krebs-Kanzow et al. (2023). While some variability among the models forced with ERA5 exists, the models commonly exhibit a higher mean JJA downward shortwave radiation at the surface (SWD) compared to HIRHAM-ERA1. Krebs-Kanzow et al. (2023) similarly reported an overestimation of SWD in ERA5 compared to ERA1. Since the differences in air temperature and SWD between ERA5 and ERA1 have implications for meltwater production and ultimately runoff, Krebs-Kanzow et al. (2023) concluded that replacing ERA1 with ERA5 forcing in an energy balance model of the Greenland Ice Sheet requires some recalibration to reproduce existing observations. In this study, we show the implications for meltwater production and extent when forcing the HIRHAM5 SMB model with ERA5 instead of ERA1 without recalibration, though the inclusion of a different albedo scheme is likely to be more important.

In addition to the distinct forcing fields in the two HIRHAM5 outputs, the approach for determining surface albedo differs. The simple surface albedo scheme in HIRHAM-ERA5 results in a lower mean JJA surface albedo in the ablation zone and lower accumulation zone compared to that in HIRHAM-ERA1, which included MODIS observations in the surface albedo scheme (Fig. 3e and f). RACMO2.3p2 is characterized by the lowest surface albedo across the entire ice sheet, while MARv3.12 and HIRHAM5-ERA1 are dominated by lower surface albedo particularly in the accumulation zone. Surprisingly, while RACMO2.3p2 and HIRHAM5-ERA1 both report incorporating MODIS

bare ice albedo observations into their surface albedo computation in the ablation zone, the resulting surface albedo differs greatly between the models. In the RCMs, the surface albedo is a crucial parameter for simulating the surface energy balance, contributing to a higher surface energy balance when the surface albedo is low. A low albedo in the ablation zone and lower accumulation zone will increase the meltwater production in these areas. Due to the high variability of air temperature, SWD, and surface albedo among the models, the albedo parameterization and radiation and temperature schemes within the models should be critically assessed using high-quality measurements to reduce the observed discrepancies in the models' estimates of meltwater production (Fig. 2). By aligning simulated melt rates more closely with observational data, we can improve the model estimates of meltwater production and, ultimately, runoff.

5.2 Limitations of ASCAT melt observations

We use the maximum extent of the snow line to mask out the ablation zone when comparing the RCMs to ASCAT due to nonphysical observations of refreezing by ASCAT in these areas (Fig. 7). This is not only due to biases in the melt classification algorithm but also because of the microwave data themselves. Although melt causes a drop in the backscatter signal, processes other than the refreezing of liquid water can counteract the decrease in the backscatter signal. Once the surface starts to melt, the surface roughness changes, which can cause an increase in the backscatter signal, potentially concealing the liquid water and melting in the backscatter signal. Since the effect of changing surface roughness on the backscatter signal is not fully understood, it is difficult to remove this bias in wet-snow detection other than by removing areas where changes in surface roughness are very pronounced. We further see this systematic misclassification

of refreeze or no melting in ASCAT when investigating the 2012 melt season. Here, ASCAT detects a lower melt extent in the late melt season compared to the simulated RCM melt extent. Since the refreezing periods identified from ASCAT data are not included in the melt season analysis, misclassifications that stem from melting being misclassified as refreezing due to changes in surface roughness are not included, potentially contributing to a smaller melt extent at the end of the melt season.

As seen in both the 2012 melt season (Fig. 8b) and the annual mean cycle of melt extent (Fig. 8a), ASCAT detects the increase in melt extent before the RCMs, which is partly explained by the preprocessing averaging done to enhance the spatial resolution. However, based on Fig. 7, we show an average offset of 10–15 d between the increase in ASCAT melt extent and RCM melt extent, which the preprocessing averaging cannot fully explain. Although it is most pronounced at the beginning of the season, we see that a similar offset also occurs when there is an increase in the melt extent in the late season. On average, the magnitude of the seasonal cycle of melt extent agrees well with that of the RCMs, suggesting that liquid water is observed earlier but to a similar extent.

Finally, the ASCAT backscatter varies due to additional factors such as specific properties of the snow, e.g., grain size and the presence of ice, due to its influence on dielectric properties and roughness geometries. Here, there are two possible approaches for improving ASCAT melt retrievals: we could use the surface properties from either the individual RCMs or in situ observations. However, there is a seasonal bias in the in situ observations and a lack of spatial coverage, making them difficult to use for ice-sheet-wide Earth observation data production. As for the RCM implementation, this would hamper the use of ASCAT data as an independent data record.

6 Conclusions

ASCAT wet-snow maps provide a valuable tool for evaluating the performance of RCMs, which is currently the only source for assessing the melt volume of the Greenland Ice Sheet on a global scale. By utilizing observations of melt extent by ASCAT, we can evaluate how well four different RCMs' melt outputs represent melt spatially and temporally across the Greenland Ice Sheet in the present day. To ensure that the RCM-modeled melt aligns well with in situ observations, we compare the RCM melt output at the PROMICE GC-net AWS. Assuming a strong correlation between 2 m air temperature and melting, we can use PROMICE air temperature measurements to assess temperature biases and determine a melting threshold in the RCMs to identify days with significant melt. Here, we find that HIRHAM5 forced with ERA5 shows potential issues with the representation of melt due to its relatively poor alignment with PROMICE air temperature measurements. HIRHAM5

forced with ERA-Interim, MARv3.12, and RACMO2p2.3 all show similar alignment with PROMICE. When comparing with ASCAT melt extent, MARv3.12 and RACMO2p2.3 show better agreement with ASCAT, but each RCM shows distinct patterns of variations. This suggests that in some cases, one RCM can represent present-day melt extent across the entire ice sheet more effectively than the ensemble mean, as the variability among the models does not appear to be random variations. For all RCMs included in this study, the onset of melt occurs more than a week later than what is observed from ASCAT, even when accounting for the averaging of satellite data. Further, we see that the regions with the largest differences in the total number of melt days across all RCMs are the SW and SE basins of the ice sheet, an indication that melt is likely not well represented in the RCMs in these areas.

The variability among modeled JJA air temperature, SWD, and surface albedo can explain the large discrepancies in modeled meltwater production (Fig. 2). Notably, we show the implications for meltwater production when running the HIRHAM5 SMB model using ERA5 instead of ERAI without any recalibration. Further, ensuring that the models accurately simulate the variability in albedo, such as by incorporating MODIS bare ice data, can lead to a more accurate representation of the surface energy balance and, consequently, meltwater production. Despite the fact that both RACMO2.3p3 and HIRHAM5-ERA5 use similar MODIS bare ice observations and have similar dynamical schemes, the surface albedo and its effects on meltwater production varied substantially between models, highlighting the need for a critical evaluation of model outputs against high-quality measurements to reduce inter-model discrepancies. In general, our analysis demonstrates the value of using independent datasets like ASCAT wet-snow maps to identify the spatiotemporal variability of RCM-simulated melt. This approach complements traditional model validation methods and intercomparison exercises, which can inform future model development to better simulate ice sheet surface melt and potentially be incorporated into future MIP efforts.

Appendix A: Comparison between RCMs and ASCAT using the baseline threshold of $0.1 \text{ mm w.e. d}^{-1}$




Figure A1. (a–d) The mean annual number of melt days modeled by the RCMs using a baseline threshold ($0.1 \text{ mm w.e. d}^{-1}$) to define days with significant melt. Pixels with $< 1 \text{ d}$ of melt on average are indicated in white, showcasing areas where melt rarely occurs. (e–h) The mean annual difference between the number of melt days in ASCAT and RCM areas above the 2007–2020 maximum snow line elevation (Fig. 4d). The red areas correspond to more melt days in ASCAT on average, and the blue areas correspond to more melt days in the RCMs on average. Melt in ASCAT is defined as ST-2A and ST-2B.

Code availability. Code is publicly available at https://github.com/anpug/4DG_meltmaps.git (Puggaard, 2025).

Data availability. ASCAT wet-snow maps are available at <https://cryoportal.enveo.at/data/> (Nagler et al., 2024) under meltmap. PROMICE GC-net AWS 2 m air temperature data are available on the GEUS Dataverse (<https://doi.org/10.22008/FK2/IW73UU>, How et al., 2022). Further, historical GC-net 2 m air temperature data are available on the GEUS Dataverse (<https://doi.org/10.22008/FK2/VVXGUT>, Steffen et al., 2022). HIRHAM5 forced with ERA-Interim is available at https://ensemblesrt3.dmi.dk/data/prudence/temp/nichan/Daily2D_GrIS/ (Langen et al., 2017), and HIRHAM5 forced with ERA5 will be made available once the paper has been published (<https://doi.org/10.11583/DTU.25568040>). RACMO3.2p2 is freely available upon request (Noël et al., 2019). MARv3.12 is available at <ftp://ftp.climato.be/fettweis/MARv3.12/Greenland> (Tedesco and Fettweis, 2020).

Author contributions. AMS and LSS framed the original study, and AP further developed the paper and wrote the first draft with the help of AMS, NH, LSS, RM, and SBS. TN, SS, and JW developed the presented ASCAT classification algorithm and provided support for the data analysis and discussion. NH processed and provided the HIRHAM5 data. All authors revised the paper.

Competing interests. At least one of the (co-)authors is a member of the editorial board of *The Cryosphere*. The peer-review process was guided by an independent editor, and the authors also have no other competing interests to declare.

Disclaimer. Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes every effort to include appropriate place names, the final responsibility lies with the authors.

Acknowledgements. The authors thank the editor and the two anonymous reviewers for their constructive comments and feedback. Anna Puggaard, Nicolaj Hansen, and Ruth Mottram are supported by the Danish State through the National Centre for Climate Research (NCKF). Furthermore, Nicolaj Hansen is supported by the Novo Nordisk Foundation project PRECISE (NNF23OC0081251). Model simulations and analysis in this publication were supported by the PROTECT project funded by the European Union's Horizon 2020 research and innovation program under grant agreement no. 869304. Thomas Nagler, Jan Wuite, Anne M. Solgaard, and Stefan Scheiblaue acknowledge support from the European Space Agency through the POLAR+ 4DGreenland (ESA contract no. 4000132139/20/I-EF) and 4DAntarctica projects (ESA contract no. 801ESA/AO/1-9570/18/I-DT). Further, Anne M. Solgaard is supported by the Programme for Monitoring of the Greenland Ice Sheet (PROMICE).

Financial support. This research has been supported by the Danish State through the National Centre for Climate Research (NCKF).

Review statement. This paper was edited by Kristin Poinar and reviewed by two anonymous referees.

References

- Abdalati, W. and Steffen, K.: Passive microwave derived snow melt regions on the Greenland Ice Sheet, *Geophys. Res. Lett.*, 22, 787–790, <https://doi.org/10.1029/95GL00433>, 1995.
- Antwerpen, R. M., Tedesco, M., Fettweis, X., Alexander, P., and van de Berg, W. J.: Assessing bare-ice albedo simulated by MAR over the Greenland ice sheet (2000–2021) and implications for meltwater production estimates, *The Cryosphere*, 16, 4185–4199, <https://doi.org/10.5194/tc-16-4185-2022>, 2022.
- Ashcraft, I. S. and Long, D. G.: Comparison of methods for melt detection over Greenland using active and passive microwave measurements, *Int. J. Remote Sens.*, 27, 2469–2488, <https://doi.org/10.1080/01431160500534465>, 2006.
- Box, J. E., Fettweis, X., Stroeve, J. C., Tedesco, M., Hall, D. K., and Steffen, K.: Greenland ice sheet albedo feedback: thermodynamics and atmospheric drivers, *The Cryosphere*, 6, 821–839, <https://doi.org/10.5194/tc-6-821-2012>, 2012.
- Brangers, I., Lievens, H., Miège, C., Demuzere, M., Brucker, L., and De Lannoy, G. J. M.: Sentinel 1 detects firm aquifers in the Greenland Ice Sheet, *Geophys. Res. Lett.*, 47, e2019GL085192, <https://doi.org/10.1029/2019GL085192>, 2020.
- Brun, E., David, P., Sudul, M., and Brunot, G.: A numerical model to simulate snow-cover stratigraphy for operational avalanche forecasting, *J. Glaciol.*, 38, 13–22, <https://doi.org/10.3189/S0022143000009552>, 1992.
- Chandler, D. M., Wadham, J. L., Lis, G. P., Cowton, T., Sole, A., Bartholomew, I., Telling, J., Nienow, P., Bagshaw, E. B., Mair, D., Vinen, S., and Hubbard, A.: Evolution of the subglacial drainage system beneath the Greenland Ice Sheet revealed by tracers, *Nat. Geosci.*, 6, 195–198, <https://doi.org/10.1038/ngeo1737>, 2013.
- Colosio, P., Tedesco, M., Ranzi, R., and Fettweis, X.: Surface melting over the Greenland ice sheet derived from enhanced resolution passive microwave brightness temperatures (1979–2019), *The Cryosphere*, 15, 2623–2646, <https://doi.org/10.5194/tc-15-2623-2021>, 2021.
- Cuffey, K. and Paterson, W. S. B.: *The Physics of Glaciers*, Butterworth-Heinemann/Elsevier, Burlington, MA, 4th edn., oCLC: ocn488732494, ISBN 9780123694614, 2010.
- Cureton, E. E.: Rank-biserial correlation, *Psychometrika*, 21, 287–290, <https://doi.org/10.1007/BF02289138>, 1956.
- Davis, J. and Goadrich, M.: The relationship between Precision-Recall and ROC curves, in: *Proceedings of the 23rd international conference on Machine learning, ICML '06*, Association for Computing Machinery, New York, NY, USA, <https://doi.org/10.1145/1143844.1143874>, 233–240, 2006.
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., Van De Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M.,

- Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, L., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., De Rosnay, P., Tavalato, C., Thépaut, J.-N., and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Q. J. Roy. Meteor. Soc.*, 137, 553–597, <https://doi.org/10.1002/qj.828>, 2011.
- Dethinne, T., Glaude, Q., Picard, G., Kittel, C., Alexander, P., Orban, A., and Fettweis, X.: Sensitivity of the MAR regional climate model snowpack to the parameterization of the assimilation of satellite-derived wet-snow masks on the Antarctic Peninsula, *The Cryosphere*, 17, 4267–4288, <https://doi.org/10.5194/tc-17-4267-2023>, 2023.
- ECMWF: IFS Documentation CY33R1 – Part IV: Physical Processes, <https://www.ecmwf.int/node/9227> (last access: 14 December 2023) 2009.
- Ettema, J., van den Broeke, M. R., van Meijgaard, E., van de Berg, W. J., Box, J. E., and Steffen, K.: Climate of the Greenland ice sheet using a high-resolution climate model – Part 1: Evaluation, *The Cryosphere*, 4, 511–527, <https://doi.org/10.5194/tc-4-511-2010>, 2010.
- Fausto, R. S., Andersen, S. B., Ahlstrøm, A. P., As, D. v., Box, J. E., Binder, D., Citterio, M., Colgan, W., Haubner, K., Hansen, K., Karlsson, N. B., Mankoff, K. D., Pedersen, A. Ø., Solgaard, A., and Vandecrux, B.: The Greenland ice sheet – snowline elevations at the end of the melt seasons from 2000 to 2017, *Geol. Surv. Den. Greenl.*, 41, 71–74, <https://doi.org/10.34194/geusb.v41.4346>, 2018.
- Fausto, R. S., van As, D., Mankoff, K. D., Vandecrux, B., Citterio, M., Ahlstrøm, A. P., Andersen, S. B., Colgan, W., Karlsson, N. B., Kjeldsen, K. K., Korsgaard, N. J., Larsen, S. H., Nielsen, S., Pedersen, A. Ø., Shields, C. L., Solgaard, A. M., and Box, J. E.: Programme for Monitoring of the Greenland Ice Sheet (PROMICE) automatic weather station data, *Earth Syst. Sci. Data*, 13, 3819–3845, <https://doi.org/10.5194/essd-13-3819-2021>, 2021.
- Fettweis, X., Tedesco, M., van den Broeke, M., and Ettema, J.: Melting trends over the Greenland ice sheet (1958–2009) from spaceborne microwave data and regional climate models, *The Cryosphere*, 5, 359–375, <https://doi.org/10.5194/tc-5-359-2011>, 2011.
- Fettweis, X., Franco, B., Tedesco, M., van Angelen, J. H., Lenaerts, J. T. M., van den Broeke, M. R., and Gallée, H.: Estimating the Greenland ice sheet surface mass balance contribution to future sea level rise using the regional atmospheric climate model MAR, *The Cryosphere*, 7, 469–489, <https://doi.org/10.5194/tc-7-469-2013>, 2013.
- Fettweis, X., Box, J. E., Agosta, C., Amory, C., Kittel, C., Lang, C., van As, D., Machguth, H., and Gallée, H.: Reconstructions of the 1900–2015 Greenland ice sheet surface mass balance using the regional climate MAR model, *The Cryosphere*, 11, 1015–1033, <https://doi.org/10.5194/tc-11-1015-2017>, 2017.
- Fettweis, X., Hofer, S., Krebs-Kanzow, U., Amory, C., Aoki, T., Berends, C. J., Born, A., Box, J. E., Delhasse, A., Fujita, K., Gierz, P., Goelzer, H., Hanna, E., Hashimoto, A., Huybrechts, P., Kapsch, M.-L., King, M. D., Kittel, C., Lang, C., Langen, P. L., Lenaerts, J. T. M., Liston, G. E., Lohmann, G., Mernild, S. H., Mikolajewicz, U., Modali, K., Mottram, R. H., Niwano, M., Noël, B., Ryan, J. C., Smith, A., Streffing, J., Tedesco, M., van de Berg, W. J., van den Broeke, M., van de Wal, R. S. W., van Kampenhou, L., Wilton, D., Wouters, B., Ziemen, F., and Zolles, T.: GrSMBMIP: intercomparison of the modelled 1980–2012 surface mass balance over the Greenland Ice Sheet, *The Cryosphere*, 14, 3935–3958, <https://doi.org/10.5194/tc-14-3935-2020>, 2020.
- Figa-Saldaña, J., Wilson, J. J., Attema, E., Gelsthorpe, R., Drinkwater, M. R., and Stoffelen, A.: The advanced scatterometer (ASCAT) on the meteorological operational (MetOp) platform: a follow on for European wind scatterometers, *Can. J. Remote Sens.*, 28, 404–412, <https://doi.org/10.5589/m02-035>, 2002.
- Forster, R. R., Box, J. E., Van Den Broeke, M. R., Miège, C., Burgess, E. W., Van Angelen, J. H., Lenaerts, J. T. M., Koenig, L. S., Paden, J., Lewis, C., Gogineni, S. P., Leuschen, C., and McConnell, J. R.: Extensive liquid meltwater storage in firn within the Greenland ice sheet, *Nat. Geosci.*, 7, 95–98, <https://doi.org/10.1038/ngeo2043>, 2014.
- Gallée, H. and Schayes, G.: Development of a three-dimensional meso- γ primitive equation model: katabatic winds simulation in the area of Terra Nova Bay, Antarctica, *Mon. Weather Rev.*, 122, 671–685, [https://doi.org/10.1175/1520-0493\(1994\)122<0671:DOATDM>2.0.CO;2](https://doi.org/10.1175/1520-0493(1994)122<0671:DOATDM>2.0.CO;2), 1994.
- Glaude, Q., Noel, B., Olesen, M., Van Den Broeke, M., Van De Berg, W. J., Mottram, R., Hansen, N., Delhasse, A., Amory, C., Kittel, C., Goelzer, H., and Fettweis, X.: A factor two difference in 21st century Greenland Ice Sheet surface mass balance projections from three regional climate models under a strong warming scenario (SSP5 8.5), *Geophys. Res. Lett.*, 51, e2024GL111902, <https://doi.org/10.1029/2024GL111902>, 2024.
- Goelzer, H., Nowicki, S., Payne, A., Larour, E., Seroussi, H., Lipscomb, W. H., Gregory, J., Abe-Ouchi, A., Shepherd, A., Simon, E., Agosta, C., Alexander, P., Aschwanden, A., Barthel, A., Calov, R., Chambers, C., Choi, Y., Cuzzone, J., Dumas, C., Edwards, T., Felikson, D., Fettweis, X., Golledge, N. R., Greve, R., Humbert, A., Huybrechts, P., Le clec'h, S., Lee, V., Leguy, G., Little, C., Lowry, D. P., Morlighem, M., Nias, I., Quiquet, A., Rückamp, M., Schlegel, N.-J., Slater, D. A., Smith, R. S., Straneo, F., Tarasov, L., van de Wal, R., and van den Broeke, M.: The future sea-level contribution of the Greenland ice sheet: a multi-model ensemble study of ISMIP6, *The Cryosphere*, 14, 3071–3096, <https://doi.org/10.5194/tc-14-3071-2020>, 2020.
- Harper, J., Humphrey, N., Pfeffer, W. T., Brown, J., and Fettweis, X.: Greenland ice-sheet contribution to sea-level rise buffered by meltwater storage in firn, *Nature*, 491, 240–243, <https://doi.org/10.1038/nature11566>, 2012.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R. J., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., De Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.: The ERA5 global reanalysis, *Q. J. Roy. Meteor. Soc.*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.
- How, P., Abermann, J., Ahlstrøm, A., Andersen, S., Box, J. E., Citterio, M., Colgan, W., Fausto, R., Karlsson, N., Jakobsen, J., Langley, K., Larsen, S., Mankoff, K., Pedersen, A.,

- Rutishauser, A., Shield, C., Solgaard, A., Van As, D., Vandecrux, B., and Wright, P.: PROMICE and GC-Net automated weather station data in Greenland, GEUS Dataverse [data set], <https://doi.org/10.22008/FK2/IW73UU>, 2022.
- Husman, S. d. R., Hu, Z., Wouters, B., Munneke, P. K., Veldhuijsen, S., and Lhermitte, S.: Remote sensing of surface melt on Antarctica: opportunities and challenges, *IEEE J. Sel. Top. Appl.*, 16, 2462–2480, <https://doi.org/10.1109/JSTARS.2022.3216953>, 2023.
- Koenig, L. S., Lampkin, D. J., Montgomery, L. N., Hamilton, S. L., Turrin, J. B., Joseph, C. A., Moutsafa, S. E., Panzer, B., Casey, K. A., Paden, J. D., Leuschen, C., and Gogineni, P.: Wintertime storage of water in buried supraglacial lakes across the Greenland Ice Sheet, *The Cryosphere*, 9, 1333–1342, <https://doi.org/10.5194/tc-9-1333-2015>, 2015.
- Krebs-Kanzow, U., Rodehacke, C. B., and Lohmann, G.: Brief communication: Surface energy balance differences over Greenland between ERA5 and ERA-Interim, *The Cryosphere*, 17, 5131–5136, <https://doi.org/10.5194/tc-17-5131-2023>, 2023.
- Langen, P. L., Mottram, R. H., Christensen, J. H., Boberg, F., Rodehacke, C. B., Stendel, M., Van As, D., Ahlstrøm, A. P., Mortensen, J., Rysgaard, S., Petersen, D., Svendsen, K. H., Aðalgeirsdóttir, G., and Cappelen, J.: Quantifying energy and mass fluxes controlling Godthåbsfjord freshwater input in a 5-km simulation (1991–2012), *J. Climate*, 28, 3694–3713, <https://doi.org/10.1175/JCLI-D-14-00271.1>, 2015.
- Langen, P. L., Fausto, R. S., Vandecrux, B., Mottram, R. H., and Box, J. E.: Liquid water flow and retention on the Greenland Ice Sheet in the regional climate model HIRHAM5: local and large-scale impacts, *Front. Earth Sci.*, 4, <https://doi.org/10.3389/feart.2016.00110>, 2017 (data available at: https://ensemblesrt3.dmi.dk/data/prudence/temp/nichan/Daily2D_GrIS/, last access: 11 July 2025).
- Lindsley, R. D. and Long, D. G.: Adapting the sir algorithm to ASCAT, in: 2010 IEEE International Geoscience and Remote Sensing Symposium, 7–12 July 2024, Athens, Greece, <https://doi.org/10.1109/IGARSS.2010.5650207>, 3402–3405, 2010.
- Long, D. G.: Polar applications of spaceborne scatterometers, *IEEE J. Sel. Top. Appl.*, 10, 2307–2320, <https://doi.org/10.1109/JSTARS.2016.2629418>, 2017.
- Long, D. G. and Drinkwater, M. R.: Greenland ice-sheet surface properties observed by the Seasat-A scatterometer at enhanced resolution, *J. Glaciol.*, 40, 213–230, <https://doi.org/10.3189/S0022143000007310>, 1994.
- Long, D., Hardin, P., and Whiting, P.: Resolution enhancement of spaceborne scatterometer data, *IEEE T. Geosci. Remote*, 31, 700–715, <https://doi.org/10.1109/36.225536>, 1993.
- Lucas Picher, P., Wulff Nielsen, M., Christensen, J. H., Aðalgeirsdóttir, G., Mottram, R., and Simonsen, S. B.: Very high resolution regional climate model simulations over Greenland: identifying added value, *J. Geophys. Res.-Atmos.*, 117, 2011JD016267, <https://doi.org/10.1029/2011JD016267>, 2012.
- Mann, H. B. and Whitney, D. R.: On a test of whether one of two random variables is stochastically larger than the other, *Ann. Math. Stat.*, 18, 50–60, <https://doi.org/10.1214/aoms/1177730491>, 1947.
- Mätzler, C.: Applications of the interaction of microwaves with the natural snow cover, *Remote Sensing Reviews*, 2, 259–387, <https://doi.org/10.1080/02757258709532086>, 1987.
- Nagler, T., Wuite, J., Rott, H., Scheiblauer, S., Puggaard, A., and Prieto, D. F.: Monitoring Ice Sheet Melt and Re-freeze Using Active Microwave Measurements, in: *IGARSS 2024–2024 IEEE International Geoscience and Remote Sensing Symposium*, 25–30 July 2010, Honolulu, Hawaii, USA, <https://doi.org/10.1109/IGARSS53475.2024.10640970>, 49–52, 2024 (data available at: <https://cryoportal.enveo.at/data/>, last access: 5 August 2025).
- Nghiem, S., Steffen, K., Kwok, R., and Tsai, W. Y.: Detection of snowmelt regions on the Greenland ice sheet using diurnal backscatter change, *J. Glaciol.*, 47, 539–547, <https://doi.org/10.3189/172756501781831738>, 2001.
- Nghiem, S. V., Hall, D. K., Mote, T. L., Tedesco, M., Alibert, M. R., Keegan, K., Shuman, C. A., DiGirolamo, N. E., and Neumann, G.: The extreme melt across the Greenland ice sheet in 2012, *Geophys. Res. Lett.*, 39, 2012GL053611, <https://doi.org/10.1029/2012GL053611>, 2012.
- Noël, B., van de Berg, W. J., Machguth, H., Lhermitte, S., Howat, I., Fettweis, X., and van den Broeke, M. R.: A daily, 1 km resolution data set of downscaled Greenland ice sheet surface mass balance (1958–2015), *The Cryosphere*, 10, 2361–2377, <https://doi.org/10.5194/tc-10-2361-2016>, 2016.
- Noël, B., van de Berg, W. J., van Wessem, J. M., van Meijgaard, E., van As, D., Lenaerts, J. T. M., Lhermitte, S., Kuipers Munneke, P., Smeets, C. J. P. P., van Uft, L. H., van de Wal, R. S. W., and van den Broeke, M. R.: Modelling the climate and surface mass balance of polar ice sheets using RACMO2 – Part 1: Greenland (1958–2016), *The Cryosphere*, 12, 811–831, <https://doi.org/10.5194/tc-12-811-2018>, 2018.
- Noël, B., Van De Berg, W. J., Lhermitte, S., and Van Den Broeke, M. R.: Rapid ablation zone expansion amplifies north Greenland mass loss, *Science Advances*, 5, eaaw0123, <https://doi.org/10.1126/sciadv.aaw0123>, 2019.
- Otosaka, I. N., Shepherd, A., Ivins, E. R., Schlegel, N.-J., Amory, C., van den Broeke, M. R., Horwath, M., Joughin, I., King, M. D., Krinner, G., Nowicki, S., Payne, A. J., Rignot, E., Scambos, T., Simon, K. M., Smith, B. E., Sørensen, L. S., Velicogna, I., Whitehouse, P. L., A, G., Agosta, C., Ahlstrøm, A. P., Blazquez, A., Colgan, W., Engdahl, M. E., Fettweis, X., Forsberg, R., Gallée, H., Gardner, A., Gilbert, L., Gourmelen, N., Groh, A., Gunter, B. C., Harig, C., Helm, V., Khan, S. A., Kittel, C., Konrad, H., Langen, P. L., Lecavalier, B. S., Liang, C.-C., Loomis, B. D., McMillan, M., Melini, D., Mernild, S. H., Mottram, R., Mouginot, J., Nilsson, J., Noël, B., Pattie, M. E., Peltier, W. R., Pie, N., Roca, M., Sasgen, I., Save, H. V., Seo, K.-W., Scheuchl, B., Schrama, E. J. O., Schröder, L., Simonsen, S. B., Slater, T., Spada, G., Sutterley, T. C., Vishwakarma, B. D., van Wessem, J. M., Wiese, D., van der Wal, W., and Wouters, B.: Mass balance of the Greenland and Antarctic ice sheets from 1992 to 2020, *Earth Syst. Sci. Data*, 15, 1597–1616, <https://doi.org/10.5194/essd-15-1597-2023>, 2023.
- Peterson, W., Birdsall, T., and Fox, W.: The theory of signal detectability, *Transactions of the IRE Professional Group on Information Theory*, 4, 171–212, <https://doi.org/10.1109/TIT.1954.1057460>, 1954.

- Picard, G., Leduc-Leballeur, M., Banwell, A. F., Brucker, L., and Macelloni, G.: The sensitivity of satellite microwave observations to liquid water in the Antarctic snowpack, *The Cryosphere*, 16, 5061–5083, <https://doi.org/10.5194/tc-16-5061-2022>, 2022.
- Puggaard, A.: *anpug/4DG_meltmaps: v1.0.0 (v1.0.0)*, Zenodo [data set], <https://doi.org/10.5281/zenodo.15480109>, 2025.
- Rae, J. G. L., Aðalgeirsdóttir, G., Edwards, T. L., Fettweis, X., Gregory, J. M., Hewitt, H. T., Lowe, J. A., Lucas-Picher, P., Mottram, R. H., Payne, A. J., Ridley, J. K., Shannon, S. R., van de Berg, W. J., van de Wal, R. S. W., and van den Broeke, M. R.: Greenland ice sheet surface mass balance: evaluating simulations and making projections with regional climate models, *The Cryosphere*, 6, 1275–1294, <https://doi.org/10.5194/tc-6-1275-2012>, 2012.
- Rignot, E. and Mouginot, J.: Ice flow in Greenland for the International Polar Year 2008–2009, *Geophys. Res. Lett.*, 39, L11501, <https://doi.org/10.1029/2012GL051634>, 2012.
- Roeckner, E., Bäuml, G., Bonaventura, L., Brokopf, R., Esch, M., Giorgetta, M., Hagemann, S., Kirchner, I., Kornblüeh, L., Manzini, E., Rhodin, A., Schlese, U., Schulzweida, U., and Tompkins, A.: The atmospheric general circulation model ECHAM 5. PART I: model description, Max Planck Institute for Meteorology Report, https://pure.mpg.de/rest/items/item_995269_2/component/file_995268/content (last access: 4 August 2025) 349, 2003.
- Smith, L. C., Chu, V. W., Yang, K., Gleason, C. J., Pitcher, L. H., Rennermalm, A. K., Legleiter, C. J., Behar, A. E., Overstreet, B. T., Moustafa, S. E., Tedesco, M., Forster, R. R., LeWinter, A. L., Finnegan, D. C., Sheng, Y., and Balog, J.: Efficient meltwater drainage through supraglacial streams and rivers on the southwest Greenland ice sheet, *P. Natl. Acad. Sci. USA*, 112, 1001–1006, <https://doi.org/10.1073/pnas.1413024112>, 2015.
- Steffen, K., Vandecrux, B., Houtz, D., Abdalati, W., Bayou, N., Box, J., Colgan, W., Espona Pernas, L., Griessinger, N., Haas-Artho, D., Heilig, A., Hubert, A., Iosifescu Enescu, I., Johnson-Amin, N., Karlsson, N., Kurup Buchholz, R., McGrath, D., Cullen, N., Naderpour, R., Molotch, N., Pedersen, A., Perren, B., Philipps, T., Plattner, G.-K., Proksch, M., Revheim, M., Særrelse, M., Schneebli, M., Sampson, K., Starkweather, S., Steffen, S., Stroeve, J., Watler, B., Winton, Ø A., Zwally, J., and Ahlstrøm, A.: GC-Net Level 1 historical automated weather station, GEUS Dataverse [data set], <https://doi.org/10.22008/FK2/VVXGUT>, 2022.
- Tedesco, M.: Snowmelt detection over the Greenland ice sheet from SSM/I brightness temperature daily variations, *Geophys. Res. Lett.*, 34, 2006GL028466, <https://doi.org/10.1029/2006GL028466>, 2007.
- Tedesco, M. and Fettweis, X.: Unprecedented atmospheric conditions (1948–2019) drive the 2019 exceptional melting season over the Greenland ice sheet, *The Cryosphere*, 14, 1209–1223, <https://doi.org/10.5194/tc-14-1209-2020>, 2020 (data available at: <ftp://ftp.climato.be/fettweis/MARv3.12/Greenland>, last access: May 2023).
- Undén, P., Rontu, L., Järvinen, H., Lynch, P., Calvo-Sanchez, J., Cats, G., Cuxart, J., Eerola, K., Fortelius, C., and García-Moya, J.: HIRLAM-5 Scientific Documentation, <https://scispace.com/pdf/hirlam-5-scientific-documentation-31lwtaucm1.pdf> (last access: 5 August 2025) 2002.
- van Angelen, J. H., Lenaerts, J. T. M., Lhermitte, S., Fettweis, X., Kuipers Munneke, P., van den Broeke, M. R., van Meijgaard, E., and Smeets, C. J. P. P.: Sensitivity of Greenland Ice Sheet surface mass balance to surface albedo parameterization: a study with a regional climate model, *The Cryosphere*, 6, 1175–1186, <https://doi.org/10.5194/tc-6-1175-2012>, 2012.
- van den Broeke, M. R., Enderlin, E. M., Howat, I. M., Kuipers Munneke, P., Noël, B. P. Y., van de Berg, W. J., van Meijgaard, E., and Wouters, B.: On the recent contribution of the Greenland ice sheet to sea level change, *The Cryosphere*, 10, 1933–1946, <https://doi.org/10.5194/tc-10-1933-2016>, 2016.
- Vandecrux, B., Box, J. E., Ahlstrøm, A. P., Andersen, S. B., Bayou, N., Colgan, W. T., Cullen, N. J., Fausto, R. S., Haas-Artho, D., Heilig, A., Houtz, D. A., How, P., Iosifescu Enescu, I., Karlsson, N. B., Kurup Buchholz, R., Mankoff, K. D., McGrath, D., Molotch, N. P., Perren, B., Revheim, M. K., Rutishauser, A., Sampson, K., Schneebli, M., Starkweather, S., Steffen, S., Weber, J., Wright, P. J., Zwally, H. J., and Steffen, K.: The historical Greenland Climate Network (GC-Net) curated and augmented level-1 dataset, *Earth Syst. Sci. Data*, 15, 5467–5489, <https://doi.org/10.5194/essd-15-5467-2023>, 2023.
- Vernon, C. L., Bamber, J. L., Box, J. E., van den Broeke, M. R., Fettweis, X., Hanna, E., and Huybrechts, P.: Surface mass balance model intercomparison for the Greenland ice sheet, *The Cryosphere*, 7, 599–614, <https://doi.org/10.5194/tc-7-599-2013>, 2013.
- WCRP Global Sea Level Budget Group: Global sea-level budget 1993–present, *Earth Syst. Sci. Data*, 10, 1551–1590, <https://doi.org/10.5194/essd-10-1551-2018>, 2018.
- Wismann, V.: Monitoring of seasonal snowmelt on Greenland with ERS scatterometer data, *IEEE T. Geosci. Remote*, 38, 1821–1826, <https://doi.org/10.1109/36.851766>, 2000.