



Investigating the impact of reanalysis snow input on an observationally calibrated snow-on-sea-ice reconstruction

Alex Cabaj¹, Paul J. Kushner², and Alek A. Petty³

¹Climate Research Division, Environment and Climate Change Canada, Toronto, ON M3H 5T4, Canada

²Department of Physics, University of Toronto, Toronto, ON M5S 1A7, Canada

³Earth System Science Interdisciplinary Center, University of Maryland, College Park, MD 20740, USA

Correspondence: Alex Cabaj (alex.cabaj@ec.gc.ca)

Received: 14 August 2024 – Discussion started: 2 September 2024

Revised: 29 April 2025 – Accepted: 22 May 2025 – Published: 13 August 2025

Abstract. A key uncertainty in reanalysis-based snow-on-sea-ice reconstructions is the choice of reanalysis product used for snowfall input. Although reanalysis products have many similarities in their precipitation output over the Arctic Ocean, they nevertheless have relative biases that impact derived snow-on-sea-ice estimates. In this study, snowfall from the European Centre for Medium-range Weather Forecasts (ECMWF) Reanalysis v5 (ERA5), the Japanese Meteorological Agency's Japanese 55-year Reanalysis (JRA-55), and NASA's Modern-Era Retrospective analysis for Research and Applications, Version 2 (MERRA-2), is used as input to the NASA Eulerian Snow On Sea Ice Model (NESOSIM). A Markov chain Monte Carlo (MCMC) approach is used to calibrate the wind packing and blowing snow parameters in NESOSIM run with these different snowfall inputs. JRA-55 shows the largest departure from the previously used values (Bayesian priors) when the MCMC calibration is run and also has the largest posterior uncertainty due to parameter uncertainties. The MCMC calibration reconciles snow depths between NESOSIM run with different reanalysis snowfall inputs but produces larger discrepancies in snow densities due to the sensitivity of snow density in NESOSIM to parameter values and the weak observational constraints on density. Regional climatologies and trends in the calibrated products are examined and compared to another reanalysis-based snow-on-sea-ice reconstruction, SnowModel-LG. NESOSIM and SnowModel-LG show close agreement in snow depth climatologies in the central Arctic Ocean region but differ more in peripheral seas. The models perform comparably when evaluated against IceBird airborne snow depth observations and in situ depth and density observations from the Multidisciplinary

plinary Drifting Observatory for the Study of Arctic Climate (MOSAIC). Trends are found to be region-dependent, and central Arctic Ocean snow depth trends are more sensitive to the choice of reanalysis input than to the choice of model.

1 Introduction

Snow on Arctic sea ice plays a key role in controlling Arctic climate and ecosystem function and is a crucial input to altimetry-derived sea ice thickness retrieval, but it is challenging to characterize consistently across the Arctic Ocean at basin scales (Webster et al., 2018). Satellite remote sensing data using, for example, depth retrievals from passive microwave data (Brucker and Markus, 2013; Rostosky et al., 2018) and altimetry-based snow depth retrievals (Lawrence et al., 2018; Kwok et al., 2020) provide basin-wide estimates of snow depth on Arctic sea ice but are subject to significant retrieval limitations and uncertainties. Airborne (MacGregor et al., 2021; Jutila et al., 2022) and in situ (Wagner et al., 2022; Radionov et al., 1997) observation campaigns and automated snow buoys (Perovich et al., 2019; Nicolaus et al., 2017) provide generally more accurate but also more localized observations. A complementary approach to estimate snow on Arctic sea ice on basin scales is through reanalysis-based snow reconstructions, in which reanalysis snowfall forces a model that simulates snow processes while accounting for sea ice concentration and drift. These reconstructions include SnowModel-LG (Liston et al., 2020), the University of Washington snow-on-sea-ice reconstruction (Blanchard-Wrigglesworth et al., 2018), and the NASA Eulerian Snow

on Sea Ice Model (NESOSIM; Petty et al., 2018), which is the focus of our study. NESOSIM has been previously used with altimetry measurements from the NASA Ice, Cloud, and land Elevation Satellite 2 (ICESat-2) to produce estimates of Arctic sea ice thickness over the winter season (Petty et al., 2020, 2023b).

Not surprisingly, reanalysis snow-on-sea-ice reconstructions are strongly sensitive to snowfall input, which depends on several factors such as atmospheric process representation in reanalysis products (e.g. microphysical processes and partitioning between solid-, liquid-, and mixed-phase precipitation), data assimilation inconsistencies, and product resolution. Reanalysis precipitation assessment for the Arctic (Behrangi et al., 2016; Boisvert et al., 2018; Barrett et al., 2020; Edel et al., 2020; Cabaj et al., 2020) is challenged by uncertainty in polar precipitation observations, especially over the Arctic Ocean. Reanalysis precipitation intercomparison work by Barrett et al. (2020) recommends that the European Centre for Medium-range Weather Forecasts (ECMWF) Reanalysis v5 (ERA5) be used to provide precipitation for sea ice thickness estimates over other contemporary reanalysis products but acknowledges that other reanalysis products investigated in that study are of similar value for that application, given the difficulty of observational validation and bias adjustment. Biases between reanalysis products can be reduced through calibration using satellite snowfall observations, but differences between products nevertheless persist, and satellite snowfall measurements themselves may be biased (Cabaj et al., 2020; Edel et al., 2020). This motivates the need for further calibration of snow-on-sea-ice reconstructions.

The purpose of this study is to improve consistency and characterize uncertainty amongst several reanalysis snowfall inputs for NESOSIM's snow-on-sea-ice reconstruction, using bias-adjusted snowfall input and automated calibration of NESOSIM's snow model parameters. We also seek to assess the variability and trends in basin-wide and regional snow on Arctic sea ice produced by NESOSIM using these newly recalibrated snow depth estimates. Our starting point is the latest version of NESOSIM, version 1.1 (v1.1; Petty et al., 2023b). In Cabaj et al. (2023), NESOSIM v1.1 free parameters for the wind packing (densification) and blowing snow (loss) processes were calibrated to snow-on-sea-ice depth and density observations using a Markov chain Monte Carlo (MCMC) approach, and uncertainty estimates for these free parameters were obtained. Several considerations motivated the development of an automated parameter calibration approach for NESOSIM. In recent upgrades to NESOSIM that included the switch to the latest ERA5 snowfall input, the free parameters were manually tuned to increase agreement with snow depths obtained from Operation IceBridge (Petty et al., 2023b). This process alluded to the potential for parameter tuning to account for forcing biases between products, as well as the potential benefits of a more automated system that could bring in different types of observations to

evaluate both the snow depth and density concurrently. Optimizing NESOSIM output is motivated by its continued use as the main snow loading input to satellite-derived sea ice thickness estimates from ICESat-2 (Petty et al., 2023a, b) and the need to better characterize the snow loading uncertainty contribution to the overall thickness uncertainty.

To better reconcile differences between NESOSIM run with different snowfall inputs and to incorporate estimates of uncertainties due to the choice of model snowfall input, in this current study, we run the MCMC optimization for NESOSIM with additional reanalysis snowfall inputs, introducing NASA's Modern-Era Retrospective analysis for Research and Applications, Version 2 (MERRA-2), and the Japanese Meteorological Agency's Japanese 55-year Reanalysis (JRA-55) to this study in addition to ERA5. This also necessitates a revisiting of the CloudSat calibration for reanalysis snowfall first performed in Cabaj et al. (2020) since a longer time record and an additional reanalysis product are used in this study. We estimate resulting snow depth and density uncertainties due to parameter uncertainty, which helps account for uncertainties due to snow input, and examine the impact of this parameter optimization on the agreement between the snow depth and density derived using these products. Then we construct consensus snow depth and density estimates that account for variability in reanalysis snowfall from the average of calibrated NESOSIM output for different reanalysis snow inputs, motivated by work combining land snow products (Mudryk et al., 2015), which demonstrates how multi-dataset approaches can help to reveal biases between datasets and facilitate the characterization of dataset uncertainties. We evaluate the consistency of the outputs across different snowfall forcing inputs, examining the climatologies, the interannual variability, and trends over the 1980–2021 time period. We compare the NESOSIM output to SnowModel-LG (Liston et al., 2020), another reanalysis-based snow-on-sea-ice model. SnowModel-LG likewise includes observation-based calibration, namely an assimilation-based bias correction to precipitation to bring modelled snow depth into agreement with ground-based and remote sensing observations, including Operation IceBridge measurements (Liston et al., 2020; Stroeve et al., 2020). We further compare the NESOSIM and SnowModel-LG outputs to independent observational datasets, namely in situ snow depth and density observations from the 2019–2020 Multi-disciplinary Drifting Observatory for the Study of Arctic Climate (MOSAIC) observational campaign (Itkin et al., 2023; Macfarlane et al., 2023) and airborne snow depth observations from April 2017 and 2019 from the Alfred Wegener Institute (AWI) IceBird campaign (Juttila et al., 2022).

2 Data products and models

2.1 Reanalysis products

Snowfall rates from the ERA5, MERRA-2, and JRA-55 reanalysis products are used as input to NESOSIM in this study. ERA-Interim is examined for reference but is not used as input to NESOSIM, as it has been superseded by ERA5. A summary of the reanalysis products used in this study is shown in Table 1, and each product is discussed in more detail in the subsections below. To format the reanalysis snowfall for use as input to NESOSIM, the snowfall rate from each reanalysis product is aggregated by day to produce daily snowfall and then regridded to the $100\text{ km} \times 100\text{ km}$ equal-area NESOSIM model grid. NESOSIM also uses 10 m wind input from reanalysis products, but for this study, ERA5 winds were used for all model runs.

2.1.1 ERA-Interim

The European Centre for Medium-range Weather Forecasts (ECMWF) Re-Analysis Project ERA-Interim (Dee et al., 2011) is widely used in studies of Arctic snow and is often used for precipitation input in snow models (Kwok and Cunningham, 2008; Petty et al., 2018; Blanchard-Wrigglesworth et al., 2018). It has been found to have high correlations and low biases with respect to observations of Arctic land precipitation (Lindsay et al., 2014). Sea ice concentration is represented as a fractional quantity for grid cells with concentrations greater than 20 %, while grid cells with less than 20 % concentration are designated as open ocean. ERA-Interim is produced using a 4DVar assimilation scheme, and it features a T255 ($\sim 79\text{ km}$) resolution spectral dynamical core. The ERA-Interim snowfall product is provided on an N128 Gaussian grid, re-gridded to a $0.75^\circ \times 0.75^\circ$ latitude/longitude grid in this study. Production of ERA-Interim has stopped as of August 2019.

2.1.2 ERA5

The ECMWF Reanalysis v5 (Hersbach et al., 2020), the successor to ERA-Interim, features many improvements, such as a finer model resolution, an updated assimilation scheme, and an improved cloud scheme, including improvements to the representation of mixed-phase clouds and ice-phase cloud microphysics (Hersbach et al., 2020). It has been found to produce more snow than ERA-Interim, especially in the Atlantic sector (Wang et al., 2019). Like ERA-Interim, ERA5 uses a 4DVar assimilation scheme. The representation of sea ice concentration is also the same as in ERA-Interim, with fractional concentration above a 20 % open-ocean threshold. In this study, the ERA5 snowfall rate product is interpolated from its native N320 Gaussian grid to a $0.25^\circ \times 0.25^\circ$ grid. Currently, ERA5 is used as the default snowfall and as the 10 m wind input for NESOSIM v1.1 (Petty et al., 2023b; Cabaj et al., 2023).

2.1.3 MERRA-2

NASA's Modern-Era Retrospective analysis for Research and Applications, Version 2 (Gelaro et al., 2017), is produced on a cubed-sphere grid with a finite-element dynamics scheme and is used in this study with its native horizontal resolution of $0.5^\circ \times 0.625^\circ$ ($\sim 55\text{ km}$). Unlike the other reanalysis products investigated in this study, which use a 4DVar assimilation scheme, MERRA-2 uses a 3DVar assimilation scheme with an incremental analysis update procedure, which applies the analysis increment as a constant term over the assimilation window instead of only correcting the initial condition, as is done conventionally for 3DVar (Gelaro et al., 2017). Sea ice is distinguished from open ocean based on a 50 % concentration threshold. MERRA-2 is known to produce more total precipitation over the Arctic compared to other reanalysis products (Barrett et al., 2020; Boisvert et al., 2018).

2.1.4 JRA-55

The Japanese Meteorological Agency's Japanese 55-year Reanalysis (Kobayashi et al., 2015) is another widely used product for Arctic snowfall estimates, and it is interpolated onto a $1.25^\circ \times 1.25^\circ$ grid from its native TL319 ($\sim 55\text{ km}$) spectral resolution. The product uses a 4DVar assimilation scheme. Sea ice is represented in JRA-55 with a binary classification based on a 55 % concentration threshold. JRA-55 has been previously used as a source of snowfall input for snow-on-sea-ice reconstructions and was investigated as an input for NESOSIM version 1.0 (v1.0; Petty et al., 2018). In comparisons of total precipitation over the Arctic Ocean, JRA-55 has been found to produce less precipitation overall than other reanalysis products (Barrett et al., 2020).

2.2 CloudSat

CloudSat was a satellite equipped with a 94 GHz Cloud Profiling Radar (CPR) instrument, which measured vertical profiles of cloud and hydrometeor reflectivity, from which the near-surface snowfall rate was retrieved (Kulie and Benkert, 2009). The satellite had an observational footprint of $1.4 \times 1.7\text{ km}$ (along and across track) and a 16 d repeat cycle. The instrument was operational from 2006 to 2023, with an interruption in 2011 due to a battery malfunction and a change to a lower orbit in 2018. In this study, surface snowfall rates from the 2C-SNOW-PROFILE product, version P1 R05 (Wood et al., 2013, 2014), are used to bias correct snowfall rates from reanalysis products by scaling the reanalysis monthly climatologies to the monthly climatology of regionally aggregated CloudSat snowfall, following the approach in Cabaj et al. (2020). CloudSat measurements from 2006 to 2016 are used in this study. CloudSat's ground track had latitudinal coverage between 82°N and 82°S . Its sampling is also limited following the 2011 battery malfunction to ob-

Table 1. Reanalysis products examined in this study. Spatial resolution refers to the regular lat–long grid used for the products in this study. To provide input to NESOSIM, all reanalysis products are regridded to the equal-area 100 km × 100 km polar grid used by the model.

Reanalysis	Spatial resolution	Time resolution	Assimilation scheme	Reference
ERA-Interim	0.75° × 0.75°	6-hourly	4DVar	Dee et al. (2011)
ERA5	0.25° × 0.25°	Hourly	4DVar	Hersbach et al. (2020)
MERRA-2	0.5° × 0.625°	Hourly	3DVar	Gelaro et al. (2017)
JRA-55	1.25° × 1.25°	3-hourly	4DVar	Kobayashi et al. (2015)

servations taken when the instrument had line-of-sight to the sun, which may introduce low biases during the later observational period (Milani and Wood, 2021). Nevertheless, CloudSat has been used extensively in studies of high-latitude precipitation and snowfall (Behrangi et al., 2016; Edel et al., 2020; Kulie et al., 2016; Milani et al., 2018). To mitigate ground clutter contamination of surface returns in this study, near-surface snowfall rate measurements are retrieved from the third vertical bin above ocean surfaces (approximately 720 m above the surface) or the fifth vertical bin above sea ice (as determined by a climatological sea ice mask at approximately 1200 m above the surface) (Wood and L’Ecuyer, 2018). Data quality flags are applied to exclude potentially contaminated observations, as described in Cabaj et al. (2020).

2.3 MOSAiC

The Multidisciplinary Drifting Observatory for the Study of Arctic Climate (MOSAiC) expedition took place in September 2019–October 2020, providing high-quality in situ observations of snow and sea ice for the duration of an entire sea ice season at a central Arctic location (Nicolaus et al., 2022). In this study, we use snow depth measurements recorded using automated snow depth probes (MagnaProbes) (Itkin et al., 2021) and bulk snow densities calculated from snow density cutters (Macfarlane et al., 2021, 2022) as independent observational datasets for comparison with model output (Macfarlane et al., 2023). Snow depth measurements were collected over regions representative of first-year and second-year ice over level ice and ridges. Snow density cutter measurements likewise were taken under a variety of snow conditions. To partly compensate for the highly localized and heterogeneous nature of the observations, we collocate the observed values with the nearest respective model grid cell and average the values by day within each grid cell. Then we calculate monthly means of these daily aggregated values for comparison with modelled values. Prior to aggregation, bulk snow densities are first calculated by calculating the height-weighted average of snow density samples measured in each snow density profile.

2.4 IceBird

The Alfred Wegener Institute (AWI) IceBird observational campaign is an airborne observational campaign for collecting measurements of sea ice thickness, and sea ice and snow properties. We make use of snow depth observations collected by airborne snow radar from April 2017 and April 2019 (Jutila et al., 2022, 2024a, b) as independent observational datasets for comparison with model output. Survey tracks cover regions over the western Arctic Ocean, around the Canadian Arctic Archipelago and Beaufort Sea regions, and they encompass both first-year and multi-year ice. For comparison with model output, we calculate the average of observed values from the transects within each coincident model grid cell. Each transect spans multiple model grid cells, although some measurements coincide with grid cells considered to be land by the models due to model resolution limitations.

2.5 NESOSIM and MCMC calibration

The NASA Eulerian Snow on Sea Ice Model (NESOSIM) produces estimates of snow depth and bulk snow density over winter Arctic (September to April) sea ice on a 100 × 100 km polar grid (Petty et al., 2018). The model is a two-layer Eulerian snow-on-sea-ice model and includes parameterized representations of snow accumulation, densification through wind packing, loss to the atmosphere and open ocean from blowing snow, and redistribution of snow due to sea ice motion. NESOSIM was initially developed to provide estimates of snow depth to enable the rapid production of Arctic sea ice thickness estimates from ICESat-2 (Petty et al., 2020, 2023b).

Several observational and reanalysis inputs are used in NESOSIM. Snowfall input for NESOSIM is provided from reanalysis products, with ERA5 being used as the default product as of v1.1 and ERA-Interim used previously as the default in v1.0. In this study, multiple reanalysis products are investigated as a source of snowfall input. Reanalysis products are also used for wind input to NESOSIM. This study uses ERA5 10 m wind as input to the model, motivated by findings that the ERA5 wind product performs relatively well with respect to observations compared to other reanalysis products in Arctic regions (Graham et al., 2019a, b). Sea ice concentration is provided by the National Oceanic and Atmo-

spheric Administration and National Snow and Ice Data Center (NOAA/NSIDC) Climate Data Record (CDR) product (Peng et al., 2013). Sea ice drift for the MCMC calibration is obtained from the low-resolution sea ice drift product of the European Organisation for the Exploitation of Meteorological Satellites (EUMETSAT) Ocean and Sea Ice Satellite Application Facility (OSI SAF; Lavergne et al., 2010). Since the OSI SAF drift product is not available for years prior to 2009, sea ice drift from the NSIDCv4 Polar Pathfinder product (Tschudi et al., 2019) was used to generate the full 1980–2021 datasets. Aside from reanalysis products, these inputs are the same as those used in previous work using NESOSIM v1.1 (Petty et al., 2020, 2023b; Cabaj et al., 2023).

Representations of snow processes in NESOSIM are highly simplified. Since NESOSIM is a two-layer model, bulk snow density in the model is represented as a weighted sum of the prescribed densities for old snow (350 kg m^{-3}) and new snow (200 kg m^{-3}). The old-snow density represents both wind slab and depth hoar (Petty et al., 2018). These prescribed values impose maximum and minimum values on the bulk density represented by the model. Snow may be redistributed between model grid cells through the action of sea ice drift, although the representation of drift represents a large-scale average, given the model resolution. Melt processes are currently not included in NESOSIM, so the model is run from September to April in each season and re-initialized on 1 September based on climatological mean snow depths scaled by the number of melt days in the previous season (Petty et al., 2018).

The wind packing and blowing snow parameters in NESOSIM are free parameters, and previous work introduced an automated calibration of these parameters using an MCMC process (Cabaj et al., 2023). At each model step and grid point, if the input wind speed exceeds a prescribed threshold speed of 5 m s^{-1} , the wind packing and blowing snow processes act on the snow in NESOSIM. Wind packing controls the amount of snow transferred between layers, decreasing the snow depth and increasing the bulk snow density as snow is transferred from the upper (less dense) layer to the lower (denser) layer. The blowing snow process acts only on the upper snow layer and decreases the snow depth in the upper layer linearly with wind speed. The blowing snow term includes an atmosphere loss term and an open-water loss term, which are prescribed separately in NESOSIM v1.1 (Petty et al., 2023b). The open-water loss term accounts for sea ice concentration, with regions of lower sea ice concentration experiencing more open-water loss. For the purpose of this study, the blowing snow term parameters are treated as a single term, as was done in previous work (Cabaj et al., 2023), with the atmospheric loss factor being 0.15 times the blowing snow parameter. The blowing snow term is exclusively a loss term and does not include redistribution. When snow is lost from a grid cell via this process, it is removed, not redistributed to another grid cell. This current study will extend previous parameter calibration work by investigating the im-

pact of using different reanalysis snowfall input products in NESOSIM.

Previous work (Cabaj et al., 2023) demonstrated successful calibration of NESOSIM's wind packing and blowing snow parameters using an MCMC process when NESOSIM was run with ERA5 snowfall. MCMC is an algorithm applied to Bayesian problems where, given prior information about the parameters and observational constraints on the parameters, posterior parameters may be obtained, which produce model output that is more closely aligned to observations, as determined by a cost function; in this case, a log likelihood function of the difference between model output and selected aggregated observations used for the calibration, weighted by the uncertainty in the observations. Using an MCMC approach for calibrating NESOSIM allows for the automated estimation of free parameters, which were previously manually estimated via comparison to observations (Petty et al., 2018). An added benefit of this approach is that it yields posterior distributions of the parameters, which provide an estimate of parameter uncertainty subject to observational constraints. The MCMC process is iterative and is conducted for NESOSIM following the approach in Cabaj et al. (2023). Further description of the NESOSIM MCMC calibration is provided in Appendix B.

2.6 SnowModel-LG

SnowModel-LG (Liston et al., 2020; Stroeve et al., 2020) is a Lagrangian snow-on-sea-ice model. Like NESOSIM, it takes snowfall input from reanalysis products. However, the representation of snow processes in SnowModel-LG is considerably more complex than in NESOSIM, with the notable inclusion of snowmelt, snowpack metamorphosis processes, and multiple snow layers (a maximum of 25 layers for the product used in this study). Output is provided with a daily temporal resolution and a spatial resolution of $25 \times 25 \text{ km}$. SnowModel-LG output has been found to compare favourably with data from several observational campaigns (Stroeve et al., 2020), although agreement depends on the region and time period of comparison.

The ERA5 and MERRA-2 reanalysis products are used to provide snowfall input to SnowModel-LG. SnowModel-LG also includes an observation-based calibration, with scaling factors applied to the reanalysis snowfall based on a correction empirically derived from Operation IceBridge snow depth measurements (Liston et al., 2020; Stroeve et al., 2020). The assimilation approach used in SnowModel-LG is consistent with optimal interpolation approaches (Liston and Hiemstra, 2008).

In this study, output from SnowModel-LG run with ERA5 and MERRA-2 input (retrieved from the National Snow and Ice Data Center; Liston et al., 2021) is used for comparison with NESOSIM. SnowModel-LG does not include the Canadian Arctic Archipelago region, so this region is not considered for the comparisons between SnowModel-LG and NE-

SOSIM in this study. Furthermore, whereas NESOSIM is initialized in September, SnowModel-LG is initialized in August and run through the melt season. For consistency, only months during which NESOSIM and SnowModel-LG data are both available will be considered in this study.

3 Investigating different reanalysis snowfall products

Here, we present a comparison of the reanalysis snowfall products used in this study as input to NESOSIM. Reanalysis snowfall products are calibrated to CloudSat following the approach from Cabaj et al. (2020), but in this study, additional products are used and a longer time series is examined, as discussed below.

3.1 Reanalysis snowfall calibration using CloudSat

Figure 1 shows regionally aggregated monthly mean snowfall rates over the ocean region in the 60–82° N latitude band from reanalysis products and CloudSat from 1980 to 2016 without and with scaling to the CloudSat monthly climatology (Cabaj et al., 2020). The scaling entails taking the monthly reanalysis snowfall rate for each month and multiplying it by a scaling factor, which consists of the CloudSat climatological monthly mean snowfall rate divided by the reanalysis climatological monthly mean snowfall rate for each respective month. The climatological means for this scaling are taken from 2006 to 2016, excluding months in 2011 when CloudSat observations are absent due to instrument malfunctions. Further details of this scaling are provided in Cabaj et al. (2020). Before the scaling is applied in Fig. 1, there is some variation between the reanalysis products, although they have similar seasonal cycles and generally coincident seasonal maxima and minima. ERA5 and MERRA-2 have relatively high snowfall compared to ERA-Interim and JRA-55. Snowfall rates from CloudSat, which are available from 2006 to 2016 with a gap in 2011, are comparable to the snowfall rates of the other products. MERRA-2 is known to be wetter compared to other reanalysis products over the Arctic when total precipitation is considered (Barrett et al., 2020; Boisvert et al., 2018). This is particularly reflected in the summer months, when MERRA-2 snowfall rates are the largest relative to the other products.

As in Cabaj et al. (2020), we bias adjust reanalysis snowfall input to climatological CloudSat snowfall for 2006–2016. CloudSat scaling improves agreement amongst the reanalyses both within and outside the 2006–2016 calibration period (Fig. 1b). Although MERRA-2's seasonal snowfall cycle is consistently of greater amplitude than those of the other products prior to 2006, the agreement of MERRA-2 with the other products is considerably improved with the scaling. JRA-55, which was not previously investigated in this context, is also brought into closer agreement with the other products using this approach. This highlights the con-

tinued benefits of this bias-adjustment approach for reconciling reanalysis snowfall products.

3.2 Snowfall comparison over ocean and sea ice for the NESOSIM domain

To apply CloudSat scaling over the NESOSIM model domain, a more regionally refined scaling approach is used. Reanalysis snowfall rates are scaled to CloudSat measurements from 60 to 82° N over four quadrants, as described in Cabaj et al. (2020). The CloudSat scaling was previously found to improve agreement in basin-averaged and regionally averaged snow depths in NESOSIM v1.0 (Cabaj et al., 2020). Some adjustments were made to the scaling for NESOSIM v1.1, which has a larger model domain (Petty et al., 2023b) extending down to 50° N compared to 60° N for NESOSIM v1.0 (Petty et al., 2018). For the current version of the model, the scaling is performed as follows.

As in previous work, the NESOSIM v1.1 model domain is divided into quadrants with longitudinal boundaries at longitudes 135, 45, 45, and 135° E, as illustrated in Fig. A1. For each quadrant within a region between latitudes of 60 and 82° N (indicated by solid lines in Fig. A1), CloudSat scaling factors are calculated by dividing the CloudSat monthly snowfall climatology by the reanalysis monthly snowfall climatology averaged over the region. The same CloudSat scaling factors previously calculated for Cabaj et al. (2020) are used, with the addition of scaling factors for JRA-55, which were not previously calculated. The scaling factors are linearly interpolated across the pole from corners at latitudes of 45° N (longitudes of 90° W, 0° E, 90° E, and 180° E, indicated by A–D in Fig. A1), consistent with the interpolation performed for version 1.0 of the model. To account for the extended model domain in NESOSIM version 1.1, the scaling factors are extrapolated southwards from the same corners (A–D) as constant values. This process creates a set of monthly gridded scaling factors that are then multiplied by the daily reanalysis snowfall rates used as input to NESOSIM v1.1, with a different set of scaling factors used for each month.

Figure 2 shows the impact of CloudSat scaling applied to NESOSIM model input for monthly climatologies of reanalysis snowfall rates over ocean (which includes both ice and open ocean, with land masked out) and over sea ice only, regionally averaged over a representative subset of the different Arctic regions shown in Fig. A1.

CloudSat scaling effectively reconciles differences between reanalysis products for the pan-Arctic ocean region in Fig. 1 during the satellite era but shows less consistency for individual regions and when ice-covered scenes are broken out. Over the ice-plus-ocean region for which the CloudSat scaling was originally developed, the CloudSat scaling reconciles differences between the products for most months in most regions. A notable exception is in the central Arctic region, where the September snowfall values are excessively




Figure 1. Monthly mean snowfall rates from reanalysis products and CloudSat, regionally averaged over the ocean region in the 60–82° N latitude band (i.e. excluding land) **(a)** without scaling to CloudSat and **(b)** scaled to the CloudSat monthly snowfall climatology. Panel **(c)** shows the difference between the reanalysis products and CloudSat for the no-scaling (dashed) and with-scaling (solid) cases from 2006 to 2016.

large for JRA-55 following the application of the CloudSat scaling. This may be because JRA-55 is biased relatively low compared to CloudSat and the other products, so the CloudSat scaling, determined using ice-plus-ocean scaling factors, greatly increases the snowfall rates, especially in the early part of the sea ice season. Furthermore, since CloudSat observations are limited to latitudes south of 82° N, the scaling factors may be less reliable over more poleward regions. Over the ice-covered region alone, the CloudSat scaling reduces inter-product consistency in some regions and months. Over sea ice, the overall inter-product spread increases in September and October in the central Arctic, in October in the Beaufort Sea, in October–November in the Chukchi Sea, in all months except January and September for the Kara Sea, in all months except April and September in the Barents Sea, and in all months except April in the East Greenland Sea. In the Kara, Barents, and East Greenland Sea regions in par-

ticular, JRA-55 and MERRA-2 are closely reconciled by the CloudSat scaling, but ERA5 is less changed by the scaling, which results in it being biased relatively low with respect to the other products.

Seasonal cycles of snowfall over sea ice may be similar to snowfall over the ice-plus-ocean region in some regions, but other regions show stark differences. Of the regions shown, the seasonal cycles and magnitudes differ considerably between the two cases in the Chukchi Sea, Barents Sea, and East Greenland Sea regions. By comparison, the differences are notably less stark in the central Arctic region, which has considerable ice cover even in the early season. In the Kara and Beaufort seas, the seasonality is similar between the two cases, although the magnitudes differ. Many of the regions show relatively low snowfall over the ocean-plus-ice region in September, but those same snowfall minima are not as prominent in the plots for the ice-covered regions. This sug-

Regional snowfall climatologies from reanalysis products before and after CloudSat scaling




Figure 2. Monthly climatologies of regionally averaged snowfall rates from reanalysis products from 1980 to 2021: over the ocean region (i.e. including both ice-covered and ice-free ocean) and over sea ice only without and with CloudSat scaling (CS) applied. Bars represent the interannual standard deviations of the climatologies. Note the differing scales on the vertical axes between regions.

gests that much of the snow that is present during the early part of the season is coincident with the presence of sea ice. This may be due to ice-covered regions having lower temperatures, which permits the presence of early-season snowfall where other regions may have rainfall. Despite these regional inconsistencies due to the limited overlap between CloudSat orbits and ice-covered regions, which would likewise adversely impact CloudSat scaling if it were restricted to ice-covered regions, we proceed with the established CloudSat scaling factors. We will return to the discussion of issues related to CloudSat scaling in Sect. 7.

Discrepancies in reanalysis input yield discrepancies in the output from NESOSIM driven by different reanalysis snowfall products. This motivates the observation-constrained calibration of NESOSIM run with different reanalysis snowfall inputs using an MCMC method, as was previously done in Cabaj et al. (2023). The following section discusses an updated calibration of NESOSIM and the impact on model-derived snow depth and density.

4 Impact of MCMC calibration on NESOSIM output

4.1 Posterior model parameters

In this study, the same MCMC approach as in Cabaj et al. (2023) is repeated but with the addition of other reanalysis snowfall inputs; MERRA-2 and JRA-55 snowfall are used in addition to that of ERA5. The MCMC process was run for 10 000 iterations for each snowfall input product. The first 5000 iterations are discarded to exclude “burn-in” values, as was done in Cabaj et al. (2023). Nevertheless, in each case, the optimal posterior parameter values did not differ significantly between the first (burn-in) and subsequent (after burn-in) sets of iterations, demonstrating robust convergence of the MCMC process.

Figure 3 shows the posterior distributions obtained from the MCMC calibrations of NESOSIM run with snowfall input from ERA5, JRA-55, and MERRA-2. The posterior distributions are aggregated from parameter values that are ac-

cepted during the MCMC process and provide both the optimal (maximum-likelihood) parameter values and estimates of the parameter uncertainties (Gelman et al., 2013). Numerical values for the optimal parameters and associated uncertainties are shown in Table 2, along with the coefficients of variation and the acceptance rates. The acceptance rate, calculated from the ratio of accepted parameters to the total number of iterations, indicates the efficiency of the MCMC process, with an optimal efficiency for a two-parameter MCMC process being approximately 23 % (Gelman et al., 2013). Coefficients of variation are calculated from the standard deviation of the posterior distribution divided by the posterior parameter value and quantify the relative spread of the posterior distribution. This provides a quantitative indication of how well-constrained the parameters are by the MCMC calibration. The posterior distribution of ERA5 is considerably narrower than the distributions of MERRA-2 and JRA-55, with the latter being noticeably broad relative to the posterior distributions of the other two products. The coefficients of variation for the wind packing parameters, as indicated in Table 2, are 15 % for ERA5, 42 % for JRA-55, and 21 % for MERRA-2. The coefficients of variation for the blowing snow parameters are 13 % for ERA5, 38 % for JRA-55, and 19 % for MERRA-2. The JRA-55 parameter distribution has a slight bimodality in both wind packing and blowing snow, although the maximum-likelihood parameter corresponds to the larger mode. The spreads of the parameters for ERA5 and MERRA-2 are more comparable, with the MERRA-2 posterior distributions being somewhat wider than those of ERA5. In terms of departure from the prior values, ERA5 has the closest blowing snow value to the prior, and MERRA-2 has the closest wind packing to the prior. JRA-55 demonstrates the largest departure from the prior parameter values overall. The acceptance rates are included primarily as an indicator of the efficiency of the MCMC process; ERA5 and MERRA-2 are relatively close to the optimal (23 %) efficiency for a two-parameter MCMC optimization (Gelman et al., 2013). The comparatively large acceptance rate for JRA-55 suggests that a larger step size could be used for the MCMC optimization for this product, but given that the NESOSIM–MCMC optimization is not excessively computationally expensive, the existing configuration is sufficient for the scope of this study. The correlation between the wind packing and blowing snow parameters may be a consequence of the processes compensating for each other, as described in Cabaj et al. (2023). The wind packing process transfers snow to the lower layer, where the blowing snow process cannot remove snow, so if wind packing is strengthened, the blowing snow process may also be strengthened to compensate and to enable additional snow depth reduction.

These results highlight the fact that model parameter tuning is heavily dependent on the forcing dataset. Care must be taken when using reanalysis-based snow-on-sea-ice reconstructions such as NESOSIM with different snow input




Figure 3. Posterior wind packing and blowing snow parameter distributions from MCMC calibration using snow input from ERA5, JRA-55, and MERRA-2. Note that the distributions have some overlap. The dark red dot indicates the prior parameter values, and the other coloured dots indicate the optimal parameter values for the three respective products. The side panels show the marginal distributions, highlighting the overlap.

datasets since model processes may be less physically representative when different inputs are used. When developing parameterizations for such model processes, it is important to consider that biases in model inputs will likewise be reflected in model parameterizations. Biases in the observations used for calibration will also impact the model output, but may be inevitable given the relative scarcity of in situ observations of snow on sea ice. Overall, MCMC can be used to objectively determine model parameters and their uncertainty given the uncertainty in the inputs.

4.2 Snow depth and density uncertainty estimates

Given the posterior parameter uncertainties provided from the MCMC calibration, uncertainty in the NESOSIM output can be calculated. Figure 4 shows the depth and density for a single representative late-decade year, with shading representing the associated MCMC-estimated uncertainty for each respective product. The uncertainty is calculated from a 100-parameter ensemble run, with the wind packing and blowing snow parameters sampled from the posterior distribution for each respective MCMC calibration, as in Cabaj et al. (2023). This represents uncertainty due to the model parameter uncertainty and therefore does not characterize all the uncertainties in the model. The day-to-day variability is quite similar between the time series, although some differences re-

Table 2. Optimal parameters from MCMC optimization for different reanalysis inputs, with the default (prior) configuration and the prescribed prior standard deviations included in the first row for reference. WP refers to wind packing and BS refers to blowing snow. MCMC-derived standard deviations are denoted by σ . CV refers to the respective coefficients of variation for each parameter. AR refers to the acceptance rate of the MCMC optimization, i.e. the percentage of iterations whose test parameter values are accepted.

Configuration	WP (s ⁻¹)	σ_{WP} (s ⁻¹)	CV _{WP}	BS (m ⁻¹)	σ_{BS} (m ⁻¹)	CV _{BS}	AR
Model default (prior)	5.8×10^{-7}	1×10^{-8}	–	2.9×10^{-7}	1×10^{-8}	–	–
MCMC–ERA5	2.05×10^{-6}	3.11×10^{-7}	15 %	4.01×10^{-7}	5.30×10^{-8}	13 %	29 %
MCMC–JRA-55	2.35×10^{-6}	9.82×10^{-7}	42 %	1.05×10^{-6}	4.01×10^{-7}	38 %	46 %
MCMC–MERRA-2	1.59×10^{-6}	3.40×10^{-7}	21 %	6.03×10^{-7}	1.12×10^{-7}	19 %	33 %

main between the products. NESOSIM with JRA-55 input shows a sharper early-season increase in snow depth compared to the other products, although the late-season snow depth is comparable to that of the other products. Snow density values for the NESOSIM–JRA-55 output are also the largest overall, reflecting its stronger wind packing. The agreement in day-to-day density variations is likely a consequence of the identical ERA5 wind inputs to each NESOSIM run since wind packing is dependent on the wind input to NESOSIM. NESOSIM–JRA-55 has the largest uncertainty in snow depth and snow density, which is consistent with the spread of the posterior parameter distributions.

The multi-product average was calculated as the average of the MCMC-calibrated output from NESOSIM for the three different reanalysis inputs. The uncertainty in the multi-product average was calculated using the standard deviation of the three 100-parameter model-run ensembles for the three reanalysis products. It thus quantifies both the uncertainty due to model parameter uncertainty and the spread from the use of different model snowfall inputs. Initially, a bootstrap sampling approach was attempted to produce potentially more robust estimates, but it was found that the bootstrap-estimated standard deviations differed from the directly calculated standard deviations by at most only 0.05 %. Hence, the standard deviation of the combined parameter ensemble was used to calculate the multi-product average uncertainty for the depth and density.

To enable more direct uncertainty comparisons, uncertainties and percent uncertainties for depth and density due to model parameter uncertainty are shown in Fig. 5. A plot of monthly interannual-averaged uncertainties from 1980 to 2021 is also available in Fig. A2. As in Fig. 4, the percent uncertainty magnitudes reflect the shape of the posterior distributions. The relative insensitivity of NESOSIM snow output to model parameter values, as was observed in Cabaj et al. (2023), persists here; the percent uncertainties are considerably smaller than the NESOSIM parameter uncertainty, represented by coefficients of variation, as discussed in Sect. 4.1. The NESOSIM–ERA5 uncertainties are relatively small compared to the other products. The percent uncertainties for all the products attain their initial maxima within approximately 15 d of when the model is initialized

despite the differing snow inputs. This further justifies the choice of 15 d as a “ramp-up” period for uncertainty in Cabaj et al. (2023).

The multi-product-average percent uncertainty is larger than the ERA5 percent uncertainty because it accounts for the inter-product differences across snowfall input products, reaching a range of 8 %–18 % snow depth uncertainty and a smaller range of 2 %–5 % uncertainty for snow density. The relatively low percent uncertainty for density may be because the density values are constrained to a relatively narrow range, with a maximum prescribed by the model. The multi-product density percent uncertainty is also notably lower than the JRA-55 density percent uncertainty, which suggests that the JRA-55 density data alone has more relative spread compared to all the ensemble data aggregated together. In particular, as seen in Fig. 4, the JRA-55 uncertainty overlaps considerably with that of ERA5 and to some extent with that of MERRA-2. Hence, there is some reduction in the standard deviation when the parameter ensembles are consolidated to construct a single inter-product standard deviation.

4.3 Impact of MCMC calibration on snow depth

Figure 6 shows basin-average monthly snow depth climatologies from NESOSIM, illustrating how re-calibrating NESOSIM parameters for each individual reanalysis forcing brings the output snow depths from NESOSIM into better consistency across the datasets. The average snow depth is lowered somewhat overall, with the multi-product average in April (27.4 cm) now very close in value to the ERA5 end-of-season value (27.3 cm). Given that in both Fig. 6a and b, ERA5 is plotted with its posterior parameters, it follows that the other products have values that more closely match the ERA5 output in Fig. 6b after the remaining two products have also been MCMC calibrated to the same target observations to which ERA5 was calibrated previously. Some of the relative biases between the products persist; JRA-55 continues to have a relatively large early-season snow depth, which is not seen in the other products, consistent with its early-season snowfall bias. Conversely, at the end of the season, JRA-55 and MERRA-2 previously both exceeded ERA5 at the end of the season, consistent with snowfall biases over




Figure 4. Single-year daily snow depth and density time series for each MCMC-calibrated NESOSIM configuration (with snowfall input from the ERA5, JRA-55, and MERRA-2 reanalysis products) and for the multi-product average. Shading denotes the uncertainty estimated based on the MCMC parameter uncertainty.

sea ice in most regions, particularly over the central Arctic. Following the MCMC calibration, JRA-55 and MERRA-2 bracket ERA5 snow depth on either side, with the multi-product average closely matching the ERA5 values.

The bars in Fig. 6 show interannual variability in the ERA5-calibrated and individually calibrated model runs, which is calculated as the standard deviation of the climatology (these quantities are also plotted separately in Fig. A3). The interannual variability reaches its seasonal peak at the beginning of the season for JRA-55 and ERA5, and at the end of the season for MERRA-2, although the seasonal cycle attains its minimum for all products in October. JRA-55 has the largest interannual variability in September, and in October onward, MERRA-2 has the largest interannual variability of all the products. MCMC calibration reconciles some of the overall spread in interannual variability between the snow depth outputs, although there is less agreement in interannual variability between JRA-55 and MERRA-2 in October following the calibration. Both JRA-55's high early-season variability and MERRA-2's high late-season variability de-

crease somewhat following the MCMC calibration, bringing them into closer agreement with the rest of the products.

4.4 Impact of MCMC calibration on snow density

Although the MCMC calibration reconciles snow depths for NESOSIM run with different snowfall inputs, the opposite is seen for bulk snow density. Figure 7 shows plots of the basin-average monthly bulk snow density climatologies before and after the calibration. The climatologies show very close agreement when the same (MCMC-ERA5) parameters are used for each NESOSIM run but differ considerably when the individually calibrated MCMC parameters are used. This is likely a consequence of how snow density is represented in the model. Since snow density in each layer of NESOSIM is fixed, bulk density is a function of the ratio of snow depths in the two layers. Hence, the bulk density in NESOSIM is strongly sensitive to the strength of the wind packing process, which transfers snow between the layers. Model runs with different snowfall inputs can still produce similar bulk densities, as long as the same wind packing parameter




Figure 5. Daily uncertainty estimates for snow depth (a, b) and density (c, d) for a single season (2018–2019) of the NESOSIM run with all the products separately (colours) and the multi-product average (black). The absolute uncertainties are shown in panels (a) and (c), and percent uncertainties are shown in panels (b) and (d). Grey shading indicates the first 15 d of the season.

and wind input are used. The slight differences between the uncalibrated snow density outputs may depend on the timing of snowfall. For example, a high-snow-accumulation event will reduce the overall bulk density in the short term, but if this accumulation occurs early in the season, more snow may subsequently be transferred to the lower layer, increasing the bulk snow density in the long term if subsequent accumulation is lower. NESOSIM driven by JRA-55 shows deep snow in the early season relative to other products, which may contribute to its high later-season snow density bias, as seen in Fig. 7. Conversely, if the wind packing parameter changes, the modelled density will shift accordingly. The inter-product density differences following MCMC calibration are consistent with the posterior parameter values shown in Fig. 3: the wind packing parameter is largest in JRA-55, which reports the highest bulk snow density value, whereas the smallest wind packing parameter is obtained in the MERRA-2 calibration, which reports the lowest density value.

The widened spread between products following the calibration also reflects the fact that the density values are relatively under-constrained by the MCMC calibration approach

due to the small number of density measurements used. Monthly climatologies of basin-averaged historical density measurements are used as observational constraints for the calibration due to a relative lack of widespread contemporary density measurements. These density observations are vastly outnumbered by the Operation IceBridge (OIB) depth observations used in the optimization, which puts more weight on the OIB measurements in the likelihood function. Hence, because the snow depth constraint is stronger, the MCMC calibration will tend to reconcile differences in snow depth while potentially introducing discrepancies in density. Some of this spread may also be related to how the wind packing and blowing snow parameters vary in tandem during the calibration, which may also be a consequence of the relatively few density measurements provided. Given that previous work (Cabaj et al., 2023) found that sea ice thickness estimates produced using NESOSIM snow input are more sensitive to snow depth than differences in snow density, we proceed with using the individually calibrated density values to produce the NESOSIM multi-product-average density despite their wider spread.




Figure 6. Basin-average monthly snow depth climatologies from NESOSIM for 1980–2021, (a) with the model run using the MCMC–ERA5 configuration for all products (i.e. using the same wind packing and blowing snow parameters) and (b) with the parameter values re-tuned to each respective reanalysis input. Bars indicate the interannual variability (standard deviation of the climatology), which is also shown separately in Fig. A3.




Figure 7. Basin-average bulk snow density monthly climatologies for 1980–2021 (a) for the MCMC–ERA5 configuration and (b) with each product calibrated separately. Bars represent the standard deviations of the climatologies, indicating interannual variability.

4.5 Regional snow-on-sea-ice climatologies

Figure 8 shows regionally averaged snow depth and density climatologies by region (with regions as defined in Meier and Stewart, 2023) from NESOSIM–MCMC output and from SnowModel-LG. Sea ice area calculated from the NOAA/NSIDC Climate Data Record (CDR) product (Peng et al., 2013) is also shown; this product is used in NESOSIM. The sea ice product used in SnowModel-LG differs in that it uses the NASA Team algorithm, whereas the CDR product uses the highest value from the NASA Team

and Bootstrap algorithms (Cavalieri et al., 1996; Peng et al., 2013). NESOSIM and SnowModel-LG snow depths agree well in the central Arctic, Beaufort Sea, and Chukchi Sea regions, but NESOSIM depths exceed those of SnowModel-LG in the Kara, Barents, and East Greenland Sea regions. Notably, NESOSIM shows the greatest snow depths in the East Greenland Sea region, as expected from the presence of the North Atlantic storm track (Webster et al., 2019), but SnowModel-LG records this as a region with much shallower snow (~ 27 cm versus ~ 72 cm in the late season). Knowing that NESOSIM's simplicity might challenge its realism

in high-latitude North Atlantic regions like the East Greenland Sea, improved observations of snow on sea ice are critical in such regions. In several peripheral seas of the Arctic Ocean, SnowModel-LG demonstrates a slight levelling-off of the snow depth in March and April (as can be seen in the Beaufort, Chukchi, and Kara seas). Conversely, snow depth in NESOSIM steadily increases in the late-season months in these same regions. This discrepancy may be due to a lack of representation of melt or other snow metamorphosis processes in NESOSIM since even at high latitudes, some melt is expected at the end of the season. However, other inter-model process differences may also contribute.

For regionally averaged snow densities from NESOSIM and SnowModel-LG, the limitations of the simple representation of snow density in NESOSIM are apparent since the density in NESOSIM does not exceed 350 kg m^{-3} , the prescribed maximum density of the model. The seasonal cycles of density in NESOSIM exhibit few regional differences. By contrast, in SnowModel-LG, snow densities and their seasonal cycles vary considerably by region. There is an early-season decline in density in several regions for SnowModel-LG that is not represented in NESOSIM, including in the Beaufort, Chukchi, and Kara seas. The densest snow in SnowModel-LG is present in the East Greenland Sea region, exceeding the maximum snow density possible in NESOSIM as early as January. This high density may explain some of the representational discrepancies for NESOSIM in this region. Because the snow cannot become as dense in NESOSIM, given equal amounts of snowfall input, lower density will yield a deeper snowpack. Nevertheless, differences in snow density representation do not entirely explain differences between SnowModel-LG and NESOSIM. In some regions where NESOSIM has a higher density (e.g. Barents Sea), it also has a deeper snowpack.

5 Comparison to MOSAiC and IceBird observations

NESOSIM (run with the ERA5-calibrated parameters (“E5config”) and individually calibrated parameters (“MCMC”)) and SnowModel-LG output is compared to MagnaProbe snow depth (Itkin et al., 2021) and snow density cutter (Macfarlane et al., 2021, 2022) observations from the 2019 to 2020 MOSAiC observational campaign (Macfarlane et al., 2023; Itkin et al., 2023) and to airborne snow radar depth observations from April 2017 and April 2019 from the AWI IceBird observational campaign (Jutila et al., 2022) in Fig. 9. The observed data are aggregated to the respective NESOSIM and SnowModel-LG model grids.

Both NESOSIM and SnowModel-LG snow depths show relatively good agreement with the gridded IceBird and MOSAiC observations, given the considerable differences in spatial and temporal sampling between the modelled and observed values. Overall root-mean-square differences are low (no greater than 10 cm for all products and observations), and

mean biases are generally relatively small (absolute value less than 5 cm for all products and observations). Overall observational correlations are somewhat better for NESOSIM than for SnowModel-LG, particularly for SnowModel-LG driven by MERRA-2. Root-mean-square differences are generally comparable between NESOSIM and SnowModel-LG for MOSAiC data, although they are somewhat lower for NESOSIM when compared to IceBird. Mean biases for snow depth are largest (in absolute value) for SnowModel-LG driven by ERA5 relative to MOSAiC and for NESOSIM driven by JRA-55 with ERA5 parameters relative to IceBird. Some of the MOSAiC snow depth observations show high values ($> 60 \text{ cm}$) that are not captured by the models; these observations tend to coincide with measurements made on or near sea ice ridges. Conversely, for some gridded measurements, the models, especially SnowModel-LG, can considerably overestimate snow depths relative to IceBird. Differences in model gridding may also contribute to differences in correlations and biases. Since SnowModel-LG has a finer grid than that of NESOSIM, observations gridded to be compared with NESOSIM are more aggregated and can have fewer extreme values, as is the case for IceBird.

For snow density, SnowModel-LG has more variability than NESOSIM, which is a consequence of the relatively simple representation of snow density in the latter model. Nevertheless, both models have similar challenges reproducing observed snow density despite the differences in snow density representation between the models, with correlations no greater than 0.23. This disagreement may be partly due to sampling bias in the snow density observations and to the overall difficulty in comparing point measurements to large-scale modelled values. Both models show a high mean bias relative to observed values, with SnowModel-LG driven by MERRA-2 having the largest mean bias, as well as the largest root-mean-square difference. Although NESOSIM has a high bias overall, some snow density measurements were found to exceed 350 kg m^{-3} (the maximum density the model can represent), highlighting another limitation of the model. Snow density observations from MOSAiC could be used in future work to guide the development of the representation of snow density in NESOSIM to help address this limitation. Several large observed density values ($> 400 \text{ kg m}^{-3}$) are not well-represented by either model.

The IceBird and MOSAiC observations have been made available recently and have not been used to calibrate either NESOSIM or SnowModel-LG. IceBird data capture some regional variability over some regions, and MOSAiC data capture seasonal variability for a single season. Nevertheless, both datasets are limited in their observational coverage, and as such, this comparison provides a limited assessment of the impact of the MCMC calibration. The NESOSIM MCMC calibration improves snow depth agreement slightly relative to IceBird observations but has mixed results relative to MOSAiC depth and density observations. In general, for snow depth, the effect of the MCMC calibration is to lower the




Figure 8. Climatologies of regionally averaged snow depth and density from MCMC-calibrated NESOSIM output and SnowModel-LG output for 1980–2021. Regional CDR sea ice area climatologies also shown. “Average” indicates the inter-product average for the three NESOSIM configurations. Climatologies from SnowModel-LG driven with ERA5 and MERRA-2 are also shown. Regions are as described in Fig. A1. Bars indicate the interannual variability in each respective climatology, which is quantified by the standard deviation of the climatology.

mean bias. This brings the model output into closer agreement with the observations when the model is biased high, as is the case for IceBird, but can degrade agreement if the bias is comparatively low, as is the case for NESOSIM driven by MERRA-2 relative to MOSAiC. The correlation is improved for all products relative to IceBird but is unchanged by the calibration relative to MOSAiC depth and density. The MCMC calibration has mixed impacts on agreement with observed density, but the overall impact of the calibration is small, likely due to sampling differences and model representational challenges, as discussed above. The improved agreement with airborne observations may be partly a consequence of similar observations (from OIB) being used as input to the MCMC calibration.

6 Trends in MCMC-calibrated NESOSIM output

Trends were calculated using a Theil–Sen trend estimator, consistent with the approach used by Mudryk et al. (2015). The Theil–Sen trend estimator produces estimates of trends by finding the median of slopes between all pairs of points in a dataset. This approach allows for the estimation of trend

uncertainty based on a chosen confidence interval; a 95 % confidence interval was chosen for this study. In the following discussion, trends are considered significant if the 95 % confidence interval does not overlap with zero. If the confidence interval overlaps with zero, we consider there to be no trend.

Basin-average trends from NESOSIM for snowfall over sea ice, snow depth, snow density, and sea ice area are shown in Fig. 10. The trends in snowfall over sea ice are not statistically significant for most products except for a decline for MERRA-2 from October onwards and a decreasing snowfall trend in January and February for JRA-55. The basin-average trends in snow depth from MCMC-calibrated NESOSIM output vary in magnitude according to the product but are all broadly similar in sign. MERRA-2 has the strongest trends in the basin average overall. The trend is found to be negative (declining snow depth) in all months except September, when the trend is significantly positive for all products, and October, when only MERRA-2 shows a (declining) trend. Snow density trends are generally similar between the products, aside from in October, when only JRA-55 shows a trend. Similarity between the snow density




Figure 9. Scatter plots with kernel density estimates of NESOSIM and SnowModel-LG output compared to observations aggregated daily to the respective model grids from IceBird airborne snow radar (April 2017 and April 2019) (Jutila et al., 2022, 2024a, b), MOSAiC MagnaProbe snow depth (2019–2020) (Itkin et al., 2021), and MOSAiC snow density cutters (2019–2020) (Macfarlane et al., 2021, 2022). Correlation (r), mean bias (MB) and root-mean-square difference (RMS) are shown. The 1 : 1 line is also indicated for reference.




Figure 10. Basin-average monthly trends from 1980 to 2021 for snowfall over sea ice from reanalysis products with CloudSat scaling applied, with MCMC-calibrated NESOSIM snow depth and density, and with CDR sea ice concentration, calculated using a Theil–Sen trend estimator for all products. “Average” denotes the multi-product average. Error bars indicate a 95 % confidence interval as given by the trend estimator; points where there are no trends (the interval overlaps the zero line) are not shown. The dashed grey lines indicate the zero line for reference. SnowModel-LG is excluded from this plot due to differences in model domains.

trends is expected since snow density in NESOSIM is less sensitive to snow input, being primarily dependent on wind speed. Since snow density in NESOSIM is limited to the range of $200\text{--}350\text{ kg m}^{-3}$, the density trends may be spuriously low, particularly towards the end of the season, when density values approach the maximum and interannual variability is low (Fig. 7). The comparatively large declining trends in MERRA-2 for depth may result from its high early-decade snowfall bias relative to the other products. Higher early-year snowfall rates in MERRA-2 can be seen in Fig. 1 and are consistent with findings on Arctic total precipitation in MERRA-2, which is likewise consistently higher in early years (Barrett et al., 2020).

Regional trends in snow depth, snow density, and sea ice area are shown in Fig. 11. Regional trends in snowfall are shown in Fig. A4; although there is regional variation in snowfall trends, most products show no trend for most months, likely due to high interannual variability in snowfall. A large and significant early-season decline is apparent in the Kara Sea region but only for the month of September for most products.

Trends in snow depth are generally stronger and more statistically significant than trends in snowfall. Many of the peripheral seas show a significant declining trend for all products from October onward. These trends are consistent with results from Webster et al. (2019), who find delays in sea ice formation, particularly in the Chukchi Sea region, and attribute declining snow-on-sea-ice trends partly to the increasingly late sea ice onset in this region. The East Greenland Sea region differs noticeably from the other regions shown, with no trend for most months except for a slight increase in November for SnowModel-LG driven by MERRA-2 and declines in April only for NESOSIM driven by MERRA-2 and JRA-55. In the central Arctic region, declines are generally seen only for products driven by MERRA-2. A slight October decline is seen for SnowModel-LG driven by ERA5, a September decline for NESOSIM driven by ERA5, and

declines in November and March for NESOSIM driven by JRA-55.

Despite the differences in the snow depth climatologies between NESOSIM and SnowModel-LG, the snow depth trends show considerably more overlap between the two models. This demonstrates how the choice of snowfall input to reanalysis-based snow-on-sea-ice reconstructions can impact the magnitude and significance of the derived snow depth trends. In several regions, the strongest declining trends are found in MERRA-2, whereas trends often tend to be smaller or absent for ERA5 for both NESOSIM and SnowModel-LG.

For snow density trends, inter-model differences tend to be larger than inter-product differences. Declining trends are largest around October–November for most products and regions except in the Barents and East Greenland seas. The East Greenland Sea region shows increases in snow density for some months and products. NESOSIM and SnowModel-LG disagree on the sign of the snow density trend for September in the Beaufort Sea and for March in the East Greenland Sea for products driven by MERRA-2. Overall, densities in SnowModel-LG tend to show large declines relative to NESOSIM. As discussed previously, NESOSIM end-of-season density trends may be spuriously small due to NESOSIM snow densities approaching their maximum towards the end of the season, although the end-of-season density trends represented in SnowModel-LG also tend to be smaller.

Sea ice area trends vary by region, but strong declines are found for at least part of the season in all regions shown. In the central Arctic and the Siberian sector, as well as in the Beaufort Sea, the largest declining trends are in the earlier months of the cold season (larger trends may be present in months outside of the NESOSIM study period). When sea ice in these regions attains its maximum extent, the trends largely vanish, suggesting a persistent cold-season cover. Towards the North Atlantic (Barents, East Greenland), larger declines are seen in later months.




Figure 11. Monthly trends for regionally averaged quantities over the 1980–2021 time period: reanalysis snowfall over sea ice, snow depth, snow density (from NESOSIM and SnowModel-LG), and sea ice area (from the Climate Data Record product). Error bars indicate a 95 % confidence interval given by the trend estimator; points where there are no trends (the interval overlaps with the zero line) are not shown.

To provide a more regional perspective on snow trends, maps of snow depth trends in NESOSIM and SnowModel-LG output are shown in Fig. 12. Corresponding snowfall trends are shown in Fig. A5. For these plots, trends were also calculated using a Theil–Sen estimator, but only grid squares containing at least 20 years of values were included to exclude spurious trends. Consistent with results from the regional monthly trend plots, there is a lack of snowfall trends over most of the Arctic basin due to the high interannual variability in Arctic snowfall relative to the magnitude of the trends. Slight increases are seen in the Barents Sea and in the Sea of Okhotsk, and decreasing trends are seen east of Greenland for all products. The depth trends are more robust, highlighting a decline in the peripheral seas consistent with the results shown in the regional plots, as well as some slight declines around Hudson Bay and the Labrador Sea. Some significant increasing depth trends north of the Beaufort Sea are found in SnowModel-LG driven by ERA5, as well as in NESOSIM driven by ERA5 and JRA-55, although the products differ regarding the existence of the increasing trend near the North Pole. The spatial pattern of increasing trends north of Greenland and the Canadian Arctic Archipelago and decreasing trends elsewhere is consistent with the pattern of spring-time trends found by other studies, including Webster et al. (2019) and Zhou et al. (2021), although the spatial extent of the significant trends differs. Some differences are expected since the other studies mentioned examine different months

and time periods. There is broad consistency, however, in the declining trend found in the Barents Sea region. The overall large declining trends in depth derived from MERRA-2 are particularly apparent in Fig. 12. ERA5 and JRA-55 agree better with each other on the spatial pattern of the snow depth trends compared to MERRA-2.

The impact of model resolution is apparent since some of the strong trends seen in the SnowModel-LG output are highly localized. There are small but significant increases in snow depth in Hudson Bay that are absent from the NESOSIM output and some increases east of Greenland towards the Fram Strait that are not apparent in the NESOSIM output. This highlights the continued need for further analysis of snow on sea ice in these regions, as well as a need for further observations to validate models in these difficult-to-characterize regions. Nevertheless, the broad patterns of trends between NESOSIM and SnowModel-LG are similar, and there is good agreement between NESOSIM and SnowModel-LG for the central Arctic and adjacent regions.

7 Discussion

The results of this study highlight the value of producing a snow-on-sea-ice product that accounts for uncertainty in model input and formulation and for sparse observations. We find that snowfall climatologies differ considerably between




Figure 12. Snow depth trend maps for March 1980–2021 from the NESOSIM–MCMC run with snowfall input from ERA5, MERRA-2, and JRA-55, and SnowModel-LG with snowfall input from ERA5 (SM-ERA5) and MERRA-2 (SM-MERRA-2). The snow depth is output from NESOSIM with parameters specific to each separate reanalysis product. The trend in the average of the output of the three NESOSIM runs is also plotted (average). Regions with no trends (not significant at a 95 % confidence interval) are shaded in grey. Note that SnowModel-LG is not provided within the Canadian Arctic Archipelago, so data from that region are absent in this map.

the ice-covered region of the Arctic Ocean and the full ice-plus-ocean region. This result is expected, given that sea ice controls atmosphere–ocean moisture and heat fluxes, which in turn influence high-latitude precipitation. In particular, cumuliiform snowfall observed by CloudSat has been shown to vary seasonally with sea ice cover (Kulie and Milani, 2017; Kulie et al., 2016). This impacts the CloudSat scaling, which performs well over the ocean basin but has more difficulty reconciling snowfall over sea ice in some cases, particularly in September when the largest basin-average accumulation takes place, as seen in Fig. 2. This was not considered in Cabaj et al. (2020). The CloudSat scaling factors applied to NESOSIM were calculated over ocean regions, including both sea-ice-free and sea-ice-covered regions, with new factors calculated for JRA-55. Although the scaling of model snowfall input to CloudSat reconciled inter-product differences in snow depth for NESOSIM v1.0, inconsistencies remain in NESOSIM v1.1 snow depth output even with the application of the CloudSat calibration. In general, the CloudSat scaling performs best in aggregate over larger regions since smaller regions may have high variability that may not necessarily be accounted for by the scaling. Re-calculating the CloudSat scaling factors masked exclusively over sea ice may not be feasible due to the relative lack of CloudSat measurements over sea ice. In regions such as the Greenland Sea where sea ice is present only in a very narrow region along the coast, CloudSat reports much less cold-season snowfall relative to the reanalysis products, which suggests that CloudSat may not be adequately sampling snowfall events in the region. This is illustrated in Fig. A6, where CloudSat fails to reproduce the climatology and interannual variability found in the reanalysis products when restricted to over-ice observations in the Greenland/Norwegian Sea region despite agreeing comparatively better with reanalysis products in other regions. As a result, constructing scaling factors using CloudSat restricted to over the ice-covered region yields excessively low (< 18 cm) basin-average snow depths. Hence, in this work, CloudSat scaling factors are calculated based on snowfall over land-free regions, including both open ocean and ice-covered ocean. Nevertheless, future work may entail some revision of the existing CloudSat scaling factors over sea ice, particularly for JRA-55. A more regionally refined calibration may be appropriate, with the caveat that aggregating CloudSat observations over smaller regions may introduce additional uncertainty due to observational under-sampling, as is likely the case with the Greenland/Norwegian Sea region. Refining the calibration using more contemporary forthcoming snowfall measurements from satellite missions such as EarthCARE (Wehr et al., 2023) and using more sophisticated calibration techniques may be other options for future work.

This study also investigates calibration of NESOSIM wind packing and blowing snow parameters using an MCMC process when different reanalysis products are used for NESOSIM snowfall input. The MCMC parameter tuning is de-

pendent on the choice of snowfall input to NESOSIM. Given the discrepancies between the NESOSIM output products prior to the MCMC calibration and the fact that they are all being calibrated to the same observational target, it is unsurprising that the posterior parameter distributions obtained from the calibration differ. This has some implications for the model physics since it suggests that the representation of the physics in the calibrated model is highly dependent on the input. Caution must be taken, then, when taking the wind packing and blowing snow parameters at face value because wide ranges of these parameters can produce physically reasonable model output. This also has implications for other reanalysis-based snow-on-sea-ice estimates, which tend to make use of a selected reanalysis product. As with NESOSIM, snow depth and density in SnowModel-LG are dependent on the choice of reanalysis input, even with the corrections applied in SnowModel-LG to the reanalysis snowfall inputs that were used (Liston et al., 2020). Using automated model parameter recalibration when changing snowfall inputs used for NESOSIM and other snow-on-sea-ice models provides an objective means to address this issue.

The MCMC calibration improves agreement in both the magnitude and the interannual variability in the snow depths output from NESOSIM forced with different reanalysis snowfall products, but it reduces agreement in snow density. This is likely a consequence of two factors: the relative lack of density constraints in the MCMC calibration and the snow density being more sensitive to the wind packing factor than to the snow inputs examined in this study. With different snow inputs, when the wind packing and blowing snow parameters were the same for all runs, there was relatively minimal variation in the density. Given that the density does not depend on the total snow depth but, rather, on the proportion of snow in each layer, one expects the density to be relatively insensitive to snow input and more sensitive to differences in the parameters. Nevertheless, the lack of density constraints in the MCMC calibration may also be an influence, since if density were more strongly constrained, the parameters would be optimized to produce output with a narrower spread in density between the products. Despite this, the estimated density uncertainty in the multi-product average is also quite low, highlighting how the densities produced by NESOSIM are limited to a relatively narrow range due to constraints imposed by the model itself. Since other models produce higher densities (as seen in the comparison with SnowModel-LG) and observations indicate the presence of denser snow than what can be produced by NESOSIM (King et al., 2020; Itkin et al., 2023; Macfarlane et al., 2023), the density assumptions in NESOSIM may need to be revisited. The matter of scale must also be considered because density measurements are highly localized, and NESOSIM represents bulk density over large regions, consistent with its coarse resolution. Overall, this result highlights the need to include additional observational density constraints in the calibration and to eventually reformulate NESOSIM's representation of density.

In regional comparisons with SnowModel-LG, NESOSIM snow depth is found to generally agree over the central Arctic region, but agreement is weaker in the peripheral seas, particularly towards the end of the sea ice season. Overall, NESOSIM snow depth tends to be biased high relative to SnowModel-LG. SnowModel-LG includes loss processes not currently captured by NESOSIM, such as snowmelt, which may contribute to the inter-product differences. NESOSIM and SnowModel-LG also have different approaches to address sea ice drift. The Lagrangian approach of SnowModel-LG allows it to track individual ice parcels, providing a more complex representation of the contribution of sea ice dynamics to the snow budget. NESOSIM presents a more simplified approach but is less computationally expensive to run. Limitations in NESOSIM's representation of snow density may also impact the agreement with SnowModel-LG; snow density in NESOSIM is limited to a maximum value of 350 kg m^{-3} , whereas it can attain larger values in SnowModel-LG. NESOSIM snow depth is biased especially high relative to SnowModel-LG in the East Greenland Sea region, where its density conversely has a low bias. SnowModel-LG includes snow grain parameterizations that are absent in NESOSIM, which allows for the representation of processes that may be essential to quantifying variations in snow density. Nevertheless, discrepancies in snow density representation are likely not the sole explanatory factor of inter-model differences since, in some regions, NESOSIM is biased high relative to SnowModel-LG even while the densities are more comparable.

Despite the differences between NESOSIM and SnowModel-LG in terms of model processes and complexity, the models generally perform comparably in comparisons with in situ MagnaProbe snow depth (Itkin et al., 2021) and density cutter (Macfarlane et al., 2021, 2022) observational data from the 2019–2020 MOSAiC observational campaign, although some larger inter-model differences are seen in comparison with April 2017 and 2019 IceBird snow radar observations (Juttila et al., 2024a, b, 2022). The highly localized nature of the in situ measurements from MOSAiC makes comparisons with comparatively coarse gridded model output challenging. Both models represent snow depth relatively well with respect to MOSAiC and IceBird (correlations upwards of 0.54 and root-mean-square difference (RMSD) $< 11 \text{ cm}$) but have more difficulty representing observed snow density (correlations < 0.23). SnowModel-LG driven by MERRA-2 is notable for having the lowest snow depth correlation and the highest root-mean-square difference relative to MOSAiC and IceBird but also the smallest mean bias compared to all other products for MOSAiC. The root-mean-square difference and mean bias for snow density are also high for this product, which is biased high overall and does not reproduce any lower density values coincident with the observations. This highlights how even with broad-scale observational adjustments such as those made to the reanalysis input to

SnowModel-LG, the use of different snow forcings can still have a large impact on model output. The MCMC calibration has mixed impacts on NESOSIM agreement with MOSAiC, and the overall magnitude of the impacts is small. However, there is slight improvement in correlations and biases relative to IceBird. The observational data used in the MCMC calibration differ considerably from MOSAiC since they include sparser observational data from multiple years and, in the case of OIB measurements, span larger spatial extents. The measurement approach of IceBird more closely resembles that of OIB, both being airborne campaigns making measurements in April. Some of the differences between observed and modelled values are likely a consequence of sampling bias. MOSAiC, for example, includes a variety of measurements made on different days, which were made with the intention of sampling various snow conditions. Thus, MOSAiC measurements show a high degree of variability compared to modelled values, especially as different observational sites change location throughout the season (Itkin et al., 2023). Given the differing scales involved, it is unsurprising that the current configurations of the models can have difficulty reproducing outlying observed values. Model simplifications likely also play a role in the discrepancies, such as in the case of NESOSIM snow density. Since snow in NESOSIM cannot be removed from the lower layer (for a given grid cell, it can only decrease as a consequence of sea ice motion), end-of-season densities are expected to approach 350 kg m^{-3} as an increasing proportion of the snow in each grid cell is old (lower-layer) snow. This yields a limited range of possible representations of snow values and a large number of snow densities near 350 kg m^{-3} , as seen in Fig. 9. Future work could explore using IceBird and MOSAiC measurements to help constrain NESOSIM model output and, in the case of MOSAiC, guide the improved representation of NESOSIM snow density. Caution would be necessary to avoid overfitting, given the spatial and temporal limitations of the observations. Nevertheless, high-quality contemporary airborne and in situ observations of snow on sea ice, such as those from IceBird and MOSAiC, provide a new path to assess and improve model representation of snow on sea ice.

There are many possibilities for further refinements to NESOSIM and to the MCMC calibration approach. NESOSIM is a comparatively low-resolution model with highly parameterized snow processes. This makes it computationally inexpensive, facilitating both MCMC optimization and the rapid production of ice thickness estimates. A key advancement presented by the MCMC calibration lies in the ease of quickly generating updated parameter estimates as new input products are introduced. Additional reanalysis products not examined in this study could be investigated as possible additional inputs to NESOSIM, especially when existing products are inevitably superseded by updated versions. The MCMC calibration itself could also be adjusted; future work could investigate the use of additional observational

constraints on snow depth and especially on snow density. High-quality contemporary snow density observations exist from several observational campaigns such as MOSAiC (Wagner et al., 2022), and although the measurements available during a single day may be highly localized, making use of these observations with an appropriate uncertainty estimate could help constrain the wind packing parameter better, yielding more representative estimates of snow on Arctic sea ice. With the inclusion of additional observational constraints, calibration of additional model parameters in NESOSIM could also be explored. Future work could also investigate impacts of a similar calibration using more complex models or of added complexity in NESOSIM. Several processes, including snow redistribution by wind, sublimation, and melt and refreeze processes, are simplified in NESOSIM. Given the warming conditions in the Arctic, it may be particularly beneficial to represent melt processes in NESOSIM. NESOSIM is run over the September–April period to exclude the melt season, but melt also occurs throughout the year, with a trend towards earlier onset (Stroeve and Notz, 2018). In the current calibration process, observed melt may hence be misrepresented in NESOSIM as a decrease in depth due to densification or blowing snow. Future work could explore parameterizations of additional processes, such as sublimation and snow redistribution by wind, to improve snow depth and density estimates across a variety of environmental conditions.

The relatively coarse resolution of NESOSIM may impact its representativeness since some snow-on-sea-ice processes operate on very small spatial scales and short timescales. The sea ice advection and divergence processes in NESOSIM represent a spatially averaged tendency of snow to be redistributed with sea ice motion but may fail to capture the small-scale effects from localized ridging and the small-scale leads often seen in observational studies (Itkin et al., 2023; Macfarlane et al., 2023). The amount of blowing snow loss due to leads has been observed to be influenced by strong winds and warm air temperatures from Arctic cyclone events, which may be challenging to capture in the current configuration of NESOSIM (Clemens-Sewall et al., 2023). The coarse time resolution also limits the model's ability to capture rapid changes in snow depth due to short-term accumulation events. In a broader modelling context, high-resolution modelling may be necessary to adequately capture small-scale processes (Lecomte et al., 2015). NESOSIM could be run at a higher resolution to take advantage of the higher resolution of available drift products to better capture the influence of sea ice motion. However, sub-grid-scale parameterization would still be necessary to better capture smaller-scale effects. The representation of sea ice differs between reanalysis products and may not align with the observational sea ice concentration used as input to NESOSIM in this work. This, in conjunction with regridding, may introduce some artefacts in regions of marginal sea ice cover such as the Greenland Sea region.

The ERA5 wind product was used in all configurations in this study to isolate the contribution of snowfall to NESOSIM since snowfall is the primary input to the NESOSIM budget. In observational comparisons in the Arctic, ERA5 has been found to perform relatively well compared to other reanalysis products, including JRA-55 and MERRA-2, which motivates the choice of ERA5 over other products (Graham et al., 2019a, b). However, the choice of reanalysis wind input may also have an impact on NESOSIM output. The wind packing and blowing snow processes take effect only when the wind speed exceeds the 5 m s^{-1} wind action threshold. If wind speeds from different input products are on opposite sides of the threshold, wind-related snow processes may take effect at a given location and time for one product and not another. The strength of the blowing snow process is also dependent on wind speed. Future work could investigate the impact of differing wind input products on NESOSIM.

Uncertainties derived from the MCMC parameter uncertainty for each product reflect the widths of the posterior distributions produced from the MCMC process. The percent uncertainties in the model depth and density output are considerably smaller than the percent uncertainty in the posterior parameters (expressed as coefficients of variation). This is consistent with the result in Cabaj et al. (2023), which highlights the relative insensitivity of NESOSIM to the model parameters. However, the snow depth and density uncertainties for NESOSIM run with MERRA-2 and JRA55 are larger than the uncertainties for NESOSIM–ERA5 alone, and likewise, the aggregated multi-product uncertainties exceed the MCMC–ERA5 values. This highlights the importance of accounting for uncertainties arising from differences in reanalysis input products. The multi-product snow depth uncertainty ranges from 8 % to 18 %, which is more reasonable compared to the $< 3 \%$ uncertainty in MCMC–ERA5 alone. The estimated snow density uncertainty is relatively small, particularly when compared to the 40 kg m^{-3} uncertainty prescribed for the ICESat-2 product (Petty et al., 2020) based on the in situ snow observations compiled and analyzed by Warren et al. (1999). The uncertainty estimated for NESOSIM in this study is likely underestimated because of the model's limited density range. Near the end of the season, the densities in many grid cells may be near the maximum 350 kg m^{-3} density value, limiting the possible density variation and thus decreasing the spread in parameter ensemble density values. This challenges the assumption of Gaussian uncertainty distributions, and it may be beneficial to revisit the calibration with non-symmetrical distributions in future work. A similar underestimate of uncertainty takes place in the early season, where uncertainties are artificially low due to the common point of initialization for all the parameter ensemble runs. However, in general, by around the 15th model day, the percent uncertainty saturates, and moreover, this saturation is observed not just for ERA5 (as in Cabaj et al., 2023) but for each individual product. Overall, the estimated uncertainties in the NESOSIM–MCMC-average prod-

uct must be treated with caution since they do not fully characterize all sources of uncertainty, but they can be used to provide a more robust estimate of uncertainty from the NESOSIM model input and calibration assumptions.

Inter-product differences in snow depth and density may have substantial impacts on estimates of sea ice thickness from sea ice altimetry measurements. For example, given representative values of lidar freeboard and representative densities of snow, ice, and water, if snow depth estimates with a 5 cm difference are used to estimate sea ice thickness, the difference in derived sea ice thickness can be as large as 30 cm (Giles et al., 2007). Thus, if trends differ between snow products, trends in derived sea ice thickness will be impacted as well. For sea ice freeboard, a snow product with a decreasing trend would impose an increasing derived ice thickness trend on top of any trend in the freeboard itself. Interannual variability in snow was found to strongly influence the sea ice volume derived from CryoSat-2 altimetry measurements (Bunzel et al., 2018). Hence, differing snow depth trends (or lack thereof) between products could lead to differing conclusions related to trends in derived sea ice thickness.

Intercomparisons of reanalysis products (and quantities derived from them) have some associated caveats. Insight into Arctic precipitation can be gained from the analysis of reanalysis precipitation trends (Boisvert et al., 2018), but caution may be necessary in their interpretation due to discontinuities in the assimilation (Barrett et al., 2020), although contemporary reanalysis systems include bias corrections that mitigate some of the issues introduced by these discontinuities. It is important to consider inter-product differences due to reanalysis inputs because differing snow depth trends between model outputs may have impacts on estimates of the sensitivity of sea ice variables to changes in snow depth. Coupled climate model simulations have found contrasting climate impacts of snow on Arctic sea ice due to competing influences on congelation sea ice growth and surface melt (Holland et al., 2021), but snow-free summers may increase sea ice melt (Webster et al., 2021). Thus, by influencing sea ice thickness, a declining snow depth trend could influence trends in atmosphere–ice heat fluxes, which in turn could influence sea ice extent and other climate variables. There are also discrepancies between the representation of sea ice cover in the three reanalysis products used, which may yield larger inter-product differences, particularly in regions with thinner sea ice (Barrett et al., 2020). Averaging multiple data products together is a well-established approach, and the development of blended land snow products motivates this study (Mudryk et al., 2015, 2018). Constructing a multi-product average using a wider range of input products and incorporating other models and observations could be of future interest.

8 Conclusions

Quantifying snow on Arctic sea ice is an ongoing challenge and existing approaches face difficulties due to spatial and temporal sampling discrepancies, relative biases, and the sparse availability of in situ validation data. Nevertheless, NESOSIM has free parameters that can be observationally calibrated for different snowfall inputs to reconcile inter-product biases. Averaging together model outputs run with different snow inputs can also account for relative differences between the products, and thus, we construct a snow-on-sea-ice product that averages the output of NESOSIM with calibrated snowfall input from ERA5, JRA-55, and MERRA-2 after calibrating each model output to depth and density observations using an MCMC process.

MCMC calibration of NESOSIM with different snowfall inputs following the approach in Cabaj et al. (2023) reconciles differences in snow depth in NESOSIM run with different reanalysis inputs but enhances differences in snow density. The posterior parameter distributions obtained from the calibration differ between the products, with JRA-55 yielding the largest values for the wind packing and blowing snow parameters and yielding posterior parameter distributions with the largest spread compared to those of the other two products.

When MCMC-calibrated regionally aggregated NESOSIM monthly climatologies from 1980 to 2021 are compared to those of SnowModel-LG, good agreement in snow depth is found in the central Arctic Ocean and nearby regions, although NESOSIM has a high bias relative to that of SnowModel-LG in more peripheral regions. Snow densities differ greatly between NESOSIM and SnowModel-LG in both magnitude and seasonality, likely as a consequence of the comparatively simpler representation of snow density in NESOSIM and the weaker constraints in the calibration, although it is challenging to ascertain accuracy in this poorly observed quantity. NESOSIM and SnowModel-LG compare similarly to airborne snow depth measurements from the 2017 and 2019 IceBird campaigns and to in situ snow depth and density measurements from the 2019–2020 MOSAiC observational campaign, and both models have challenges representing snow density in particular. The discrepancies may be partly due to insufficient representation of physical processes in models but also likely result from differences in spatial scales and in timing between the modelled and observed values, as well as measurement sampling biases.

Trends in MCMC-calibrated NESOSIM run with the different products over the 1980–2021 time period broadly agree, with decreasing trends being the strongest for NESOSIM run with MERRA-2 snow input. Basin-average snow depth on Arctic sea ice from NESOSIM is declining in most months and for most products except in September, where there is a slight increasing trend, and trends are not statistically significant in October except for in MERRA-2, where there is a modest decline. Regional snow depth trends vary in

magnitude and statistical significance, but most regions show a declining trend in snow depth on sea ice over the mid-to-late season. In regions where climatologies disagree between models, snow depth trends between models can sometimes show more agreement. In other cases, the choice of reanalysis snow input can still greatly impact the magnitude and statistical significance of snow depth trends, and thus, trends derived from reanalysis-based reconstructions of snow on sea ice must be treated with caution. In general, when using reanalysis-based reconstructions of snow on sea ice, the impact of reanalysis input must be accounted for since changing the reanalysis input may yield less representative model processes.

The uncertainties from MCMC–ERA5 are low relative to other products, and combining uncertainties from the MCMC calibrations for additional reanalysis products yields a more reasonable estimate of basin-average snow depth uncertainty, accounting for uncertainties due to model parameter calibration and the choice of reanalysis input. Estimates of snow density uncertainty remain relatively low, likely due to the implicit constraints on snow density imposed by the model since the fixed density values in each layer impose minimum and maximum density values.

Overall, the findings in this study motivate the continued need for widespread in situ observations of snow on Arctic sea ice, particularly for snow density. In the meantime, however, synthesizing existing models and observations can help provide best-guess estimates of snow on Arctic sea ice. The consensus snow depth product produced in this work incorporates uncertainties from both reanalysis and parameter uncertainties, albeit limited by the simplicity of NESOSIM. Future work in synthesizing models and observations could entail incorporating additional observations and reanalysis products or possibly applying similar calibration and blending approaches to other snow-on-sea-ice products.

Appendix A: Supporting figures




Figure A1. Map of the NESOSIM v1.1 model domain, with the NSIDC-defined regions (Meier and Stewart, 2023) investigated in this study identified. The lower (60° N) and upper (82° N) limits of the latitude band from which CloudSat measurements were aggregated for the CloudSat climatology scaling are indicated with dark-grey contours. The quadrants of the domain used for the CloudSat scaling as applied to the NESOSIM reanalysis input are likewise indicated. Dashed lines indicate the bounds of the model domain for the previous version of NESOSIM, and A–D indicate the corner points from which the CloudSat scaling factors are linearly interpolated.




Figure A2. Basin-average monthly climatologies (1980–2021) of estimated uncertainties for NESOSIM snow depth and density, shown for all products separately and for the multi-product average.




Figure A3. Interannual variability (IAV) of snow depth for NESOSIM, as shown in the bars in Fig. 6 (a) with MCMC–ERA5 calibration and (b) with all products calibrated. This value is calculated as the standard deviation over the time period for the monthly mean for each given month.




Figure A4. Monthly trends for regionally averaged reanalysis snowfall over sea ice quantities over the 1980–2021 time period. Bars indicate 95 % confidence intervals.




Figure A5. Snowfall trend maps over ocean for March 1980–2021 for ERA5, MERRA-2, JRA-55 and the average of the three reanalysis products (average). Snowfall is provided for each respective reanalysis product regridded to the NESOSIM domain. Regions with no trends (not significant at a 95 % confidence interval) are shaded in grey. Snowfall trends over land-covered regions are not shown.




Figure A6. Monthly climatologies (left column) and interannual variability (IAV, right column) of snowfall over sea ice in the 60–82° N latitude band within the four quadrants (East Siberian/Chukchi Sea, Kara/Laptev Sea, Canadian Arctic Archipelago, and Greenland/Norwegian Sea), which are illustrated on the map in Fig. A1. CloudSat fails to adequately represent the monthly climatology in the Greenland/Norwegian Sea region.

Appendix B: MCMC calibration

The MCMC calibration is carried out in this work following the approach in Cabaj et al. (2023) and is described as follows:

1. Begin with a model run with prior parameter values a_0 and observed values y and calculate the log likelihood $\log(p(y|a_0))$. The prior parameter values are associated with prior parameter distributions $p(a_0)$ for which the mean is a_0 and the uncertainty is a prescribed prior parameter uncertainty value. These prior values are given in Table 2. In the following iterative loop, set $a_{\text{current}} = a_0$.
2. For each subsequent step in the Markov chain:
 - a. Choose new parameters a_{test} that are a small step from a_{current} , with the step chosen from $p(a_0)$: a normal distribution centered at a_0 whose standard deviation is determined by the prior parameter uncertainty.
 - b. Calculate the new test log likelihood function $\log(p(y|a_{\text{test}}))$.
 - c. Calculate the log likelihood difference $R = \log(p(y|a_{\text{test}})) - \log(p(y|a_{\text{current}}))$. If $R > \log(U(0, 1))$ (where $U(0, 1)$ is chosen from a uniform distribution between 0 and 1), then the new parameters are accepted and $a_{\text{current}} := a_{\text{test}}$.

As in Cabaj et al. (2023), the observations used for the calibration of NESOSIM are snow depth measurements from the median of airborne Operation IceBridge (OIB) measurements (Petty et al., 2020) and from Cold Regions Research and Engineering Laboratory (CRREL)-Dartmouth buoys (Perovich et al., 2019) and historical snow density measurements from Soviet drifting stations (Radionov et al., 1997; Mallett et al., 2022). OIB measurements are available exclusively in March and April and represent the majority of the observations used for calibrating the parameters. Basin-averaged monthly climatologies are used for the drifting station and buoy measurements, and OIB measurements are aggregated to daily averages over the NESOSIM model grid. This aggregation helps to mitigate the impact of observational biases due to the relatively sparse and infrequent observations in these datasets.

Log likelihood is used to reduce the number of exponential operations calculated and thus reduce computational costs. The log likelihood function used in this study is the same as that used in Cabaj et al. (2023), shown below:

$$L = -\frac{1}{2} \sum_{i=1}^M \frac{(h_{N,i} - h_{O,i})^2}{u_{h_O}^2} - \frac{1}{2} \sum_{j=1}^8 \frac{(\langle \rho_{N,j} \rangle - \langle \rho_{d,j} \rangle)^2}{\langle u_{\rho_{d,j}} \rangle^2} - \frac{1}{2} \sum_{k=1}^8 \frac{(\langle h_{N,k} \rangle - \langle h_{b,k} \rangle)^2}{\langle u_{h_{b,k}} \rangle^2}. \quad (\text{B1})$$

Here, M denotes the number of grid points with Operation IceBridge snow depth measurements, $h_{N,i}$ denotes NESOSIM snow depth output values for a given grid point, and $h_{O,i}$ denotes corresponding OIB snow depth measurements aggregated to a single grid point for a single day. u_{h_O} denotes OIB observational uncertainty. ρ denotes snow density, with subscripts N for NESOSIM and d for drifting stations, respectively, and with $u_{\rho_{d,j}}$ denoting the corresponding uncertainty. h_b denotes CRREL-Dartmouth buoy depth measurements, with corresponding uncertainties $u_{h_{b,k}}$. Angle brackets denote basin-averaged monthly climatologies, and the indices j and k denote months from September to April. The observational uncertainties also account for estimated errors in representativeness in each term.

The acceptance step in the MCMC algorithm allows for the avoidance of local maxima, and posterior parameter distributions are obtained from the distributions of accepted parameters. As in Cabaj et al. (2023), all distributions (the prior parameter distribution, the likelihood function, and the posterior distribution) are assumed to be Gaussian. The modes of the posterior distributions provide optimal values for the parameters. Parameter uncertainty can be estimated from the spread of the posterior distributions. This parameter uncertainty may be propagated through the model to provide estimates of model uncertainty due to parameter uncertainty. Additional parameters may also be calibrated using the MCMC process, but in previous work, limitations were found due to observations not providing sufficiently strong constraints for the optimization to provide suitable optimal parameter values (Cabaj et al., 2023).

To enable NESOSIM to be run with MCMC parameter calibration, the model was modified to keep model output in memory, minimizing the number of file I/O operations and providing a 20 % speedup for MCMC model runs (Cabaj et al., 2023). The NESOSIM-MCMC code was also adapted to enable the calibration to be run with different reanalysis snowfall input products for this study. This highlights the versatility of NESOSIM as a model well-suited to observational calibration.

Code and data availability. The NASA Eulerian Snow On Sea Ice Model (Petty et al., 2018) is available at <https://doi.org/10.5281/zenodo.6342069> (Petty and Cabaj, 2022), modified for MCMC at <https://doi.org/10.5281/zenodo.7644948> (Cabaj and Petty, 2023). The final calibrated multi-product average with uncertainties, including the individually calibrated model runs and the Cloud-Sat scaling factors, is provided at <https://doi.org/10.5281/zenodo.13307800> (Cabaj et al., 2024). SnowModel-LG (Liston et al., 2020; Stroeve et al., 2020) output was obtained from the NSIDC at <https://doi.org/10.5067/27A0P5M6LZBI> (Liston et al., 2021). Forcing data for NESOSIM, including atmospheric input from ECMWF ERA5 (Hersbach et al., 2020; Hersbach et al., 2023, <https://doi.org/10.24381/cds.adbb2d47>), NOAA/NSIDC sea ice concentration at <https://doi.org/10.7265/EFMZ-2T65> (Meier et al., 2021), EUMETSAT OSI SAF sea ice drift (Laverne et al., 2010;

EUMETSAT Ocean and Sea Ice Satellite Application Facility, 2017, https://doi.org/10.15770/EUM_SAF_OSI_NRT_2007, and NSIDCv4 Polar Pathfinder sea ice drift at <https://doi.org/10.5067/INAWUW07QH7B> (Tschudi et al., 2019), as well as processed Operation IceBridge data (Petty et al., 2023b), is available on Zenodo at <https://doi.org/10.5281/zenodo.7051062> (Petty, 2021). Additional forcing data for NESOSIM were regridded from MERRA-2 (Gelaro et al., 2017) obtained from NASA GES DISC at <https://doi.org/10.5067/7MCPBJ41Y0K6> (Global Modeling and Assimilation Office (GMAO), 2015) and from JRA-55 (Kobayashi et al., 2015) obtained from the NCAR RDA at <https://doi.org/10.5065/D6HH6H41> (Japan Meteorological Agency, 2013). The CloudSat 2C-SNOW-PROFILE product (Wood et al., 2013, 2014) was obtained from <https://www.cloudsat.cira.colostate.edu/data-products/2c-snow-profile> (Wood and L'Ecuyer, 2018). In addition to the processed Operation IceBridge data mentioned above, the NESOSIM MCMC calibration also makes use of processed snow input from Soviet drifting stations (Mallett et al., 2022; Radionov et al., 1997) and from CRREL-Dartmouth buoy observations (Perovich et al., 2022). AWI IceBird data were obtained from Jutila et al. (2022), Jutila et al. (2024a, <https://doi.org/10.1594/PANGAEA.966059>), and Jutila et al. (2024b, <https://doi.org/10.1594/PANGAEA.966009>). MOSAiC MagnaProbe snow depth data (Itkin et al., 2021) were obtained from <https://doi.org/10.1594/PANGAEA.937781>, and MOSAiC snow density cutter data were obtained from <https://doi.org/10.1594/PANGAEA.935934> (Macfarlane et al., 2021) and <https://doi.org/10.1594/PANGAEA.940214> (Macfarlane et al., 2022).

Supplement. The supplement related to this article is available online at <https://doi.org/10.5194/tc-19-3033-2025-supplement>.

Author contributions. AC designed this study, with input from PJK and AAP. AC ran the model simulations based on model code originally developed by AAP (with subsequent contributions from AC). AC conducted the analysis in the article, with input from all co-authors. AC prepared the paper with contributions from all co-authors.

Competing interests. The contact author has declared that none of the authors has any competing interests.

Disclaimer. Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes every effort to include appropriate place names, the final responsibility lies with the authors.

Acknowledgements. Alex Cabaj and Paul J. Kushner conducted this study with support from the Canadian Space Agency Earth System Science, Data Analyses Fund, grant no. 16SAUSSNOW. Alek A. Petty gratefully acknowledges support from NASA under grant no. 80NSSC23K1253, awarded by the Cryospheric Sciences program (solicitation NNH22ZDA001N-ICESAT2). We thank all those

who contributed to MOSAiC (Nixdorf et al., 2021), as well as those who contributed to IceBird, the CRREL-Dartmouth buoy program, Operation IceBridge, and all other observational campaigns that provided the data used in this work. The authors would also like to thank Walt Meier for his helpful feedback on a precursor to this paper.

Financial support. This research has been supported by the Canadian Space Agency Earth System Science Data Analyses Fund (grant no. 16SAUSSNOW) and NASA through the Cryospheric Sciences program (grant no. 80NSSC23K125, solicitation NNH22ZDA001N-ICESAT2).

Review statement. This paper was edited by John Yackel and reviewed by Tian Tian and two anonymous referees.

References

- Barrett, A. P., Stroeve, J. C., and Serreze, M. C.: Arctic Ocean Precipitation From Atmospheric Reanalyses and Comparisons With North Pole Drifting Station Records, *J. Geophys. Res.-Oceans*, 125, e2019JC015415, <https://doi.org/10.1029/2019JC015415>, 2020.
- Behrangi, A., Christensen, M., Richardson, M., Lebsock, M., Stephens, G., Huffman, G. J., Bolvin, D., Adler, R. F., Gardner, A., Lambriksen, B., and Fetzer, E.: Status of High-Latitude Precipitation Estimates from Observations and Reanalyses, *J. Geophys. Res.-Atmos.*, 121, 4468–4486, <https://doi.org/10.1002/2015JD024546>, 2016.
- Blanchard-Wrigglesworth, E., Webster, M. A., Farrell, S. L., and Bitz, C. M.: Reconstruction of Snow on Arctic Sea Ice, *J. Geophys. Res.-Oceans*, 123, 3588–3602, <https://doi.org/10.1002/2017JC013364>, 2018.
- Boisvert, L. N., Webster, M. A., Petty, A. A., Markus, T., Bromwich, D. H., and Cullather, R. I.: Intercomparison of Precipitation Estimates over the Arctic Ocean and Its Peripheral Seas from Reanalyses, *J. Climate*, 31, 8441–8462, <https://doi.org/10.1175/JCLI-D-18-0125.1>, 2018.
- Brucker, L. and Markus, T.: Arctic-scale assessment of satellite passive microwave-derived snow depth on sea ice using Operation IceBridge airborne data, *J. Geophys. Res.-Oceans*, 118, 2892–2905, <https://doi.org/10.1002/jgrc.20228>, 2013.
- Bunzel, F., Notz, D., and Pedersen, L. T.: Retrievals of Arctic Sea-Ice Volume and Its Trend Significantly Affected by Inter-annual Snow Variability, *Geophys. Res. Lett.*, 45, 11751–11759, <https://doi.org/10.1029/2018GL078867>, 2018.
- Cabaj, A. and Petty, A.: NESOSIM with MCMC calibration, Version v1.1-mcmc, Zenodo [code], <https://doi.org/10.5281/zenodo.7644948>, 2023.
- Cabaj, A., Kushner, P. J., Fletcher, C. G., Howell, S., and Petty, A. A.: Constraining Reanalysis Snowfall Over the Arctic Ocean Using CloudSat Observations, *Geophys. Res. Lett.*, 47, e2019GL086426, <https://doi.org/10.1029/2019GL086426>, 2020.
- Cabaj, A., Kushner, P. J., and Petty, A. A.: Automated Calibration of a Snow-On-Sea-Ice Model, *Earth and Space Science*, 10, e2022EA002655, <https://doi.org/10.1029/2022EA002655>, 2023.

- Cabaj, A., Petty, A. A., and Kushner, P. J.: NESOSIM-MCMC Multi-Reanalysis-Average Product with Uncertainty Estimates, Zenodo [data set], <https://doi.org/10.5281/zenodo.13307800>, 2024.
- Cavalieri, D. J., Parkinson, C. L., Gloersen, P., and Zwally, H. J.: Sea Ice Concentrations from Nimbus-7 SMMR and DMSP SSM/I-SSMIS Passive Microwave Data, Version 1, NASA National Snow and Ice Data Center Distributed Active Archive Center [data set], Boulder, Colorado, USA, <https://doi.org/10.5067/8GQ8LZQVL0VL>, 1996.
- Clemens-Sewall, D., Polashenski, C., Frey, M. M., Cox, C. J., Granskog, M. A., Macfarlane, A. R., Fons, S. W., Schmale, J., Hutchings, J. K., von Albedyll, L., Arndt, S., Schneebeil, M., and Perovich, D.: Snow Loss Into Leads in Arctic Sea Ice: Minimal in Typical Wintertime Conditions, but High During a Warm and Windy Snowfall Event, *Geophys. Res. Lett.*, 50, e2023GL102816, <https://doi.org/10.1029/2023GL102816>, 2023.
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, I., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J.-N. and Vitart, F.: The ERA-Interim Reanalysis: Configuration and Performance of the Data Assimilation System, *Q. J. Roy. Meteor. Soc.*, 137, 553–597, <https://doi.org/10.1002/qj.828>, 2011.
- Edel, L., Claud, C., Genthon, C., Palermé, C., Wood, N., L'Ecuyer, T., and Bromwich, D.: Arctic Snowfall from Cloud-Sat Observations and Reanalyses, *J. Climate*, 33, 2093–2109, <https://doi.org/10.1175/JCLI-D-19-0105.1>, 2020.
- EUMETSAT Ocean and Sea Ice Satellite Application Facility: Global Low Resolution Sea Ice Drift, EUMETSAT Data Centre [data set], https://doi.org/10.15770/EUM_SAF_OSI_NRT_2007, 2017.
- Gelaro, R., McCarty, W., Suárez, M. J., Todling, R., Molod, A., Takacs, L., Randles, C. A., Darmenov, A., Bosilovich, M. G., Reichle, R., Wargan, K., Coy, L., Cullather, R., Draper, C., Akella, S., Buchard, V., Conaty, A., da Silva, A. M., Gu, W., Kim, G.-K., Koster, R., Lucchesi, R., Merkova, D., Nielsen, J. E., Partyka, G., Pawson, S., Putman, W., Rienecker, M., Schubert, S. D., Sienkiewicz, M., and Zhao, B.: The Modern-Era Retrospective Analysis for Research and Applications, Version 2 (MERRA-2), *J. Climate*, 30, 5419–5454, <https://doi.org/10.1175/JCLI-D-16-0758.1>, 2017.
- Gelman, A., Carlin, J. B., Stern, H. S., Dunson, D. B., Vehtari, A., and Rubin, D. B.: Bayesian Data Analysis, CRC Texts in Statistical Science, CRC Press, Boca Raton, 3rd edn., ISBN: 9781439840955, 2013.
- Giles, K., Laxon, S., Wingham, D., Wallis, D., Krabill, W., Leuschen, C., McAdoo, D., Manizade, S., and Raney, R.: Combined airborne laser and radar altimeter measurements over the Fram Strait in May 2002, *Remote Sens. Environ.*, 111, 182–194, <https://doi.org/10.1016/j.rse.2007.02.037>, 2007.
- Global Modeling and Assimilation Office (GMAO): MERRA-2 tavg1_2d_flux_Nx: 2d,1-Hourly,Time-Averaged,Single-Level,Assimilation,Surface Flux Diagnostics V5.12.4, Goddard Earth Sciences Data and Information Services Center (GES DISC) [data set], Greenbelt, MD, USA, <https://doi.org/10.5067/7MCPBJ41YOK6>, 2015.
- Graham, R. M., Cohen, L., Ritzhaupt, N., Segger, B., Graversen, R. G., Rinke, A., Walden, V. P., Granskog, M. A., and Hudson, S. R.: Evaluation of Six Atmospheric Reanalyses over Arctic Sea Ice from Winter to Early Summer, *J. Climate*, 32, 4121–4143, <https://doi.org/10.1175/JCLI-D-18-0643.1>, 2019a.
- Graham, R. M., Hudson, S. R., and Maturilli, M.: Improved Performance of ERA5 in Arctic Gateway Relative to Four Global Atmospheric Reanalyses, *Geophys. Res. Lett.*, 46, 6138–6147, <https://doi.org/10.1029/2019GL082781>, 2019b.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.-N.: The ERA5 Global Reanalysis, *Q. J. Roy. Meteor. Soc.*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.
- Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey, C., Radu, R., Rozum, I., Schepers, D., Simmons, A., Soci, C., Dee, D., and Thépaut, J.-N.: ERA5 hourly data on single levels from 1940 to present, Copernicus Climate Change Service (C3S) Climate Data Store (CDS) [data set], <https://doi.org/10.24381/cds.adbb2d47>, 2023.
- Holland, M. M., Clemens-Sewall, D., Landrum, L., Light, B., Perovich, D., Polashenski, C., Smith, M., and Webster, M.: The influence of snow on sea ice as assessed from simulations of CESM2, *The Cryosphere*, 15, 4981–4998, <https://doi.org/10.5194/tc-15-4981-2021>, 2021.
- Itkin, P., Webster, M., Hendricks, S., Oggier, M., Jaggi, M., Ricker, R., Arndt, S., Divine, D. V., von Albedyll, L., Raphael, I., Rohde, J., and Liston, G. E.: Magnaprobe snow and melt pond depth measurements from the 2019–2020 MOSAiC expedition, PANGAEA [data set], <https://doi.org/10.1594/PANGAEA.937781>, 2021.
- Itkin, P., Hendricks, S., Webster, M., von Albedyll, L., Arndt, S., Divine, D., Jaggi, M., Oggier, M., Raphael, I., Ricker, R., Rohde, J., Schneebeil, M., and Liston, G. E.: Sea ice and snow characteristics from year-long transects at the MOSAiC Central Observatory, *Elementa: Science of the Anthropocene*, 11, 00048, <https://doi.org/10.1525/elementa.2022.00048>, 2023.
- Japan Meteorological Agency: JRA-55: Japanese 55-year Reanalysis, Daily 3-Hourly and 6-Hourly Data, Research Data Archive at the National Center for Atmospheric Research, Computational and Information Systems Laboratory [data set], <https://doi.org/10.5065/D6HH6H41>, 2013.
- Jutila, A., Hendricks, S., Ricker, R., von Albedyll, L., Krumpén, T., and Haas, C.: Retrieval and parameterisation of sea-ice bulk density from airborne multi-sensor measurements, *The Cryosphere*, 16, 259–275, <https://doi.org/10.5194/tc-16-259-2022>, 2022.
- Jutila, A., Hendricks, S., Ricker, R., von Albedyll, L., and Haas, C.: Airborne sea ice parameters during aircraft flight P6_217_ICEBIRD_2019_1904051001, Version 2, PANGAEA [data set], <https://doi.org/10.1594/PANGAEA.966059>, 2024a.

- Jutila, A., Hendricks, S., Ricker, R., von Albedyll, L., and Haas, C.: Airborne sea ice parameters during the PAMARCMIP2017 campaign in the Arctic Ocean, Version 2, PANGAEA [data set], <https://doi.org/10.1594/PANGAEA.966009>, 2024b.
- King, J., Howell, S., Brady, M., Toose, P., Derksen, C., Haas, C., and Beckers, J.: Local-scale variability of snow density on Arctic sea ice, *The Cryosphere*, 14, 4323–4339, <https://doi.org/10.5194/tc-14-4323-2020>, 2020.
- Kobayashi, S., Ota, Y., Harada, Y., Ebata, A., Moriya, M., Onoda, H., Onogi, K., Kamahori, H., Kobayashi, C., Endo, H., Miyaoka, K., and Takahashi, K.: The JRA-55 Reanalysis: General Specifications and Basic Characteristics, *J. Meteorol. Soc. Jpn., Ser. II*, 93, 5–48, <https://doi.org/10.2151/jmsj.2015-001>, 2015.
- Kulie, M. S. and Bennartz, R.: Utilizing Spaceborne Radars to Retrieve Dry Snowfall, *J. Appl. Meteorol. Clim.*, 48, 2564–2580, <https://doi.org/10.1175/2009JAMC2193.1>, 2009.
- Kulie, M. S. and Milani, L.: Seasonal Variability of Shallow Cumuliform Snowfall: A CloudSat Perspective, *Q. J. Roy. Meteor. Soc.*, 144, 329–343, <https://doi.org/10.1002/qj.3222>, 2017.
- Kulie, M. S., Milani, L., Wood, N. B., Tushaus, S. A., Bennartz, R., and L'Ecuier, T. S.: A Shallow Cumuliform Snowfall Census Using Spaceborne Radar, *J. Hydrometeorol.*, 17, 1261–1279, <https://doi.org/10.1175/JHM-D-15-0123.1>, 2016.
- Kwok, R. and Cunningham, G. F.: ICESat over Arctic sea ice: Estimation of snow depth and ice thickness, *J. Geophys. Res.-Oceans*, 113, C08010, <https://doi.org/10.1029/2008JC004753>, 2008.
- Kwok, R., Kacimi, S., Webster, M. A., Kurtz, N. T., and Petty, A. A.: Arctic Snow Depth and Sea Ice Thickness From ICESat-2 and CryoSat-2 Freeboards: A First Examination, *J. Geophys. Res.-Oceans*, 125, e2019JC016008, <https://doi.org/10.1029/2019JC016008>, 2020.
- Lavergne, T., Eastwood, S., Teffah, Z., Schyberg, H., and Breivik, L.-A.: Sea Ice Motion from Low-Resolution Satellite Sensors: An Alternative Method and Its Validation in the Arctic, *J. Geophys. Res.-Oceans*, 115, C10032, <https://doi.org/10.1029/2009JC005958>, 2010.
- Lawrence, I. R., Tsamados, M. C., Stroeve, J. C., Armitage, T. W. K., and Ridout, A. L.: Estimating snow depth over Arctic sea ice from calibrated dual-frequency radar freeboards, *The Cryosphere*, 12, 3551–3564, <https://doi.org/10.5194/tc-12-3551-2018>, 2018.
- Lecomte, O., Fichet, T., Flocco, D., Schroeder, D., and Vancoppenolle, M.: Interactions between wind-blown snow redistribution and melt ponds in a coupled ocean–sea ice model, *Ocean Model.*, 87, 67–80, <https://doi.org/10.1016/j.ocemod.2014.12.003>, 2015.
- Lindsay, R., Wensnahan, M., Schweiger, A., and Zhang, J.: Evaluation of Seven Different Atmospheric Reanalysis Products in the Arctic, *J. Climate*, 27, 2588–2606, <https://doi.org/10.1175/JCLI-D-13-00014.1>, 2014.
- Liston, G. E. and Hiemstra, C. A.: A Simple Data Assimilation System for Complex Snow Distributions (SnowAssim), *J. Hydrometeorol.*, 9, 989–1004, <https://doi.org/10.1175/2008JHM871.1>, 2008.
- Liston, G. E., Itkin, P., Stroeve, J., Tschudi, M., Stewart, J. S., Pedersen, S. H., Reinking, A. K., and Elder, K.: A Lagrangian Snow-Evolution System for Sea-Ice Applications (SnowModel-LG): Part I – Model Description, *J. Geophys. Res.-Oceans*, 125, e2019JC015913, <https://doi.org/10.1029/2019JC015913>, 2020.
- Liston, G. E., Stroeve, J., and Itkin, P.: Lagrangian Snow Distributions for Sea-Ice Applications, Version 1, NASA National Snow and Ice Data Center Distributed Active Archive Center [data set], Boulder, Colorado, USA, <https://doi.org/10.5067/27A0P5M6LZBI>, 2021.
- Macfarlane, A. R., Schneebeil, M., Dadic, R., Wagner, D. N., Arndt, S., Clemens-Sewall, D., Hämmerle, S., Hannula, H.-R., Jaggi, M., Kolabutin, N., Krampe, D., Lehning, M., Matero, I., Nicolaus, M., Oggier, M., Pirazzini, R., Polashenski, C., Raphael, I., Regnery, J., Shimanchuck, E., Smith, M. M., and Tavri, A.: Snowpit raw data collected during the MOSAiC expedition, PANGAEA [data set], <https://doi.org/10.1594/PANGAEA.935934>, 2021.
- Macfarlane, A. R., Schneebeil, M., Dadic, R., Wagner, D. N., Arndt, S., Clemens-Sewall, D., Hämmerle, S., Hannula, H.-R., Jaggi, M., Kolabutin, N., Krampe, D., Lehning, M., Matero, I., Nicolaus, M., Oggier, M., Pirazzini, R., Polashenski, C., Raphael, I., Regnery, J., Shimanchuck, E., Smith, M. M., and Tavri, A.: Snowpit snow density cutter profiles measured during the MOSAiC expedition, PANGAEA [data set], <https://doi.org/10.1594/PANGAEA.940214>, 2022.
- Macfarlane, A. R., Schneebeil, M., Dadic, R., Tavri, A., Immerz, A., Polashenski, C., Krampe, D., Clemens-Sewall, D., Wagner, D. N., Perovich, D. K., Henna-Reetta, H., Raphael, I., Matero, I., Regnery, J., Smith, M. M., Nicolaus, M., Jaggi, M., Oggier, M., Webster, M. A., Lehning, M., Kolabutin, N., Itkin, P., Naderpour, R., Pirazzini, R., Hämmerle, S., Arndt, S., and Fons, S.: A Database of Snow on Sea Ice in the Central Arctic Collected during the MOSAiC expedition, *Scientific Data*, 10, 398, <https://doi.org/10.1038/s41597-023-02273-1>, 2023.
- MacGregor, J. A., Boisvert, L. N., Medley, B., Petty, A. A., Harbeck, J. P., Bell, R. E., Blair, J. B., Blanchard-Wrigglesworth, E., Buckley, E. M., Christoffersen, M. S., Cochran, J. R., Csathó, B. M., De Marco, E. L., Dominguez, R. T., Fahnestock, M. A., Farrell, S. L., Gogineni, S. P., Greenbaum, J. S., Hansen, C. M., Hofton, M. A., Holt, J. W., Jezek, K. C., Koenig, L. S., Kurtz, N. T., Kwok, R., Larsen, C. F., Leuschen, C. J., Locke, C. D., Manizade, S. S., Martin, S., Neumann, T. A., Nowicki, S. M., Paden, J. D., Richter-Menge, J. A., Rignot, E. J., Rodríguez-Morales, F., Siegfried, M. R., Smith, B. E., Sonntag, J. G., Studinger, M., Tinto, K. J., Truffer, M., Wagner, T. P., Woods, J. E., Young, D. A., and Yungel, J. K.: The Scientific Legacy of NASA's Operation IceBridge, *Rev. Geophys.*, 59, e2020RG000712, <https://doi.org/10.1029/2020RG000712>, 2021.
- Mallett, R. D. C., Stroeve, J. C., Tsamados, M., Willatt, R., Newman, T., Nandan, V., Landy, J. C., Itkin, P., Oggier, M., Jaggi, M., and Perovich, D. K.: Sub-Kilometre Scale Distribution of Snow Depth on Arctic Sea Ice from Soviet Drifting Stations, *J. Glaciol.*, 68, 1–13, <https://doi.org/10.1017/jog.2022.18>, 2022.
- Meier, W., Fetterer, F., Windnagel, A., and Stewart, S.: NOAA/NSIDC Climate Data Record of Passive Microwave Sea Ice Concentration, Version 4, National Snow and Ice Data Center [data set], Boulder, Colorado, USA, <https://doi.org/10.7265/EFMZ-2T65>, 2021.
- Meier, W. N. and Stewart, J. S.: NSIDC Land, Ocean, Coast, Ice, and Sea Ice Region Masks, NSIDC Spe-

- cial Report No. 25, National Snow and Ice Data Center, Boulder, CO, USA, <https://nsidc.org/sites/default/files/documents/technical-reference/nsidc-special-report-25.pdf> (last access: 15 June 2024), 2023.
- Milani, L. and Wood, N. B.: Biases in CloudSat Falling Snow Estimates Resulting from Daylight-Only Operations, *Remote Sensing*, 13, 2041, <https://doi.org/10.3390/rs13112041>, 2021.
- Milani, L., Kulie, M. S., Casella, D., Dietrich, S., L'Ecuyer, T. S., Panegrossi, G., Porcù, F., Sanò, P., and Wood, N. B.: CloudSat snowfall estimates over Antarctica and the Southern Ocean: An assessment of independent retrieval methodologies and multi-year snowfall analysis, *Atmos. Res.*, 213, 121–135, <https://doi.org/10.1016/j.atmosres.2018.05.015>, 2018.
- Mudryk, L. R., Derksen, C., Kushner, P. J., and Brown, R.: Characterization of Northern Hemisphere Snow Water Equivalent Datasets, 1981–2010, *J. Climate*, 28, 8037–8051, <https://doi.org/10.1175/JCLI-D-15-0229.1>, 2015.
- Mudryk, L. R., Derksen, C., Howell, S., Laliberté, F., Thackeray, C., Sospedra-Alfonso, R., Vionnet, V., Kushner, P. J., and Brown, R.: Canadian snow and sea ice: historical trends and projections, *The Cryosphere*, 12, 1157–1176, <https://doi.org/10.5194/tc-12-1157-2018>, 2018.
- Nicolaus, M., Hoppmann, M., Arndt, S., Hendricks, S., Katlein, C., König-Langlo, G., Nicolaus, A., Rossmann, L., Schiller, M., Schwegmann, S., Langevin, D., and Bartsch, A.: Snow Height and Air Temperature on Sea Ice from Snow Buoy Measurements, PANGAEA [data set], <https://doi.org/10.1594/PANGAEA.875638>, 2017.
- Nicolaus, M., Perovich, D. K., Spreen, G., Granskog, M. A., von Albedyll, L., Angelopoulos, M., Anhaus, P., Arndt, S., Belter, H. J., Bessonov, V., Birnbaum, G., Brauchle, J., Calmer, R., Cardellach, E., Cheng, B., Clemens-Sewall, D., Dadic, R., Damm, E., de Boer, G., Demir, O., Dethloff, K., Divine, D. V., Fong, A. A., Fons, S., Frey, M. M., Fuchs, N., Gabarró, C., Gerland, S., Goessling, H. F., Gradinger, R., Haapala, J., Haas, C., Hamilton, J., Hannula, H.-R., Hendricks, S., Herber, A., Heuzé, C., Hoppmann, M., Høyland, K. V., Huntemann, M., Hutchings, J. K., Hwang, B., Itkin, P., Jacobi, H.-W., Jaggi, M., Jutila, A., Kaleschke, L., Katlein, C., Kolabutin, N., Krampe, D., Kristensen, S. S., Krumpen, T., Kurtz, N., Lampert, A., Lange, B. A., Lei, R., Light, B., Linhardt, F., Liston, G. E., Loose, B., Macfarlane, A. R., Mahmud, M., Matero, I. O., Maus, S., Morgenstern, A., Naderpour, R., Nandan, V., Niubom, A., Oggier, M., Oppelt, N., Pätzold, F., Perron, C., Petrovsky, T., Pirazzini, R., Polashenski, C., Rabe, B., Raphael, I. A., Regnery, J., Rex, M., Ricker, R., Riemann-Campe, K., Rinke, A., Rohde, J., Salganik, E., Scharien, R. K., Schiller, M., Schneebeli, M., Semmling, M., Shimanchuk, E., Shupe, M. D., Smith, M. M., Smolyanitsky, V., Sokolov, V., Stanton, T., Stroeve, J., Thielke, L., Timofeeva, A., Tonboe, R. T., Tavri, A., Tsamados, M., Wagner, D. N., Watkins, D., Webster, M., and Wendisch, M.: Overview of the MOSAiC expedition: Snow and sea ice, *Elementa: Science of the Anthropocene*, 10, 000046, <https://doi.org/10.1525/elementa.2021.000046>, 2022.
- Nixdorf, U., Dethloff, K., Rex, M., Shupe, M., Sommerfeld, A., Perovich, D. K., Nicolaus, M., Heuzé, C., Rabe, B., Loose, B., Damm, E., Gradinger, R., Fong, A., Maslowski, W., Rinke, A., Kwok, R., Spreen, G., Wendisch, M., Herber, A., Hirsekorn, M., Mohaupt, V., Frickenhaus, S., Immerz, A., Weiss-Tuider, K., König, B., Menedoht, D., Regnery, J., Gerchow, P., Ransby, D., Krumpen, T., Morgenstern, A., Haas, C., Kanzow, T., Rack, F. R., Saitzev, V., Sokolov, V., Makarov, A., Schwarze, S., Wunderlich, T., Wurr, K., and Boetius, A.: MOSAiC Extended Acknowledgement, Zenodo, <https://doi.org/10.5281/zenodo.5541624>, 2021.
- Peng, G., Meier, W. N., Scott, D. J., and Savoie, M. H.: A long-term and reproducible passive microwave sea ice concentration data record for climate studies and monitoring, *Earth Syst. Sci. Data*, 5, 311–318, <https://doi.org/10.5194/essd-5-311-2013>, 2013.
- Perovich, D. K., Richter-Menge, J. A., and Polashenski, C.: Observing and Understanding Climate Change: Monitoring the Mass Balance, Motion, and Thickness of Arctic Sea Ice, CRREL-Dartmouth [data set], <http://imb-crrel-dartmouth.org> (last access: 30 July 2025), 2019.
- Perovich, D. K., Richter-Menge, J. A., and Polashenski, C.: Observing and Understanding Climate Change: Monitoring the Mass Balance, Motion, and Thickness of Arctic Sea Ice, CRREL-Dartmouth [data set], <http://imb-crrel-dartmouth.org> (last access: 30 July 2025), 2022.
- Petty, A.: NASA Eulerian Snow On Sea Ice Model Version 1.1 (NESOSIMv1.1) data: 1980 - 2022, Version v1.1, Zenodo [data set], <https://doi.org/10.5281/zenodo.7051062>, 2021.
- Petty, A. and Cabaj, A.: akpetty/NESOSIM, Version v1.1.1, Zenodo [code], <https://doi.org/10.5281/zenodo.6342069>, 2022.
- Petty, A. A., Webster, M., Boisvert, L., and Markus, T.: The NASA Eulerian Snow on Sea Ice Model (NESOSIM) v1.0: initial model development and analysis, *Geosci. Model Dev.*, 11, 4577–4602, <https://doi.org/10.5194/gmd-11-4577-2018>, 2018.
- Petty, A. A., Kurtz, N. T., Kwok, R., Markus, T., and Neumann, T. A.: Winter Arctic Sea Ice Thickness From ICESat-2 Freeboards, *J. Geophys. Res.-Oceans*, 125, e2019JC015764, <https://doi.org/10.1029/2019JC015764>, 2020.
- Petty, A. A., Kurtz, N., Kwok, R., Markus, T., Neumann, T. A., and Keeney, N.: ICESat-2 L4 Monthly Gridded Sea Ice Thickness, Version 3, National Snow and Ice Data Center (NSIDC) [data set], <https://doi.org/10.5067/ZCSU8Y5U1BQW>, 2023a.
- Petty, A. A., Keeney, N., Cabaj, A., Kushner, P., and Bagnardi, M.: Winter Arctic sea ice thickness from ICESat-2: upgrades to freeboard and snow loading estimates and an assessment of the first three winters of data collection, *The Cryosphere*, 17, 127–156, <https://doi.org/10.5194/tc-17-127-2023>, 2023b.
- Radionov, V. F., Bryazgin, N. N., and Alexandrov, E. I.: The Snow Cover of the Arctic Basin, APL-UW TR 9701, 98 pp., Washington University Applied Physics Laboratory, <https://apps.dtic.mil/sti/tr/pdf/ADA327057.pdf> (last access: 1 July 2024), 1997.
- Rostosky, P., Spreen, G., Farrell, S. L., Frost, T., Heygster, G., and Melsheimer, C.: Snow Depth Retrieval on Arctic Sea Ice From Passive Microwave Radiometers—Improvements and Extensions to Multiyear Ice Using Lower Frequencies, *J. Geophys. Res.-Oceans*, 123, 7120–7138, <https://doi.org/10.1029/2018JC014028>, 2018.
- Stroeve, J. and Notz, D.: Changing state of Arctic sea ice across all seasons, *Environ. Res. Lett.*, 13, 103001, <https://doi.org/10.1088/1748-9326/aade56>, 2018.
- Stroeve, J., Liston, G. E., Buzzard, S., Zhou, L., Mallett, R., Barrett, A., Tschudi, M., Tsamados, M., Itkin, P., and Stewart, J. S.: A Lagrangian Snow Evolution System for Sea Ice Applications (SnowModel-LG): Part II – Analyses, *J. Geophys. Res.-Oceans*,

- 125, e2019JC015900, <https://doi.org/10.1029/2019JC015900>, 2020.
- Tschudi, M., Meier, W. N., Stewart, J. S., Fowler, C., and Maslanik, J.: Polar Pathfinder Daily 25 km EASE-Grid Sea Ice Motion Vectors, Version 4, NASA National Snow and Ice Data Center Distributed Active Archive Center [data set], Boulder, Colorado, USA, <https://doi.org/10.5067/INAWUWO7QH7B>, 2019.
- Wagner, D. N., Shupe, M. D., Cox, C., Persson, O. G., Uttal, T., Frey, M. M., Kirchaessner, A., Schneebeli, M., Jaggi, M., Macfarlane, A. R., Itkin, P., Arndt, S., Hendricks, S., Krampe, D., Nicolaus, M., Ricker, R., Regnery, J., Kolabutin, N., Shimanshuck, E., Oggier, M., Raphael, I., Stroeve, J., and Lehnig, M.: Snowfall and snow accumulation during the MOSAiC winter and spring seasons, *The Cryosphere*, 16, 2373–2402, <https://doi.org/10.5194/tc-16-2373-2022>, 2022.
- Wang, C., Graham, R. M., Wang, K., Gerland, S., and Granskog, M. A.: Comparison of ERA5 and ERA-Interim near-surface air temperature, snowfall and precipitation over Arctic sea ice: effects on sea ice thermodynamics and evolution, *The Cryosphere*, 13, 1661–1679, <https://doi.org/10.5194/tc-13-1661-2019>, 2019.
- Warren, S. G., Rigor, I. G., Untersteiner, N., Radionov, V. F., Bryazgin, N. N., Aleksandrov, Y. I., and Colony, R.: Snow Depth on Arctic Sea Ice, *J. Climate*, 12, 1814–1829, [https://doi.org/10.1175/1520-0442\(1999\)012<1814:SDOASI>2.0.CO;2](https://doi.org/10.1175/1520-0442(1999)012<1814:SDOASI>2.0.CO;2), 1999.
- Webster, M. A., Gerland, S., Holland, M., Hunke, E., Kwok, R., Lecomte, O., Massom, R., Perovich, D. K., and Sturm, M.: Snow in the Changing Sea-Ice Systems, *Nat. Clim. Change*, 8, 946–953, <https://doi.org/10.1038/s41558-018-0286-7>, 2018.
- Webster, M. A., Parker, C., Boisvert, L., and Kwok, R.: The role of cyclone activity in snow accumulation on Arctic sea ice, *Nat. Commun.*, 10, 1–12, <https://doi.org/10.1038/s41467-019-13299-8>, 2019.
- Webster, M. A., DuVivier, A. K., Holland, M. M., and Bailey, D. A.: Snow on Arctic Sea Ice in a Warming Climate as Simulated in CESM, *J. Geophys. Res.-Oceans*, 126, e2020JC016308, <https://doi.org/10.1029/2020JC016308>, 2021.
- Wehr, T., Kubota, T., Tzeremes, G., Wallace, K., Nakatsuka, H., Ohno, Y., Koopman, R., Rusli, S., Kikuchi, M., Eisinger, M., Tanaka, T., Taga, M., Deghaye, P., Tomita, E., and Bernaerts, D.: The EarthCARE mission – science and system overview, *Atmos. Meas. Tech.*, 16, 3581–3608, <https://doi.org/10.5194/amt-16-3581-2023>, 2023.
- Wood, N. B. and L'Ecuyer, T. S.: Level 2C Snow Profile Process Description and Interface Control Document, <https://www.cloudsat.cira.colostate.edu/data-products/2c-snow-profile> (last access: 1 February 2024), 2018.
- Wood, N. B., L'Ecuyer, T. S., Bliven, F. L., and Stephens, G. L.: Characterization of video disdrometer uncertainties and impacts on estimates of snowfall rate and radar reflectivity, *Atmos. Meas. Tech.*, 6, 3635–3648, <https://doi.org/10.5194/amt-6-3635-2013>, 2013.
- Wood, N. B., L'Ecuyer, T. S., Heymsfield, A. J., Stephens, G. L., Hudak, D. R., and Rodriguez, P.: Estimating Snow Microphysical Properties Using Collocated Multisensor Observations, *J. Geophys. Res.-Atmos.*, 119, 8941–8961, <https://doi.org/10.1002/2013JD021303>, 2014.
- Zhou, L., Stroeve, J., Xu, S., Petty, A., Tilling, R., Winstrup, M., Rostosky, P., Lawrence, I. R., Liston, G. E., Ridout, A., Tsamados, M., and Nandan, V.: Inter-comparison of snow depth over Arctic sea ice from reanalysis reconstructions and satellite retrieval, *The Cryosphere*, 15, 345–367, <https://doi.org/10.5194/tc-15-345-2021>, 2021.