



Southern Ocean sea-ice leads: first insights into regional lead patterns, seasonality, and trends, 2003–2023

Umesh Dubey, Sascha Willmes, and Günther Heinemann

Department of Environmental Meteorology, Trier University, Trier, Germany

Correspondence: Umesh Dubey (dubey@uni-trier.de) and Sascha Willmes (willmes@uni-trier.de)

Received: 17 February 2025 – Discussion started: 19 March 2025

Revised: 2 June 2025 – Accepted: 18 June 2025 – Published: 5 September 2025

Abstract. Sea-ice leads play a key role in the climate system by facilitating heat and moisture exchanges between the ocean and atmosphere, as well as by providing essential habitats for marine life. This study presents new insights from a gap-filled monthly dataset on sea-ice leads in the Southern Ocean and a first comprehensive analysis of spatial patterns, seasonal variability, and long-term trends of winter-time (April to September) sea-ice leads over a 21-year period (2003–2023). Our findings reveal that leads are ubiquitous in the Southern Ocean and show distinct spatial patterns with maximum lead frequencies close to the coastline, over the shelf break, and close to seafloor ridges and peaks. We see a strong seasonal variability in lead occurrence, with lead frequencies peaking in mid-winter. Weak but significant trends in lead frequencies are shown for the presented period for individual regions and months. Rather small changes in lead occurrence over the 21 years suggest stable winter-time sea-ice compactness despite the observed strong fluctuations and recent anomalies in sea-ice extent. Expanding upon previous work of lead detection in Antarctic sea ice, this study provides first results on the long-term regional, seasonal, and inter-annual variability of sea-ice leads in the Southern Ocean and can thereby contribute to an improved understanding of air–sea–ice–ocean interactions in the climate system. It also underscores the need for further investigation into the individual contributions of atmospheric and oceanic drivers to sea-ice lead formation in the Antarctic.

1 Introduction

The Southern Ocean is a key component of the climate system, and its sea-ice cover is characterized by a dynamic pattern of leads – narrow, elongated openings in the pack ice. Understanding the temporal variability and spatial distribution of leads is essential due to their role in facilitating significant exchanges of heat and moisture between the relatively warm ocean and cold winter atmosphere (Alam and Curry, 1995; Marcq and Weiss, 2012; Lüpkes et al., 2008; Heinemann et al., 2022). In addition to their impact on ocean–sea–ice–atmosphere interactions, leads are valuable diagnostic parameters for sea-ice drift (Spreen et al., 2017; Kwok et al., 2013) and provide habitats for marine mammals and birds (Stirling, 1997). In the context of the sea-ice–albedo feedback (Curry et al., 1995; Nihashi and Cavalieri, 2006), a larger fraction of leads can accelerate sea-ice thinning through an increased uptake of solar radiation, mainly during late winter. Ice-free areas including sea-ice leads have also been identified as sources of methane emissions (Kort et al., 2012; Damm et al., 2015).

Leads are not uniformly distributed in the sea ice. Patterns in their spatial distribution can be influenced by atmospheric forcing in the short term, such as that due to wind speed and wind divergence, and by ocean processes in the long term (Willmes et al., 2023). These leads, forming and evolving due to complex interactions between atmospheric and oceanic processes, represent an important component for understanding climatic variability and trends in polar regions (Wang et al., 2016; Zhang, 2014; Rheinlaender et al., 2022). Hence, studying lead dynamics is relevant for gaining better insight into the feedback mechanisms within the Southern Ocean climate system.

While distinct patterns in the spatial distribution of leads have recently been identified in both polar regions (Reiser et al., 2019; Willmes et al., 2023), the spatial and temporal variability and trends in wintertime sea-ice lead dynamics for individual regions in the Southern Ocean are yet to be determined. The potential of using thermal-infrared satellite data for the detection of sea-ice leads has been demonstrated for the Southern Ocean in a previous study (Reiser et al., 2020). Their lead detection relies on identifying significant positive ice-surface temperature anomalies from the Moderate Resolution Imaging Spectroradiometer (MODIS) on-board NASA's Aqua and Terra satellites and the associated ice-surface temperature dataset (MOD29/MYD29; Hall and Riggs, 2015), which includes the MODIS cloud mask to flag pixels affected by cloud cover. The method of Reiser et al. (2020) thereby detects the presence of leads in both hemispheres during the winter months (here: April to September) and also identifies false lead detections due to surface-temperature anomalies that result from deficiencies in the MODIS cloud mask. As a result, leads can be detected on a daily basis at a spatial resolution of 1 km²; however, due to high cloud fractions in the Antarctic as compared to the Arctic, the daily lead maps in the Southern Ocean suffer from extended data gaps that need to be dealt with when it comes to investigating the long-term regional lead patterns, seasonality, and trends.

The primary objectives of this study are therefore to deduce a continuous and gap-free monthly lead dataset from the daily lead data obtained from Reiser et al. (2020) and to use this dataset to infer the spatial distribution of sea-ice leads, quantify temporal trends, and compare regional differences for the months of April to September (hereafter referred to as “winter”) in the 21 years from 2003 to 2023. By addressing these objectives, this research contributes to a deeper understanding of the mechanisms driving sea-ice dynamics and the associated climatic implications. By enhancing our understanding of sea-ice leads in the Southern Ocean and their relationship with atmospheric and oceanic forcing, this work aims to contribute a more comprehensive picture of air–sea–ice–ocean interactions. The presented data can provide valuable contributions for improving sea ice/ocean components in climate models and forecasting systems, which are essential for predicting future changes in polar environments and their global implications.

The structure of the study is as follows. The data and methods used to derive the monthly lead dataset and to conduct a preliminary analysis are described in Sect. 2. Following this, in Sect. 3, we identify regions with the highest frequency of leads and show the predominant spatial and temporal lead patterns in wintertime sea ice. We examine trends and anomalies across various regions in the Southern Ocean and discuss our results in Sect. 4. Finally, we conclude our work in Sect. 5.

2 Data and methods

2.1 Sea-ice lead data

The detection of sea-ice leads from thermal-infrared satellite data is based on identifying significant positive spatial surface temperature anomalies. This method, as described by Reiser et al. (2020), classifies each pixel into one of four classes: sea ice, clouds, artefacts, and leads. We use these daily lead maps on a 1 km² resolution grid to calculate monthly lead frequencies (LFs) in the months of April to September for the years 2003 to 2023. The monthly aggregated maps are the main database for the presented analysis in this study. Daily sea-ice concentration from passive-microwave data from the University of Bremen (Spren et al., 2008) is used to define a mask by applying a sea-ice concentration threshold of 15 %, which is a common value to determine the sea-ice extent. For the months of April to June 2012, due to the gap in AMSR-E and AMSR2 data between October 2011 and June 2012, the mean sea-ice concentration as derived from the period with available AMSR data is used. It is important to note that our dataset focuses on small-scale leads and does not capture large-scale polynyas. Areas being classified as open water in the passive microwave sea-ice concentration data will not be detected as leads in our dataset. For studies on polynya areas, we suggest the inclusion of different datasets, e.g. Lin et al. (2024).

2.2 Retrieval of monthly lead frequency

The LF is a temporally integrated quantity that represents the number of days a pixel is covered by a lead during a specified period, relative to the number of available clear-sky (sea-ice or lead) observations (Willmes and Heinemann, 2016). This aggregation provides monthly, annual (winter), and total LF values for the Southern Ocean. The retrieval of monthly LFs involves several steps, starting with the daily lead maps, as shown in panel (a) of Fig. 1. As the daily lead maps are derived from satellite thermal-infrared data, valid lead detections are only available for clear-sky conditions. This means that cloud cover introduces gaps in the data, and the uncertainty in monthly LF calculations increases with the number of cloud gaps in the monthly stack. An example from July 2004 is shown here to demonstrate how daily data are influenced by cloud cover and the resulting data gaps.

In Fig. 1b, the raw monthly LF (LF_r) is calculated by averaging daily lead maps over a month. The number of data gaps in the monthly stack, represented by the data gap frequency (dgF), influences the confidence of the applied integration. Potential biases in LF_r calculations due to high dgF values need to be taken into consideration. This is addressed by applying an LF correction scheme together with an uncertainty measure for the monthly LF values. Therefore, we first derive basic statistical quantities of monthly LF_r, i.e. the long-term mean (mLF_r) and standard deviation (sdLF_r). Then, a cor-




Figure 1. Retrieval of monthly lead frequency (LF) from daily lead maps: (a) Daily lead maps for July 2004 with categories cloud, sea ice, lead, artefact, (b) raw monthly lead frequency (LF_r) and data gap frequency (dgF) derived from daily data, and (c) corrected monthly lead frequency (LF) and the potential integration error (pIE) derived from the overall mean LF_r (mLF_r), the standard deviation of LF_r (sdLF_r), and monthly dgF.

rection method is applied to obtain the monthly corrected LF (see Fig. 1c), which essentially represents a weighted smoothing of LF values based on the number of data gaps (dgF) per pixel (see Eqs. 1 and 2).

$$LF = LF_r - dgF \cdot (LF_r - mLF_r) \pm pIE \quad (1)$$

$$pIE = dgF \cdot sdLF_r \quad (2)$$

By this, we assume large LF_r deviations from the long-term mean to be potentially biased by data gaps and the associated error. The correction offset becomes larger with an increasing LF_r anomaly and higher dgF values and always pushes LF towards the climatological mean. This approach seeks to prevent over-interpretation of large deviations from the long-term mean, while enhancing confidence in the observed signals, i.e. LF anomalies.

We also define the absolute potential integration error (pIE) to measure the associated uncertainty in LF due to a combination of dgF and the local standard deviation of LF_r (sdLF_r) from the monthly climatology. A larger local sdLF_r and a higher dgF will, in this context, increase the representation error and thereby the uncertainty. The pIE represents a potential absolute LF error and will here be used to constrain an analysis of spatial–temporal LF patterns to pixels and months when the observed LF anomaly (LFA) was

larger than the associated pIE (hereafter referred to as the LFA > pIE criterion, with LFA < pIE indicating low-quality LF data). It thereby serves as a metric to exclude low-quality LF data in quantitative investigations (e.g. Sect. 3.2). The obtained monthly LF anomalies are independent from anomalies in cloud cover and the associated data gaps (see Appendix, Fig. A1), which is important to make sure that the obtained LF variability is not a function of available clear-sky data.

2.3 Auxiliary data

For a preliminary evaluation of confidence in the identified lead patterns and to assess the influence of ocean dynamics on lead formation, we use ocean surface current data provided by the European Centre for Medium-Range Weather Forecasts (ECMWF) Ocean Reanalysis System 5 (ORAS5) dataset (Zuo et al., 2019). Surface currents can play a crucial role in modulating lead formation by influencing sea-ice drift, mechanical stress, and ice deformation. ORAS5 provides global ocean reanalysis products with monthly surface current velocity data at a spatial resolution of 0.25°. These data include estimates of zonal and meridional current velocities, offering insights into large-scale ocean current velocity

patterns. For this study, we use ORAS5 surface current data with the climatological mean of LF to calculate the maximum coincident percentile exceedance (CPE) as a measure of spatial similarity. The CPE represents a simple measure of spatial coincidence by comparing the exceedance probability of two variables in a spatial field. This approach is inspired by Willmes et al. (2023), where it was used to analyse spatial patterns of leads in the Arctic Ocean. Here, CPE is used to identify areas where ocean currents modulate lead formation and variability.

The bathymetry data used in this study are derived from the International Bathymetric Chart of the Southern Ocean (IBCSO) version 2.0 (Dorschel et al., 2022), a digital bathymetric model for the area south of 50° S in the Southern Ocean. It has a horizontal resolution of 500 m. Similarly to the use of ORAS5 data, we use bathymetric gradients and mean LF to compute CPE and thereby highlight regions where bathymetry is obviously related to the formation of leads. The results of this comparison are shown in Sect. 3.8.

3 Results

3.1 Spatial patterns of sea-ice leads in the Southern Ocean

Figure 2 shows the long-term mean of the corrected LF (mean LF) over the 2003–2023 period, which essentially represents an updated and improved version of the lead climatology shown and discussed in Reiser et al. (2019). This visualization highlights the spatial patterns in lead frequency across the Southern Ocean. The LF value indicates how often, on average, a lead was present during wintertime at a specific position in the 21-year period.

The most prominent spatial lead patterns are found in the Weddell and Ross seas along the shelf break with values exceeding 0.25, which means that leads were present on more than 25 % of days during winter. Lead hotspots often follow linear patterns, with a prominent line of increased LF running close to the Antarctic coastline. In regions such as the Weddell Sea, Prydz Bay, and Ross Sea, further areas of concentrated lead activity are found along the continental shelf break and deep sea ridges, such as Maud Rise and Gunnerus Ridge. Close to Maud Rise we see what is generally referred to as a warm-water “Halo” around Maud Rise and represents a ring-like structure of high LF values at the location where the Maud Rise polynya frequently forms (note that larger polynyas are not included in this dataset). The presence of bathymetric features, such as ridges and troughs, align with these regions of increased LF, which confirms the findings by Reiser et al. (2019). This agreement suggests that the seafloor topography plays a crucial role in the formation and persistence of leads. The distinct band of increased LF at the coastline and at the edge of fast-ice regions (Fraser et al., 2021) likely reflects the interaction between ocean currents, wind

patterns, and ice dynamics, particularly in areas where the ice is thinner or more mobile (Wang et al., 2023). The associated overall mean potential integration error (mpIE) in Fig. 2b identifies areas that are more prone to data gaps and potential errors in the monthly LF integration. The mpIE apparently becomes generally larger towards the marginal ice zone (MIZ) and in areas with increased sdLF_r. It represents the potential error in the integrated monthly LF and does not exceed values of 0.07 in most regions of the inner sea-ice pack (south of the MIZ).

3.2 Inter-annual variability of leads

Figure 3 shows annual wintertime LF anomalies in the Southern Ocean from 2003 to 2023, with respect to the overall mean LF, from April to September, 2003–2023. Areas that do not meet the LFA > pIE criterion are highlighted in yellow (low-quality). In all of the presented years, a mixture of positive and negative anomalies is found, with their magnitude and regional differentiation changing from year to year. Also, the total area of low-quality data (LFA < pIE, yellow) is subject to inter-annual changes. The years 2021–2023 exhibit notably larger low-quality LF areas compared to the previous years. These years are obviously affected by more data gaps in a broader region south of the MIZ. No fixed pattern in LF anomalies can be found. It is possible, however, to identify some years with remarkable regional lead anomalies. For example, in 2006 there were substantially more leads in the Weddell Sea than in the previous and following years, with anomaly values partly exceeding 0.05, which means that leads were found here during roughly 10 more days of the 182 d period. Lead occurrence in the Ross and Amundsen seas was especially strong in 2004, 2012, 2013, and 2016. LF anomalies after 2020 seem to be generally larger than in previous years. This comes, however, with increased pIE and associated low-quality LF areas. There is no year with an overall positive or negative anomaly, but both occur in different regions of the Southern Ocean in each of the presented years.

3.3 Seasonal variability of leads

The seasonal evolution of the monthly mean LF anomalies is given in Fig. 4. These anomalies provide insight into the general seasonal variability of sea-ice leads in the Southern Ocean. In April, when new sea ice starts to consolidate, negative and close-to-zero anomalies indicate that leads are less abundant as compared to the total winter mean in most of the regions. In the southern Weddell Sea and in coastal regions leads seem to be more frequent in April on average. From May to June, leads also become more frequent in the Ross and Bellingshausen–Amundsen seas. In June, the maximum occurrence of leads south of the MIZ is observed, i.e. leads are more abundant in June than in any other month of the winter season. In mid-winter, July, and August, sea ice




Figure 2. (a) Overall mean of corrected lead frequency (mLF) and (b) overall mean absolute potential integration error (mpIE) for the months of April to September in the period from 2003 to 2023.

seems to consolidate, and leads form predominantly in the marginal ice zones, while in August LF is found to increase in the southwestern Ross Sea on average. In September, the overall lead activity seems to cease. However, in contrast, the coastal regions in the East Antarctic, with a pronounced positive anomaly north of the Dronning Maud and Enderby land, are characterized by an increased LF during this time of the year. The fraction of areas covered by low-quality data ($LFA < pIE$) increases throughout the season. In August, almost the entire Eastern Antarctic sea-ice region is affected, except the coastal regions. In September, confidence in anomalies is limited to the southern Weddell Sea (negative) and the pronounced positive anomaly north of Dronning Maud and Enderby land mentioned above. Overall, the seasonal cycle of lead occurrence in the Southern Ocean can be summarized as increasing from April to June and then decreasing again until September.

3.4 Seasonal and sectoral variability of leads

The time series plots in Fig. 5 illustrate the inter-annual variability of monthly LF in the Southern Ocean and its sub-regions, i.e. the Weddell Sea, Ross Sea, Indian Ocean, Pacific Ocean, and Bellingshausen–Amundsen seas from 2003 to 2023. Each plot represents monthly mean LF values (distinguished by different colours) for April to September. Data points without colour fill in the markers indicate months in which the valid LF data coverage (i.e. $LFA > pIE$) for the LF mean in the specific region is lower than 50 %, which means that the data quality is low.

A strong seasonal variability is confirmed here with low monthly mean LF in April (blue) and high LF in June (red).

The LF variability across all sectors suggests that significant anomalies occur primarily on a monthly basis. The time series reveals mostly increasing LF values over the 21-year period, mainly in the Southern Ocean, Weddell Sea, and Indian Ocean. It has to be noted, however, that a large part of this observed increase is attributed to an increase in low-quality data points during the last 3 years of the observed period, which does not allow us to infer a trend in LF. While low quality is not necessarily the cause for increased LF (low-quality data points are also found in months with low LF; see also Fig. A1), it still reduces confidence in a retrieval of linear trends from the presented time series. In the above mentioned sectors, a weak but obvious LF increase can also be determined when the low-quality data points are not taken into account (for a detailed trend analysis, see Sect. 3.6). Some periods stand out due to pronounced deviations of the monthly mean LF values. Most obvious is the two-year period of 2012 and 2013 with substantially higher monthly mean LF in the Ross Sea in all months. Also, June 2012 and May 2012 stand out with high LF in the Bellingshausen–Amundsen seas. In the Indian and Pacific Ocean sectors, low-quality data points are more frequent, which is a result of their rather narrow area and proximity to the MIZ with increased data gaps (high pIE).

3.5 Temporal evolution of lead frequency anomalies

The regional and temporal lead variability is given in more detail in latitude-averaged monthly LF anomalies, categorized into 5° longitude bins, covering the period from April 2003 to September 2023 as presented in Fig. 6. The analysis of meridional and zonal lead patterns between 60




Figure 3. Annual lead frequency anomalies of wintertime (April to September) lead frequency with respect to the 21-year (2003–2023) climatological mean. Areas that do not meet the lead frequency anomaly (LFA) > the potential integration error (PIE) criterion are highlighted in yellow.

and 90° S as shown in Fig. 6a provides a comprehensive view on how lead dynamics vary across longitudes in the Southern Ocean over time (the coastlines below Fig. 6a represent the corresponding positions along the longitudes). The most notable aspect is the obvious presence of “strong lead events” and “low lead events”, which are highlighted by pronounced positive anomalies (red areas) and negative anomalies (blue areas), respectively. Four of these events have been selected and are illustrated as monthly LF anomaly maps in Fig. 5b–e. Low-quality data (yellow) are mostly present in the Pacific

and Indian Sectors (0–135° E) and increase substantially after 2020, accompanied by an overall increase of LF. Some examples for the pronounced regional anomalies are depicted by arrows. The map for April 2006 (Fig. 6b) shows a significant positive anomaly, with values reaching up to 0.1 over the Weddell Sea at around 45° W. In June 2008 (Fig. 6c), a negative LF anomaly is observed along the coastal area in the Cosmonaut Sea near 50° E longitude in the Indian Ocean. The map for June 2012 (Fig. 6d) reveals a notable positive LF anomaly of more than 0.1 along the Dronning Maud coast of




Figure 4. Seasonal variability of monthly mean lead frequency anomalies from April to September, 2003–2023. These anomalies are calculated relative to the 21-year climatological mean. Areas that do not meet the $LFA > pIE$ criterion are highlighted in yellow.




Figure 5. Inter-annual variability of monthly mean lead frequency (April to September) in the Southern Ocean and its sub-regions from 2003 to 2023. Sub-regions are identified in the inset map. Individual months are colour-coded as follows: April (blue), May (green), June (red), July (orange), August (brown), and September (black). Month dots without colour fill indicate low-quality ($LFA < pIE$) data point coverage above 50 %. Note the different scales on the y axes for monthly LF.

the East Antarctic, particularly between 0 to 45° E longitude. A positive LF anomaly of less than 0.12 is evident in the Bellingshausen–Amundsen seas during May 2021 (Fig. 6e). These maps just exemplarily show the spatial patterns of lead anomalies on the monthly scale. As Fig. 6a reveals, more of these events can be identified for different regions throughout the 21-year period.

The variability of lead dynamics shown here indicates how monthly anomalies are superimposed to the previously described general seasonal, inter-annual, and long-term spatial patterns. Regional monthly anomalies provide a good basis to identify atmospheric drivers for sea-ice break-up and lead dynamics (see Sect. 4: Discussion).

3.6 Trend analysis of leads

Figure 7 shows the spatial variability of the overall LF change in the 21-year period as derived from the slope of a linear regression using annual LF values based on different data configurations. Figure 7a indicates the overall change in LF if low-quality data ($LFA < pIE$) are replaced by the climatological mean. Figure 7b shows the LF change but without replacing low-quality LF data. Figure 7c and d show the LF change for both alternatives but only for where the derived slope is significant at the 5 % level using a two-tailed test (note different colour scales). In both configurations, with filled mean and all data, the same spatial patterns can be identified, with mostly positive trends that amount to a total LF change of up to 0.04 (i.e. an increase of approximately 8 open lead days during winter in the period 2003 to 2023). A decreasing number of winter lead days is found only in the southern Weddell Sea, east and west of the Antarctic peninsula, in the central Ross Sea. If only significant trends are considered (Fig. 7c and d), the general spatial patterns with mostly positive trends in the majority of regions and decreasing LF in the southern Weddell and central Ross seas remain consistent. The total LF change for the 21-year period is, however, only about half as pronounced if only significant trends are considered.

3.7 Leads in coastal areas

As leads are especially abundant in coastal regions and at fast-ice edges (flaw-leads), we also want to focus on these areas to highlight coastal region lead dynamics based on the presented dataset. Figure 8 illustrates the spatial patterns of mean LF across various regions of the East and West Antarctic during the period from 2003 to 2023. Box 8 is an exception as it is not a coastal region but lies over the pronounced band of high LF over the Weddell Sea shelf break. The spatial LF patterns clearly reveal several sea-ice features that are typical for coastal regions. Particularly noticeable is the pronounced band of flaw-leads that is found right in front of the ice shelves, but also at the edge of fast-ice areas (Fraser et al., 2021), which become visible here through low LF values in

the used dataset (see for example the red boxes 2, 3, and 4 in Fig. 8a). The LF values in these regions generally correspond to areas of frequent fast-ice occurrence (see Fraser et al., 2021), with low LF often indicating the presence of stable fast ice. Fast-ice, flaw-leads, and polynyas can also be recognized in the West Antarctic (Fig. 8b). The presence of leads can affect the dynamics of fast ice by altering local wind and ocean currents, which can lead to changes in ice thickness and extent, and fast ice can also influence the formation of leads by providing a barrier that modifies the stress distribution in the surrounding ice cover, potentially leading to fractures and the opening of leads (Kim et al., 2018). Overall, the spatial patterns across the East and West Antarctic highlight pronounced regional differences in lead occurrence and the presence of fast-ice areas. Coastal polynyas, which are regions of persistent open water or thin ice, are mainly driven dynamically by wind systems, such as cold winds from the ice shelf or ice sheet (Paul et al., 2015), causing high heat energy fluxes compared to the surrounding pack ice (Marcq and Weiss, 2012). The continuous exchange of heat and moisture in these regions further influences local sea-ice dynamics, potentially maintaining higher lead frequencies. As a result, coastal areas generally exhibit higher LF values than the inner ice-pack, the shelf breaks, and some other bathymetry hot spots (e.g. Maud Rise Halo) excluded. The observational data summarized in Fig. 8 are useful to expand upon observations of long-term polynya areas, for example Lin et al. (2024).

Figures 9 and 10 show the seasonal and inter-annual variability of mean monthly LF in the sub-regions indicated by red boxes in Fig. 8 during the 2003–2023 winter seasons. The heat maps (left) highlight regional monthly changes in LF and indicate some pronounced lead events, e.g. June 2012 and 2013 in the Droning Maud region, low LF at the coast of Enderby land in April between 2005 and 2007, high LF in July 2018 at Terre Adélie land, and generally an overall increase in LF during September after 2020 in most of the regions. As in Fig. 5, this obvious increase is accompanied by a stronger presence of low-quality ($LFA < pIE$) data. The dots in the presented heat maps are representative of months in which the regional coverage of these low-quality LF data exceeded 50 % of the box area. The time series plots (middle) indicate the inter-annual LF variability for each sub-region and highlight again the pronounced increase in LF after 2020. This increase was only found in the Eastern Weddell Sea (boxes 1–4), while in the West Antarctic, no significant changes are notable in the 21-year period. The strong presence of low-quality data points ($LFA < pIE$ above 50 %) is here indicated by dots without colour fill. The mean seasonal changes in LF per region (right) offer a detailed overview of regional differences in the seasonality of lead dynamics. While the mean LF increases in boxes 1 and 2 (Dronning Maud and Enderby) from April to September, it shows differing evolutions in the other presented regions. Box 8 (WeddellSlope) stands out here, with a strong drop




Figure 6. (a) Latitude-averaged (60 to 90°S) monthly lead frequency (LF) anomalies in 5° longitude bins from April 2003 to September 2023. Areas that do not meet the LFA > pIE criterion are highlighted in yellow. (b–e) Monthly LF anomaly maps of selected years and months from panel (a) are shown. Strong lead anomalies can be seen, for example, in (b) April 2006, (d) June 2012, and (e) May 2021. Anomalies are calculated relative to the climatological monthly mean.

in average LF in July and low values also in August and September.

3.8 Spatial lead patterns in context with surface currents and ocean depth

Here, we aim to evaluate the spatial similarity in local maxima of the climatological mean LF (see Fig. 2a) to ocean surface currents and bathymetry gradients. We therefore use the CPE (see Sect. 2.3) as a metric of comparison, which represents the common maximum percentile that is exceeded in both datasets at a given grid point. High values in CPE indi-

cate areas where both datasets show high values coincidentally.

Figure 11a illustrates the spatial patterns of CPE between mean LF and mean ORAS5 surface current velocity in wintertime 2003–2023. The map highlights regions where CPE values exceed high percentiles in both datasets. High CPE values (of more than 80 %) are observed mainly in regions of the Antarctic Slope Current (ASC) and close to the MIZ, particularly along the Weddell Sea, Ross Sea, and Indian Ocean sectors. The highest values are concentrated near the Antarctic coastal zones, where the mean surface currents and LF are




Figure 7. Lead frequency (LF) change from 2003 to 2023 according to linear trend in annual LF values for (a) where low-quality LF data ($LFA < pIE$) are replaced by the overall mean LF per pixel, (b) all LF data including low-quality LF, (c) and (d) only for where the slope of the linear trend in (a) and (b), respectively, is significant at the 5 % level according to a two-tailed test.

most pronounced. The CPE map also highlights regions close to seafloor ridges and peaks (Maud Rise, Gunnerus Ridge, Iselin Bank, etc.), which are both characterized by high LF and strong surface currents at the same time. This is demonstrated even better in Fig. 11b, which shows the relationship between mean LF and bathymetric gradients derived from IBCSO version 2.0 data. High CPE values are predominantly observed in areas where significant changes in ocean depth occur, such as along the continental shelf break and near prominent underwater features, including seamounts, ridges, and troughs (see Fig. A2b). For example, the Weddell Sea, Ross Sea, and Prydz Bay regions display strong CPE values, underscoring the critical role of underwater topography in modulating the occurrence of sea-ice leads as discussed in Reiser et al. (2019) for the Antarctic and in Willmes et al. (2023) for the Arctic.

4 Discussion

The analysis of monthly LF anomalies from 2003 to 2023 reveals distinct temporal and spatial variations across the Southern Ocean. Figure 6 indicates that certain regions and time periods exhibit significant deviations in LF from the

climatological mean. These anomalies complement the seasonal variability of monthly mean lead frequency anomalies shown in Fig. 4 and can be used to get a better insight into the large-scale drivers of sea-ice break-up and lead dynamics (Kimura and Wakatsuchi, 2011). We hypothesize that the presented monthly anomalies in regional lead frequencies are driven by a combination of large-scale atmospheric and oceanic processes. However, the relative importance of those two components is yet to be estimated. In this study, we therefore present first insights into this new lead dataset to describe the spatial and temporal variability of the Southern Ocean lead dynamics.

Our findings support the argument that sea-ice leads are widely distributed across the Southern Ocean, with particularly high frequencies of occurrence along the coastline, over the continental shelf break and ridges (see Fig. 2a), which underlines the findings of Reiser et al. (2019) and extends upon their results. In many of these regions, especially near the coast, these patterns are caused by flaw-leads and polynyas, which are opening mainly due to wind-driven processes and thereby promote new ice formation and the production of cold and dense shelf water, a precursor to Antarctic bottom water, which represents a key element of the global cli-




Figure 8. Mean LF for wintertime period 2003–2023 in coastal regions for (a) the East Antarctic and (b) the West Antarctic. Red boxes indicate sub-regions used for the regional analysis of individual inter-annual and seasonal lead variabilities in Figs. 9 and 10.

mate system (Goosse and Fichefet, 1999; Jacobs and Weiss, 1998; Silvano, 2018). Therefore, knowledge about the seasonal, inter-annual, and spatial variability of lead dynamics and their drivers is one of the crucial fields to put forward our understanding of global ocean circulation and climate (Bindoff, 2000; Jacobs, 2004). The presented dataset does not include the presence of polynyas, which were large enough to cause a signal passive microwave data. However, it can be very well combined with long-term polynya observations as given, for example, in Lin et al. (2024).

The occurrence and variability of polynyas in the Southern Ocean and the consequent new ice production and salt release were investigated in many previous studies (e.g. Golledge et al., 2025; Macdonald et al., 2023; Tamura et al., 2008; Tamura et al., 2016). However, these investigations mostly did not take small-scale leads within the pack ice into account, which is mainly because satellite data were used that did not allow the resolution of sea-ice surface features at the km scale or below. The data we present here show that leads are ubiquitous in the Southern Ocean pack ice during wintertime. This supports the need for more detailed information about sea-ice dynamics within the pack ice to improve our understanding of ocean–sea-ice–atmosphere interactions and climate prediction in the future (Mohrmann et al., 2021).

The continental slope plays a key role in shaping spatial lead patterns. One of the possible explanations for this feature is ice divergence in these regions, driven by the ASC and tidal flows (Stewart et al., 2019; Heil et al., 2008; Hutchings et al., 2012). Also, regions such as Maud Rise (Beckmann et al., 2001) or Gunnerus Ridge exhibit enhanced tidal currents, which can increase sea-ice velocity and divergence. In addition to tidal flows, the thermal variability at Maud Rise is significantly influenced by advection and mesoscale eddies (Gülk et al., 2023). These processes underscore the interplay between bathymetry, ocean circulation, and lead formation in this region. Our observations provide evidence that this also facilitates lead formation in these regions and over seafloor ridges in general. The opening of leads during winter triggers strong energy exchange processes that can alter both the stratification of the upper ocean (Venables and Meredith, 2014; Cohan, 2021) and the atmospheric boundary layer (Heinemann et al., 2022; Marcq and Weiss, 2012). In this context, the role of leads in shaping the feedback between sea ice, atmosphere, and ocean must not be neglected and requires a robust identification of forcings, for which we think the presented lead climatology will yield a useful basis.

In light of the observed trends (e.g. Schroeter, 2023; Eayrs et al., 2021) and recent changes in Antarctic sea ice (Suryawanshi et al., 2023; Turner et al., 2016), a detailed




Figure 9. Mean monthly LF per month and year (left) with open dots indicating months with an LFA < pIE pixel coverage above 50 %, times series of monthly LF with blue open dots indicating months with an LFA < pIE pixel coverage > 50 % (middle), and mean seasonal LF change (right) with its standard deviations (vertical lines) for each month from April to September, 2003–2023. Sub-figures (a)–(d) indicate sub-regions of the East Antarctic (red boxes) shown in Fig. 8a.

look at what has happened within the pack ice has become increasingly important and has made sea-ice lead dynamics receive increasing attention. Our results indicate weak but significant trends in wintertime lead frequencies in the period between 2003 and 2023, which are mostly positive (Figs. 5 and 7). We do not find a signal in the presented lead climatology that can be associated with the observed negative sea-ice extent anomalies after 2015 (Zhang et al., 2022). Likewise, the observed minimum Antarctic sea-ice extent in July 2023 (Ionita, 2024) is not accompanied by more leads in the remaining pack ice. This demonstrates that sea-ice compactness during winter, at least south of the MIZ, seems not to be affected by the processes responsible for the overall sea-ice decline in the Southern Ocean since 2016. Although the detected LF changes in the 21-year period are small, Fig. 7 exhibits some spatial and seasonal contrasts in the sign of the

slope, which might point to a weak, but spatially distinguishable contrast in lead changes over the last 21 years. The LF trends presented here exclude the contribution of low-quality data (LFA < pIE) to provide high confidence for the observed change signal. If the significant increase in LF in certain regions – linked to larger areas of low-quality data after 2020 – is indeed real, then substantial changes in lead dynamics may have occurred since. For this to be confirmed, however, a more in-depth analysis with additional datasets would be necessary. LF after 2020 tends to be larger compared to previous years (Fig. 6a) in almost all of the observed regions. This observation coincides, however, with higher pIE values and the associated decrease in confidence of monthly LF values, due to an increase in data gaps. While a direct analysis of detailed cloud cover changes is beyond the scope of this paper, it is important to acknowledge the impact of clouds




Figure 10. Same as in Fig. 9 but for sub-regions of the West Antarctic shown in Fig. 8b.

on lead detection using thermal-infrared satellite data. For example, Goessling et al. (2025) examine the trend in ERA5 low cloud cover from 2013 to 2022 and identify an increasing coverage of low clouds in the Weddell Sea in the last couple of years. Detected changes in monthly LF may also partly be attributed to changes in shelf ice edges and associated local modifications in the presence of open water areas that have occurred since 2002. The lead maps used in this study are based on the land mask that is provided with the MODIS ice surface temperature product (see Reiser et al., 2020).

For the pan-Antarctic sea-ice extent as a broader quantity, the connections to atmospheric and oceanic drivers were demonstrated in a wide range of published research papers. For example, changes in atmospheric circulation patterns, such as the Southern Annular Mode (SAM) and the El Niño–Southern Oscillation (ENSO), significantly influence the extent of Antarctic sea ice as well as the position of the MIZ (Yuan, 2004; Hall and Visbeck, 2002; Maksym, 2012). These modes affect wind strength and di-

rection, which in turn impact sea-ice dynamics and presumably also the opening/closing of leads. In addition, sea surface temperature (SST) anomalies in subtropical regions, such as the Indian Ocean Dipole (IOD), show a clear impact on the compactness, drift patterns, and trends of sea ice in the Southern Ocean (Blanchard-Wrigglesworth et al., 2021). SST anomalies propagate through atmospheric circulation patterns, which influence sea-ice transport and temperature advection (Yu, 2023). As far as trends are concerned, the atmosphere is generally considered the primary driver, while the ocean is important for the seasonality of the trend patterns (Holland and Kwok, 2012; Hobbs et al., 2016). However, in general, previous investigations of large-scale modes and their impact on sea ice are mostly constrained to sea-ice extent or low-resolution sea-ice concentration. Therefore, the dataset we present here offers a new opportunity and supplement to existing datasets to conduct thorough investigations of the relationship between high-resolution sea-ice compactness in the Southern Ocean and its large-scale drivers.




Figure 11. The maximum coincident percentile exceedance (CPE) for overall mean LF and (a) ORAS5 surface current velocity (m s^{-1}) and (b) IBCSO2 bathymetry gradient (m). Values indicate the maximum percentile per pixel that is exceeded coincidentally in both datasets.

5 Conclusions

In this study, we present a new dataset on the spatial distribution and temporal variability of wintertime sea-ice leads in the Southern Ocean for the months of April to September for the 21-year period from 2003 to 2023. The data include monthly lead frequencies and their potential integration error on a per-pixel basis at 1 km^2 spatial resolution. Monthly and annual mean lead frequencies, along with their anomalies and trends are derived to provide first insights into long-term lead dynamics. The results reveal pronounced spatial patterns in the long-term sea-ice lead climatology, with higher lead frequencies observed along the Antarctic coastline, particularly in the Weddell and Ross seas, over continental shelf breaks and bathymetric features, such as ridges. This provides strong evidence for a significant role of ocean currents in shaping lead dynamics in the Southern Ocean. On the monthly scale, we can identify a pronounced, but regionally different, seasonal cycle in lead dynamics and the presence of strong monthly anomalies in different regions across the Southern Ocean. We suggest these anomalies to be potentially attributed to atmospheric forcing. We find small but significant long-term trends in lead frequencies in the Southern Ocean with regionally differing signs of change. The fact that no strong trend is found indicates relative stability in wintertime sea-ice compactness in areas south of the MIZ, while the overall sea-ice extent in the Antarctic has shown substantial variability and recent declines, particularly post-2016. This work provides initial results for future investigations into the complex pan-Antarctic interplay between sea ice, ocean, and atmosphere. The presented data can help to better estimate and untangle the individual contribution of atmospheric and oceanic forcings on sea-ice lead dynamics in the Southern Ocean.

Appendix A

A1 LF anomaly variability with respect to dgF

Figure A1 presents the distribution of LF anomalies as a function of the coincident dgF anomaly class. The figure highlights the independence of the LF variability from dgF anomalies.




Figure A1. Range of LF anomalies with respect to corrected monthly mean LF as a function of the coincident dgF anomaly: median (circles), 1 (blue) and 2 (grey) standard deviations.

A2 ORAS5 current velocity and ocean depth

Figure A2a represents the mean surface current velocity for the Southern Ocean during the winter months (April–September), 2003–2023. The highest current velocities are observed near the Antarctic Circumpolar Current and along the continental shelf. Figure A2b shows bathymetry data from the IBCSO version 2 dataset. The alignment of high LF occurrences with these features emphasizes the critical role of underwater topography in shaping spatial LF patterns (see Sect. 3.8).




Figure A2. (a) Overall mean of ORAS5 surface current velocity for 2003–2023 (April to September) and (b) IBCSO2 ocean depth (Dorschel et al., 2022).

Data availability. Monthly lead frequencies (LFs) and potential integration error (pIE) data for the period April 2003 to September 2023 are available in NetCDF format on PANGAEA (AntLeads: Monthly wintertime sea-ice lead maps for the Antarctic, April–September, 2003–2023, <https://doi.org/10.1594/PANGAEA.977634>) (Dubey et al., 2025). ORAS5 data (Zuo et al., 2019) were downloaded from the Copernicus Climate Data Store (<https://cds.climate.copernicus.eu/>, last access: 13 December 2024), and version 2.0 of the International Bathymetric Chart of the Southern Ocean (IBCSO, Dorschel et al., 2022) was acquired from the General Bathymetric Chart of the Ocean (GEBCO) gridded bathymetric datasets (<https://gebcoset.net>, last access: 13 December 2024; Dorschel et al., 2022).

Author contributions. UD and SW processed and analysed the data and drafted the main script. The final version was prepared with contributions from all co-authors including GH. All authors have read and agreed to the submitted version of the manuscript.

Competing interests. The contact author has declared that none of the authors has any competing interests.

Disclaimer. Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes every effort to include appropriate place names, the final responsibility lies with the authors.

Acknowledgements. All processing was done in Python3. The authors want to thank the ECMWF and GEBCO for providing the data used in this study.

Financial support. The research was funded by the Deutsche Forschungsgemeinschaft (DFG) in the framework of the priority programme “Antarctic Research with comparative investigations in Arctic ice areas” (SPP1158) under grant WI 3314/6-1. The publication was funded by the Open Access Fund of Trier University and the German Research Foundation (DFG).

Review statement. This paper was edited by Lars Kaleschke and reviewed by Alexander Fraser and one anonymous referee.

References

Alam, A. and Curry, J.: Lead-induced atmospheric circulations, *J. Geophys. Res.*, 100, 4643, <https://doi.org/10.1029/94jc02562>, 1995.

Beckmann, A., Timmermann, R., and Pereira, A.: The effect of flow at Maud Rise on the sea-ice cover numerical experiments, *Ocean*

Dynam., 52, 11–25, <https://doi.org/10.1007/s10236-001-8173-5>, 2001.

Bindoff, N.: Bottom Water Formation and Polynyas in Adélie Land, Antarctica, University of Tasmania, <https://doi.org/10.26749/RSTPP.133.3.51>, 2000.

Blanchard-Wrigglesworth, E., Roach, L., Donohoe, A., and Ding, Q.: Impact of winds and Southern Ocean SSTs on Antarctic sea ice trends and variability, *J. Climate*, 34, 949–965, <https://doi.org/10.1175/JCLI-D-20-0386.1>, 2021.

Cohanin, K.: Dynamics of Eddies Generated by Sea Ice Leads, *J. Phys. Oceanogr.*, 51, 3071–3092, <https://doi.org/10.1175/JPO-D-20-0169.1>, 2021.

Curry, J. A., Schramm, J. L., and Ebert, E. E.: Sea Ice-Albedo Climate Feedback Mechanism, *J. Climate*, 8, 240–247, [https://doi.org/10.1175/1520-0442\(1995\)008<0240:siacfm>2.0.co;2](https://doi.org/10.1175/1520-0442(1995)008<0240:siacfm>2.0.co;2), 1995.

Damm, E., Rudels, B., Schauer, U., Mau, S., and Dieckmann, G.: Methane excess in Arctic surface water triggered by sea ice formation and melting, *Scientific Reports*, 5, 16179, <https://doi.org/10.1038/srep16179>, 2015.

Dorschel, B., Hehemann, L., Viquerat, S., Warnke, F., Dreutter, S., Tenberge, Y. S., Accettella, D., An, L., Barrios, F., Bazhenova, E., Black, J., Bohoyo, F., Davey, C., De Santis, L., Dotti, C. E., Fremand, A. C., Fretwell, P. T., Gales, J. A., Gao, J., Gasperini, L., Greenbaum, J. S., Jencks, J. H., Hogan, K., Hong, J. K., Jakobsson, M., Jensen, L., Kool, J., Larin, S., Larter, R. D., Leitchenkov, G., Loubrieu, B., Mackay, K., Mayer, L., Millan, R., Morlighem, M., Navidad, F., Nitsche, F. O., Nogi, Y., Pertuisot, C., Post, A. L., Pritchard, H. D., Purser, A., Rebesco, M., Rignot, E., Roberts, J. L., Rovere, M., Ryzhov, I., Sauli, C., Schmitt, T., Silvano, A., Smith, J., Snaith, H., Tate, A. J., Tinto, K., Vandenbossche, P., Weatherall, P., Wintersteller, P., Yang, C., Zhang, T., and Arndt, J. E.: The International Bathymetric Chart of the Southern Ocean Version 2, *Scientific Data*, 9, 275, <https://doi.org/10.1038/s41597-022-01366-7>, 2022.

Dubey, U., Willmes, S., and Heinemann, G.: AntLeads: Monthly wintertime sea-ice lead maps for the Antarctic, April–September, 2003–2023, PANGAEA [data set], <https://doi.org/10.1594/PANGAEA.977634>, 2025.

Eayrs, C., Li, X., Raphael, M., and Holland, D.: Rapid decline in Antarctic sea ice in recent years hints at future change, *Nat. Geosci.*, 14, 460–464, <https://doi.org/10.1038/s41561-021-00768-3>, 2021.

Fraser, A. D., Massom, R. A., Handcock, M. S., Reid, P., Ohshima, K. I., Raphael, M. N., Cartwright, J., Klekociuk, A. R., Wang, Z., and Porter-Smith, R.: Eighteen-year record of circum-Antarctic landfast-sea-ice distribution allows detailed baseline characterisation and reveals trends and variability, *The Cryosphere*, 15, 5061–5077, <https://doi.org/10.5194/tc-15-5061-2021>, 2021.

Goessling, H. F., Rackow, T., and Jung, T.: Recent global temperature surge intensified by record-low planetary albedo, *Science*, 387, 68–73, <https://doi.org/10.1126/science.adq7280>, 2025.

Golledge, N. R., Keller, E. D., Gossart, A., Malyarenko, A., Bahamondes-Dominguez, A., Krapp, M., Jendersie, S., Lowry, D. P., Alevropoulos-Borrill, A., and Notz, D.: Antarctic coastal polynyas in the global climate system, *Nature Reviews Earth & Environment*, 6, 126–139, <https://doi.org/10.1038/s43017-024-00634-x>, 2025.

- Goosse, H. and Fichefet, T.: Importance of ice-ocean interactions for the global ocean circulation: A model study, *J. Geophys. Res.*, 104, 23337–23355, <https://doi.org/10.1029/1999JC900215>, 1999.
- Gülk, B., Roquet, F., Naveira Garabato, A. C., Narayanan, A., Rousset, C., and Madec, G.: Variability and Remote Controls of the Warm-Water Halo and Taylor Cap at Maud Rise, *J. Geophys. Res.-Oceans*, 128, e2022JC019517, <https://doi.org/10.1029/2022jc019517>, 2023.
- Hall, A. and Visbeck, M.: Synchronous Variability in the Southern Hemisphere Atmosphere, Sea Ice, and Ocean Resulting from the Annular Mode, *J. Climate*, 15, 3043–3057, [https://doi.org/10.1175/1520-0442\(2002\)015<3043:SVITSH>2.0.CO;2](https://doi.org/10.1175/1520-0442(2002)015<3043:SVITSH>2.0.CO;2), 2002.
- Hall, D. and Riggs, G.: MODIS/Terra Sea Ice Extent 5-Min L2 Swath 1km, Version 6, NASA National Snow and Ice Data Center Distributed Active Archive Center [data set], <https://doi.org/10.5067/MODIS/MOD29.006>, 2015.
- Heil, P., Hutchings, J. K., Worby, A. P., Johansson, M., Lau-niainen, J., Haas, C., and Hibler, W. D.: Tidal forcing on sea-ice drift and deformation in the western Weddell Sea in early austral summer, 2004, *Deep-Sea Res. Pt. II*, 55, 943–962, <https://doi.org/10.1016/j.dsr2.2007.12.026>, 2008.
- Heinemann, G., Schefczyk, L., Willmes, S., and Shupe, M.: Evaluation of simulations of near-surface variables using the regional climate model CCLM for the MOSAiC winter period, *Elementa*, 10, 33, <https://doi.org/10.1525/elementa.2022.00033>, 2022.
- Hobbs, W., Massom, R., Stammerjohn, S., Reid, P., Williams, G., and Meier, W.: A review of recent changes in Southern Ocean sea ice, their drivers and forcings, *Global Planet. Change*, 143, 228–250, <https://doi.org/10.1016/j.gloplacha.2016.06.008>, 2016.
- Holland, P. and Kwok, R.: Wind-driven trends in Antarctic sea-ice drift, *Nat. Geosci.*, 5, 872–875, <https://doi.org/10.1038/ngeo1627>, 2012.
- Hutchings, J., Heil, P., Steer, A., and Hibler, W.: Subsynoptic scale spatial variability of sea ice deformation in the western Weddell Sea during early summer, *J. Geophys. Res.*, 117, C01002, <https://doi.org/10.1029/2011jc006961>, 2012.
- Ionita, M.: Large-scale drivers of the exceptionally low winter Antarctic sea ice extent in 2023, *Front. Earth Sci.*, 12, 1333706, <https://doi.org/10.3389/feart.2024.1333706>, 2024.
- Jacobs, S.: Bottom water production and its links with the thermohaline circulation, *Antarct. Sci.*, 16, 427–437, <https://doi.org/10.1017/S095410200400224X>, 2004.
- Jacobs, S. and Weiss, R.: Ocean, Ice, and Atmosphere: Interactions at the Antarctic Continental Margin, *Arct. Antarct. Alpine Res.*, 31, 329–332, <https://doi.org/10.2307/1552267>, 1998.
- Kim, S., Saenz, B., Scanniello, J., Daly, K., and Ainley, D.: Local climatology of fast ice in McMurdo Sound, Antarctica, *Antarct. Sci.*, 30, 125–142, <https://doi.org/10.1017/s0954102017000578>, 2018.
- Kimura, N. and Wakatsuchi, M.: Large-scale processes governing the seasonal variability of the Antarctic sea ice, *Tellus A*, 63, 828–840, <https://doi.org/10.1111/j.1600-0870.2011.00526.x>, 2011.
- Kort, E., Wofsy, S., Daube, B., Diao, M., Elkins, J., Gao, R., Hints, E., Hurst, D., Jimenez, R., Moore, F., Spackman, J., and Zondlo, M.: Atmospheric observations of Arctic Ocean methane emissions up to 82 degrees north, *Nat. Geosci.*, 5, 318–321, <https://doi.org/10.1038/ngeo1452>, 2012.
- Kwok, R., Spreen, G., and Pang, S.: Arctic sea ice circulation and drift speed: Decadal trends and ocean currents, *J. Geophys. Res.-Oceans*, 118, 2408–2425, <https://doi.org/10.1002/jgrc.20191>, 2013.
- Lin, Y., Nakayama, Y., Liang, K., Huang, Y., Chen, D., and Yang, Q.: A dataset of the daily edge of each polynya in the Antarctic, *Scientific Data*, 11, 1006, <https://doi.org/10.1038/s41597-024-03848-2>, 2024.
- Lüpkes, C., Vihma, T., Birnbaum, G., and Wacker, U.: Influence of leads in sea ice on the temperature of the atmospheric boundary layer during polar night, *Geophys. Res. Lett.*, 35, L03805, <https://doi.org/10.1029/2007gl032461>, 2008.
- Macdonald, G. J., Ackley, S. F., Mestas-Núñez, A. M., and Blanco-Cabanillas, A.: Evolution of the dynamics, area, and ice production of the Amundsen Sea Polynya, Antarctica, 2016–2021, *The Cryosphere*, 17, 457–476, <https://doi.org/10.5194/tc-17-457-2023>, 2023.
- Maksym, T.: Antarctic sea ice-A polar opposite?, *Oceanography*, 25, 140–151, <https://doi.org/10.5670/OCEANOGRAPHY.2012.88>, 2012.
- Marcq, S. and Weiss, J.: Influence of sea ice lead-width distribution on turbulent heat transfer between the ocean and the atmosphere, *The Cryosphere*, 6, 143–156, <https://doi.org/10.5194/tc-6-143-2012>, 2012.
- Mohrmann, M., Heuzé, C., and Swart, S.: Southern Ocean polynyas in CMIP6 models, *The Cryosphere*, 15, 4281–4313, <https://doi.org/10.5194/tc-15-4281-2021>, 2021.
- Nihashi, S. and Cavalieri, D. J.: Observational evidence of a hemispheric-wide ice-ocean albedo feedback effect on Antarctic sea-ice decay, *J. Geophys. Res.-Oceans*, 111, C12001, <https://doi.org/10.1029/2005jc003447>, 2006.
- Paul, S., Willmes, S., and Heinemann, G.: Long-term coastal-polynya dynamics in the southern Weddell Sea from MODIS thermal-infrared imagery, *The Cryosphere*, 9, 2027–2041, <https://doi.org/10.5194/tc-9-2027-2015>, 2015.
- Reiser, F., Willmes, S., Hausmann, U., and Heinemann, G.: Predominant Sea Ice Fracture Zones Around Antarctica and Their Relation to Bathymetric Features, *Geophys. Res. Lett.*, 46, 12117–12124, <https://doi.org/10.1029/2019GL084624>, 2019.
- Reiser, F., Willmes, S., and Heinemann, G.: A New Algorithm for Daily Sea Ice Lead Identification in the Arctic and Antarctic Winter from Thermal-Infrared Satellite Imagery, *Remote Sensing*, 12, 1957, <https://doi.org/10.3390/rs12121957>, 2020.
- Rheinlaender, J., Davy, R., Ólason, E., Rampal, P., Spensberger, C., Williams, T., Korosov, A., and Spengler, T.: Driving Mechanisms of an Extreme Winter Sea Ice Breakup Event in the Beaufort Sea, *Geophys. Res. Lett.*, 49, e2022GL099024, <https://doi.org/10.1029/2022GL099024>, 2022.
- Schroeter, S., O’Kane, T. J., and Sandery, P. A.: Antarctic sea ice regime shift associated with decreasing zonal symmetry in the Southern Annular Mode, *The Cryosphere*, 17, 701–717, <https://doi.org/10.5194/tc-17-701-2023>, 2023.
- Silvano, A.: Freshening by glacial meltwater enhances melting of ice shelves and reduces formation of Antarctic Bottom Water, *Science Advances*, 4, <https://doi.org/10.1126/sciadv.aap9467>, 2018.

- Spreen, G., Kaleschke, L., and Heygster, G.: Sea ice remote sensing using AMSR-E 89-GHz channels, *J. Geophys. Res.-Oceans*, 113, C02S03, <https://doi.org/10.1029/2005JC003384>, 2008.
- Spreen, G., Kwok, R., Menemenlis, D., and Nguyen, A. T.: Sea-ice deformation in a coupled ocean–sea-ice model and in satellite remote sensing data, *The Cryosphere*, 11, 1553–1573, <https://doi.org/10.5194/tc-11-1553-2017>, 2017.
- Stewart, A. L., Klocker, A., and Menemenlis, D.: Acceleration and Overturning of the Antarctic Slope Current by Winds, Eddies, and Tides, *J. Phys. Oceanogr.*, 49, 2043–2074, <https://doi.org/10.1175/jpo-d-18-0221.1>, 2019.
- Stirling, I.: The importance of polynyas, ice edges, and leads to marine mammals and birds, *J. Marine Syst.*, 10, 9–21, [https://doi.org/10.1016/S0924-7963\(96\)00054-1](https://doi.org/10.1016/S0924-7963(96)00054-1), 1997.
- Suryawanshi, K., Jena, B., Bajish, C., and Anilkumar, N.: Recent Decline in Antarctic Sea Ice Cover From 2016 to 2022: Insights From Satellite Observations, Argo Floats, and Model Reanalysis, *Tellus A*, 75, 193–212, <https://doi.org/10.16993/tellusa.3222>, 2023.
- Tamura, T., Ohshima, K. I., and Nihashi, S.: Mapping of sea ice production for Antarctic coastal polynyas, *Geophys. Res. Lett.*, 35, L07606, <https://doi.org/10.1029/2007gl032903>, 2008.
- Tamura, T., Ohshima, K., Fraser, A., and Williams, G.: Sea ice production variability in Antarctic coastal polynyas, *J. Geophys. Res.-Oceans*, 121, 2967–2979, <https://doi.org/10.1002/2015JC011537>, 2016.
- Turner, J., Hosking, J. S., Marshall, G. J., Phillips, T., and Bracegirdle, T. J.: Antarctic sea ice increase consistent with intrinsic variability of the Amundsen Sea Low, *Clim. Dynam.*, 46, 2391–2402, <https://doi.org/10.1007/s00382-015-2708-9>, 2016.
- Venables, H. and Meredith, M.: Feedbacks between ice cover, ocean stratification, and heat content in Ryder Bay, western Antarctic Peninsula, *J. Geophys. Res.*, 119, 5323–5336, <https://doi.org/10.1002/2013JC009669>, 2014.
- Wang, Q., Danilov, S., Jung, T., Kaleschke, L., and Wernecke, A.: Sea ice leads in the Arctic Ocean: Model assessment, interannual variability and trends, *Geophys. Res. Lett.*, 43, 7019–7027, <https://doi.org/10.1002/2016GL068696>, 2016.
- Wang, Y., Ji, Q., Pang, X., Qu, M., Cha, M., Zhang, F., Yan, Z., and He, B.: Distribution Characteristics and Influencing Factors of Sea Ice Leads in the Weddell Sea, Antarctica, *Remote Sensing*, 15, 5568, <https://doi.org/10.3390/rs15235568>, 2023.
- Willmes, S. and Heinemann, G.: Sea-ice wintertime lead frequencies and regional characteristics in the Arctic, *Remote Sensing*, 8, 4, <https://doi.org/10.3390/rs8010004>, 2016.
- Willmes, S., Heinemann, G., and Schnaase, F.: Patterns of wintertime Arctic sea-ice leads and their relation to winds and ocean currents, *The Cryosphere*, 17, 3291–3308, <https://doi.org/10.5194/tc-17-3291-2023>, 2023.
- Yu, L.: Antarctic sea ice variability linked to three dipole modes of subtropical sea surface temperature in the Southern Hemisphere, *Int. J. Climatol.*, 43, 6131–6149, <https://doi.org/10.1002/joc.8195>, 2023.
- Yuan, X.: ENSO-related impacts on Antarctic sea ice: a synthesis of phenomenon and mechanisms, *Antarct. Sci.*, 16, 415–425, <https://doi.org/10.1017/S0954102004002238>, 2004.
- Zhang, J.: Modeling the Impact of Wind Intensification on Antarctic Sea Ice Volume, *J. Climate*, 27, 202–214, <https://doi.org/10.1175/JCLI-D-12-00139.1>, 2014.
- Zhang, L., Delworth, T., Yang, X., Zeng, F., Lu, F., Morioka, Y., and Bushuk, M.: The relative role of the subsurface Southern Ocean in driving negative Antarctic Sea ice extent anomalies in 2016–2021, *Communications Earth & Environment*, 3, 302, <https://doi.org/10.1038/s43247-022-00624-1>, 2022.
- Zuo, H., Balmaseda, M. A., Tietsche, S., Mogensen, K., and Mayer, M.: The ECMWF operational ensemble reanalysis–analysis system for ocean and sea ice: a description of the system and assessment, *Ocean Sci.*, 15, 779–808, <https://doi.org/10.5194/os-15-779-2019>, 2019.