

Abstract

Fluctuations in terrestrial surface freshwater flux to Sermilik Fjord, SE Greenland, were simulated and analyzed. SnowModel, a state-of-the-art snow-evolution, snow and ice melt, and runoff modeling system, was used to simulate the temporal and spatial terrestrial runoff distribution to the fjord based on observed meteorological data (1999–2008) from stations located on and around the Greenland Ice Sheet (GrIS). Simulated runoff was compared and verified against independent glacier catchment runoff observations (1999–2005). Modeled runoff to Sermilik Fjord was highly variable, ranging from $2.9 \times 10^9 \text{ m}^3 \text{ y}^{-1}$ in 1999 to $5.9 \times 10^9 \text{ m}^3 \text{ y}^{-1}$ in 2005. The uneven spatial runoff distribution produced an areally-averaged annual maximum runoff at the Helheim glacier terminus of more than 3.8 m w.e. The sub-catchment runoff of the Helheim glacier region accounted for 25% of the total runoff to Sermilik Fjord. The runoff distribution from the different sub-catchments suggested a strong influence from the spatial variation in glacier coverage. To assess the Sermilik Fjord freshwater flux, simulated terrestrial runoff and net precipitation (precipitation minus evaporation and sublimation) for the fjord area were combined with satellite-derived ice discharge and subglacial geothermal and frictional melting due to basal ice motion. A terrestrial freshwater flux of $\sim 40.4 \times 10^9 \text{ m}^3 \text{ y}^{-1}$ was found for Sermilik Fjord, with an 11% contribution originated from surface runoff. For the Helheim glacier sub-catchment only 4% of the flux originated from terrestrial surface runoff.

1 Introduction

Global atmospheric temperatures showed a warming trend since the 1970s, and all years during the present century (2001–2008) have been among the warmest since the beginning of instrumental records (Allison et al., 2009). Surface air temperature observations reveal the strongest increase occurring over Northern Hemisphere land areas (40–70° N) since 1970s; an increase almost twice the global average rate (IPCC, 2007), accompanied by a $\sim 1\%$ decade⁻¹ increase in precipitation (ACIA, 2005).

Freshwater flux to Sermilik Fjord, SE Greenland

S. H. Mernild et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Freshwater flux to Sermilik Fjord, SE Greenland

S. H. Mernild et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



There is clear evidence of increased melting of the GrIS and marginal glaciers in Greenland since the mid-1990s (e.g., Mote, 2007; Tedesco, 2007; Knudsen and Hasholt, 2008; Steffen et al., 2008), and rapid mass loss has been observed and simulated (e.g., Hanna et al., 2008; Mernild et al., 2008a; Allison et al., 2009; Khan et al., 2010). In a warming climate, we would expect an accelerating freshwater flux: ice discharge (calving) and runoff to the ocean, manifested by, e.g., decreasing ocean salinity, and increasing global eustatic sea level rise (e.g., ACIA, 2005; Box et al., 2006; IPCC, 2007). A few freshwater runoff measurements are available for Greenland from 1990s at the Sermilik Research Station, Sermilik Fjord (65° N), SE Greenland, and at the Zackenberg Research Station, Young Sund/Tyroler Fjord (74° N), NE Greenland, and shorter term measurements are available from Kangerlussuaq (Søndre Strømfjord), Kangerlussuaq Fjord (67° N), W Greenland, since 2007, and from Kobberfjord (64° N), W Greenland, since 2008 (e.g., Mernild and Hasholt, 2006, 2009; Jensen and Rasch, 2009). These data series are important tools for assessing and quantifying the impact of climate change and variability on freshwater runoff from glaciated landscapes such as Greenland.

The first documentation of glaciers in the Sermilik Fjord catchment basin was carried out in 1933, and in 1970 the Sermilik Research Station was established close to the Mittivakkat Glacier to study the control of climate on a low-arctic (Born and Böcher, 2001), partly glaciated landscape. An automated terrestrial monitoring program was initiated for the Mittivakkat Glacier catchment in 1993, which presents today the longest continuous monitoring program in E Greenland. Data on observed climate conditions have been presented by Mernild et al. (2008b) and Jakobsen et al. (2008). Seasonal and annual observations on the Mittivakkat Glacier include: winter, summer, and net mass-balance (Knudsen and Hasholt, 2004, 2008), freshwater runoff (e.g., Hasholt, 1980; Hasholt and Mernild, 2004, 2008), solute export (Yde et al., 2010), and sediment transport (Hasholt and Walling, 1992; Buskamp and Hasholt, 1996; Hasholt and Mernild, 2008). Modeling studies for this region include seasonal and annual climate processes (Mernild and Liston, 2010), snow cover distribution (Hasholt et al.,

2003; Mernild et al., 2006, 2008a), glacier surface mass-balance (Mernild et al., 2006, 2008a), and runoff (Mernild and Hasholt, 2006; Mernild et al., 2008a).

This collection of extensive observations and model results from the Mittivakkat Glacier catchment was used to simulate the terrestrial surface runoff for the Sermilik Fjord. Not only runoff but also ice discharge from e.g., the Helheim Glacier (one of the most conspicuous calving outlet glaciers from the GrIS) at the innermost part of the fjord, seems to be an important source of freshwater for both, the Sermilik Fjord and the Irminger Sea. We present a 10-year time series (1999–2008) of freshwater flux to the Sermilik Fjord in order to assess variability and trend thereof due to changes in air temperature, net precipitation (hereafter referred to as precipitation), and ice dynamics. In particular, we address the simulated temporal and spatial distribution of terrestrial surface freshwater runoff to the fjord and also on a sub-catchment scale. The runoff was simulated in SnowModel (Liston and Elder, 2006a; Mernild et al., 2006), based on in situ meteorological data within the Sermilik Fjord area. Runoff was initially simulated for the Mittivakkat Glacier catchment area of $\sim 18 \text{ km}^2$ and tested against observed runoff data from the Mittivakkat Glacier catchment outlet which is the only place in the Sermilik Fjord where runoff is observed. The simulated runoff was verified (bias corrected) against runoff observations, before runoff simulations were scaled up to the entire Sermilik Fjord catchment area. The following objectives are addressed: (1) assess the performance of SnowModel by comparing simulated runoff against observed runoff for the Mittivakkat Glacier catchment; (2) simulate the spatial runoff variability and quantify whether the annual freshwater runoff to the Sermilik Fjord has been increasing throughout the simulation period; (3) compare simulated runoff with observed Helheim ice discharge to illustrate the respective distribution from each freshwater source; and (4) merge different sources of freshwater input, e.g., simulated runoff, simulated precipitation over the fjord with satellite-derived ice discharge and geothermal and frictional melting due to basal ice motion to quantify the freshwater flux to the Sermilik Fjord.

Freshwater flux to Sermilik Fjord, SE Greenland

S. H. Mernild et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2 Study area

2.1 Physical settings and climate

The Sermilik Fjord catchment (58 045 km²) is located on the east coast of Greenland (65° N; 37° W), connected to the Irminger Sea (Fig. 1a). The fjord is 1103 km² in area, 85 km in length, and the largest fjord system in SE Greenland. The catchment drains a part of the GrIS, including the Helheim, Fenris, and Midgård Glaciers (the three major outlet glaciers in Sermilik Fjord catchment), and marginal glaciers, among these the Mittivakkat Glacier on Ammassalik Island (see Figs. 1c and 3a for location), where long-term monitoring of climate, mass-balance, and runoff was observed (Mernild and Hasholt, 2006; Knudsen and Hasholt, 2008). The Sermilik Fjord catchment ranges in elevation from sea level to ~2900 m a.s.l. The lower parts of the terrain (elevation below 700–1000 m a.s.l.) are dominated by exposed bedrock, sporadic thin soil layers, and sparse vegetation. Landscapes above 700–1000 m a.s.l. are mostly covered by glaciers and the GrIS (Fig. 1c). For the purposes of this study, the Sermilik Fjord catchment has been divided into 7 sub-catchments, each draining into specific parts of the fjord. These areas also represent characteristic variations in glacier ice coverage from approximately 10% (area 2) to 87% (area 4) (Figs. 1c and 4f).

The Sermilik Fjord region represents one of the most humid parts of Greenland (the zone of largest annual precipitation is located ~200 km further south). The mean annual air temperature (MAAT) for the full catchment including the GrIS was -4.8°C (1999–2008), varying from around freezing point at the near coastal stations (Tasiilaq and Coast Stations) to around -19°C on the GrIS (Station NASA-SE) (Fig. 2a). The total annual simulated precipitation (corrected after Allerup et al., 1998, 2000; for further information about the detailed precipitation correction procedures, see Mernild et al., 2009, 2010a) varied from ~1200 to ~1800 mm w.eq. y⁻¹ within the catchment. Our simulated precipitation values were in the same order of magnitude compared to values from Ettema et al. (2009). Mean annual catchment relative humidity and wind speed were 83% and 4.1 m s⁻¹, respectively. Sermilik Fjord is dominated by seasonal

Freshwater flux to Sermilik Fjord, SE Greenland

S. H. Mernild et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and local-to-regional variations in climate. During summer, the low lying coastal areas, on, e.g., Ammassalik Island (approximately below 300 m a.s.l.), are influenced by air temperature inversions which are common in Arctic coastal landscapes due to the effect of thermal differences between land and ocean (e.g., Kozo, 1982; Weick and Rouse, 1991; Mernild and Liston, 2010). The climate and its seasonal variability are illustrated in Fig. 2b using positive summer air temperature lapse rates in the near coastal areas. Apart from this temperature inversion in the lower lying near coastal areas during summer, meteorological data from all seven stations in the Sermilik Fjord catchment (Fig. 1) showed constantly negative mean monthly lapse rates, very similar to the high-elevation GrIS air temperature lapse rates (Fig. 2a) (e.g., Steffen and Box, 2001; Mernild et al., 2009).

3 Model description and satellite data

3.1 SnowModel and model simulations

SnowModel (Liston and Elder, 2006a), is a spatially-distributed snow-evolution, ice melt, and runoff modeling system useful in all landscapes, climates, and conditions where snow and ice play an important role in hydrological cycling (Mernild et al., 2006; Mernild and Liston, 2010). For a detailed description of SnowModel, including its sub-programs: *MicroMet*, *EnBal*, *SnowPack*, *SnowTran-3D*, and *SnowAssim* see Liston and Elder (2006a,b), Liston et al. (2008), Liston and Hiemstra (2008), and Mernild and Liston (2010). SnowModel is a surface model simulating first-order effects of atmospheric forcing on snow, glacier ice, and runoff, but processes related to glacier dynamics are not included.

3.2 Input data, model verification, and uncertainties

Meteorological data of air temperature, relative humidity, wind speed, wind direction, and precipitation were obtained from seven meteorological stations at different elevations within the simulation domain (Fig. 1b). Four stations were located on the GrIS,

Freshwater flux to Sermilik Fjord, SE Greenland

S. H. Mernild et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and three outside the GrIS in the coastal region (Table 1). Simulations were performed on a one-day time step, based on lapse rates, generated using observed data from all seven stations (Fig. 2b). The simulation period spans from 1 September 1998 through 31 August 2009, whereas output values were simulated for calendar years (1999–2008). Air temperature and precipitation data are summarized in Fig. 2a.

The Greenland topographic data at 625-m resolution from Bamber et al. (2001) was used with the image-derived correction published by Scambos and Haran (2002), and interpolated to a 500-m grid increment covering a 400.5 by 300.5 km simulation domain for the Sermilik Fjord catchment (Fig. 1a). The location of the Sermilik Fjord coast line, GrIS terminus, and marginal glaciers were estimated by using Moderate Resolution Imaging Spectroradiometer (MODIS) satellite images (observed on 30 August 2009). User-defined constants for SnowModel are shown in Mernild et al. (2009), and parameter definitions are given in Liston and Sturm (1998).

SnowModel simulated runoff was tested and verified (bias corrected) against observed runoff from the outlet of the Mittivakkat Glacier catchment for the period 1999–2005 (Fig. 2). The cumulative simulated runoff was initially underestimated by 34–43%, averaging 38% according to runoff observations. Therefore, a linear regression ($r^2=0.95$; where r^2 is the explained variance) was used for verification as shown in Fig. 3a. The verified cumulative annual Mittivakkat Glacier runoff is illustrated in Fig. 3b. This underestimation of runoff is expected to be due to: (1) englacial and subglacial water flow to neighboring glacier sub-catchments; (2) uncertainties associated with model inputs; (3) unrepresented or poorly-represented processes in SnowModel; and (4) uncertainties related to runoff observations (see below). A dye trace study on the Mittivakkat Glacier by Mernild (2006) confirmed englacial and subglacial water flow through crevasses, moulins, and tunnels to neighboring glacier sub-catchments, and modeling studies further indicated that the location of the northern catchment divide on the Mittivakkat Glacier itself may vary because of crevasse and tunnel development due to, e.g., glacier dynamic activity. Therefore, the glacier may deliver and, in this case, receive freshwater water across this divide from neighboring glacier sub-catchments.

**Freshwater flux to
Sermilik Fjord,
SE Greenland**

S. H. Mernild et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The assumed accuracy of single outlet discharge measurements is within 5–10%, whereas calculated stage-discharge values might deviate up to 25% from simultaneous manual measurements. However, long-term discharges (monthly and annual) are typically accurate within approximately 5–15% (Hasholt et al., 2006; Mernild and Hasholt, 2009).

Statistical analysis from previous SnowModel studies on snow distributions, snow and glacier melt, and runoff from marginal glaciers in Greenland and the GrlS (e.g., Mernild et al., 2006, 2009), along with uncertainties in observed discharge used for verification, indicates that simulated influx of runoff to the Sermilik Fjord might be influenced by the same order of uncertainties: We assume an error of 10–25%. This includes uncertainties related to not using routines for air temperature inversion in low lying near coastal areas (Mernild and Liston, 2010) and the associated influence on snow and glacier ice melt and glacier mass-balance simulations; unfortunately no radiosonde data exist for the inner part of the Sermilik Fjord.

3.3 Satellite-derived ice discharge

Time series of ice flux to the calving front, which we term the ice discharge, for Helheim Glacier, Midgård Glacier, and Ferris Glacier were calculated from observed average surface velocity, glacier width and estimated ice thickness. Speeds were measured from automated Repeat-Image Feature Tracking (RIFT) using pairs of orthorectified images from: (1) Landsat 7 Enhanced Thematic Mapper Plus (panchromatic band) distributed by the United States Geological Survey; (2) visible to near-infrared bands of the Advanced Spaceborne Thermal Emissivity and reflection Radiometer (ASTER) distributed by the NASA Land Processes Distributed Active Archive (LP DAAC); and (3) SPOT-5 panchromatic images distributed through the SPIRIT program. Landsat and ASTER images have a pixel resolution of 15 m and the 5 m SPOT-5 images were down-sampled to 15 m for RIFT processing. The Multi-Image/Multi-Chip (MIMC) RIFT algorithm (Ahn and Howat, 2010) was used to measure surface displacements every 100 m. Individual displacement vectors were then averaged over a 1-km wide band

Freshwater flux to Sermilik Fjord, SE Greenland

S. H. Mernild et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



near the front the glacier to provide a time series of average velocity at the ice front. Due to the high speeds observed, we assumed that speed was constant with ice depth. Averaged glacier width over the region of velocity sampling was obtained from 15-m Landsat imagery.

Ice thickness for Helheim Glacier was obtained in 2001 and 2008 by the University of Kansas Coherent Radar Depth Sounder (CoRDS) (Gogineni et al., 2001; Howat et al., 2005). For Fenris and Midgård glaciers, for which no thickness data are available, ice thickness at the start of the time series was estimated from the height of the calving front assuming a grounded ice front at hydrostatic equilibrium and densities of ice and sea water of 910 and 1027 kg m³, respectively. We assume an error of ±50 m in this thickness estimate. Changes in ice thickness through time were then measured from repeat ASTER digital elevation models produced by the LP DAAC and vertically co-registered using tie points over ice-free terrain. These data have a relative error of ±10 m in the vertical (Fujisada et al., 2005).

3.4 Geothermal and frictional melting due to basal ice motion

The upper-bounds for the melt water generated through melting at the ice bed due to: (a) geothermal heating; and (b) frictional heating due to basal ice motion were estimated. For (a), it was liberally assumed that the bed was at the melting temperature over all regions with surface elevations below 1200 m a.s.l., and area of 2300 km². A typical geothermal heat flux of 0.05 W m⁻² gives a basal melt rate of 5 mm w.eq. y⁻¹ (Cuffey and Paterson, 2010, p. 118) for ice at the melting temperature, totaling approximately 0.01 × 10⁹ m³ y⁻¹ produced by geothermal heating over this area, which was 2 orders of magnitude less than the contributions from runoff and ice discharge, and can therefore be ignored (Table 2).

For (b), the maximum rate of basal melt due to frictional heating caused by ice sliding over the bed is Eq. (1):

$$m = (tb \cdot ub) / (\rho \cdot Lf) \quad (1)$$

Freshwater flux to Sermilik Fjord, SE Greenland

S. H. Mernild et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



where, τ_b is the basal shear stress, u_b is sliding speed, ρ is the ice density, and L_f is the latent heat. Again, it was assumed that the bed was at the melting temperature over the drainage area with surface elevations below 1200 m a.s.l. We also assumed that 100% of the ice motion needed to maintain mass balance (i.e., the balance velocity) was accomplished through basal sliding in this region. Balance velocities were obtained from Bamber et al. (2000). Finally, we assumed that the basal drag was equal to the driving stress, which we calculated from the ice thickness and surface elevation maps from Bamber et al. (2000, 2001). From this we obtain a total melt volume rate of approximately $0.5 \times 10^9 \text{ m}^3 \text{ y}^{-1}$, which is approximately 1% of the average total freshwater flux (Table 2).

4 Results and discussion

4.1 Terrestrial surface runoff to Sermilik Fjord

Annual (1999–2008) cumulative simulated runoff from all seven sub-catchments, and from the entire catchment, to Sermilik Fjord is shown in Fig. 4b. Total runoff to Sermilik Fjord for the modeled decade averaged $4.6 \times 10^9 \text{ m}^3 \text{ y}^{-1}$, from a minimum of $2.9 \times 10^9 \text{ m}^3 \text{ y}^{-1}$ in 1999 to a maximum of $5.9 \times 10^9 \text{ m}^3 \text{ y}^{-1}$ in 2005; these values were expected to be among the highest since the simulation period included the warmest years since the beginning of instrumental records. For the simulation period, data showed an average insignificant increase in runoff of $1.0 \times 10^9 \text{ m}^3$ ($r^2=0.14$, $p<0.25$; where p level of significance) (Fig. 4b), due to both increasing annual precipitation ($r^2=0.13$) and increasing mean annual summer air temperature (June through August) ($r^2=0.32$). An increase in precipitation for the Sermilik Fjord catchment of $\sim 15\%$ decade $^{-1}$, which was above the average increase of $\sim 1\%$ decade $^{-1}$ estimated by ACIA (2005).

For the simulation period, 2007 showed the largest satellite-derived GrIS cumulative melt extent followed by 2005 (Steffen et al., 2008), but also the largest melt index –

Freshwater flux to Sermilik Fjord, SE Greenland

S. H. Mernild et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



defined as the melting area times the number of melting days for areas above 2000 m in elevation (Tedesco, 2007) – followed by 2005 as the fourth highest for the simulation period. The largest amount of modeled runoff to the Sermilik Fjord occurred in 2005, and not in 2007 (Fig. 4b). This discrepancy between the GrIS melting conditions and the Sermilik Fjord runoff conditions was due to a record high annual precipitation for 2005 of ~ 1800 mm w.eq. y^{-1} combined with the second highest mean annual summer air temperatures of 2.2°C (influencing the melting snow and ice conditions) (Fig. 2a). The record high 2005 precipitation combined with the relatively high percentage of rain ($\sim 65\%$ of the total annual precipitation) was the reason why less precipitation accumulated as snow during winter, and more streamed directly into the fjord as runoff. The connection between snow melting, melt water retention and refreezing within the snowpack, and runoff is described, e.g., in Hanna et al. (2008), Mernild et al. (2009) related to the variation in annual snow accumulation/precipitation.

Weather conditions for the Sermilik Fjord are strongly affected by low pressure systems, especially the associated wind and precipitation which varies significantly due to year-to-year changes in the storm tracks. Most low pressure centers affecting Greenland arrive from directions between south and west, steered by an upper level cyclone, the “polar vortex”. During winter these are normally centered over the Canadian Cold Pole and during summers they are less pronounced and centered over the Arctic Ocean (Hansen et al., 2008). This pressure system is of utmost importance for the specific weather conditions at a given position in the southeastern Greenland sector. Therefore, it should be kept in mind, even though maximum melting conditions occurred for the GrIS as in 2007, local variability in precipitation can be the reason for annual runoff peaks, as illustrated for the Sermilik Fjord catchment for 2005.

On a sub-catchment scale, the interannual runoff variability generally followed the variability of the overall runoff to the fjord, showing lowest runoff values in 1999 and highest values in 2005 (Fig. 4b). In Fig. 4b the spatial distribution of runoff to the Sermilik Fjord is illustrated, displaying that sub-area 7 contributed, on average, the lowest annual runoff volume of $0.4 \times 10^9 \text{ m}^3 \text{ y}^{-1}$, and sub-area 4 the highest value of

Freshwater flux to Sermilik Fjord, SE Greenland

S. H. Mernild et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

5 $1.4 \times 10^9 \text{ m}^3 \text{ y}^{-1}$. Besides the general effect of precipitation and summer air temperatures on runoff from all sub-catchments, both the percentage of glacier cover and its hypsometry within each sub-catchment strongly influenced simulated runoff within the region. Generally, sub-catchments showing high fractions of glacier cover and glaciers at low elevations show stronger positive runoff effects during years with high temperatures (Fig. 4c).

10 The Sermilik Fjord accumulated catchment and sub-catchment runoff (1999–2008) are illustrated in Fig. 4c, showing an overall cumulative freshwater runoff volume of $46.0 \times 10^9 \text{ m}^3$. The lowest cumulative runoff contribution occurred from sub-area 7, with a total of $3.5 \times 10^9 \text{ m}^3$, which equalled about 8% of the overall freshwater runoff to the fjord. The highest contribution of $10.4 \times 10^9 \text{ m}^3$ came from sub-area 4 – the Helheim sub-catchment –, which equalled about 25% of the overall runoff. The percentage of cumulative freshwater runoff from the other sub-areas (area 1–3 and 5–6) averaged from 9% to 17% of the overall runoff (Fig. 4d). Obviously sub-catchments with the greatest glacier coverage, combined with the highest percentage of glaciers at low elevations, were the sub-areas where the greatest freshwater runoff contribution to the fjord occurred, and vice versa. In Fig. 4e the differences between sub-catchments 4 and 7 in glacier cover (km^2) and in glacier cover (%) within each 100-m elevation interval are shown. Area 4 was the sub-catchment having the greatest glacier area of 910 km^2 from where runoff occurred: $\sim 30\%$ of the area was located below 500–600 m a.s.l. Area 7 was, however, a sub-catchment having a glacier cover of only 65 km^2 of which only $\sim 10\%$ was found below 500–600 m a.s.l. Areas 1–3 and 5–6 represent a mixture of the main characteristics found in areas 4 and 7. Even though areas 3 and 6 both have a relatively high glacier cover of 67% and 79%, respectively, the cumulative runoff only accounted for 14% and 9% of the overall runoff to the fjord (Fig. 4d). The reason for these relatively low percentages of runoff values from sub-area 3 and 6 were due to the high elevated glacier cover within each sub-catchment. For sub-area 5 the situation was however different: In area 5 the amount of runoff accounted for 17% of the overall runoff to Sermilik Fjord, even though the sub-catchment

Freshwater flux to Sermilik Fjord, SE Greenland

S. H. Mernild et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

areal and the glacier cover area were relative low. The reason for the relatively high runoff volume from sub-area 5 was because of the low elevated glacier cover in the sub-area.

In Fig. 5 the spatial distributions of annual cumulative runoff to Sermilik Fjord are illustrated for 1999 through 2008. Those parts of the fjord catchment exhibiting glaciers covering low altitudes, e.g., both marginal glaciers and the Helheim glacier terminus showed the highest simulated runoff values. At the Helheim glacier terminus the areally-averaged annual maximum runoff ranged from ~ 1.8 m.w.eq. in 2003 to more than ~ 3.8 m.w.eq. in 2007. Simulated runoff values which seemed to be in line with previously published values, e.g., by Ettama et al. (2008), and consistent with previous GrIS runoff simulations by Mernild et al. (2009). The amount of simulated runoff decreased with increasing altitude, on average by ~ 250 mm w.eq. 100 m^{-1} from the ice margin all the way to the boundary where runoff occurs (Fig. 5); for Jakobshavn drainage area, W Greenland (69° N), the value was similar with ~ 220 mm w.eq. 100 m^{-1} (Mernild et al., 2010b). On the GrIS within the Sermilik Fjord catchment (for a latitude range of $65\text{--}66^\circ \text{ N}$) this annual runoff boundary line was located about 25–40 km from the GrIS terminus at an elevation of 1140 m a.s.l. to 1600 m a.s.l., averaging 1150(± 140) m a.s.l.

4.2 Freshwater flux to Sermilik Fjord

To account for the freshwater flux to the Sermilik Fjord, not only terrestrial surface runoff needs to be addressed, but also: (1) ice discharge influenced by GrIS dynamical processes (as described by Howat et al., 2005, 2008) and temperature of near-coastal ocean currents (Holland et al., 2008; Straneo et al., 2010); (2) seasonal changes in internal drainage system due to melting; (3) runoff from subglacial geothermal melting and frictional melting due to basal ice motion; (4) submarine melting at tidewater glacier margins; and (5) precipitation (rain and snow accumulation on sea ice) at the Sermilik Fjord surface area. Unfortunately, seasonal changes in internal drainage system was omitted, due to missing data (values probably insignificant related to the overall

Freshwater flux to Sermilik Fjord, SE Greenland

S. H. Mernild et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



terrestrial freshwater flux budget to the Sermilik Fjord), while submarine melting at tidewater glacier margins was integrated in the ice discharge values. Contributions of ice discharge from minor GrIS outlet glaciers, e.g., glaciers located in Johan Petersens Fjord were ignored due to lack of available data, while ice discharge from the three major outlet glaciers: the Helheim Glacier, Fenris Glacier, and Midgård Glacier at the innermost part of the fjord were included (Figs. 6 and 7). Based on satellite-derived observations, the Helheim average ice discharge for the period 1999–2008, was estimated to be $25.9 \pm 2.6 \times 10^9 \text{ m}^3 \text{ w.eq. y}^{-1}$ (Table 2), and for the Fenris and Midgård Glaciers ice discharge were 2.5 ± 0.5 and $5.5 \pm 1.0 \times 10^9 \text{ m}^3 \text{ w.eq. y}^{-1}$, respectively (Table 2 and Fig. 6). In Fig. 6 an example of variations in surface ice velocity, ice thickness, and ice discharge for the Helheim Glacier, Fenris Glacier, and Midgård Glacier are illustrated, showing substantial variations in velocity, ice thickness, and discharge, with a general increase in velocity and discharge after 2002 and peaking in 2005 and 2006. Due to both decreased ice velocity and thickness, ice discharge at Helheim Glacier decreased to earlier levels by 2007 (Howat et al., 2007). However, for the simulation period, ice discharge from the three major outlet glaciers showed an average significant increase of $13.2 \times 10^9 \text{ m}^3 \text{ w.eq.}$ ($r^2=0.49$; $p<0.01$) (Fig. 7). In Table 2 mean annual ice discharge values were combined with annual SnowModel simulated precipitation at the Sermilik Fjord surface area, terrestrial surface runoff, and subglacial geothermal and frictional melting, to deduce the freshwater flux: (1) from the Helheim Glacier catchment, but also; (2) to the entire Sermilik Fjord. Calculations showed a freshwater flux averaging $26.9 \times 10^9 \text{ m}^3 \text{ y}^{-1}$ from the Helheim Glacier sub-catchment, and a flux of $40.0 \times 10^9 \text{ m}^3 \text{ y}^{-1}$ to the Sermilik Fjord for 1999–2008. The suggested freshwater flux entering the Sermilik Fjord of $40.4 \times 10^9 \text{ m}^3 \text{ y}^{-1}$, equaled 5% of the freshwater flux of $786 \times 10^9 \text{ m}^3 \text{ y}^{-1}$ (1995–2007) from the entire GrIS to the ocean (Mernild et al., 2009). For the Helheim Glacier catchment, around $25.9 \times 10^9 \text{ m}^3 \text{ y}^{-1}$ (96%) originated from satellite-derived ice discharge, and $1.0 \times 10^9 \text{ m}^3 \text{ y}^{-1}$ (4%) from SnowModel simulated runoff. For Sermilik Fjord $4.6 \times 10^9 \text{ m}^3 \text{ y}^{-1}$ (11%) out of a freshwater flux of $40.4 \times 10^9 \text{ m}^3 \text{ y}^{-1}$, originated from the SnowModel simulated runoff, $1.4 \times 10^9 \text{ m}^3 \text{ y}^{-1}$

Freshwater flux to Sermilik Fjord, SE Greenland

S. H. Mernild et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(3%) from the SnowModel simulated precipitation at the Sermilik Fjord surface area, $33.9 \times 10^9 \text{ m}^3 \text{ y}^{-1}$ (85%) from ice discharge, and $0.5 \times 10^9 \text{ m}^3 \text{ y}^{-1}$ (1%) from subglacial frictional melting due to basal ice motion (Table 2). In general, ice discharge represented 65%, 14%, and 6% of the total freshwater flux from Helheim, Midgård, and Fenris glaciers, respectively (Table 2).

The Helheim Glacier represents one of the major outlet glaciers from the GrIS, which dominated 65% of the freshwater flux into Sermilik Fjord. Runoff only forms a minor part (11%) of the overall freshwater flux to the fjord. Similar conditions were found for the Jakobshavn drainage area (2000–2007), where 7% of the average annual freshwater flux originated from surface runoff (Mernild et al., 2010b). For two of the major GrIS outlet glacier sub-catchments (Helheim and Jakobshavn) it can be concluded that runoff was a minor contributor to the freshwater flux, which was highly dominated by ice discharge. For both Helheim and Jakobshavn, it should be kept in mind that the freshwater flux to the Sermilik Fjord and the Illulissat Fjord, respectively, does not include values for a melting internal drainage system through out runoff season. The freshwater flux should be seen as first estimate, even though, omitted freshwater inputs from changes in englacial and subglacial internal drainage system due to melting only counts for a minor part of the overall flux to the fjords.

5 Summary and conclusion

The amount of freshwater runoff reaching the ocean from marginal glaciers, the GrIS, and ice free landscapes depends on the precipitation and storage changes in reservoirs of ice, snow, and water on land. In many places around Greenland, glaciers calve directly into the sea and the overall flux of freshwater from specific catchments, e.g., the Sermilik Fjord catchment, will be influenced by or even dominated by a discharge of calving ice. At the Sermilik Fjord catchment, 85% of the average annual freshwater flux of $40.4 \times 10^9 \text{ m}^3 \text{ y}^{-1}$ (1999–2008) originated from ice discharge, 11% from terrestrial surface runoff, 3% from net precipitation at the fjord area, and 1% from subglacial

Freshwater flux to Sermilik Fjord, SE Greenland

S. H. Mernild et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



frictional melting. In period of a warmer climate, as for example during the recent decade an increase in runoff ($r^2=0.14$) and ice discharge ($r^2=0.49$) occurred. The Sermilik Fjord increasing runoff was caused by increasing mean annual summer air temperature and precipitation: even though maximum melting conditions occurred in 2007, local variability in precipitation can be the reason for annual runoff peaks, as illustrated for the Sermilik Fjord for 2005.

Acknowledgements. Special thanks to the Danish Meteorological Institute, the Cooperative Institute for Research in Environmental Sciences (CIRES), University of Colorado at Boulder, the Department of Geography and Geology, University of Copenhagen, and the Geological Survey of Denmark and Greenland (GEUS) for use of observed meteorological data. Further, thanks to the Department of Geography and Geology, University of Copenhagen, for use of observed runoff data for verification. Thanks are also given to Byrd Polar Research Center, Ohio State University, for use of satellite-derived ice discharge. This work was supported by grants from the Climate Change Prediction Program within the US Department of Energy's Office of Science. Los Alamos National Laboratory is operated under the auspices of the National Nuclear Security Administration of the U.S. Department of Energy under Contract No. DE-AC52-06NA25396.

References

- ACIA: Arctic Climate Impact Assessment, Cambridge University Press, 1042 pp., 2005.
- Ahn, Y. and Howat, I. M.: Efficient, automated glacier surface velocity measurements from repeat images using multi-image/multi-chip (MIMC) feature tracking, IEEE T. Geosci. Remote, submitted, 2010.
- Allerup, P., Madsen, H., and Vejen, F.: Estimating true precipitation in Arctic areas, Proc. Nordic Hydrological Conf., Helsinki, Finland, Nordic Hydrological Programme Rep. 44, 1–9, 1998.
- Allerup, P., Madsen, H., and Vejen, F.: Correction of precipitation based on off-site weather information. Atmos. Res., 53, 231–250, 2000.
- Allison, I., Bindoff, N. L., Bindshadler, R. A., Cox, P. M., de Noblet, N., England, M. H., Francis, J. E., Gruber, N., Haywood, A. M., Karoly, D. J., Kaser, G., Le Quéré, C., Lenton, T. M., Mann, M. E., McNeil, B. I., Pitman, A. J., Rahmstorf, S., Rignot, E., Schellnhuber, H. J.,

Freshwater flux to Sermilik Fjord, SE Greenland

S. H. Mernild et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Freshwater flux to
Sermilik Fjord,
SE Greenland**S. H. Mernild et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Schneider, S. H., Sherwood, S. C., Somerville, R. C. J., Steffen, K., Steig, E. J., Visbeck, M., and Weaver, A. J.: The Copenhagen Diagnosis: Updating the World on the Latest Climate Science, The University of New South Wales Climate Change Research Centre (CCRC), Sydney, Australia, 60 pp., 2009.

5 Bamber, J., Ekholm, S., and Krabill, W.: A new, high-resolution digital elevation model of Greenland fully validated with airborne laser altimeter data, *J. Geophys. Res.*, 106, 6733–6746, 2001.

Bamber J., Hardy, R. J., and Joughin, I.: An analysis of balance velocities over the Greenland Ice Sheet and comparison with synthetic aperture radar interferometry, *J. Glaciol.*, 46(152), 67–74, 2000.

10 Born, E. W. and Böcher, J.: The Ecology of Greenland, Nuuk, Ministry of Environment and Natural Resources, Greenland, 429 pp., 2001.

Box, J., Bromwich, D. H., and Veenhuis, B. A., et al.: Greenland Ice Sheet surface mass balance variability (1998–2004) from calibrated polar MM5 output, *J. Climate*, 19, 2783–2800, 2006.

15 Buskamp, R. and Hasholt, B.: Coarse bed load transport in a glacial valley, Sermilik, South East Greenland, *Z. Geomorphol.*, 40, 349–358, 1996.

Cuffey, K. M. and Paterson, W. S. B.: The Physics of Glaciers, 4th edn., Butterworth-Heinemann, Oxford, 704 pp., 2010.

20 Ettema, J., van den Broeke, M., van Meijgaard, E., van de Berg, M. J., Bamber, J. L., Box, J. E., and Bales, R. C.: Higher surface mass balance of the Greenland Ice Sheet revealed by high resolution climate modeling, *Geophys. Res. Lett.*, 36, L12501, doi:10.1029/2009GL038110, 2009.

Gogineni, S., Tammana, D., Braaten, D., et al.: Coherent radar ice thickness measurements over Greenland Ice Sheet, *J. Geophys. Res.*, 106, 33761–33772, 2001.

25 Hanna, E., Huybrechts, P., Steffen, K., et al.: Increased runoff from melt from the Greenland Ice Sheet: a response to global warming, *J. Climate*, 21, 331–341, 2008.

Hansen, B. U., Sigsgaard, C., Rasmussen, L., Cappelen, J., Hinkler, J., Mernild, S. H., Petersen, D., Tamstorf, M., Rasch, M., and Hasholt, B.: Present day climate at Zackenberg, *Adv. Ecol. Res.*, 40, 115–153, 2008.

30 Hasholt, B.: Morphological and hydrological possibilities for the development of water power at Angmassalik – a case study of applied physical geography, *Geogr. Tidsskr.*, 80, 57–62, 1980.

**Freshwater flux to
Sermilik Fjord,
SE Greenland**

S. H. Mernild et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Hasholt, B., Bobrovitskaya, N., Bogen, J., McNamara, J., Mernild, S. H., Milbourn, D., and Walling, D. E.: Sediment transport to the Arctic Ocean and adjoining cold oceans, *Nord. Hydrol.*, 37(4–5), 413–432, 2006.

Hasholt, B., Liston, G. E., and Knudsen, N. T.: Snow-distribution modelling in the Ammassalik Region, South East Greenland, *Nord. Hydrol.*, 34(1/2), 1–16, 2003.

Hasholt, B. and Mernild, S. H.: Estimation of water balance in and around the Mittivakkat Glacier basin, Ammassalik Island, Southeast Greenland, in: *Northern Research Basins Water Balance*, edited by: Kane, D. L. and Yang, D., Wallingford, UK, IAHS, 129–142, publications no. 290, ISBN 1-901502-82-1, 2004.

Hasholt, B. and Mernild, S. H.: Hydrology, sediment transport, and water resources of Ammassalik Island, SE Greenland, *Geogr. Tidsskr.*, 108(1), 73–95, 2008.

Hasholt, B., Mernild, S. H., Sigsgaard, C., Elberling, B., Petersen, D., Jakobsen, B. H., Hansen, B. U., Hinkler, J., and Søgaard, H.: Hydrology and transport of sediment and solutes at Zackenberg, in: *High-Arctic Ecosystem Dynamics in a Changing Climate: Ten Years of Monitoring and Research at Zackenberg Research Station, Northeast Greenland*, edited by: Meltofte, H., Christensen, T. R., Elberling, B., Forchhammer, M. C., and Rasch, M.: *Hydrology and transport of sediment and solutes at Zackenberg*, *Adv. Ecol. Res.*, 40, 111–149, 2008.

Hasholt, B. and Walling, D. E.: Use of caesium-137 to investigate sediment sources and sediment delivery in a small glacierized mountain drainage basin in eastern Greenland, in: *Erosion, Debris Flows and Environment in Mountain Regions*, IAHS, Publ. 209, 87–99, 1992.

Holland, D. M., Thomas, R. H., de Young, B., Ribergaard, M. H., and Lyberth, B.: Acceleration of Jakobshavn Isbræ triggered by warm subsurface ocean waters, *Nat. Geosci.*, 1(10), 659–664, 2008.

Howat, I. M., Joughin, I., Tulaczyk, S., and Gogineni, S.: Rapid retreat and acceleration of Helheim Glacier, East Greenland, *Geophys. Res. Lett.*, 32, L22502, doi:10.1029/2005GL024737, 2005.

Howat, I. M., Joughin, I., and Scambos, T.: Rapid changes in ice discharge from Greenland outlet glaciers, *Science*, 315, p. 1559, doi:10.1126/science.1138478, 2007 (advanced publication in *Science Express* on 8 February 2007).

IPCC: Summary for policymakers, in: *Climate Change 2007, The Physical Science Basis, Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M.,

Freshwater flux to Sermilik Fjord, SE Greenland

S. H. Mernild et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Averyt, K. B., Tignor, M., and Miller, H. L., Cambridge University Press, Cambridge, UK and New York, USA, 2007.

Jakobsen, B. H., Fredskild, B., and Pedersen, J. B. T.: Holocene changes in climate and vegetation in the Ammassalik area, East Greenland, recorded in lake sediment and soil profiles, *Geogr. Tidsskr.*, 108(1), 21–50, 2008.

Jensen, L. M. and Rasch, M. (Eds.): Zackenbergt Ecological Research Operations, 14th Annual Report 2008, National Environmental Research Institute, Aarhus University, Denmark, 116 pp., 2009.

Jujisada, H., Bailey, G. B., Kelly, G. G., Hara, S., and Abrams, M. J.: ASTER DEM performance, *IEEE T. Geosci. Remote*, 43(12), 2707–2714, 2005.

Kozo, T. L.: An observational study of sea breezes along the Alaskan Beaufort Sea Coast: Part I, *J. Appl. Meteorol.*, 21, 891–905, 1982.

Khan, S. A., Wahr, J., Bevis, M., Velicogna, I., and Kendrick, E.: Spread of ice mass into Northwest Greenland observed, *Geophys. Res. Lett.*, 37, L06501, doi:10.1029/2010GL042460, 2010.

Knudsen, N. T. and Hasholt, B.: Mass balance observations at the Mittivakkat Gletscher, Southeast Greenland 1995–2002, *Nord. Hydrol.*, 35(4–5), 381–390, 2004.

Knudsen, N. T. and Hasholt, B.: Mass balance observations at Mittivakkat Glacier, Ammassalik Island, Southeast Greenland 1995–2006, *Geogr. Tidsskr.*, 108(1), 111–120, 2008.

Liston, G. E. and Hiemstra, C. A.: A simple data assimilation system for complex snow distributions (SnowAssim), *J. Hydrometeorol.*, 9, 989–1004, 2008.

Liston, G. E. and Elder, K.: A distributed snow-evolution modeling system (SnowModel), *J. Hydrometeorol.*, 7, 1259–1276, 2006a.

Liston, G. E. and Elder, K.: A meteorological distribution system for high-resolution terrestrial modeling (MicroMet), *J. Hydrometeorol.*, 7, 217–234, 2006b.

Liston, G. E. and Hiemstra, C. A.: A simple data assimilation system for complex snow distributions (SnowAssim), *J. Hydrometeorol.*, 9, 989–1004, 2008.

Liston, G. E. and Sturm, M.: A snow-transport model for complex terrain, *J. Glaciol.*, 44, 498–516, 1998.

Mernild, S. H.: The internal drainage system of the lower Mittivakkat Glacier, Ammassalik Island, SE Greenland, *Geogr. Tidsskr.*, 106(1), 13–24, 2006.

Mernild, S. H., Hansen, B. U., Jakobsen, B. H., and Hasholt, B.: Climatic conditions at the Mittivakkat Glacier catchment (1994–2006), Ammassalik Island, SE Greenland, and in a 109

Freshwater flux to Sermilik Fjord, SE Greenland

S. H. Mernild et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



years term perspective (1898–2006), *Geogr. Tidsskr.*, 108(1), 51–72, 2008b.

Mernild, S. H. and Hasholt, B.: Climatic control on river discharge simulations, Mittivakkat Glacier catchment, Ammassalik Island, SE Greenland, *Nord. Hydrol.*, 37(4–5), 327–346, 2006.

5 Mernild, S. H. and Hasholt, B.: Observed runoff, jökulhlaups, and suspended sediment load from the Greenland Ice Sheet at Kangerlussuaq, West Greenland, for 2007 and 2008, *J. Glaciol.*, 55(193), 855–858, 2009.

Mernild, S. H. and Liston, G. E.: The influence of air temperature inversion on snow melt and glacier surface mass-balance simulations, SW Ammassalik Island, SE Greenland, *J. Appl. Meteorol. Clim.*, 49(1), 47–67, 2010.

10 Mernild, S. H., Liston, G. E., Hasholt, B., and Knudsen, N. T.: Snow distribution and melt modeling for Mittivakkat Glacier, Ammassalik Island, SE Greenland, *J. Hydrometeorol.*, 7, 808–824, 2006.

15 Mernild, S. H., Liston, G. E., Hiemstra, C. A., Steffen, K., Hanna, E., and Christensen, J. H.: Greenland Ice Sheet surface mass-balance modeling and freshwater flux for 2007, and in a 1995–2007 perspective, *Hydrol. Process.*, 23(17), 2470–2484, doi:10.1002/hyp.7354, 2009.

Mernild, S. H., Liston, G. E., Kane, D. L., Hasholt, B., and Knudsen, N. T.: Spatial snow distribution, runoff, and mass balance modelling for entire Mittivakkat Glacier (1998–2006), Ammassalik Island, SE Greenland, *Geogr. Tidsskr.*, 108(1), 121–136, 2008a.

20 Mernild, S. H., Liston, G. E., Hiemstra, C. A., and Christensen, J. H.: Greenland Ice Sheet surface mass-balance modeling in a 131-year perspective 1950–2080, *J. Hydrometeorol.*, 11(1), 3–25, 2010a.

25 Mernild, S. H., Liston, G. E., Steffen, K., and Chylek, P.: Meltwater flux and runoff modeling in the ablation area of the Jakobshavn Isbræ, West Greenland, *J. Glaciol.*, 56(195), 20–32, 2010b.

Mote, T. L.: Greenland surface melt trends 1973–2007: evidence of a large increase in 2007, *Geophys. Res. Lett.*, 34, L22507, doi:10.1029/2007GL031976, 2007.

30 Rysgaard, S. and Glud, R. N. (Eds.): Carbon cycling in Arctic marine ecosystems: case study young sound, *Meddr. om Grønland, Bioscience*, 58, 24–43, 2007.

Scambos, T. and Haran, T.: An image-enhanced DEM of the Greenland ice sheet, *Ann. Glaciol.*, 34, 291–298, 2002.

- Steffen, K. and Box, J.: Surface climatology of the Greenland ice sheet: Greenland Climate Network 1995–1999, *J. Geophys. Res.*, 106, 33951–33964, doi:10.1029/2001JD900161, 2001.
- Steffen, K., Clark, P. U., Cogley, J. G., Holland, D., Marshall, S., Rignot, E., and Thomas, R.: Rapid changes in glacier and ice sheet and their impacts on sea level, in: *Abrupt Climate Change, a Report by the US Climate Change Science Program and the Subcommittee on Global Change Research*, US Geological Survey, reston, VA, 60–142, 2008.
- 5 Straneo, F., Hamilton, G. S., Sutherland, D. A., Stearne, L. A., Davidson, F., Hammill, M. O., Stenson, G. B., and Rosing-Asvid, A.: Rapid circulation of warm subtropical waters in a major glacial fjord in East Greenland, *Nat. Geosci.*, 3(3), 182–186, 2010.
- 10 Tedesco, M.: A new record in 2007 for melting in Greenland, *EOS T. Am. Geophys. Un.*, 88(39), 383 pp., 2007.
- Velicogna, I. and Wahr, J.: Acceleration of Greenland ice mass loss in spring 2004, *Nature*, 443, 329–331, 2006.
- Weick, E. J. and Rouse, W. R.: Advection in the coastal Hudson Bay Lowlands, Canada, II. Impact of atmospheric divergence on the surface energy balance, *Arctic Alpine Res.*, 23, 338–348, 1991.
- 15 Yde, J. C., Knudsen, N. T., Hasholt, B., and Mernild, S. H.: Chemical weathering and solute export in Southeast Greenland, *Chem. Geol.*, in review, 2010.

Freshwater flux to Sermilik Fjord, SE GreenlandS. H. Mernild et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Freshwater flux to Sermilik Fjord, SE Greenland

S. H. Mernild et al.

Table 1. Meteorological input data for the Sermilik Fjord simulation based on meteorological station data on the GrIS: KULU and NASA-SE (provided by University of Colorado at Boulder) and TAS_L and TAS_U (by Geological Survey of Denmark and Greenland (GEUS)), and from the outside the GrIS: Station Tasiilaq (by Danish Meteorological Institute (DMI)) and Station Coast and Station Nunatak (by University of Copenhagen, Department of Geography and Geology). The abbreviations indicate: (Ta) air temperature, (Rh) relative humidity, (Ws) wind speed, (Wd) wind direction, and (P) precipitation. For station locations see Fig. 1b.

Meteorological station	Location	Grid	Elevation (m a.s.l.)	Data period	Parameters	Operated by
KULU	GrIS	65°45′ N; 39°36′ W	880	20 Jun 1999–13 Sep 2000	Ta, Rh, Ws, and Wd	University of Colorado at Boulder
NASA-SE	GrIS	66°28′ N; 42°30′ W	2390	1 Jan 1999–25 May 2005	Ta, Rh, Ws, and Wd	University of Colorado at Boulder
TAS_L	GrIS	65°38′ N; 38°54′ W	280–270	26 Jun 2006–31 Dec 2008	Ta, Rh, Ws, and Wd	GEUS
TAS_U	GrIS	65°42′ N; 38°52′ W	590–580	16 Apr 2004–31 Dec 2008	Ta, Rh, Ws, and Wd	GEUS
Tasiilaq	Outside GrIS	65°36′ N; 37°38′ W	44	1 Sep 1998–31 Aug 2009	Ta, Rh, Ws, Wd, and P	DMI
Coast	Outside GrIS	65°41′ N; 37°55′ W	25	1 Jan 1999–31 Dec 2008	Ta, Rh, Ws, Wd, and P	University of Copenhagen
Nunatak	Outside GrIS	65°42′ N; 37°49′ W	515	1 Jan 1999–31 Dec 2008	Ta, Rh, Ws, Wd, and P	University of Copenhagen

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Freshwater flux to Sermilik Fjord, SE Greenland

S. H. Mernild et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 2. Freshwater flux from the Helheim Glacier catchment and to the Sermilik Fjord based on SnowModel simulated freshwater runoff, precipitation at the Sermilik Fjord surface area, subglacial geothermal melting and subglacial frictional melting due to basal ice motion, and satellite-derived ice discharge from the Helheim Glacier, Fenris Glacier, and Midgård Glacier (the three major outlet glaciers in Sermilik Fjord catchment). Ice discharge is recalculated from Gt y^{-1} to $10^9 \text{ m}^3 \text{ y}^{-1}$ water equivalent. Mean freshwater flux to the Sermilik Fjord is calculated based on mean values from each input component.

	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	Mean
SnowModel simulated runoff from the Helheim Glacier catchment (sub-area 4), $10^9 \text{ m}^3 \text{ y}^{-1}$	0.7	1.1	0.8	1.1	0.7	1.2	1.3	1.1	1.3	1.0	1.0 (4%)
Satellite-derived ice discharge from the Helheim Glacier including error, $10^9 \text{ m}^3 \text{ y}^{-1}$	21.7±3.3	22.7±2.5	21.8±2.2	26.8±2.3	28.1±3.8	28.0±2.5	33.8±2.8	25.9±1.8	25.9±2.1	24.3±2.8	25.9±2.6 (96%)
Freshwater flux from the Helheim Glacier catchment, $10^9 \text{ m}^3 \text{ y}^{-1}$	22.4	23.8	22.6	27.9	28.8	29.2	35.1	27.0	27.2	25.3	26.9
SnowModel simulated terrestrial runoff to the Sermilik Fjord, $10^9 \text{ m}^3 \text{ y}^{-1}$	2.9	5.1	4.1	5.2	3.5	5.2	5.9	5.1	4.8	4.2	4.6 (11%)
SnowModel simulated precipitation (rain and snow accumulation on sea ice) at the Sermilik Fjord surface area (1103 km ²), $10^9 \text{ m}^3 \text{ y}^{-1}$	0.9	1.2	1.7	1.7	1.4	1.4	1.8	1.4	1.5	1.3	1.4 (3%)
Subglacial geothermal melting, $10^9 \text{ m}^3 \text{ y}^{-1}$	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01 (>0.03%)
Subglacial frictional melting due to basal ice motion, $10^9 \text{ m}^3 \text{ y}^{-1}$	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5 (1%)
Satellite-derived ice discharge from the Helheim Glacier including error, $10^9 \text{ m}^3 \text{ y}^{-1}$	21.7±3.3 (97%)	22.7±2.5 (95%)	21.8±2.2 (96%)	26.8±2.3 (96%)	28.1±3.8 (98%)	28.0±2.5 (96%)	33.8±2.8 (96%)	25.9±1.8 (96%)	25.9±2.1 (95%)	24.3±2.8 (96%)	25.9±2.6 (65%)
Satellite-derived ice discharge from the Fenris Glacier including error, $10^9 \text{ m}^3 \text{ y}^{-1}$	–	2.3±0.5	2.5±0.5	2.6±0.5	2.8±0.5	2.9±0.5	2.8±0.5	2.2±0.5	2.5±0.5	2.4±0.5	2.5±0.5 (6%) 33.9 (85%)
Satellite estimated ice discharge from the Midgård Glacier including error, $10^9 \text{ m}^3 \text{ y}^{-1}$	–	3.7±0.7	3.9±0.8	3.6±0.7	4.2±0.8	6.1±1.0	7.9±1.2	7.3±1.2	7.5±1.2	–	5.5±1.0 (14%)
Freshwater flux to the Sermilik Fjord, $10^9 \text{ m}^3 \text{ y}^{-1}$	–	28.7	28.8	33.0	35.1	37.0	44.5	35.4	35.9	–	40.4




Fig. 1. (a) Greenland, including the Sermilik Fjord simulation domain ($120\,350\text{ km}^2$) and area of interest ($18\,750\text{ km}^2$) in South East Greenland; (b) Sermilik Fjord with topography (300-m contour interval) and meteorological stations (See Table 1 for station information); and (c) MODIS satellite-derived surface characteristics (30 August 2009), including catchment ($58\,045\text{ km}^2$), watershed divide, and locations of GrIS outlet glaciers.

Freshwater flux to Sermilik Fjord, SE Greenland

S. H. Mernild et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Freshwater flux to Sermilik Fjord, SE Greenland

S. H. Mernild et al.




Fig. 2. (a) Time series of observed mean monthly air temperature from coastal (Station Tasiilaq, Coast, Nunatak, TAS_U, TAS_L, and KULU) and GrIS stations (NASA-SE). Observed cumulative monthly precipitation from coastal stations (Tasiilaq; light color, and Coast and Nunatak; dark colors) are illustrated; and (b) mean monthly air temperature lapse rates for all the meteorological stations in the simulation domain, for Station Coast and Nunatak (classified as coastal region; Mernild et al., 2006), and for different areas around the GrIS are illustrated (Mernild et al., 2008, 2009).

Freshwater flux to Sermilik Fjord, SE Greenland

S. H. Mernild et al.




Fig. 3. (a) Observed and simulated Mittivakkat Glacier catchment runoff from 1999–2005. The linear regression was used for verification of the Sermilik Fjord simulated runoff. Be aware that the annual observed runoff periods are different. The inset figure indicates the general location of the Mittivakkat Glacier catchment (black polygon) within sub-catchment area 1 (for a general location of the sub-catchments see Fig. 4a) inside the Sermilik Fjord catchment; and **(b)** observed, simulated, and verified runoff from the Mittivakkat Glacier catchment from 1999–2005.




Fig. 4. (a) Area of interest including the seven sub-catchments for the Sermilik Fjord (simulated in River Tools); (b) time series (1999–2008) of annual sub-catchment simulated runoff and annual cumulative runoff, including trend line (linear) for cumulative runoff; (c) cumulative sub-catchment runoff and overall runoff; (d) percentage of sub-catchment runoff of total runoff; (e) glacier cover distribution in percentage and square kilometer within the elevations from where runoff occurred for the sub-catchment with the lowest cumulative runoff (sub-catchment 7) and the highest (sub-catchment 4); and (f) the percentage of sub-catchment glacier cover within the area of interest.

Freshwater flux to Sermilik Fjord, SE Greenland

S. H. Mernild et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Freshwater flux to Sermilik Fjord, SE Greenland

S. H. Mernild et al.




Fig. 5. Spatial simulated annual cumulative runoff within the area of interest to the Sermilik Fjord for 1999–2008.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Freshwater flux to Sermilik Fjord, SE Greenland

S. H. Mernild et al.




Fig. 6. Satellite-derived time series for the period 1998–2008 of ice velocity, ice thickness, and ice discharge at a stationary flux gate within 3 km of the most-retreated front position of the three major outlet glaciers: Helheim Glacier, Fenris Glacier, and Midgård Glacier draining into the Sermilik Fjord. Ice thickness observations (denoted by vertical error bars) were linearly interpolated, denoted by the curve, and multiplied by constant glacier width and observations of ice velocity and the ratio of water and ice densities (0.91) to obtain discharge of water. Ice discharge data for Helheim Glacier reported here are updated from Howat et al. (2005, 2007).

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)





Fig. 7. Average monthly ice discharge observations (closed circle) and linearly-interpolated values (open circle) for Helheim Glacier, Fenris Glacier, and Midgård Glacier, and sum of the three glaciers for the period 1999 through 2008.

Freshwater flux to Sermilik Fjord, SE Greenland

S. H. Mernild et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

