

This discussion paper is/has been under review for the journal The Cryosphere (TC).  
Please refer to the corresponding final paper in TC if available.

## Comment on “100-year mass changes in the Swiss Alps linked to the Atlantic Multidecadal Oscillation” by Matthias Huss et al. (2010)

P. W. Leclercq, R. S. W. van de Wal, and J. Oerlemans

IMAU, Universiteit Utrecht, Utrecht, The Netherlands

Received: 25 October 2010 – Accepted: 7 November 2010 – Published: 17 November 2010

Correspondence to: P. W. Leclercq (p.w.leclercq@uu.nl)

Published by Copernicus Publications on behalf of the European Geosciences Union.

2475

### Abstract

The paper by Huss et al. (2010) presents a comprehensive set of 100-year specific mass balance series for 30 Swiss glaciers. In the second part of the paper, the authors relate the fluctuations in alpine glacier specific mass balance to climatic changes attributed to the Atlantic Multidecadal Oscillation (AMO). We believe that the specific mass balance is not the appropriate measure to interpret climatic fluctuations. Due to the dynamic response of glaciers to changes in their climatic forcing, the importance of short-term climatic oscillations is overestimated. Taking the changes in glacier geometry into account, the AMO related climate variations are far less important to the recent mass loss than the trend caused by the gradual warming over the past century.

### 1 Introduction

Huss et al. (2010) base the modeled mass balance series for the 100-year period 1908–2008 on in-situ measurements and ice volume changes derived from sequential Digital Elevation Models (DEMs) of the studied glaciers (with a total of 3 to 9 DEMs per glacier). These measurements constrain a degree-day mass balance model that is used to compute daily mass balance values for the studied glaciers on a 25×25 m grid. For each year Huss et al. (2010) calculate the specific mass balance by dividing the total mass balance of the glacier surface by the glacier area of that year. They show that the specific mass balance of the Swiss glaciers was mostly negative for the last century, leading to a total volume loss of 14 km<sup>3</sup> for the 30 glaciers from 1908 to 2008.

In the second part of the paper, the authors link the specific mass balance anomalies to climatic fluctuations and show a statistical correspondence between the AMO and the mean mass balance anomalies of the 30 glaciers. They fit a combination of a sinusoidal and a linear trend to the specific mass balance anomaly and conclude that about half of the loss of ice mass over the most recent decade can be attributed to the 65-year period variation that is superimposed on the negative linear trend.

2476




Supplementary material related to this article is available online at:  
<http://www.the-cryosphere-discuss.net/4/2475/2010/tcd-4-2475-2010-supplement.pdf>.

5 *Acknowledgements.* We would like to thank M. van de Broeke and F. Paul for their useful comments.

## References

- Adhikari, S. and Huybrechts, P.: Numerical modelling of historical front variations and the 21st-century evolution of Glacier AX010, Nepal Himalaya, *Ann. Glaciol.*, 50, 27–34, 2009. 2477
- Brugger, K. A.: The non-synchronous response of Rabots Glaciär and Storglaciären, northern Sweden, to recent climate change: a comparative study, *Ann. Glaciol.*, 46, 275–282, 2007. 2477
- 10 Elsberg, D. H., Harrison, W. D., Echelmeyer, K. A., and Krimmel, R. M.: Quantifying the effects of climate and surface change on glacier mass balance, *J. Glaciol.*, 47, 159, 649–658, 2001. 2477
- 15 Harrison, W. D., Cox, L. H., Hock, R., March, R. S., and Pettit, E. C.: Implications for the dynamic health of a glacier from comparison of conventional and reference-surface balances, *Ann. Glaciol.*, 50, 25–30, 2009. 2477
- Huss, M., Hock, R., Bauder, A., and Funk, M.: 100-year mass changes in the Swiss Alps linked to the Atlantic Multidecadal Oscillation, *Geophys. Res. Lett.*, 37, doi:10.1029/2010GL042616, 2010. 2476, 2477, 2478
- 20 Jóhannesson, T., Raymond, C., and Waddington, E.: Time-scale for adjustment of glaciers to changes in mass balance, *J. Glaciol.*, 35, 355–369, 1989. 2477
- Machgut, H., Paul, F., Hoelzle, M., and Haerberli, W.: Distributed glacier mass balance modelling as an important component of modern multi-level glacier monitoring, *Ann. Glaciol.*, 43, 335–343, 2006. 2478
- 25 Nemeč, J., Huybrechts, P., Rybak, O., and Oerlemans, J.: Reconstruction of the annual balance of Vadret da Morteratsch, Switzerland, since 1865, *Ann. Glaciol.*, 50, 126, 2009. 2477, 2478
- Oerlemans, J.: Climate sensitivity of Franz Josef Glacier, New Zealand, as revealed by numerical modelling, *Arctic Alpine Res.*, 29, 233–239, 1997. 2477
- 2479
- Oerlemans, J.: *Glaciers and Climate Change*, AA Balkema Publishers, 2001. 2481
- Oerlemans, J.: *Minimal Glaciers Models*, Igitur, Utrecht Publishing & Archiving Services, Universiteitsbibliotheek Utrecht, 2008. 2477
- 5 Paul, F.: The influence of changes in glacier extent and surface elevation on modeled mass balance, *The Cryosphere Discuss.*, 4, 737–766, doi:10.5194/tcd-4-737-2010, 2010. 2477, 2478
- Paul, F., Escher-Vetter, H., and Machgut, H.: Comparison of mass balances for Vernagtferner obtained from direct measurements and distributed modeling, *Ann. Glaciol.*, 50, 169–177, 2009. 2478
- 10 Zemp, M., Haerberli, W., Hoelzle, M., and Paul, F.: Alpine glaciers to disappear within decades?, *Geophys. Res. Lett.*, 33, L13504, doi:10.1029/2006GL026319, 2006. 2477



**Fig. 1.** Schematic response of the mass balance (blue) and glacier area (black) to an idealized climate change (red). Due to the slow change in glacier area the mass balance acts as a high-pass filter (cf. Oerlemans, 2001, Section 9.7).