



This discussion paper is/has been under review for the journal The Cryosphere (TC).
Please refer to the corresponding final paper in TC if available.

A numerical study of glacier advance over deforming till

G. J.-M. C. Leysinger Vieli^{1,3} and G. H. Gudmundsson²

¹Versuchsanstalt für Wasserbau, Hydrologie und Glaziologie (VAW), ETH Zürich, Switzerland

²British Antarctic Survey, Natural Environment Research Council, Cambridge, UK

³Durham University, Department of Geography, Durham, UK

Received: 18 June 2010 – Accepted: 21 June 2010 – Published: 30 June 2010

Correspondence to: G. J.-M. C. Leysinger Vieli (g.j.m.c.leysinger-vieli@durham.ac.uk)

Published by Copernicus Publications on behalf of the European Geosciences Union.

TCD

4, 823–864, 2010

Glacier advance over deforming till

G. J.-M. C. Leysinger
Vieli and
G. H. Gudmundsson

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Abstract

The advance of a glacier over a deforming sediment layer is analysed numerically. We treat this problem as a contact problem involving two slowly-deforming viscous bodies. The surface evolution of both bodies, and that of the contact interface between them, 5 is followed in time. Using various different till rheologies we show how the mode of advance depends on the relative effective viscosities of ice and till. Three modes of advances are observed: 1) overriding, where the glacier advances through ice deformation only and without deforming the sediment; 2) plug-flow, where the sediment is 10 strongly deformed, the ice moves forward as a block and a bulge is built in front of the glacier; and 3) mixed-flow, where the glacier overrides deforming sediment. An 15 inverse depth-age relationship is obtained for a glacier advance by both overriding and mixed-flow. An additional model experiment, using a till with near plastic rheology, shows that the contrast in effective viscosity between ice and till is the single most important model parameter defining the mode of advance and the resulting thickness 20 distribution of the till. Furthermore, the model calculations imply that measurements of sediment thickness and sediment deformation taken close to the glacier front significantly overestimate the average sediment thickness and displacement due to sediment deformation. Given sufficiently large contrast in effective viscosity between ice and till, a sediment bulge is formed in front of the glacier. During glacier advance, the bulge 25 quickly reaches a steady state form strongly resembling single-crested push moraines. Inspection of particle paths within the sediment bulge, shows the material particle of the till to travel at different speed to that of the bulge itself, and the push moraine to advance as a form-conserving non-linear wave.

1 Introduction

25 The advance of glaciers has been observed by measurements of the snout position and records of glacier mass-balance (Oerlemans, 1989; Haeberli and Beniston, 1998; Oer-

Glacier advance over deforming till

G. J.-M. C. Leysinger
Vieli and
G. H. Gudmundsson

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



lemans, 2001), by geomorphological observations of the glacier forefield deformation, as well as by using traces in the landscape and historical glacier records to interpret the glacier fluctuations in the past (e.g., Drewry, 1986; Benn and Evans, 1998; Haeberli and Beniston, 1998; Oerlemans, 2001). Observed length changes have been used to extract climate signals (e.g., Boulton, 1986; Oerlemans, 2005; Beedle et al., 2009).

Glaciers advance by: 1) ice deformation; 2) deformation of the glacier bed; 3) sliding at the ice-bed interface; or 4) a combination of these processes (e.g., Benn and Evans, 1998). A glacier advance by ice deformation only (ice frozen to bed) and no surface melt would lead to an advance by “overriding”, which means that the advancing glacier snout rolls its surface over the glacier forefield, potentially giving rise to inverse depth-age relationship. For an advance by 2) or 3) only, the ice mass is moving as a block (“plug-flow”). A glacier advance over a deformable till can lead to the formation of “push moraines”, where the glacier appears to be bulldozing over and through the ice-marginal sediments (Boulton, 1986; Van der Wateren, 1995; Benn and Evans, 1998; Bennett, 2001; Motyka and Echelmeyer, 2003).

In contrast to the larger number of observational studies on the advance of glaciers over deforming substrate, there is a conspicuous lack of any numerical studies. Although a variety of numerical models have been used for calculating temporal changes in front position (e.g., Greuell, 1992; Schlosser, 1997; Schmeits and Oerlemans, 1997; Wallinga and Van de Wal, 1998; Oerlemans, 2001; Smedt and Pattyn, 2003), none of these models attempt to describe the details of ice flow, or the mechanical interactions between ice and till in the vicinity of the glacier terminus. In this area assumptions commonly used in flow modelling of large ice masses, such as the shallow ice approximation (SIA) model (Hutter, 1983) do not apply. Furthermore, we are not aware of any numerical work where the time-dependent deformation of both the ice and the till is calculated in a fully coupled way using the full-set of the momentum equations.

In this paper we aim to answer the question of how the front of a glacier residing on a sediment advances and to investigate the details of the flow at an advancing glacier front, by using a two dimensional finite-element model designed for this purpose. This

Glacier advance over deforming till

G. J.-M. C. Leysinger
Vieli and
G. H. Gudmundsson

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



study extends and complements a previous study of ours (Leysinger Vieli and Gudmundsson, 2004) focusing on the advance of glacier over non-deforming substrate.

Our modelling approach is generic in its nature and we do not try to simulate the advance of any one particular glacier. Rather, we are interested in the overall characteristics of the flow of ice and till, and the mechanical interaction between advancing glacier and the underlying till. Of special interest is the flow regime in the vicinity of the snout where assumption of most previous models break down. We aim at answering how till deformation is induced by the flow of a glacier, how the thickness distribution of till changes as a glacier advances over it, and where till deforms primarily through shearing as opposed to horizontal compression. We study numerically the formation of push moraines and suggest that push moraines may be considered to be a type of form-conserving non-linear till waves.

2 Modelling the advance of a glacier

For isotropic ice a viscous rheology formulated in Glen's flow law is commonly used (Glen, 1955). For subglacial till various different rheological models have been proposed (e.g., Fowler, 2002; Clarke, 2005). The rheology of the till has been suggested to behave like a linear or slightly nonlinear viscous fluid (Alley et al., 1986, 1987, 1989; Boulton and Hindmarsh, 1987; Alley, 1989; MacAyeal, 1989, 1992), where the strain rates $\dot{\epsilon}$ are related to the effective stress τ through

$$20 \quad \dot{\epsilon} \propto \tau^m, \quad (1)$$

with a stress exponent m in the range of $1 \leq m \leq 5$ (e.g., Kamb, 2001), such as $m=1.33$ as fitted by Boulton and Hindmarsh (1987) for the till beneath Breidamerkurjökull or $m=3.04$ as optimised by Gudmundsson (2007) for the subglacial till of Rutford Ice Stream. It has also been pointed out that since till is a granular medium its rheological behaviour should fall within the range commonly exhibited by such materials (Iverson et al., 1998; Tulaczyk et al., 2000a,b; Truffer et al., 2000; Kamb, 2001; Kavanaugh and

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Clarke, 2006). Therefore, it should show Coulomb-plastic rheology (termed “triboplastic” by Kamb, 2001), with a yield stress which is controlled by intergranular friction and which depends only slightly, if at all, on the strain rate (Kamb, 1991; Iverson et al., 1998).

5 A perfectly plastic till rheology can be obtained by letting the flow exponent m in Eq. (1) go to infinity. For stresses below the *yield stress* (τ_0), deformation is negligible and for stresses exceeding the yield stress the deformation would be infinitely large. As a result, the stress in the sediment never exceeds the yield stress (Paterson, 1994, p. 98; Van der Veen, 1999, p. 145). Kamb (2001) suggests that till rheology, as observed for ice streams B, C and D (now Whillans, Kamb and Bindschadler), is best described as “imperfect triboplastic”, because of an appreciable strain rate ($\dot{\epsilon}$) dependence on the stress (σ), which he expresses in a highly nonlinear flow law with an exponent $m \approx 40 \pm 20$. Here we use the term *near perfect plastic* to refer to such a highly nonlinear rheology.

15 As pointed out by Fowler (2002, 2003) a rigid/perfectly plastic material can deform over long time-scales as if viscous. More recently the controversy is dealt with by investigating constitutive models of subglacial sediments which allow for different deformation under different conditions (e.g., Sane et al., 2008; Altuhafi et al., 2009). Here we consider cases corresponding to both non-linear viscous and near perfect plastic rheology.

20

2.1 Model description

25 A time-dependent two-dimensional numerical plane-strain model is used to study the advance of a glacier over a layer of deforming till. The model treats the glacier and the underlying till as two independent slowly-deforming viscous bodies. The surface evolutions of both bodies is followed with time and contacts between the bodies automatically determined. In each time step nodes that enter in or out of contact are found and the boundary conditions changed accordingly. Unstructured grids and automated remeshing is employed to limit element distortion. The full set of the equilibrium

Glacier advance over deforming till

G. J.-M. C. Leysinger
Vieli and
G. H. Gudmundsson

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

equations is solved, allowing for accurate estimates of stress and strain.

2.1.1 Ice rheology

Glen's flow law (Glen, 1955; Steineman, 1958)

$$\dot{\epsilon}_{ij} = A \tau^{n-1} \sigma_{ij}^{(d)}, \quad (2)$$

5 is used to describe the relationship between deviatoric stresses and strain rates in the ice as well as in the sediment layer. Here $\dot{\epsilon}_{ij}$ are the components of the strain rate tensor, and $\sigma_{ij}^{(d)}$ are the components of the deviatoric stress tensor

$$\sigma_{ij}^{(d)} = \sigma_{ij} - \frac{1}{3} \delta_{ij} \sigma_{kk}, \quad (3)$$

10 where σ_{ij} are the components of the stress tensor and δ_{ij} is the Kronecker delta. The effective stress τ is defined by

$$\tau^2 = \frac{1}{2} \sigma_{ij}^{(d)} \sigma_{ij}^{(d)}. \quad (4)$$

15 The rate factor A and the stress exponent n are material parameters. For the ice body the flow law exponent is chosen as $n=3$. Values for other parameters are given below. All parameters are made non-dimensional by the use of appropriate scalings (see Sect. 2.1.4 and Appendix A).

2.1.2 Till rheology

The till is treated in a similar way as the ice, namely as an incompressible non-linear viscous medium (Glen's flow law, Eq. 2). The rate factor and the stress exponent of the till are denoted by A' and m , respectively. The till density was set to $\rho_s=2200 \text{ kg m}^{-3}$.

20 It is one of the interesting facts of glaciological research that despite decades of work on this subject no general consensus has emerged on the most appropriate numerical

TCD

4, 823–864, 2010

Glacier advance over deforming till

G. J.-M. C. Leysinger
Vieli and
G. H. Gudmundsson

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



values for the modelling parameter m . Values for m ranging from 1 to infinity are commonly used (e.g., Iverson et al., 1998; Kamb, 2001; Clarke, 2005). Here we bracket the range of possible rheologies by using a wide range of values for the rate factor (A') and the stress exponent (m).

5 The till rheology is determined by a number of factors and processes that do not concern us here. We do not consider in detail how the particular numerical values of the rheological parameters of the till are related to grain size, pore water pressure, etc. which is an important question in itself but entirely unrelated to our modelling work.

2.1.3 Model geometry and mass balance distribution

10 The general model geometry is depicted schematically in Fig. 1. The glacier is everywhere in contact with a deformable till layer. The till layer extends sufficiently far to the front of the initial glacier geometry to ensure that while advancing the glacier remains in contact with the till. Initially, the thickness of the till is uniform and set to about 1/20th of the maximum ice thickness. Where the ice is in contact with the till no differential motion between ice and till is allowed. No mass is added or extracted to the sediments 15 in the course of the calculations. In all our experiments the geometry of the base of the till layer is kept fixed, with a steep upper part (45°) and a lower part sloped at 5° as shown in Fig. 1.

We chose a height-dependent mass balance function such as

$$20 \quad \dot{b}(z) = \begin{cases} a_{\text{acc}}(z_s - \text{ELA}) & \text{when } z_s > \text{ELA} \\ 0 & \text{when } z_s \leq \text{ELA} \end{cases} \quad (5)$$

where ELA is the equilibrium line altitude, located at a height of 0.5 dimensionless units (defined below) and z_s is the surface height of the glacier. Above the ELA a positive mass balance gradient a_{acc} of 0.0117 (non-dimensional units) is used. Note that this mass-balance distribution includes altitude-mass balance feedback in the accumulation 25 area only. The use of this particular form of the surface mass-balance distributing is motivated by the requirement that the overall mass balance must remain positive

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

in order to ensure continued advance. Setting the mass-balance to zero for $z_s \leq \text{ELA}$ simplifies our analysis of the movements of material particles in the vicinity of the snout.

An initial glacier geometry was generated for a hard bed by running a SIA model. Once the glacier had advanced to a total length of three times the size of the accumulation area, we switched from using the SIA model and did the rest of the run using the full-system model. Our conclusions are not affected by the details of the starting geometry. The use of SIA in the initial modelling stage is based on practical considerations and a desire to limit computational times involved.

2.1.4 Normalisation

10 We introduce non-dimensional variables for the space coordinates $\{X, Z\}$, the components of the velocity vector $\{U, W\}$, the time $\{T\}$ and the effective stress $\{T_{\text{eff}}\}$ defined as $\{x/[h], z/[h], u/[u_s], w/[u_s], t/([h]/[u_s]), \tau/[\tau_d]\}$, where $[h]$ is a typical mean thickness of a glacier, $[u_s]$ a typical mean horizontal surface velocity and $[\tau_d] = \rho g [h] \sin(\alpha_s)$ the driving stress assuming the same surface slope $\alpha_s = 5^\circ$ as the main basal slope. As

15 we are scaling with the extended inclined ice slab (see Appendix A) $[h]$ corresponds to the thickness of the ice slab and $[u_s]$ to the analytical solution for the surface velocity (Eq. A1). The non-dimensional ice thickness H and mean ice thickness $[H]$ are defined as $h_o/[h]$ and $[h_o]/[h]$, respectively, where h_o is the local and $[h_o]$ the mean ice thickness of the observed or modelled glacier.

20 2.1.5 Numerical solution technique

The model calculations were performed with the commercial finite-element (FE) program MARC (2000) tailored for use in glaciology by Gudmundsson (1999). The code solves the full set of momentum equations and has been used extensively for flow modelling of alpine and of grounded tidewater glaciers (Gudmundsson, 1999; Vieli et al., 2001; Leysinger Vieli and Gudmundsson, 2004). We refer to this model as the *full system* model (FS). The transient evolution of the surface and the ice-sediment interface is

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Glacier advance over deforming till

G. J.-M. C. Leysinger
Vieli and
G. H. Gudmundsson

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

followed using a mixed Lagrange-Euler approach. As the surface evolves, automated remeshing is used when needed to limit mesh distortion (Leysinger and Gudmundsson, 2000; Leysinger Vieli, 2003). The remeshing strategy (Leysinger Vieli, 2003) uses a cubic spline interpolation to reposition nodal points along all singular surfaces. It 5 has been verified that the remeshing algorithm preserves the total volume accurately and that the model results are not affected by smaller time steps or more refined mesh. The FE code has been verified by comparing its numerical results with transient analytical solutions (Gudmundsson, 1997; Adalgeirsdóttir et al., 2000; Raymond et al., 2003; Gudmundsson, 2007) and by comparing the time-dependent response for a linear flow 10 law (Newtonian flow) with the analytical solution (ISMIP-HOM Benchmark experiment F, Pattyn et al., 2008).

2.1.6 Model parameters

We consider two end-member cases of different modes of advance where a) the till is much stiffer than the ice and all the deformation takes place in the ice, and b) the opposite situation where ice is much stiffer than the till and all deformation takes place within 15 the till layer. These different limiting cases of modes of advance can be generated in our numerical model simply by varying the contrast in effective viscosity between ice and till.

After non-dimensionalisation, the results of the numerical runs depend only on the 20 ratio of the rheological parameters. We keep the rate factor A and the stress exponent n fixed and vary the corresponding rheological parameters A' and m of the till. The non-dimensional rate factor for the ice is $A=2$ (Eq. A4), this follows directly from the scalings used, when we set $n=3$. For a soft, highly deforming sediment the ratio between the rate factor A in the ice and A' in the sediment is small ($\ll 1$) and for a highly rigid 25 sediment this ratio is large ($\gg 1$). For the till, values of both $m=3$ and $m=40$ were used. The value $m=40$ is used to simulate a nearly perfect plastic rheology as suggested by Kamb (2001).

To be able to get similar effective viscosities ($\eta=1/2A'\tau^{m-1}$) in the sediment at given stresses with different m we define B for the till as

$$B = A'^{-1/m}. \quad (6)$$

and rewrite Glen's flow law (Eq. 2) as

$$5 \quad \dot{\epsilon}_{ij} = \left(\frac{\tau}{B} \right)^{m-1} \frac{\sigma_{ij}^{(d)}}{B}. \quad (7)$$

Note that the material parameter B has the dimensions of stress, and that increasing B has the effect of decreasing the strain rates for a given state of stress. We refer to B as the *hardness factor* while A is the *rate factor*. For $m \rightarrow \infty$, Eq. (7) shows that the rheology of till corresponds to a perfectly plastic rheology with B being the yield stress

10 τ_0 . Hence, by adopting a high stress exponent such as $m=40$, the sediment rheology goes from a nonlinear viscous to a near perfect plastic one. For such a high stress exponent the till will only deform significantly in regions where the effective stress τ is larger than B . The values used in the model experiments are shown in Table 1.

3 Results

15 As already mentioned, the model is formulated in dimensionless variables and therefore the results shown are all dimensionless. One can easily go from non-dimensional variables to dimensional ones. Our non-dimensional vertical space coordinate Z extends from 0 to 1, as the extend of the grid in z has been chosen to be the same as the thickness $[h]$ of the infinitely extended inclined ice slab (see Appendix A).

20 For example, choosing a rate factor of $A=2.06 \times 10^{-15} \text{ s}^{-1} (\text{kPa})^{-3}$ for temperate ice (Hubbard et al., 1998; Gudmundsson, 1999; Albrecht et al., 2000), a mean thickness $[h]=1000 \text{ m}$ for the ice slab and a slope of $\alpha=5^\circ$ results in a mean horizontal surface velocity of $[u_s]=2A/(n+1)(\rho g \sin(\alpha))^n [h]^{n+1}=1.5665 \times 10^4 \text{ m a}^{-1}$. For this example

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

one non-dimensionless unit in length, velocity and in time, corresponds to 1000 m, $1.5665 \times 10^4 \text{ m a}^{-1}$ and 0.0638 years, respectively. Note that since our non-dimensional vertical space coordinate Z extends from 0 to 1, the mean dimensionless thickness of the modelled glacier [H] is approximately 0.2 Z (e.g. see Figs. 3, 10, 13 and 14), which corresponds to a mean glacier thickness [h_0]=200 m for our example.

3.1 Experiment A: relatively stiff and moderately non-linear till

We start our discussion of the numerical results by considering a case where the till is relatively stiff in comparison to the ice. We use $B=8$, which is a high value as compared to the mean effective basal stress of unity, and $m=3$. This experiment illustrates how 10 glaciers advance over stiff till, and serves as a reference case for the set of experiments described later where the advance gives rise to significant till deformation.

The front advance is shown in Fig. 2a as a series of snapshots depicting the longitudinal surface profiles at different times. The thickness of the till at different times is also plotted in Fig. 2a. As expected, due to the relative stiffness of the till in this experiment, 15 no change in till thickness is visible in the figure.

The surface profile at $t=0$ in Fig. 2a corresponds to the last profile obtained with the SIA model, and the starting profile of the full-system (FS) model. As Fig. 2a shows, the surface profile of the SIA model is markedly different from the one obtained with the FS model. The SIA model calculates velocity from local slope and thickness, and predicts 20 infinitely large velocities for vertical slopes. The FS model includes the effects of horizontal deviatoric stresses on the flow and gives a much more accurate description of the flow field in the vicinity of glacier terminus. Figure 2a shows how the surface slope initially increases, finally resulting in an overhanging snout.

Figure 2b–d show how various material particles (dots in the figures) within the ice 25 move as the front advances. In Fig. 2b positions of selected material particles at the surface, at the base, and within the glacier are shown at different points in time. Due to the relative stiffness of the till, all material particles at the ice-sediment interface stay fixed at their respective locations, and the ice-sediment interface remains parallel to the

[Title Page](#)
[Abstract](#) [Introduction](#)
[Conclusions](#) [References](#)
[Tables](#) [Figures](#)
[◀](#) [▶](#)
[◀](#) [▶](#)
[Back](#) [Close](#)
[Full Screen / Esc](#)

[Printer-friendly Version](#)
[Interactive Discussion](#)

Glacier advance over deforming till

G. J.-M. C. Leysinger
Vieli and
G. H. Gudmundsson

[Title Page](#)

[Abstract](#) [Introduction](#)

[Conclusions](#) [References](#)

[Tables](#) [Figures](#)

◀

▶

◀

▶

[Back](#) [Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



bed (see Fig. 2b–d). Material particles within the ice, however, gradually move down-slope eventually passing the position of the initial glacier front. As the front advances, ice particles that previously were located at the glacier surface come in contact with the till and form a new basal layer of ice giving rise to depth-age inversion.

Figure 3 shows vertical profiles of the non-dimensional horizontal velocity U and effective stress T_{eff} at three different positions along the glacier for the overriding case. At each location the effective stress generally increases with depth. Note that at the snout the effective stress reaches its maximum value at the interface between the ice and the till, and not at the base of the till as is the case further upstream.

3.2 Experiment B: relatively soft and moderately non-linear till

We now consider the situation where the till is considerably softer than the ice by setting $B=0.35$. In this experiment the glacier advances primarily through deformation of the underlying sediment and without any significant deformation of the ice.

In Fig. 4a, glacier and sediment surface profiles are shown at different times. Clearly visible is the formation of a sediment bulge in front of the glacier. The amplitude of the bulge as a function of distance is shown in Fig. 4b. The geometry and the positions of a number of material points within the ice and at the till interface are shown in Fig. 4c–e and the corresponding profiles in Fig. 4a are labelled c, d, and e. Further details of the sediment deformation are given in Figs. 5 and 6.

One of the most interesting results of the experiment is the formation of the sediment bulge in front of the glacier snout. After an initial period of rapid growth the amplitude of the bulge approaches a steady state value (see Fig. 4a). Following the movement of material points within the till (see Figs. 5 and 6) reveals that till particles within the bulge travel at different speed to that of the bulge itself. Thus, the bulge is an example of a till wave. Till particles get incorporated into the till wave for a limited period of time, and are then left behind as the wave travels further. The distribution of horizontal and vertical velocity, and of the effective stress is shown in Fig. 7a–c, respectively. This figure can be compared directly to Fig. 8a–c depicting results from experiment A. Because of the

Glacier advance over deforming till

G. J.-M. C. Leysinger
Vieli and
G. H. Gudmundsson

[Title Page](#)

[Abstract](#) [Introduction](#)

[Conclusions](#) [References](#)

[Tables](#) [Figures](#)

◀

▶

◀

▶

[Back](#) [Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



relative stiffness of the ice, the velocity field of the glacier is close to being constant everywhere, and almost the entire forward motion is due to till deformation (Fig. 7a). The vertical velocity distribution within the till bulge (Fig. 7b) shows how the part of the bulge in contact with the ice is being pressed downward, resulting in negative vertical velocities, while the down-glacier side of the bulge moves upwards. The stresses are transferred into the sediment bulge and therefore contour lines of the effective stress, Fig. 7c, are almost continuous across ice-sediment interface.

The vertical velocity distribution within the sediment bulge is shown in detail in Fig. 9. As the figure shows, the ice in the frontal area presses the till downwards and forwards at the same time. Note that the ice velocity is larger than that of any of the material particles in the till bulge. With time the glacier therefore overrides the till particles previously within the bulge, while at the same time, due to the upward motion in the frontal region, new till particles are incorporated into the bulge.

From the ratio of basal to surface velocity in Fig. 10b we see that at the glacier front the flow is dominated by sediment deformation ($U_b/U_s \approx 1$), and that the contribution of the ice to the total deformation increases in upstream direction. The increase in till deformation towards the glacier front is also seen in the vertical profiles of the horizontal velocity in Fig. 10c–e and in Table 2. The observed distribution of the effective stress in the three vertical profiles is similar to the one obtained for experiment A with the exception of the frontal profile, where the highest shear stress is found at the sediment base (Fig. 10e) and not at the ice-sediment interface.

3.3 Experiment C: moderately stiff and moderately non-linear till

We set $B=2^{-1/3} \approx 0.79$, giving an equal rate factor within the ice and the till. The advance now follows through a mixture of both ice and till deformation (see Figs. 11a–e and 12). In this experiment, flow features that were only seen either in experiment A or B act in combination. As the ice advances over the till, material particles previously at the surface come in contact with the till and form a new basal ice layer. At the same time the till deforms and we see a till wave propagating with the snout in a similar

fashion as seen in experiment B (Fig. 11a–e).

Detailed view of the till deformation is given Fig. 12. The advance of the glacier causes shearing within the till, combined with a horizontal compression in the snout region. As in experiment B a sediment bulge is formed, but due to the larger stiffness of the till the bulge is about 8 times smaller.

The ratio of basal to surface velocity as a function of distance is shown in Fig. 13b. The ratio increases sharply towards the snout implying that measurements of till deformation and surface velocity in that area are not reliable estimates of the glacier wide contribution of basal motion to the forward motion of a glacier. The effective stress distribution shown in Fig. 13c–e also reveals that the magnitude and distribution of stresses within the till in the terminus area (Fig. 13e) differ strongly from those found further up-glacier. In the terminus area, the stresses in the till decrease with depth rather than increasing as is the case elsewhere.

4 Nonlinear viscous vs. near perfect plastic rheology

In the above experiments glacier advance over deforming till was modelled using a stress exponent $m=3$ (Eq. 7). Here we repeat those experiments for a near perfect plastic till rheology by choosing a flow law exponent $m=40$ (Kamb, 2001). We performed a number of experiments using parameter sets that produced both relatively stiff and moderately soft till rheologies.

The main effect of increasing the value of m is to sharpen the contrast between those regions where the effective stress, τ , in the till is either similar to or smaller than the hardness factor B (see Eq. 7). Where $\tau>B$ the till deforms readily, but hardly at all where $\tau<B$. We found in all our experiments that the highest effective stresses in the till were reached in the terminus area. One of the consequences of increasing the value of m from 3 to 40, while keeping B fixed, was therefore to raise the ratio between basal motion and surface velocity in those areas where $\tau>B$ (see Fig. 14). For example, for $B=0.42$ (Fig. 14c–e) at approx. $1/20[H]$ behind the glacier front 62% of the horizontal

Glacier advance over deforming till

G. J.-M. C. Leysinger
Vieli and
G. H. Gudmundsson

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Glacier advance over deforming till

G. J.-M. C. Leysinger
Vieli and
G. H. Gudmundsson

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)
[Back](#)[▶](#)
[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

deformation is due to till deformation, 33% at $2/3[H]$ and no till deformation is taking place at only 5 times the mean ice thickness, $[H]$, behind the front (Table 2).

By increasing the hardness factor B from 0.42 to 0.44, the value used in experiment B described above, the point of transition from non-deforming to deforming till migrates further towards the snout. The contribution of sediment deformation for $B=0.44$ to the total horizontal deformation decreased to 56% at $1/20[H]$, 23% at $2/3[H]$ and 0% at $5[H]$ behind the glacier front (Table 2).

The till thickness generally increases towards the terminus. This can be seen clearly in Fig. 14c–e where the till thickness is indicated by a horizontal dotted line (note different vertical scales). Note that since the initial prescribed till thickness was everywhere uniform, this thickening of the till in the terminus area is solely caused by the flow of the glacier and the mechanical interaction between the ice and the till. Note also that since we do not describe in the model any sediment movements due to action of water and/or erosion, all sediment deformation is solely due to shearing through the till column and horizontal compression towards the terminus.

The formation of a sediment bulge in front of the snout was also observed for $m=40$ just as it was for $m=3$. Although the details of the form and the shape of the bulge change as modelling parameters such as m and B are varied, the genesis of this bulge, hence, does not depend on the exact parameter values chosen and is, in this sense, a robust modelling feature.

5 Discussion

Our model was not designed at the outset to investigate the formation of push moraines. As illustrated by the findings of a study of the large-scale push moraine of Taku Glacier, Alaska, by Kuriger et al. (2006), the shape and form of the sediment bulge forming in front of the glacier terminus, however, bears strong similarities to those of push moraines. Furthermore, we find the value of the till hardness factor B to be the single most important modelling parameter affecting the mode of till deformation. This

agrees favourably with Kuriger et al. (2006) finding that decreased effective frictional resistance of the sediment is the key parameter affecting rates of internal sediment deformation.

Numerous descriptions of the morphology of push/squeeze moraines and conceptual models of their formation are found in the literature (e.g., Price, 1970; Kälin, 1971; Meyer, 1983; Van der Wateren, 1995; Winkler and Nesje, 1999; Bennett, 2001; Evans and Hiemstra, 2005; Kuriger et al., 2006; Evans, 2009). To our knowledge the model presented here is, however, the first numerical model of glacier advance over till that produces a feature closely resembling push moraines as they are observed.

The process of advance producing a push moraine is often observed to be a seasonal process with advance in late winter while the till is frozen and squeezing out of water-soaked till from beneath the glacier during the early summer melt period (e.g., Price, 1970; Boulton, 1986; Evans and Hiemstra, 2005). By varying the hardness of the sediment seasonally our model should be able to reproduce such seasonal push/squeeze moraines. Truffer et al. (2009) inferred from field measurements and numerical modelling that the mode of glacier advance changes on seasonal timescales on Taku Glacier, Alaska, from internal deformation (winter) to plug-flow (summer), depending only on small changes in the basal boundary condition.

The numerical modelling experiments show how a glacier advancing over a soft till causes shearing within the till that, depending on the value of the stress exponent m , gives rise to spatially highly variable rates of basal motion (see Figs. 13b and 14b). In all our modelling experiments, the highest ratio of basal motion to surface velocity was found in the terminus area. Measurements of basal motion in that area are therefore potentially not representative for a glacier as a whole. An example for such a measurement site is the tunnel system dug in the ice about 1 to 2 m above the glacier sole and approximately 24 to 44 m behind the terminus of Breidamerkurjökull, Iceland, described by Boulton (1979). By assuming no slip at the glacier sole, the displacement due to sediment deformation at this site was between 95% and 80% of the forward movement of the glacier (Boulton and Hindmarsh, 1987). If we assume an average

Glacier advance over deforming till

G. J.-M. C. Leysinger
Vieli and
G. H. Gudmundsson

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

thickness of 400 m for Breidamerkurjökull (Björnsson et al., 2001; using slope as in our model), then the position behind the terminus is about 1/16 to 1/9 of the typical mean ice thickness [h_0] behind the glacier front. Their measured deformation, especially at the higher end, is in the same range as obtained in this study for the plug-flow case (experiment B).

5 Mechanical interaction between ice and till alone is sufficient to give rise to till thickening with time in the terminus area. The thickening is caused by combined action of shearing through the till column and horizontal compression towards the terminus. No other potential modes of sediment transport, i.e. suspended sediment transport or 10 entrainment of sediment into the ice, were described in the model. Including erosion and/or hydraulic transport would modify long-term evolution of the sediment thickness (e.g., Motyka et al., 2006). Similarly, the moraine structure in front of the terminus would be modified by pro-glacial fluvial and eolian processes.

If sufficiently soft, the till is extruded from underneath the glacier towards the terminus. 15 This happens through direct vertical mechanical action of the ice on the underlying sediments. Such moraine forming by extrusion flow has been observed in nature and discussed in the literature (e.g., Price, 1970; Sharp, 1984; Evans and Hiemstra, 2005). The shape and the position of the bulge as depicted for example in Figs. 5 and 6 resembles push/squeeze moraines (e.g., Price, 1970; Evans and Hiemstra, 2005; 20 Evans, 2009). What has been touched on by Van der Wateren (1995) but possibly not been fully realised since, and is also suggested by our modelling, is the possibility that the push moraines are a form of non-linear shape-conserving waves where horizontal movement of individual till particles is insignificant in comparison to the total distance travelled by the moraine. We also emphasise that the formation of the push moraine 25 does not require the terminus to be in a steady state for any significant period of time. In fact in our modelling approach the terminus is never in steady state but advances continuously.

Where the till is too stiff to be significantly deformed, the glacier advances through “overriding”. In nature examples of this type of advance can be found in advancing

Glacier advance over deforming till

G. J.-M. C. Leysinger
Vieli and
G. H. Gudmundsson

[Title Page](#)

[Abstract](#) [Introduction](#)

[Conclusions](#) [References](#)

[Tables](#) [Figures](#)

◀

▶

◀

▶

[Back](#) [Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



glaciers in sub-/polar regions where melting is negligible (e.g. Glaciers on Axel Heiberg Island, Canadian Arctic Archipelago and in the Dry Valleys, Antarctica). A possible example of a glacier advance by overriding is Crusoe Glacier from Axel Heiberg Island where the folding structure revealed in a photo taken of the west front by Alean (2008)

5 could be interpreted as the inverse layering obtained from overriding. Another example of advance by overriding are active rock glaciers, which have insignificant mass loss through surface ablation. On the other hand, for glaciers with surface melting in alpine environments a realistic shape of an advancing glacier front would be less steep. Previous model calculations for the evolution of advancing glaciers residing on a bedrock 10 with a realistic mass balance distribution, done by Leysinger Vieli and Gudmundsson (2004), indeed show that the front geometry is less curved than the one with zero mass balance (Fig. 15).

Raising the flow exponent m from 3 to 40 changes the vertical distribution of the horizontal velocities within the sediment. For this near perfect plastic rheology the 15 velocity increase is largest close to the sediment bed (Fig. 14d), whereas for $m=3$ the increase is approximately linear over depth. The main effect of increasing the value of the stress factor, however, is to increase the relative contrast in the contribution of basal motion to the overall forward motion between different areas. The result is a more pronounced spatial variation in distribution of basal velocities.

20 6 Conclusions

By modelling the advance of a glacier over a deforming sediment layer numerically as a contact problem involving two viscous bodies, we have shown how push moraines can be formed. During glacier advance, till particles enter and then subsequently leave the push moraine. After an initial phase the size and the shape of the push moraine 25 does not change with time. The formation of the push moraine is a “robust” modelling result in the sense that its genesis does not depend on a fortuitous choice of parameters but occurs in the model for a wide range of parameter values.

Glacier advance over deforming till

G. J.-M. C. Leysinger
Vieli and
G. H. Gudmundsson

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Although we only modelled a glacier in its advancing stage, the formation of the moraine and analysis of the trajectories of the till particles suggest that such a push moraine would also form during a stationary stage or even during a retreat phase.

Despite the model parameters describing the till rheology (m and B), as well as the initial till thickness distribution, to be spatially uniform, resulting distribution of basal motion is highly non-uniform. Highest rates of basal motion, as compared to surface velocity, are found in the vicinity of the terminus area. Rates of basal motion measured close to a glacier terminus are not reliable estimates of basal motion a few ice thicknesses further up-glacier.

10 Appendix A

Scaling of the flow law

A For an infinitely extended inclined ice slab with the thickness [h] and the slope α the analytical solution for the surface velocity, due to internal deformation, is given by

$$15 \quad [u_d] = \frac{2A}{n+1} [\tau]^n [h], \quad (A1)$$

where the basal shear stress is given by $\tau = \rho g [h] \sin(\alpha)$.

To get dimensionless variables we scale dimensional quantities with corresponding quantities for the infinite extended inclined slab, that is: lengths with the thickness [h], velocities with the deformation velocity $[u_d]$ (Eq. A1), stresses with the basal shear stress $[\tau]$ and strain rates with $[u_d]/[h]$. Glen's flow law (Eq. 2) is now written as

$$20 \quad \dot{\epsilon}_{ij} \frac{[h]}{[u_d]} = A \tau^{n-1} \sigma_{ij}^{(d)} \frac{[h]}{[u_d]} \\ = \frac{\tau^{n-1} \sigma_{ij}^{(d)}}{\frac{2}{n+1} [\tau]^n}. \quad (A2)$$

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

The term on the left hand side of Eq. (A2) is the non-dimensional strain tensor \dot{E}_{ij} . The non-dimensional stress tensor and effective stress are given by $\Sigma_{ij}^{(d)} = \sigma_{ij}^{(d)} / [\tau]$ and $T_{\text{eff}} = \tau / [\tau]$, respectively. Equation (A2) can be rewritten as

$$\dot{E}_{ij} = \frac{n+1}{2} T_{\text{eff}}^{n-1} \Sigma_{ij}^{(d)}, \quad (\text{A3})$$

5 which is the non-dimensional formulation of Glen's flow law (Eq. 2). In this formulation the non-dimensional rate factor is now given by

$$A = (n+1)/2. \quad (\text{A4})$$

Acknowledgements. This study was financed by ETH internal research grant number TH-11/99-2 and supported by the Section of Glaciology at VAW ETH and by the Royal Society

10 BP Dorothy Hodgkin Fellowship, and by the British Antarctic Survey, PSPE program. We would like to thank M. Lüthi for his helpful python programs to extract data from the output files of the commercial FE software MARC and A. Vieli for many constructive discussions.

References

15 Adalgeirsdóttir, G., Gudmundsson, G. H., and Björnsson, H.: The response of a glacier to a surface disturbance, a case study on Vatnajökull ice cap, *Ann. Glaciol.*, 31, 104–110, 2000. 831

16 Albrecht, O., Jansson, P., and Blatter, H.: Modelling glacier response to measured mass balance forcing, *Ann. Glaciol.*, 31, 91–96, 2000. 832

20 Alean, J.: Glaciers online, Axel Heiberg Island, Crusoe Glacier, http://www.swisseduc.ch/glaciers/axel_heiberg/crusoe_glacier/crusoe_front_west/index-en.html?id=2, last access: 18 June 2010, 2008. 840

25 Alley, R. B.: Water-pressure coupling of sliding and bed deformation: I. Water system, *J. Glaciol.*, 35, 108–118, 1989. 826

Alley, R. B., Blankenship, D. D., Bentley, C. R., and Rooney, S. T.: Deformation of till beneath Ice Stream B, West Antarctica, *Nature*, 322, 57–59, 1986. 826

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Glacier advance over deforming till

G. J.-M. C. Leysinger
Vieli and
G. H. Gudmundsson

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Alley, R. B., Blankenship, D. D., Rooney, S. T., and Bentley, C. R.: Till beneath Ice Stream B 4. A coupled ice-till flow model, *J. Geophys. Res.*, 92, 8931–8940, 1987. 826

Alley, R. B., Blankenship, D. D., Rooney, S. T., and Bentley, C. R.: Water-pressure coupling of sliding and bed deformation: III. application to ice stream B, Antarctica, *J. Glaciol.*, 35, 130–139, 1989. 826

Altuhafi, F. N., Baudet, B. A., and Sammonds, P.: On the time-dependent behaviour of glacial sediments: a geotechnical approach, *Quaternary Sci. Rev.*, 28, 693–707, 2009. 827

Beedle, M. J., Menounos, B., Luckman, B. H., and Wheate, R.: Annual push moraines as climate proxy, *Geophys. Res. Lett.*, 36, L20501, doi:10.1029/2009GL039533, 2009. 825

Benn, D. and Evans, D.: *Glaciers and Glaciation*, Arnold, a member of the Hodder Headline Group, 338 Euston Road, London NW1 3BH, 1998. 825

Bennett, M. R.: The morphology, structural evolution and significance of push moraines, *Earth-Sci. Rev.*, 53, 197–236, 2001. 825, 838

Björnsson, H., Pálsson, F., and Gudmundsson, S.: Jökulsárlón at Breidamerkursandur, Vatnajökull, Iceland: 20th century changes and future outlook, *Jokull*, 50, 1–18, 2001. 839

Boulton, G. S.: Processes of glacier erosion on different substrata, *J. Glaciol.*, 23, 15–37, 1979. 838

Boulton, G. S.: Push-moraines and glacier-contact fans in marine and terrestrial environments, *Sedimentology*, 33, 677–698, 1986. 825, 838

Boulton, G. S. and Hindmarsh, R. C. A.: Sediment deformation beneath glaciers: rheology and geological consequences, *J. Geophys. Res.*, 92, 9059–9082, 1987. 826, 838

Clarke, G. C. K.: Subglacial processes, *Annu. Rev. Earth Pl. Sc.*, 33, 247–276, doi:10.1146/annurev.earth.33.092203.122621, 2005. 826, 829

Drewry, D.: *Glacial Geologic Processes*, Edward Arnold Publishers, London, 1986. 825

Evans, D. J. A.: Controlled moraines: origins, characteristics and palaeoglaciological implications, *Quaternary Sci. Rev.*, 28, 183–208, 2009. 838, 839

Evans, D. J. A. and Hiemstra, J. F.: Till deposition by glacier submarginal, incremental thickening, *Earth Surf. Proc. Land.*, 30, 1633–1662, 2005. 838, 839

Fowler, A. C.: Rheology of subglacial till, *J. Glaciol.*, 48, 631–632, 2002. 826, 827

Fowler, A. C.: On the rheology of till, *Ann. Glaciol.*, 37, 55–59, 2003. 827

Glen, J. W.: The creep of polycrystalline ice, *Proc. R. Soc. Lon. Ser.-A*, 228, 519–538, 1955. 826, 828

Greuell, W.: Hintereisferner, Austria: mass-balance reconstruction and numerical modelling of

Glacier advance over deforming till

G. J.-M. C. Leysinger
Vieli and
G. H. Gudmundsson

[Title Page](#)

[Abstract](#) [Introduction](#)

[Conclusions](#) [References](#)

[Tables](#) [Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



the historical length variations, *J. Glaciol.*, 38, 233–244, 1992. 825

Gudmundsson, G. H.: Basal-flow characteristics of a linear flow sliding frictionless over small bedrock undulations, *J. Glaciol.*, 43, 71–79, 1997. 831

Gudmundsson, G. H.: A three-dimensional numerical model of the confluence area of Unter-aargletscher, Bernese Alps, Switzerland, *J. Glaciol.*, 45, 219–230, 1999. 830, 832

Gudmundsson, G. H.: Tides and the flow of Rutford Ice Stream, West Antarctica, *J. Geophys. Res.*, 112(F4), F04007, doi:10.1029/2006JF000731, 2007. 826, 831

Haeberli, W. and Beniston, M.: Climate change and its impact on glaciers and permafrost in the Alps, *Ambio*, 27, 258–265, 1998. 824, 825

Hubbard, A., Blatter, H., Nienow, P., Mair, D., and Hubbard, B.: Comparison of three-dimensional model for glacier flow with field data from Haut Glacier d'Arolla, Switzerland, *J. Glaciol.*, 44, 368–378, 1998. 832

Hutter, K.: *Theoretical Glaciology; Material Science of Ice and the Mechanics of Glaciers and Ice Sheets*, D. Reidel Publishing Company/Tokyo, Terra Scientific Publishing Company, 1983. 825

Iverson, N. R., Hooyer, T. S., and Baker, R. W.: Ring-shear studies of till deformation: coulomb-plastic behavior and distributed strain in glacier beds, *J. Glaciol.*, 44, 634–642, 1998. 826, 827, 829

Kälin, M.: The active push moraine of the Thompson Glacier, Axel Heiberg Island, Canadian Arctic Archipelago, Canada, Ph.D. thesis, ETH Zürich, Nr. 4671, 68 pp., 1971. 838

Kamb, B.: Rheological nonlinearity and flow instability in the deforming bed mechanism of ice stream motion, *J. Geophys. Res.*, 96, 16585–16595, 1991. 827

Kamb, B.: Basal zone of the West Antarctic Ice Streams and its role in lubrication of their rapid motion, in: *The West Antarctic Ice Sheet. Behavior and Environment*, edited by: Alley, R. B. and Bindschadler, R. A., vol. 77 of *Antarctic Research Series*, 157–199, American Geophysical Union, 2001. 826, 827, 829, 831, 836

Kavanaugh, J. L. and Clarke, G. K. C.: Discrimination of the flow law for subglacial sediment using in situ measurements and an interpretation model, *J. Geophys. Res.*, 111, F01002, doi:10.1029/2005JF000346, 2006. 826

Kuriger, E., Truffer, M., Motyka, R. J., and Bucki, A. K.: Episodic reactivation of large-scale push moraines in front of the advancing Taku Glacier, Alaska, *J. Geophys. Res.*, 111, F01009, doi:10.1029/2005JF000385, 2006. 837, 838

Leysinger, G. J.-M. C. and Gudmundsson, G. H.: Are higher-order numerical models needed

Glacier advance over deforming till

G. J.-M. C. Leysinger
Vieli and
G. H. Gudmundsson

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Price, R. J.: Moraines at Fjallsjökull, Iceland, *Arctic Alpine Res.*, 2, 27–42, 1970. 838, 839

Raymond, M., Gudmundsson, G. H., and Funk, M.: Non-linear finite-amplitude transfer of basal perturbations to a glacier, in: *Geophysical Research Abstracts*, EGS-AGU-EUG Joint Assembly 2003, 5, abstract, 2003. 831

5 Sane, S. M., Desai, C. S., Jenson, J. W., Contractor, D. N., Carlson, A. E., and Clark, P. U.: Disturbed State constitutive modeling of two Pleistocene tills, *Quaternary Sci. Rev.*, 27, 267–283, 2008. 827

10 Schlosser, E.: Numerical simulation of fluctuations of Hintereisferner, Ötztal Alps, since AD 1850, *Ann. Glaciol.*, 24, 199–202, 1997. 825

Schmeits, M. J. and Oerlemans, J.: Simulation of the historical variations in length of Unterer Grindelwaldgletscher, Switzerland, *J. Glaciol.*, 43, 152–164, 1997. 825

15 Sharp, M. J.: Annual moraine ridges at Skalafellsjökull, South-East Iceland, *J. Glaciol.*, 30, 82–93, 1984. 839

Smedt, B. D. and Pattyn, F.: Numerical modelling of historical front variations and dynamic response of Sofiyevskiy glacier, Altai mountains, Russia, *Ann. Glaciol.*, 37, 143–149, 2003. 825

20 Steineman, S.: Experimentelle Untersuchungen zur Plastizität von Eis, *Geotechnische Serie Nr. 10, Beiträge zur Geologie der Schweiz.*, Kommissionsverlag Kümmerli & Frey AG, Geographischer Verlag, Bern, 1958. 828

Truffer, M., Harrison, W. D., and Echelmeyer, K. A.: Glacier motion dominated by processes deep in underlying till, *J. Glaciol.*, 46, 213–221, 2000. 826

25 Truffer, M., Motyka, R. J., Hekkers, M., Howat, I. M., and King, M. A.: Terminus dynamics at an advancing glacier: Taku Glacier, Alaska, *J. Glaciol.*, 55, 1052–1060, 2009. 838

Tulaczyk, S., Kamb, W. B., and Engelhardt, H. F.: Basal mechanics of ice stream B, West Antarctica 1. Till mechanics, *J. Geophys. Res.*, 105, 463–481, 2000a. 826

30 Tulaczyk, S., Kamb, W. B., and Engelhardt, H. F.: Basal mechanics of ice stream B, West Antarctica 2. Undrained plastic bed model, *J. Geophys. Res.*, 105, 483–494, 2000b. 826

Van der Veen, C. J.: *Fundamentals of Glacier Dynamics*, A. A. Balkema, Rotterdam/Brookfield, 1999. 827

Van der Wateren, D. F. M.: Structural geology and sedimentology of push moraines: processes of soft sediment deformation in a glacial environment and the distribution of glaciotectonic styles, *Mededelingen Rijks Geologische Dienst* 54, Rijks Geologische Dienst, Harlem,

Glacier advance over deforming till

G. J.-M. C. Leysinger
Vieli and
G. H. Gudmundsson

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



1995. 825, 838, 839

Vieli, A., Funk, M., and Blatter, H.: Flow dynamics of tidewater glaciers: a numerical modelling approach, *J. Glaciol.*, 47, 595–606, 2001. 830

Wallinga, J. and Van de Wal, R. S. W.: Sensitivity of Rhonegletscher, Switzerland, to climate change: experiments with a one-dimensional flowline model, *J. Glaciol.*, 44, 383–393, 1998. 825

5 Winkler, S. and Nesje, A.: Moraine formation at an advancing temperate glacier: Brigidalsbreen, Western Norway, *Geogr. Ann.*, 34, 40–45, 1999. 838

Discussion Paper | Discussion Paper

TCD

4, 823–864, 2010

Glacier advance over deforming till

G. J.-M. C. Leysinger
Vieli and
G. H. Gudmundsson

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Table 1. Flow parameters used in the sediment layer for the different experiments.

Stress exponent (m)	Hardness factor (B)	Till characteristics	Type of advance
3	8	Relatively stiff and moderately non-linear till	overriding
3	0.35	Relatively soft and moderately non-linear till	plug-flow
3	0.79	Moderately stiff and moderately non-linear till	mixed
40	8	Relatively stiff and strongly non-linear till	overriding
40	0.42	Moderately soft and strongly non-linear till	mixed
40	0.44	Moderately stiff and strongly non-linear till	mixed

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Glacier advance over deforming till

G. J.-M. C. Leysinger
Vieli and
G. H. Gudmundsson

Table 2. Contribution of sediment deformation at positions A, B, C in percentage to the total deformation.

Type of advance	Flow exponent <i>m</i>	Hardness factor <i>B</i>	Sediment deformation (%)		
			A	B	C
Mixed-flow	3	0.79	11.7	23.0	42.4
	40	0.44	0.0	23.2	56.2
	40	0.42	0.0	33.3	62.2
Plug-flow	3	0.35	57.2	80.5	94.4

[Title Page](#) | [Abstract](#) | [Introduction](#)
[Conclusions](#) | [Tables](#) | [Figures](#)
[◀](#) | [▶](#)
[◀](#) | [▶](#)
[Back](#) | [Close](#)
[Full Screen / Esc](#)

[Printer-friendly Version](#)
[Interactive Discussion](#)



Glacier advance over deforming till

G. J.-M. C. Leysinger
Vieli and
G. H. Gudmundsson




Fig. 1. Schematic view of the glacier considered in the two-layer model. The ice body (white) resides on a sediment layer (grey) which extends into the glacier forefield. The inclination of the upper part is 45° and of 5° for the lower part.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Glacier advance over deforming till


G. J.-M. C. Leysinger
Vieli and
G. H. Gudmundsson

Fig. 2. Experiment A: Relatively stiff and moderately non-linear till ($m=3$, $B=8$): (a) glacier surfaces are shown at time intervals of approximately two dimensionless time units. Selected front positions and surface shapes marked (b), (c) and (d) are shown in more detail in subfigures (b–d). (b–d) snapshots of the front showing the surface, four selected internal layers and the ice-sediment interface at dimensionless times T equal to 0 (b), 10 (c) and 20 (d). The dots indicate material particles that can be followed throughout the three pictures. The connecting lines illustrate how material layers (fine solid and dashed lines) with initial shape as in (b) become deformed with time. The arrow marks the initial position of the glacier front.




Fig. 3. Experiment A: Relatively stiff and moderately non-linear till ($m=3$, $B=8$): **(b-d)** vertical profiles of the non-dimensional horizontal velocity U (lines with crosses) and effective stress T_{eff} (lines with dots) are shown at a dimensionless time $T=15.4$ for the three locations A, B and C shown in **(a)**. Note that the vertical scale changes for each profile.




Fig. 4. Experiment B: Relatively soft and moderately non-linear till ($m=3$, $B=0.35$): **(a)** several glacier surfaces are shown at approximately two dimensionless time unit intervals. Selected front positions and surface shapes marked **(c)**, **(d)** and **(e)** are shown in more detail in sub-figures **(c–e)**. **(b)** shows the maximum height H_b of the sediment bulge as a function of its horizontal position. Note how, after the initial build up phase, the bulge reaches a constant thickness. **(c–e)** snapshots of the front showing the surface, four selected internal layers and the ice-sediment interface at dimensionless times T equal to 0 **(c)**, 16 **(d)** and 32 **(e)**. The dots indicate material particles and can be followed throughout the three pictures. The connecting lines illustrate how material layers (fine solid and dashed lines) with initial shape as in **(c)** become deformed with time.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 5. Experiment B: Relatively soft and moderately non-linear till ($m=3$, $B=0.35$): Deformation of originally parallel internal sediment layers are shown for dimensionless times T equal to 0 (a), 8 (b) and 16 (c). The dots represent material particles throughout the pictures. The topmost layer shows the strongest deformation due to the uplifting of sediment needed to form the bulge (b-d). After a while the bulge is stable in amplitude and shape, and only a small uplift of the sediment takes place at the down-facing side of the bulge (c). Three former vertical straight lines (solid) illustrate how such lines deform with time.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 6. Experiment B: Relatively soft and moderately non-linear till ($m=3$, $B=0.35$): Three snapshots (**a–c**) at intervals of two dimensionless time units T show the evolution of five internal sediment layers which are parallel at the time $T=15$ (a), when the sediment bulge had an almost stable slope. Sediment uplift takes place at the transition from the down-facing bulge slope to the undisturbed sediment layer (b–c). A vertical straight line in (a) is followed throughout (b–c).

Glacier advance over deforming till

G. J.-M. C. Leysinger
 Vieli and
 G. H. Gudmundsson




Fig. 7. Experiment B: Relatively soft and moderately non-linear till ($m=3$, $B=0.35$): Snapshots at dimensionless time $T=15$ showing the distribution of horizontal velocity **(a)**, vertical velocity **(b)** and effective stress **(c)** in the frontal part of the glacier. In (b) the dashed line represents the zero velocity contour line and the symbol + means positive vertical velocities, that is upward movement.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Glacier advance over deforming till


G. J.-M. C. Leysinger
Vieli and
G. H. Gudmundsson

Fig. 8. Experiment A: Relatively stiff and moderately non-linear till ($m=3$, $B=8$): Snapshots at dimensionless time $T=15.4$ showing the distribution of horizontal velocity (a), vertical velocity (b) and effective stress (c) in the frontal part of the glacier. Note that the effective stress in the sediment is decreasing distal of the glacier front.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Glacier advance over deforming till

G. J.-M. C. Leysinger
Vieli and
G. H. Gudmundsson




Fig. 9. Experiment B: Relatively soft and moderately non-linear till ($m=3$, $B=0.35$): Velocity vectors of the sediment bulge are shown at the dimensionless time $T=15$. Velocity vectors are mostly parallel to the bed. Sediment uplift takes place at the downfacing side of the bulge.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Glacier advance over deforming till


G. J.-M. C. Leysinger
Vieli and
G. H. Gudmundsson

Fig. 10. Experiment B: Relatively soft and moderately non-linear till ($m=3$, $B=0.35$): (c–e) vertical profiles of the non-dimensional horizontal velocity U (lines with crosses) and effective stress T_{eff} (lines with dots) are shown at a dimensionless time $T=15$ for the three locations A, B and C shown in (a). Note that the vertical scale changes for each profile. (b) shows the ratio between the horizontal velocity at the ice-sediment interface U_b to the one at the glacier surface U_s .

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Glacier advance over deforming till


G. J.-M. C. Leysinger
Vieli and
G. H. Gudmundsson

Fig. 11. Experiment C: Moderately stiff and moderately non-linear till ($m=3$, $B=0.79$): **(a)** glacier surfaces are shown at time intervals of approximately two dimensionless time units. Selected front positions marked c, d and e are shown in more detail in subfigures **(c–e)**. **(b)** shows the maximum height H_b of the sediment bulge with its horizontal position. Note that the bulge is significantly smaller than in Fig. 4. **(c–e)** snapshots of the front showing the surface, four selected internal layers and the sediment surface at dimensionless times T equal to 0 **(c)**, 8 **(d)** and 14 **(e)**. The dots indicate material particles that can be followed throughout the three pictures. The connecting lines illustrate how material layers (fine solid and dashed lines) with initial shape as in **(c)** become deformed with time. The arrow marks the initial position of the glacier front.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Glacier advance over deforming till


G. J.-M. C. Leysinger
Vieli and
G. H. Gudmundsson

Fig. 12. Experiment C: Moderately stiff and moderately non-linear till ($m=3$, $B=0.79$): Sediment layers which were originally parallel to the bed (a) are deformed and form a wave to keep the sediment bulge in position (b–c). The snapshots were taken at dimensionless times T equal to 0 (a), 8 (b) and 14 (c). Three vertical straight lines in (a) are followed throughout (b–c).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Glacier advance over deforming till


G. J.-M. C. Leysinger
Vieli and
G. H. Gudmundsson

Fig. 13. Experiment C: Moderately stiff and moderately non-linear till ($m=3$, $B=0.79$): (c–e) vertical profiles of the non-dimensional horizontal velocity U (lines with crosses) and effective stress T_{eff} (lines with dots) are shown at a dimensionless time $T=15$ for the three locations A, B and C shown in (a). Note that the vertical scale changes for each profile. (b) shows the ratio between the horizontal velocity at the ice-sediment interface U_b to the one at the glacier surface U_s .

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Glacier advance over deforming till

G. J.-M. C. Leysinger
Vieli and
G. H. Gudmundsson




Fig. 14. Moderately stiff and highly non-linear till ($m=40$, $B=0.42$): **(c–e)** vertical profiles of the horizontal velocity (lines with crosses) and effective stress (lines with dots) are shown at a dimensionless time $T=15$ for the three locations A, B and C shown in **(a)**. **(b)** shows the ratio between the horizontal velocity at the ice-sediment interface U_b to the one at the glacier surface U_s .

[Title Page](#) | [Abstract](#) | [Introduction](#)
[Conclusions](#) | [References](#)
[Tables](#) | [Figures](#)

[◀](#) | [▶](#)
[◀](#) | [▶](#)
[Back](#) | [Close](#)
[Full Screen / Esc](#)

[Printer-friendly Version](#)
[Interactive Discussion](#)

Glacier advance over deforming till

G. J.-M. C. Leysinger
Vieli and
G. H. Gudmundsson




Fig. 15. Glacier advance calculated with a altitude-dependent mass balance distribution over the whole glacier as used in Leysinger Vieli and Gudmundsson (2004) and shown using the same scale as in Fig. 2a.

[Title Page](#)
[Abstract](#) [Introduction](#)
[Conclusions](#) [References](#)
[Tables](#) [Figures](#)
[◀](#) [▶](#)
[◀](#) [▶](#)
[Back](#) [Close](#)
[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)