The Cryosphere Discuss., 4, C1388–C1389, 2010 www.the-cryosphere-discuss.net/4/C1388/2010/ © Author(s) 2010. This work is distributed under the Creative Commons Attribute 3.0 License.



TCD 4, C1388–C1389, 2010

> Interactive Comment

Interactive comment on "Present and LGM permafrost from climate simulations: contribution of statistical downscaling" by G. Levavasseur et al.

K. Saito

ksaito@iarc.uaf.edu

Received and published: 20 December 2010

Interesting paper that attempts to utilize the outputs of the GCMs that do not sufficiently resolve the sub-surface thermal regimes or archive those results. It is unfortunate that I was advised of this paper just a few days before the comment deadline, and have not had much time to read through the paper. I thought still it might be worth to compare the results with our effort conducted as a sub-project of PMIP2, in which a simpler statistical method was employed to map the sub-surface thermal regimes (permafrost, seasonally-frozen ground, no freezing, etc.) for 0, 6 and 21ka from the freeze and thaw indexes derived from the PMIP2 near-surface air temperature outputs.





If you have a chance please refer to our letter paper. You might find it interesting and/or useful: Kazuyuki Saito, Sergei Marchenko, Vladimir Romanovsky, Nancy Bigelow, Kenji Yoshikawa, and John Walsh (2009): Thermally-Conditioned Paleo-Permafrost Variations from Global Climate Modeling. SOLA, 2009, Vol. 5, 101-104, doi:10.2151/sola.2009-026.

Please also note the supplement to this comment: http://www.the-cryosphere-discuss.net/4/C1388/2010/tcd-4-C1388-2010supplement.pdf

Interactive comment on The Cryosphere Discuss., 4, 2233, 2010.

TCD

4, C1388–C1389, 2010

Interactive Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

