

**GPR-derived  
accumulation and  
compaction of snow**

N. C. Kruetzmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



# Snow accumulation and compaction derived from GPR data near Ross Island, Antarctica

N. C. Kruetzmann<sup>1,2</sup>, W. Rack<sup>2</sup>, A. J. McDonald<sup>1</sup>, and S. E. George<sup>3</sup>

<sup>1</sup>Department of Physics & Astronomy, University of Canterbury, Private Bag 4800, Christchurch 8140, New Zealand

<sup>2</sup>Gateway Antarctica, University of Canterbury, Private Bag 4800, Christchurch 8140, New Zealand

<sup>3</sup>Antarctic Climate and Ecosystems Cooperative Research Centre, University of Tasmania, Private Bag 80, Hobart, Tasmania 7001, Australia

Received: 10 December 2010 – Accepted: 16 December 2010 – Published: 5 January 2011

Correspondence to: N. C. Kruetzmann (nikolai.kruetzmann@pg.canterbury.ac.nz)

Published by Copernicus Publications on behalf of the European Geosciences Union.

## Abstract

We present a new method of using ground penetrating radar (GPR) for estimating snow accumulation and compaction rates in Antarctica. We process 500 MHz data to produce radargrams with unambiguous reflection horizons that can be observed and tracked in repeat GPR measurements made one year apart. Our processing methodology is a deterministic deconvolution via the Fourier domain using an estimate of the emitted waveform from direct measurement. At two measurement sites near Scott Base, Antarctica, point measurements of average accumulation from snow pits and firn cores are extrapolated to a larger area by identifying a dateable dust layer in the radargrams. Over an 800 m × 800 m area on the McMurdo Ice Shelf (77°45' S, 167°17' E) the average accumulation is found to be  $269 \pm 9 \text{ kg m}^{-2} \text{ a}^{-1}$ . The accumulation over an area of 400 m × 400 m in the dry snow zone on Ross Island (77°40' S, 167°11' E, 350 m a.s.l.) is found to be higher ( $404 \pm 22 \text{ kg m}^{-2} \text{ a}^{-1}$ ) and shows increased variability related to undulating terrain. Compaction of snow between 2 m and 13 m depth is estimated at both sites by tracking several internal reflection horizons along the radar profiles and calculating the average change in separation of horizon pairs from one year to the next. The derived compaction rates range from  $7 \text{ cm m}^{-1}$  at a depth of two metres, down to no measurable compaction at 13 m depth, and are similar to published values from point measurements.

## 1 Introduction

In recent decades, satellite altimeters have been used to estimate and monitor the Antarctic mass balance by measuring surface height changes around Antarctica (Wingham et al., 1998; Davis and Ferguson, 2004; Nguyen and Herring, 2005). The variability in the mass of polar ice sheets has important implications for sea-level rise and the global radiation balance (Davis et al., 2005). Key uncertainties for determining snow accumulation from changes in surface elevation are snow density and compaction

TCD

5, 1–39, 2011

## GPR-derived accumulation and compaction of snow

N. C. Kruetzmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(Arthern and Wingham, 1998) and the spatial variability thereof (Drinkwater et al., 2001). These uncertainties can only be quantified by means of ground truthing.

The amount of snow compaction near the surface is related to mechanical settling during and immediately after accumulation, the overburden pressure by additional snow deposition, and the complex mechanism of temperature metamorphosis (van den Broeke, 2008), while melt metamorphosis is mostly restricted to coastal areas. A change in surface height as measured by satellite altimeters, therefore does not necessarily reflect a mass imbalance but may instead be caused by meteorological conditions affecting snow compaction. Additionally, densification processes can change the snow morphology, which affects the height retrieval from reflected radar waveforms such as those from the CryoSat-2 radar altimeter (Wingham et al., 2006). The aim of this study is to examine the utility of ground penetrating radar (GPR) measurements in deriving accumulation and compaction rates at two sites in, and close to, the dry-snow zone in Antarctica to reduce these uncertainties.

Studies which measure compaction of snow are rare. Recently, Arthern et al. (2010) presented an experimental setup for measuring snow compaction down to three distinct depth levels with very high temporal resolution. Zwally and Li (2002) developed a compaction model which shows good agreement with point measurements of compaction rates. Their results show that compaction of dry snow is a continuous process whose seasonal variability depends largely on temperature. In this paper we investigate the feasibility of measuring compaction rates over larger areas using a ground based radar system. We believe that our GPR based methodology is complementary to the point measurements made by Arthern et al. (2010), in that it can measure compaction over larger regions with a higher vertical resolution, albeit on a longer time scale.

Radar has been utilised for glaciological analysis of ice thickness and the detection of internal layers for many years (e.g., Bentley et al., 1979; Bogorodsky et al., 1985; Arcone et al., 1995; Eisen et al., 2003; Rotschky et al., 2006). Distinct reflections seen in radar recordings can sometimes be associated with accumulation or melt events (Eisen et al., 2004; Helm et al., 2007; Dunse et al., 2008). Analysis of internal layers

## GPR-derived accumulation and compaction of snow

N. C. Kruetzmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



---

**GPR-derived  
accumulation and  
compaction of snow**N. C. Kruezmman et al.

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

in ice and snow with commercial GPR systems for estimating accumulation has been demonstrated by Arcone et al. (2004), Dunse et al. (2008), and Heilig et al. (2009), amongst others. Here we apply a more rigorous processing methodology based on deconvolution. Our study includes snow density and accumulation information derived from snow pits and firn cores. This is used to reference our processed GPR data and expand the point measurements to larger areas. Additionally, we show that it is possible to acquire estimates of the compaction rates of dry snow by tracking internal horizons in GPR data and comparing layer separations in different years.

The research was carried out at three land ice sites, the locations of which are described in Sect. 2. Section 3 details the deterministic Fourier deconvolution processing scheme used for enhancing the weak contrast of the GPR data. In Sect. 4, the accumulation and compaction estimates are presented and results are discussed in Sect. 5. Section 6 summarises our findings.

## 2 Data acquisition

Ground penetrating radar data were acquired on the McMurdo Ice Shelf and on Ross Island, Antarctica (Fig. 1a and b), in November 2008 and 2009. A Sensors & Software pulseEKKO PRO GPR system emitting at a nominal frequency of 500 MHz was used to acquire reflection profiles of the subsurface. According to the manufacturer's specifications, the system has an effective isotropically radiated power (EIRP) just below 10 mW (Sensors & Software Inc., personal communication). With the settings used for this study the pulse repetition frequency (PRF) lies between 50 kHz and 60 kHz and the system has a duty cycle of 0.02%. A frequency analysis shows that the system actually operates at an effective centre frequency of about 620 MHz (see Fig. 2b), which is equivalent to an approximate wavelength of 0.35 m in dry snow, assuming a density of 500 kg m<sup>-3</sup>. Similarly, the bandwidth of the system can be estimated from Fig. 2b to be  $\Delta f \approx 550$  MHz. Following Rial et al. (2009), the system has a theoretical resolution  $\Delta_v = c / (2 \cdot \Delta f \cdot \sqrt{\epsilon_r}) \approx 0.19$  m, where  $\epsilon_r$  is the relative dielectric permittivity of snow (see Sect. 3) at a density of 500 kg m<sup>-3</sup>.

---

**GPR-derived  
accumulation and  
compaction of snow**N. C. Kruetzmann et al.

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

The measurement sites were located within 30 km of New Zealand's Scott Base. Stake farms on land ice were established in 2008 at three locations in different climatic settings and named L1, L2, and L3 (Fig. 1b). The stake farms were setup to measure snow accumulation over a one year time period and to allow repeat GPR measurements along the same profiles. At L1 and L2, 81 stakes were installed on a regular 800 m × 800 m grid at 100 m intervals. The distances between stakes and the regularity of the grid were established with centimetre accuracy using a laser distance meter. The layout of a site is illustrated in Fig. 1c. As the study was conducted within a validation experiment for CryoSat-2, the sites were oriented along anticipated satellite ground tracks. The stake in the southwest corner of each farm is labelled A1. The stake in the northeast is I9, with numbers increasing from west to east. For topographic and safety reasons, site L3 was reduced to a 400 m × 400 m grid of 25 stakes. Only odd numbered labels were used in this case to maintain consistent nomenclature for the corners. In the following, directions are relative to grid-north/east, unless stated otherwise. In addition to the stake farms, a profile of 20 stakes was established between L1 and L2, with a separation of approximately 1.5 km between the stakes.

The transmit-receive system and the recording equipment were pulled along the grid lines on plastic sleds at a slow walking pace. The GPR data were acquired with a sampling interval of 0.1 ns. Radar shots were triggered at regular intervals with an odometer wheel. Using a 135 ns time window in 2008 allowed us to record one trace every 5 cm. In 2009 we attempted to image deeper reflections by using a time window of 205 ns. Due to system limitations, this required an increased horizontal step-size of 7 cm. An additional radar profile was recorded along a line between L1 and L2 using a skidoo, with a recording step-size of 0.4 m and a time window of 180 ns.

Based on previous studies, L1 is located in an area of low accumulation and frequent summer melting (Heine, 1967), on an almost stationary part of the ice shelf. L2, situated in Windless Bight, features considerably higher accumulation rates. Based on Scott Base temperature records we expected occasional summer melting at this site. However, no melt layers were observed in the upper eight metres of snow. GPS

measurements corrected with base station data from Scott Base yielded an ice shelf movement of 58 m towards the southwest between November 2008 and October 2009. The third test site, L3, is located on the western slopes of Mt. Erebus (Ross Island), at an altitude of approximately 350 m a.s.l., in the dry snow zone and on undulating terrain. Here, GPS measurements show that this area had moved 3.4 m towards the Erebus Glacier Tongue. In both years, density profiles were taken in at least one snow pit at each site. Densities were measured by weighing known volumes of snow. Additionally, firn cores were drilled and logged in 2009 to obtain snow density profiles up to eight metres deep. L1 did not display coherent layers in radar or density profiles in either year. The irregular and low accumulation, high wind, and frequent summer melting at this site probably prevent the formation of clear stratification in snow pits and GPR images. Consequently, the data from this site are not discussed further.

### 3 GPR data processing methodology

In many cases the processing of GPR data has been adapted from the processing of seismic recordings. One frequently used procedure is to calculate the envelope of the received signal via the Hilbert Transform (e.g., Taner et al., 1979). However, this does not completely remove the effects of the carrier signal. Deconvolution, the common remedy to this problem in seismics, has been shown to be more difficult for GPR data (e.g., Turner, 1994; Irving and Knight, 2003). Two key reasons for this difficulty are dispersion of the emitted waveform and its non-minimum-phase character (Belina et al., 2009). The former causes changes in the shape of the radar wave as it travels through the medium, which makes the task of removing one specific waveform inaccurate. The latter relates to the energy distribution of the waveform emitted by most commercial GPR systems, which has its maximum close to the centre of the pulse rather than being frontloaded. This can lead to non-convergent deconvolution operators. Recently, Xia et al. (2004) and Belina et al. (2009) successfully tested deconvolution techniques for GPR recordings on low-dispersion soils. Additionally, Spikes et al. (2004) used a “spiking deconvolution” for simplifying firn radar profiles, though they do not give any

## GPR-derived accumulation and compaction of snow

N. C. Kruetzmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



detail on how this processing was applied. In this study we use a similar method, the deterministic Fourier deconvolution, to analyse internal radar reflections of dry Antarctic snow, which is also a low-dispersion material.

A common assumption in analysing GPR data is that the distribution of dielectric contrasts in the ground is random. This assumption is also known as the whiteness hypothesis (Ulrych, 1999), because a (successfully) recovered reflectivity profile is expected to have a spectrum that is similar to that of white noise. While the whiteness hypothesis is widely accepted in a geological context, though sometimes modified to a “blueness hypothesis” (Walden and Hosken, 1985; Ulrych, 1999), it is not immediately evident that it should also apply to the reflectivity structure of stratified snow. Snow deposited in different weather conditions will have variable permittivity based on the thermodynamic properties of the environment at the time, and thus successive layers may be correlated. Nevertheless, the small-scale details of the contrast between these layers are still likely to be random in nature, even if there is some correlation. Hence, the assumption that the spectrum of the output of the deconvolution should be at least *whiter* than the recorded radargram is likely to be true.

The GPR data recorded for this study can therefore be assumed to measure a medium which consists of well-defined layers with variable dielectric properties. The conductivity of dry snow is very small and the imaginary part of the dielectric permittivity can be neglected (Eisen et al., 2003). The real part of the relative dielectric permittivity,  $\epsilon'_r$ , of dry snow can be related to its density,  $\rho$  (in  $\text{g}/\text{cm}^3$ ), using the empirical formula (Kovacs et al., 1995):

$$\epsilon'_r(\rho) = (1 + 0.845 \cdot \rho)^2 \quad (1)$$

The succession of snow layers can be thought of as a reflectivity profile,  $r(t)$ . The emitted signal is partially reflected at each interface between these layers, and the intensity of the reflection depends on the magnitude of the dielectric gradient at each interface. Therefore, the reflectivity profile can be directly related to snow density variations (Eisen et al., 2008). The received radar signal,  $s(t)$ , can be described as

## GPR-derived accumulation and compaction of snow

N. C. Kruetzmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the convolution of an emitted waveform,  $e(t)$ , with  $r(t)$ :

$$s(t) = e(t) * r(t) \quad (2)$$

Fourier deconvolution provides a method to remove the effect of the high frequency carrier wave and to recover the reflectivity profile from the radargram. According to the convolution theorem, the Fourier transform (FT) of the convolution of two functions is equal to the point-wise multiplication of the Fourier transforms of the individual functions (Bracewell, 2003):

$$\text{FT}(s) = \text{FT}(e * r) = \text{FT}(e) \cdot \text{FT}(r) \quad (3)$$

Thus, if the emitted waveform can be estimated, dividing the FT of the recorded trace by the FT of the waveform results in the FT of the reflectivity profile. An inverse Fourier transform (IFT) can then be used to derive the reflectivity profile:

$$\text{FT}(r) = \frac{\text{FT}(s)}{\text{FT}(e)} \Leftrightarrow r = \text{IFT} \left( \frac{\text{FT}(s)}{\text{FT}(e)} \right) \quad (4)$$

If the source waveform is known and does not change with time (i.e. depth in the medium), this is referred to as deterministic deconvolution (Yilmaz, 1987), since it is a well defined mathematical problem that has a single solution. To avoid the problem of non-convergent filter functions mentioned above, we perform the deconvolution in the frequency-domain.

To successfully use Eq. (4) for recovering  $r(t)$ , we measured the emitted waveform (see Fig. 2) by holding the transmitting and receiving antennae above the ground, directly facing each other. To avoid interference between the air-wave and the ground-reflection, the transmitter and receiver were placed 2.3m above the surface, and 2m apart.

Figure 2a shows the returned signal as a function of time for 100 superimposed shots. Two distinct returns are observed, the air-wave that travels directly from the transmitter to the receiver, and the ground-reflection. The small variation between

## GPR-derived accumulation and compaction of snow

N. C. Kruetzmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





shots displayed in Fig. 2a illustrates that the pulse emitted by the system is transmitted at a consistent phase and can be assumed to have the same form from one shot to another. We use an integrated version of the air-wave as our estimate of the emitted waveform for the deconvolution process. The frequency spectrum of the averaged trace, Fig. 2b, shows that the centre frequency of the pulse lies close to 620 MHz. Before performing the division in Eq. (4), the spectrum of the emitted wave is whitened by ten percent of the largest spectral peak in order to avoid over-amplification of frequencies of small amplitude (Yilmaz, 1987).

An overview of the processing steps and the order in which they are applied is given in the following list:

1. Dewow – a DC-correction to remove a localised offset caused by receiver saturation (Sensors & Software Inc., 2006).
2. Linear gain – compensates for the loss of signal power due to the spherical spreading of the wave as it propagates through the medium.
3. Fourier deconvolution – removes the effects of the carrier wave.
4. Low-pass filter – applied in the frequency domain with a 1550 MHz cut-off based on the spectrum of the emitted wave.
5. Background subtraction – to remove interference due to ringing. This distorts the radargram in the first 6 to 8 ns which are therefore not analysed.
6. Envelope calculation using the Hilbert transform.
7. Integration (stacking) – used to improve the signal-to-noise ratio.

Step (7) also allows us to match the horizontal sampling intervals of the data from the two seasons. After ten-fold stacking of the data from 2008 and seven-fold stacking of data from 2009, the horizontal sampling of both data sets reduces to approximately 0.5 m.

## GPR-derived accumulation and compaction of snow

N. C. Kruetzmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



---

**GPR-derived  
accumulation and  
compaction of snow**N. C. Kruetzmann et al.

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Figure 3 illustrates the benefits of the Fourier deconvolution scheme for a representative 200 m long radar profile from the L2 site. The simple application of dewow and gain in Fig. 3a shows that the carrier wave is still present in the data. After applying all processing steps except deconvolution, a reduced number of distinct reflections can be clearly identified in Fig. 3b. Including the deconvolution step (Fig. 3c) significantly sharpens these horizons. Comparison of the different panels in Fig. 3 shows that, by compressing the oscillating carrier wave to a single peak, the deconvolution results in a more focussed radargram.

A more stringent test of the quality of the deconvolution processing methodology is to analyse the autocorrelation of traces before and after processing. Theoretically, deconvolution should significantly reduce the autocorrelation time and flatten the tail of the autocorrelation function (ACF) of a radar trace (Ulrych, 1999). Figure 4 shows the ACF of a trace before and after deconvolution, and demonstrates that there is a decrease in autocorrelation time after deconvolution (first minimum is shifted from 0.7 ns to 0.5 ns) and that the tail has been flattened. Note that the ACF is calculated using a truncated version of the trace that does not include the first 8 ns, for reasons mentioned above.

The new GPR processing scheme is used as a basis to identify and track internal reflections in the radargrams. The tracking was performed by initially selecting from the limited number of visible horizons observed after processing (Fig. 3c) using the KINGDOM Suite 8.2 software. From a starting point somewhere within the radargram determined by the user, the program follows the reflection peak from trace to trace. The vertical guide window within which the algorithm searches for the amplitude maximum was set to 2 ns. This corresponds to approximately 0.2 m in the vertical, which is close to the estimated resolution limit. Sometimes a reflection is weak or very blurred in some part of the profile and the tracking algorithm may not be able to find the reflection or may follow the wrong path. In this case the selection is corrected manually until automatic tracking can resume.

The tracked internal reflection horizons are used for extrapolating accumulation measurements from snow pits and firn cores to larger areas and for deriving compaction rates as a function of depth at the sites L2 and L3. In order to analyse the GPR profiles it is necessary to establish a time-depth relationship. Using the density information from the snow pits and firn cores to estimate  $\varepsilon'_r$  in Eq. (4), we calculate the velocity,  $v$ , of the radar signal in snow:

$$v = \frac{c}{\sqrt{\varepsilon'_r}} \quad (5)$$

where  $c$  is the speed of light in vacuum. Using this analysis, the two-way-travel time (TWT) from a radargram can be converted to depth and vice versa, allowing accumulation estimates to be derived from the GPR data. In the following, reflection-depth is converted to water equivalent by using a depth-density relationship determined from a firn core at the respective site.

## 4 Results

### 4.1 Accumulation estimates from stake farms, snow pits, and firn cores

Stake farm and firn core measurements provided independent sources of accumulation data (Table 1). The stake farms at L2 and L3 were installed in November 2008 and revisited in November 2009. Recordings of the stake heights above the snow in both years allow the measurement of one year of accumulation in these areas using the conversion detailed in Takahashi and Kameda (2007). Taking the mean and standard deviation of the snow depths at the 81 stakes from L2 gives an average reduction in stake height of  $60.2 \pm 8.1$  cm or, using the snow densities recorded in a snow pit,  $224 \pm 21$  kg m<sup>-2</sup> a<sup>-1</sup>. At L3, only 25 stakes were installed. The measurement from one stake was omitted because of a likely error in the height recording. The average

## GPR-derived accumulation and compaction of snow

N. C. Kruezmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



decrease in stake height measured at the remaining 24 stakes was  $70.1 \pm 11.1$  cm, equivalent to  $304 \pm 83 \text{ kg m}^{-2} \text{ a}^{-1}$  of accumulation.

A previously identified dust layer (Dunbar et al., 2009) originating from a severe storm (with maximum southerly wind speeds exceeding  $55 \text{ m s}^{-1}$ ) that occurred on 16 May 2004 (Xiao et al., 2008) served as a reference point for dating. In 2008, we found the dust layer at a depth of 2.93 m in a snow pit near stake C3 (not shown) at L2. Using the dust layer for dating, we calculate an average annual accumulation of  $251 \text{ kg m}^{-2} \text{ a}^{-1}$ . In 2009, we were unable to clearly identify the dust layer in a snow pit. However, a firn core was drilled one metre east of the stake G7, to a depth of 8.46 m. The core contained an unusually coarse grained low-density layer at a depth of 3.42 m, which is likely to be related to the dust layer found previously. Similarly, considering this as a marker for the storm in May 2004 gives an accumulation rate of  $245 \text{ kg m}^{-2} \text{ a}^{-1}$ . Figure 5a shows the density profile of the firn core and some of the snow pit data from 2009.

At L3, no particularly distinct features could be found in a two metre snow pit in either year (not shown). Slight changes in snow grain size and hardness at about one metre depth are indicative of the previous year's summer layer, but it was not possible to determine this with any certainty. Generally, the snow at L3 was found to be very homogeneous, with a slightly higher average density than at L2. In 2009, we drilled a firn core down to about 7.5 m near stake C3 at L3. The resulting density profile is shown in Fig. 5b. At 5.3 m depth we were able to identify a dust layer similar to that found at L2 in the previous year. Assuming that this is also associated with the May 2004 storm gives an estimated average accumulation of  $437 \text{ kg m}^{-2} \text{ a}^{-1}$ . This value is considerably higher than the one measured by the stake farm and reflects a high inter-annual variability in snow accumulation.

The firn core data can be used to determine a depth-density relationship for the two sites, as a reference for converting the vertical scale of the radargrams to depth. Following Alley et al. (1982), we determine an empirical depth( $d$ )-density( $\rho$ ) relationship for both sites by fitting an exponential model of the form  $\rho(d) = \rho_i - a \cdot e^{-c \cdot d}$  to the firn

## GPR-derived accumulation and compaction of snow

N. C. Kruezmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



core data, where  $\rho_i=917 \text{ kg m}^{-3}$  is the density of ice, and  $a$  and  $c$  are the constants to be fitted. For the upper 0.6 m at L2, where the snow was loose and density difficult to measure by coring, snow pit data were used in order to obtain a fitted depth-density curve. The core at L3 was drilled two weeks after the snow pit had been recorded, as weather conditions prevented earlier attempts to return to the site. The core was started at the previous surface level by removing the fresh snow. However, the additional overburden will have caused some densification. Therefore, the top 1.9 m of the firn core were replaced by snow pit data when determining the depth-density relationship. The resultant equations for L2 and L3 are shown in Fig. 5a and b, respectively. As the radar profiles extend below the maximum depth of the firn cores, we also use these equations to extrapolate the density to greater depths in the following analysis. Integrating these empirical relationships allows the estimation of the total snow mass to a certain depth, which can then be used to determine the mean column density, and therefore the TWT, down to this depth.

## 4.2 Accumulation estimates from GPR measurements

The snow pit logged at L2 in 2008 is approximately located at the centre of Fig. 3c. The yellow arrow corresponds to the dust layer depth and coincides with a strongly undulating horizon in the radar profile. Therefore, it seems likely that this reflection is related to the storm event in May 2004. Figure 6a and b shows processed radargrams from line E4 to E6 at L2, recorded in 2008 and 2009, respectively. The undulating horizon, as well as several other reflections, are observed throughout the entire survey grid in both years (not shown). The result of tracking the dust layer horizon, and eight other distinct reflections in the radar lines in Fig. 6a and b, is shown in Fig. 6c and d, respectively. The tracked reflection horizons are numbered with roman numerals from top to bottom for easier referencing. The vertical scale in Fig. 6d is shifted by 4.7 ns to align the first horizon (I) in both years to facilitate comparison. The yellow line (II) in Fig. 6c is the reflection we associate with the dust horizon. Comparing its path with

## GPR-derived accumulation and compaction of snow

N. C. Kruezmman et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## GPR-derived accumulation and compaction of snow

N. C. Kruetzmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the neighbouring horizons further illustrates that it is unusually variable; its standard deviation from the linear trend is 0.65 ns, as opposed to 0.30 ns, 0.44 ns, and 0.34 ns for the red (I), purple (III), and black (IV) horizons, respectively. Assuming that the association of the undulating horizon with the dust storm is correct, we can calculate the accumulation over the whole grid since May 2004. Tracking this reflection along all 18 grid lines gives an average TWT of  $27.5 \pm 0.8$  ns, which is equivalent to an average accumulation of approximately  $269 \pm 9 \text{ kg m}^{-2} \text{ a}^{-1}$ . Note that the error term is the standard error of the measured depths, reflecting the geophysical variability of the accumulation over the whole site rather than measurement error. Figure 7a illustrates the variability in this reflection's depth over the L2 area. While there is no particularly distinct pattern, it appears that accumulation is slightly higher in the north-east part of the grid.

Figure 8 displays a radargram of the C-line at L3 from 2009. The location of the firn core is indicated by the red box. Using the dust layer depth (5.3 m) and the densities from the firn core at stake C3, the expected TWT to the dust layer is calculated to be 49 ns (yellow arrow in Fig. 8). Comparing this with the radargram shows that there is a clear reflection horizon at approximately this TWT (tracked in yellow in Fig. 8). Analogous to the approach at L2, we consider this as a marker for the May 2004 dust storm. Accordingly, the average accumulation over this site is calculated to be  $404 \pm 22 \text{ kg m}^{-2} \text{ a}^{-1}$ .

Examination of Fig. 8 shows that the internal horizons are noticeably deeper in the middle of the profile, indicating more accumulation at the centre of the grid than at the edges. This inhomogeneity is probably related to local topography since L3 is situated on sloping terrain. The vertical difference between the lowest (I1) and the highest point (A9) is almost 20 m, with the terrain slope increasing towards the south and west, i.e. the gradient of the surface is smaller on the left hand side of Fig. 8 than on the right. Previous work by Vaughan et al. (1999) found that there is a direct correlation between surface slope and accumulation-related dipping of internal horizons. Accordingly, the dip in the reflections can be explained by local differences in accumulation related to

changes in the surface slope. Hence, the accumulation rate calculated above needs to be considered as a large scale average for the whole area, rather than an accurate estimate at any particular location. The interpolated accumulation grid for L3 (Fig. 7b), calculated from the tracked dust layer reflection, illustrates the overall pattern. The central dip in the terrain clearly captures more snow than the surrounding areas.

The decreasing trend in the depth of the dust horizon from north to south observed at L2 (Fig. 7a) can be followed south along a GPR transect (Fig. 9) that was recorded as an extension of the line going from I1 to A1. Along this profile, the internal horizons gradually migrate upwards (not shown). After about 14 km, the horizon we associate with the dust layer becomes too indistinct to be reliably identified. At this point, the dust layer reflection is found at a depth of about 1.9 m, which is equivalent to a reduced average accumulation of approximately  $165 \text{ kg m}^{-2} \text{ a}^{-1}$  (see Fig. 9). However, this has to be considered a crude estimate, since it assumes the same average snow density between the surface and the dust layer as at L2. A qualitatively similar trend was previously reported by both Heine (1967) and the McMurdo Ice Shelf Project (McCrae, 1984).

Unfortunately, we were unable to reliably date other layers within the snow pit or firn core profiles and associate them with discrete reflections in the radargram. However, due to the high precision of the system, the other apparent horizons – or rather their vertical separation – can be used to estimate compaction, which is the focus of the next section.

### 4.3 Snow compaction

Most of the apparent internal horizons found in the 2008 GPR data can also be identified in the following year's record. Figure 6c shows nine reflections tracked between stakes E4 and E6 at L2 in 2008, and the same horizons tracked in the 2009 data, are presented in Fig. 6d. One might expect that compaction, caused by temperature metamorphosis and the additional overburden on the surface, will reduce the separation between horizon pairs. This is confirmed by the poor alignment of the bottom horizon

## GPR-derived accumulation and compaction of snow

N. C. Kruezmman et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(IX) in Fig. 6d compared to Fig. 6c. Clearly, the total separation between the top (I) and the bottom (IX) reflection has been reduced. Assuming that the same horizons have been identified, calculating the average distance between two successive horizons in 2008 and comparing this layer thickness with the separation of the same horizon pair in the 2009 data, allows us to estimate the compaction of the snow in the intervening time period.

Analysis suggests that not all of the horizons shown in Fig. 6c and d are suitable for compaction calculations, since one has to be confident that the same pair of reflections are tracked in both years. The high variability of the undulating horizon for example, does not allow it to be tracked reliably enough in both years to be confident that the exact same reflection was selected. Similarly, the relatively strong double-reflection visible between 41 ns and 46 ns in 2008 (Fig. 6a) and between 45 ns and 49 ns in 2009 (Fig. 6b) may be easy to recognise visually, but the noisy nature of both horizons makes them unreliable subjects for the automated tracking algorithm. In order to establish which of the reflections are likely to be reliably tracked in both years, we calculate the correlation between each horizon in 2008 and its counterpart in 2009. Only those horizons that show a correlation greater than 0.5 are used for compaction calculations. To ensure accuracy, we also require a minimum average TWT difference between horizons of 10 ns ( $\Leftrightarrow$ 1 m vertical separation) and additionally that the correlation of the TWT differences in 2008 and 2009 is greater than 0.5.

For example, the average separation of horizon III and horizon IV in 2008 is 18.7 ns (Fig. 6c), but only 18.4 ns in 2009 (Fig. 6d). This means that the snow in between these two reflections was compressed by about 1.6% during the intervening time period. Using the density information from the firn core, this translates into a compaction of about  $1.6 \text{ cm m}^{-1}$ . The same calculations for five other horizon pairs result in the compaction diagram shown in Fig. 10. Clearly, the compaction decreases with depth. The horizontal error bars are the standard errors of the respective reflections' depths along the whole line added in quadrature. The vertical scale in Fig. 10 relates to the depth of the horizons in 2008, i.e. before the compaction has occurred. Since the time between the

## GPR-derived accumulation and compaction of snow

N. C. Kruezmman et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





acquisition of the GPR datasets was nearly one year (355 days), the values in Fig. 10 can be considered estimates of annual compaction rates.

The conversion of TWT difference between two horizons to a physical separation is performed by calculating the mean depth between the two horizons in 2009, determining the density at this depth with the formula shown in Fig. 5a and then using this density to estimate the radar velocity between the two horizons via Eqs. (1) and (5). Note that the density of the snow between two horizons is determined from the TWT in 2009 (post-compaction) and assumed to be the same in 2008 (pre-compaction), as the empirical depth-density relationship is purely based on data from 2009. Therefore, we likely underestimate the radar velocity, and hence the distance, between the horizons in 2008, making our compaction rates conservative estimates. For example, assuming a constant density profile with time (Sorge's law; Bader, 1954), the average snow density in the layer initially between 5 to 10 m depth at L2 would increase by 2.5% between 2008 and 2009. This translates to an underestimation of the wave velocity in 2008 by approximately 1% and an increase in the compaction rates of about  $0.5 \text{ cm m}^{-1}$ .

The apparent expansion observed below 10 m in Fig. 10 is probably an indicator of the error in our measurements. The actual amount of compaction over a one year time period at this depth is probably too small to be measured with our system. Furthermore, below 8.4 m the conversion from TWT to depth is based solely on an extrapolation of the exponential fit to the density data from the firn core, causing additional uncertainties in the calculations.

The same calculations were performed for five more lines at L2 (A2–I2, A5–I5, A7–I7, B1–B9, and H1–H9). The results are summarised in Fig. 11a. The vertical error bars indicate the average separation of the two horizons in 2008. Figure 11a shows that the variability in layer depths and compaction between these six lines are small and thus tracking was not repeated for the remaining lines. The thicker, black data points present the average compaction between each of the horizon pairs and the connecting line illustrates the decreasing trend.

## GPR-derived accumulation and compaction of snow

N. C. Kruezmman et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Using the same methodology on seven internal horizons at L3 allows the calculation of an equivalent diagram shown in Fig. 11b. At this site, the variability of the compaction rates and the average depths is significantly higher. Therefore, the calculations were performed for all grid lines, except for A9–I9 due to corrupt data. The larger spread is in accordance with the observed spatial variability in the accumulation pattern and the resulting dipping of the internal horizons (see Figs. 7b and 8). Nevertheless, the mean compaction rate (black line) shows a clear trend of reduced compaction with depth, similar to that observed at L2.

## 5 Discussion

The deterministic Fourier deconvolution scheme suggested here can be used to remove the effects of the carrier signal of any GPR data of dry firn, provided that it was recorded with a system that has a very stable output. The result is a more focussed radargram with improved contrast. This type of processing allows the identification and precise tracking of weak internal reflection horizons associated with density variations. The processing also facilitates recognition of the same horizons in follow-up surveys, as it removes the ubiquitous apparent reflections found in the raw data, most of which are associated with the carrier frequency. Our methodology also has the potential to work successfully for recordings in areas that are subject to sporadic melt events, although one should keep in mind that an essential assumption of the deconvolution is that there is no – or only very little – frequency dependent absorption and dispersion in the medium.

Table 1 summarizes our results relating to accumulation. At both L2 and L3, the measurements derived from the stake farms showed significantly lower results than the combined firn core and GPR measurements. This discrepancy is probably largely due to temporal variability. If the readout of the stake depths had occurred one week later, the measurements at both sites would have been noticeably higher due to the occurrence of a high precipitation event from 13 to 15 November 2009. The accumulation

rates derived from the snow pit and firn core observations should therefore be considered more reliable, since they consider a longer time period.

We now compare our measurements with previous studies in the area. The average accumulation at L2 – a site that is relatively typical for coastal areas – was found to be  $269 \pm 9 \text{ kg m}^{-2} \text{ a}^{-1}$ , which is much lower than the value reported by Heine (1967) at the closest station ( $510 \text{ kg m}^{-2} \text{ a}^{-1}$  at station 200, about 6 km north east of our site). However, there is a strong accumulation gradient in this area and L2 is close to the  $320 \text{ kg m}^{-2} \text{ a}^{-1}$  contour suggested by McCrae (1984), a value which still lies above our estimate. Generally, the conversion of TWT to depth based on measured densities is critical for our method and a potential source of error. A bias in the density data would lead to over- or under-estimation of the wave velocity in the snow and therefore the accumulation estimates. However, good agreement between core- and snow pit densities from the two different seasons at both sites (not shown) allows us to be confident in the density measurements. As the dust layer is a reliable reference point for dating, the difference between our results and previous studies could indicate an overall reduction in annual accumulation or high natural variability in this area. Moreover, it is important to note that historic accumulation derived from stake measurements are very likely to be underestimates, as the consideration of snow compaction – as suggested by Takahashi and Kameda (2007) – is not reported. Additionally, the gradient in the accumulation map for L2 (Fig. 7a) is found to continue along the transect toward L1 (Fig. 9). The southernmost point up to which we were able to track the dust horizon lies relatively close to the location of a 20 m firn core analysed in Dunbar et al. (2009). They estimate an accumulation rate of  $53 \pm 20 \text{ cm}$  of snow per year. If we assume an average density of  $600 \text{ kg m}^{-3}$  for the whole length of their core, this corresponds to  $318 \pm 120 \text{ kg m}^{-2} \text{ a}^{-1}$ . Again, our estimate from the tracked reflection lies significantly lower at  $165 \text{ kg m}^{-2} \text{ a}^{-1}$ . However, the two points are approximately 3 km apart and the conversion from TWT to depth uses the average column density down to the dust layer determined at L2, which is likely an underestimate for this location.

## GPR-derived accumulation and compaction of snow

N. C. Kruezmann et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## GPR-derived accumulation and compaction of snow

N. C. Kruezmman et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Only one reflection horizon could be associated with a layer in the snow stratigraphy acquired via simple glaciological tools and visible observation. However, such difficulties in linking snow pit and radar observations are quite common (Harper and Bradford, 2003) and are due to the limited resolution of our density profiles (Eisen et al., 2003).

Furthermore, the study sites investigated here were located in areas of low dielectric contrast. The snow was dry and homogeneous and therefore contained few major dielectric gradients, such as might be caused by occasional melt layers (e.g., Dunse et al. 2008). Nevertheless, since we were able to re-identify many reflection horizons in the following season, these correspond to actual density variations. Accurate identification of the origin of the radar reflections would require high-resolution data from e.g. dielectric profiling, as suggested by Eisen et al. (2004) and Hawley et al. (2008).

The extrapolation of the point measurements to the whole areas of our test sites (Fig. 7), show that there can be high variability of the accumulation rate even on a small scale (hundreds of metres), particularly at L3. This underlines the importance of accumulation measurements with high spatial resolution to obtain representative values of the surface mass balance and may also explain some of the differences we observe with previous studies.

The location of the various horizons at the crossover points of the grid is very consistent between perpendicular profiles. In most cases, the same reflection tracked along different profiles can be found within two time samples ( $2 \times 0.1 \text{ ns} \leftrightarrow 4 \text{ cm}$ ) at the point of intersection. The concurrence of the horizon depths is equally high for all reflections at both L2 and L3, showing that the precision of the processed radar data is much higher than the theoretical resolution might suggest. This high precision allows us to estimate snow compaction with depth from changes in the separations of internal horizons from one year to the next.

At both L2 and L3, the average compaction measured between 2 and 13 m depth over a one year time period ranges from  $0 \text{ cm m}^{-1}$  to  $7 \text{ cm m}^{-1}$ . The sudden drop in compaction observed below about 4 m at both sites (see Fig. 11) could be related to a change in the compaction mechanism. For example, fresh snow largely compacts via

settling, but above a density of  $550 \text{ kg m}^{-3}$  sintering is usually considered to become dominant (Maeno, 1982; van den Broeke, 2008). However, the depth at which we observe this change in compaction rate is too shallow. According to the density profiles in Fig. 5, the  $550 \text{ kg m}^{-3}$  level lies around 6 m depth at both sites. Incidentally, the latter value compares well with the model estimate by van den Broeke (2008), who gives a range of 5 to 8 m depth for the  $550 \text{ kg m}^{-3}$  level in this area. Therefore, the observed step in compaction rate may not solely be caused by a change in the compaction mechanism, but more measurements would be required to investigate the origin of this discrepancy. Similarly, the origin of the dip in compaction rate in between 7.5 m and 8.5 m at L2 (Fig. 11a) could be a result of above average snow density at this depth, possibly related to one or two particularly warm summers. However, we do not have sufficient data to test whether this is a measurement error or a subsurface feature.

High resolution measurements of densification rates to directly compare with our results are sparse. As we cannot identify the 2008 surface in the radargrams from 2009, it is not possible to calculate the total compaction of the whole snow column. However, total compaction between 5 m and 10 m depth can be estimated from Fig. 11 and compared to the measurements from Arthern et al. (2010). Summing up the average compaction (black lines in Fig. 11a and b) between 5 m and 10 m, gives a total compaction of approximately 4.3 cm at L2 (assuming zero compaction for the last 30 cm, where the results are negative) and 5.5 cm at L3, over a time period of 355 days. Using the daily rates given in Table 3 of Arthern et al. (2010) to calculate the total compaction over the same time span and depth range gives 5.7 cm for their “Berkner Island” site, the only site at which their strainmeters worked throughout the whole trial time. While our results qualitatively agree with the “Berkner Island” data, the other sites detailed in Arthern et al. (2010) show considerably higher compaction rates. As mentioned above, one reason for this could be that we do not take into account the change in snow density due to compaction and its effect on the velocity of the radar signal, since this would require additional density data for 2008. Ultimately, the discrepancies between Arthern et al. (2010) and our results could also be due to a difference in climatic conditions,

**GPR-derived  
accumulation and  
compaction of snow**

N. C. Kruetzmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



since most of their sites are located in regions with lower mean annual temperature, lower latitude, higher elevation and higher annual accumulation. Therefore, considerable differences in the compaction behaviour of the snow could be expected.

Our results above show that internal reflectors found in GPR data can be used for estimating compaction rates in dry snow if combined with sufficient complementary density information. However, while a large part of Antarctica can be considered to be dry snow, estimating densification is probably much more difficult in percolation regions, due to the more complex dynamics of snow morphology with regular melt water intrusions (Parry et al., 2007).

## 6 Conclusion and outlook

We have established a method of deconvolution for GPR data via the Fourier domain, using an estimate of the emitted waveform from direct measurement. Applying the deterministic Fourier deconvolution to GPR data from the dry snow zone results in a radar-gram with more narrow and distinct horizons which are more easily distinguished from the background of volume scattering in the low-contrast medium. While we believe that the approach has the potential to be successful for recordings in areas that are subject to sporadic melt events, the assumption that there is (almost) no frequency dependent absorption or dispersion in the medium limits its applicability to data acquired under such conditions.

Using the thusly processed radargrams from two measurement sites near Scott Base, Ross Island, Antarctica, we were able to extrapolate point measurements of average accumulation from a snow pit at site L2 ( $251 \text{ kg m}^{-2} \text{ a}^{-1}$ ) and a firn core at site L3 ( $437 \text{ kg m}^{-2} \text{ a}^{-1}$ ), to a larger area by identifying a dateable dust layer from the firn profiles with a reflection in the radargram. From the GPR data the extrapolated average accumulation over the  $800 \text{ m} \times 800 \text{ m}$  site on the ice shelf in Windless Bight (L2) was found to be  $269 \pm 9 \text{ kg m}^{-2} \text{ a}^{-1}$ . The  $400 \text{ m} \times 400 \text{ m}$  grid on Ross Island showed higher variability in the internal horizons with an overall average accumulation

## GPR-derived accumulation and compaction of snow

N. C. Kruetzmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



of  $404 \pm 22 \text{ kg m}^{-2} \text{ a}^{-1}$ . Furthermore, we were able to identify a decreasing accumulation trend along a 14 km long GPR transect heading south from L2. At the southernmost point, the accumulation was about  $165 \text{ kg m}^{-2} \text{ a}^{-1}$ . Stake farm readings at both sites, maintained over approximately a one year time period, resulted in accumulation estimates of  $224 \pm 21 \text{ kg m}^{-2} \text{ a}^{-1}$  at L2 and  $304 \pm 83 \text{ kg m}^{-2} \text{ a}^{-1}$  at L3. The discrepancy between these values and the combined firn core and GPR measurements is probably due to the short time period spanned by the stake observations and the high temporal variability of precipitation events.

Additionally, we were able to estimate compaction rates down to 13 m depth from GPR measurements for the first time, by comparing vertical separations of internal reflection horizons from one year to the next. This technique might have implications for the validation of the CryoSat-2 satellite altimeter which consistently measures the surface height of ice sheets and shelves, in order to monitor the polar mass balance. Approximations of compaction rates are vital for accurately estimating the local mass balance of an ice sheet, since compaction can lead to an apparent decrease in surface height, without a corresponding change in mass.

Our results show that high-frequency GPR can be used to monitor the accumulation and compaction of snow in situ. Frequent repetition of our measurements over a longer time period, combined with high resolution dielectric profiles of firn cores, could therefore be used to establish a more detailed representation of time-dependent firn densification for the validation of current firn densification models. The suggested method is complementary to point measurements of compaction with a very high time resolution, such as those suggested by Arthern et al. (2010), in the sense that our method has higher vertical resolution and can be used for estimating compaction rate averages over larger areas in an efficient and non-invasive manner. Additionally, using a setup similar to the one presented in Heilig et al. (2009) and a higher frequency system it might be possible to improve the time resolution of compaction data from GPR measurements in the future.

**GPR-derived  
accumulation and  
compaction of snow**

N. C. Kruezmman et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



*Acknowledgements.* This work was supported by Antarctica New Zealand under the field event K053 “Cryosphere Remote Sensing” and by scholarships from the Christchurch City Council and the University of Canterbury for Nikolai Krueztmann. The participation of Mette Riger-Kusk in the field work in 2008 is greatly acknowledged. The measurements were conducted within the ESA CryoSat-2 calibration and validation activities for project AOCRY2CAL-4512.

## References

- Alley, R. B., Bolzan, J. F., and Whillans, I. M.: Polar firn densification and grain growth, *Ann. Glaciol.*, 3, 7–11, 1982.
- Arcone, S. A., Lawson, D., and Delaney, A.: Short-pulse wavelet recovery and resolution of dielectric contrasts within englacial and basal ice of Matanuska Glacier, Alaska, USA, *J. Glaciol.*, 41, 68–86, 1995.
- Arcone, S. A., Spikes, V. B., Hamilton, G. S., and Mayewski, P. A.: Stratigraphic continuity in 400 MHz short-pulse radar profiles of firn in West Antarctica, *Ann. Glaciol.*, 39, 195–200, 2004.
- Arthern, R. J. and Wingham, D. J.: The natural fluctuations of firn densification and their effect on the geodetic determination of ice sheet mass balance, *Climatic Change*, 40(4), 605–624, 1998.
- Arthern, R. J., Vaughan, D. G., Rankin, A. M., Mulvaney, R., and Thomas, E. R.: In situ measurements of Antarctic snow compaction compared with predictions of models, *J. Geophys. Res.*, 115, F03011, doi:10.1029/2009JF001306, 2010.
- Bader, H.: Sorge’s law of densification of snow on high polar glaciers, *J. Glaciol.*, 2(15), 319–323, 1954.
- Bentley, C. R., Clough, J. W., Jezek, K. C., and Shabtaie, S.: Ice-thickness patterns and the dynamics of the Ross Ice Shelf, Antarctica, *J. Glaciol.*, 24(90), 287–294, 1979.
- Belina, F. A., Dafflon, B., Tronicke, J., and Holliger, K.: Enhancing the vertical resolution of surface georadar data, *J. Appl. Geophys.*, 68, 26–35, doi:10.1016/j.jappgeo.2008.08.011, 2009.
- Bogorodsky, V. V., Bentley, C. R., and Gudmandsen, P. E.: *Radioglaciology*, D. Reidel Publishing Company, Norwell, Massachusetts, ISBN 90-277-1893-9, 1985.

## GPR-derived accumulation and compaction of snow

N. C. Krueztmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





- Bracewell, R. N.: Fourier analysis and imaging, Kluwer Academic/Plenum Publishers, New York, New York, ISBN 0-306-48187-1, 2003.
- Davis, C. H. and Ferguson, A. C.: Elevation change of the antarctic ice sheet, 1995–2000, from ERS-2 satellite radar altimetry, *IEEE T. Geosci. Remote*, 42(11), 2437–2445, 2004.
- 5 Davis, C. H., Li, Y., McConnell, J. R., Frey, M. M., and Hanna, E.: Snowfall-driven growth in east antarctic ice sheet mitigates recent sea-level rise, *Science*, 308, 1898–1901, 2005.
- Drinkwater, M. R., Long, D. G., and Bingham, A. W.: Greenland snow accumulation estimates from satellite radar scatterometer data, *J. Geophys. Res.*, 106(D24), 33935–33950, 2001.
- 10 Dunbar, G. B., Bertler, N. A. N., and McKay, R. M.: Sediment flux through the McMurdo ice shelf in windless bight, *Antarctica, Global Planet. Change*, 69, 87–93, doi:10.1016/j.gloplacha.2009.05.007, 2009.
- Dunse, T., Eisen, O., Helm, V., Rack, W., Steinhage, D., and Parry, V.: Characteristics and small-scale variability of GPR signals and their relation to snow accumulation in Greenland's percolation zone, *J. Glaciol.*, 54(185), 333–342, 2008.
- 15 Eisen, O., Wilhelms, F., Nixdorf, U., and Miller, H.: Revealing the nature of radar reflections in ice: DEP-based FDTD forward modeling, *Geophys. Res. Lett.*, 30(5), 1218, doi:10.1029/2002GL016403, 2003.
- Eisen, O., Nixdorf, U., Wilhelms, F., and Miller, H.: Age estimates of isochronous reflection horizons by combining ice core, survey, and synthetic radar data, *J. Geophys. Res.*, 109, B04106, doi:10.1029/2003JB002858, 2004.
- 20 Eisen, O., Frezzotti, M., Genthon, C., Isaksson, E., Magand, O., van den Broeke, M. R., Dixon, D. A., Ekaykin, A., Holmlund, P., Kameda, T., Karlo, L., Kaspari, S., Lipenkov, V. Y., Oerter, H., Takahashi, S., and Vaughan, D. G.: Ground-based measurements of spatial and temporal variability of snow accumulation in East Antarctica, *Rev. Geophys.*, 46, RG2001, doi:10.1029/2006RG000218, 2008.
- 25 Harper, J. T. and Bradford, J. H.: Snow stratigraphy over a uniform depositional surface: spatial variability and measurement tools, *Cold Reg. Sci. Technol.*, 37, 289–298, doi:10.1016/S0165-232X(03)00071-5, 2003.
- Helm, V., Rack, W., Cullen, R., Nienow, P., Mair, D., Parry, V., and Wingham, D. J.: Winter accumulation in the percolation zone of Greenland measured by airborne radar altimeter, *Geophys. Res. Lett.*, 34(6), L06501, doi:10.1029/2006GL029185, 2007.
- 30 Hawley, R. L., Brandt, O., Morris, E. M., Kohler, J., Shepherd, A. P., and Wingham, D. J.: Techniques for measuring high-resolution firn density profiles: case study from Kongsvegen,

---

## GPR-derived accumulation and compaction of snow

N. C. Kruetzmann et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



---

**GPR-derived  
accumulation and  
compaction of snow**N. C. Kruetzmann et al.

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Svalbard, J. Glaciol., 54(186), 463–468, 2008.
- Heilig, A., Schneebeli, M., and Eisen, O.: Upward-looking ground-penetrating radar for monitoring snowpack stratigraphy, *Cold Reg. Sci. Technol.*, 59, 152–162, doi:10.1016/j.coldregions.2009.07.008, 2009.
- 5 Heine, A. J.: The McMurdo ice shelf antarctica – a preliminary report, *New Zeal. J. Geol. Geop.*, 10(2), 474–478, 1967.
- Irving, J. D. and Knight, R. J.: Removal of wavelet dispersion from ground-penetrating radar data, *Geophysics*, 68(3), 960–970, doi:10.1190/1.1581068, 2003.
- Kovacs, A., Gow, A. J., and Morey, R. M.: The in-situ dielectric constant of polar firn revisited, 10 *Cold Reg. Sci. Technol.*, 23, 245–256, 1995.
- Maeno, N.: Densification rates of snow at polar glaciers, *Mem. Natl. Ins. Polar Res., Special Issue*, 24, 48–61, 1982.
- McCrae, I. R.: A summary of glaciological measurements made between 1960 and 1984 on the McMurdo ice shelf, Antarctica: a report submitted to the Antarctic Division of D.S.I.R., 15 *Auckland, Dept. of Theoretical and Applied Mechanics: University of Auckland*, 1984.
- Nguyen, A. T. and Herring, T. A.: Analysis of ICESat data using Kalman filter and kriging to study height changes in East Antarctica, *Geophys. Res. Lett.*, 32, L23S03, doi:10.1029/2005GL024272, 2005.
- Parry, V., Nienow, P., Mair, D., Scott, J., Hubbard, B., Steffen, K., and Wingham, D.: Investigations of meltwater refreezing and density variations in the snowpack and firn within the percolation zone of the Greenland ice sheet, *Ann. Glaciol.*, 46, 61–68, 2007.
- Rial, F. I., Henrique, L., Pereira, M., and Armesto, J.: Waveform analysis of UWB GPR antennas, *Sensors*, 9, 1454–1470, doi:10.3390/s90301454, 2009.
- Rotschky, G., Rack, W., Dierking, W., and Oerter, H.: Retrieving snowpack properties and accumulation estimates from a combination of SAR and scatterometer measurements, *IEEE T. Geosci. Remote*, 44(4), 943–956, 2006.
- 25 *Sensors & Software Inc.: pulseEKKO PRO – User’s Guide*, Mississauga, Ontario, Canada, 2006.
- Spikes, V. B., Hamilton, G. S., Arcone, S. A., Kaspari, S., and Mayewski, P. A.: Variability in accumulation rates from GPR profiling on the West Antarctic plateau, *Ann. Glaciol.*, 39, 238–244, 2004.
- 30 *Taner, M. T., Koehler, F., and Sheriff, R. E.: Complex seismic trace analysis*, *Geophysics*, 44(6), 1041–1063, 1979.

- Takahashi, S. and Kameda, T.: Snow density for measuring surface mass balance using the stake method, *J. Glaciol.*, 53(183), 677–680, 2007.
- Turner, G.: Subsurface radar propagation deconvolution, *Geophysics*, 59(2), 215–223, 1994.
- Ulrych, T. J.: The whiteness hypothesis: reflectivity, inversion, chaos, and Enders, *Geophysics*, 64, 1512–1523, 1999.
- van den Broeke, M.: Depth and density of the antarctic firn layer, *Arct. Antarct. Alp. Res.*, 40(2), 432–438, 2008.
- Vaughan, D. G., Corr, F. J., Doake, C. S. M., and Waddington, E. D.: Distortion of isochronous layers in ice revealed by ground-penetrating radar, *Nature*, 398, 323–326, 1999.
- Walden, A. T. and Hosken, J. W. J.: An investigation of the spectral properties of primary reflection coefficients, *Geophys. Prospect.*, 33, 400–435, 1985.
- Wingham, D. J., Ridout, A. J., Scharroo, R., Arthern, R. J., and Shum, C. K.: Antarctic elevation change from 1992 to 1996, *Science*, 282, 456–458, 1998.
- Wingham, D. J., Francis, C. R., Baker, S., Bouzinac, C., Brockley, D., Cullen, R., de Chateau-Thierry, P., Laxon, S. W., Mallow, U., Mavrocordatos, C., Phalippou, L., Ratier, G., Rey, L., Rostan, F., Viau, P., and Wallis, D. W.: CryoSat: a mission to determine the fluctuations in Earth's land and marine ice fields, *Adv. Space Res.*, 37, 841–871, 2006.
- Xia, J., Franseen, E. K., Miller, R. D., and Weis, T. V.: Application of deterministic deconvolution of ground-penetrating radar data in a study of carbonate strata, *J. Appl. Geophys.*, 56, 213–229, 2004.
- Xiao, Q. N., Kuo, Y. H., Ma, Z. Z., Huang, W., Huang, X. Y., Zhang, X., Barker, D. M., Michalakes, J., and Dudhia, J.: Application of an adiabatic WRF adjoint to the investigation of the May 2004 McMurdo, Antarctica, severe wind event, *Mon. Weather Rev.*, 136(10), 3696–3713, doi:10.1175/2008MWR2235.1, 2008.
- Yilmaz, Ö.: *Seismic Data Processing*, Society of Exploration Geophysicists, Tulsa, OK, USA, 1987.
- Zwally, H. J. and Li, J.: Seasonal and interannual variations of firn densification and ice-sheet surface elevation at the Greenland summit, *J. Glaciol.*, 48(161), 199–207, 2002.

---

**GPR-derived  
accumulation and  
compaction of snow**

---

N. C. Kruetzmann et al.

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## GPR-derived accumulation and compaction of snow

N. C. Kruetzmann et al.

**Table 1.** Summary of accumulation measurements at L2 (77°45′ S, 167°17′ E) and L3 (77°40′ S, 167°11′ E).

Site	observation period	years	accumulation kg m <sup>-2</sup> a <sup>-1</sup>	measurement technique
L2 – 81 stakes	13.11.08–12.11.09	1	224 ± 21	stake reading
L2 – at stake C3	May 2004–Nov 2008	4.5	251	snow pit/dust layer
L2 – at stake G7	May 2004–Nov 2009	5.5	245	firn core/dust layer
L2	May 2004–Nov 2008	4.5	269 ± 9	GPR/dust layer
L3 – 24 stakes	18.11.08–9.11.09	0.97	304 ± 83	stake reading
L3 – at stake C3	May 2004–Nov 2009	5.5	437	firn core/dust layer
L3	May 2004–Nov 2009	5.5	404 ± 22	GPR/dust layer
14 km transect	May 2004–Nov 2008	4.5	270 to 165	GPR/dust layer

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back


Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion






**Fig. 1.** (a) The Ross Sea region. (b) Measurement area in the Western Ross Sea region and corner points of stake farms on the McMurdo Ice Shelf (L1 and L2) and on Ross Island (L3). The circles between L1 and L2 indicate the locations of additional stakes installed for accumulation and radar measurements. (c) Outline of the measurement grid and numbering of the stake farms (Envisat ASAR image courtesy ESA).



**Fig. 2. (a)** One hundred superimposed shots recorded with facing antennae. The air-wave and the ground reflection can be clearly distinguished. **(b)** The spectrum of the averaged air-wave shows the centre frequency to be around 620 MHz.

## GPR-derived accumulation and compaction of snow

N. C. Kruezmann et al.



**Fig. 3.** Radar profile at L2 between the stakes C2 and C4 (2008) after applying **(a)** dewow and gain filters, **(b)** the full processing excluding- and **(c)** including deconvolution. The yellow arrow in **(c)** indicates the depth and location of the dust layer identified in the snow pit. The vertical scale of the image is exaggerated, representing 200 m in the horizontal direction and about 13 m vertically.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**GPR-derived  
accumulation and  
compaction of snow**

N. C. Kruetzmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures




Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



**Fig. 4.** ACF of one radar trace at L2 before (dashed line) and after deconvolution (solid line).






**Fig. 5.** Snow density profiles in 2009 from snow pits (grey shaded area) and firn cores at sites **(a)** L2 and **(b)** L3. In both cases, the snow pit was logged according to visually identified stratigraphy, while the core was largely logged in 10 cm intervals.



## GPR-derived accumulation and compaction of snow

N. C. Kruetzmann et al.



**Fig. 7.** Accumulation maps for (a) L2 and (b) L3, based on the reflection horizon associated with the dust layer tracked along each line. A four metre interpolation grid resolution was used in both cases.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



**GPR-derived  
accumulation and  
compaction of snow**

N. C. Kruezmann et al.



**Fig. 8.** Processed radargram from line C1 to C9 at site L3 (2009). The red box marks the approximate location of the firn core; the yellow arrow indicates the expected TWT to the dust layer.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back


Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Fig. 9.** Accumulation along a transect from L2 to L1, estimated by tracking the dust layer reflection for 14 km.

**GPR-derived  
accumulation and  
compaction of snow**

N. C. Kruetzmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**GPR-derived  
accumulation and  
compaction of snow**

N. C. Kruetzmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶


Back

Close

Full Screen / Esc

Printer-friendly Version


Interactive Discussion



**Fig. 10.** Snow compaction along the line from stake E1 to E9 at L2, calculated from the change in the average reflection horizon separations between Fig. 6c and d. Horizontal error bars represent the sum of the standard errors of the two corresponding horizons' depths.

**GPR-derived  
accumulation and  
compaction of snow**

N. C. Kruezmann et al.



**Fig. 11.** Snow compaction vs. depth for (a) L2 and (b) L3. (a) shows the compaction calculated from six lines at L2, while (b) includes all lines from L3 except one. The vertical axis represents depth before compaction has occurred. The horizontal error bars are the same as in Fig. 10 and the vertical error bars illustrate the thickness of the layers before compaction. The black line represents the average compaction for each horizon pair.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

