



## Abstract

Rising sea levels and increased surface melting of the Greenland ice sheet have heightened the need for direct observations of meltwater release from the ice edge to ocean. Buoyant sediment plumes that develop in fjords downstream of outlet glaciers are controlled by numerous factors, including meltwater runoff. Here, Moderate Resolution Imaging Spectroradiometer (MODIS) satellite imagery is used to average surface suspended sediment concentration (SSC) in fjords around ~80 % of Greenland from 2000–2009. Spatial and temporal patterns in SSC are compared with positive-degree-days (PDD), a proxy for surface melting, from the Polar MM5 regional climate model. Over this decade significant geographic covariance occurred between ice sheet PDD and fjord SSC, with outlet type (land- vs. marine-terminating glaciers) also important. In general, high SSC is associated with high PDD and/or a high proportion of land-terminating glaciers. Unlike previous site-specific studies of the Watson River plume at Kangerlussuaq, temporal covariance is low, suggesting that plume dimensions best capture interannual runoff dynamics whereas SSC allows assessment of long-term conditions across much broader fjord environments around the ice sheet. Remote sensing of both plume characteristics thus offers a viable approach for observing spatial and temporal patterns of meltwater release exiting the Greenland ice sheet to the global ocean.

## 1 Introduction

The Greenland ice sheet is undergoing increasing melt intensity and extent (Mote, 2007; Tedesco, 2007; Bhattacharya et al., 2009) in response to warming air temperatures (Hanna et al., 2008; Tedesco et al., 2008; Box et al., 2009). Ice mass loss has accelerated in the last decade (Chen et al., 2006; Rignot et al., 2008; Wouters et al., 2008; Allison et al., 2009; Velicogna, 2009), with increasing accumulation in the ice sheet interior (Box et al., 2006; Burgess et al., 2010) exceeded by losses in the

## Hydrologic controls on coastal suspended sediment plumes

V. W. Chu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Hydrologic controls on coastal suspended sediment plumes

V. W. Chu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

marginal ablation zone (Luthcke et al., 2006; Ettema et al., 2009; Zwally et al., 2011). Losses are exponentially higher at the margin (van den Broeke et al., 2008) with rapid thinning of near-coastal outlet glaciers (Krabill et al., 2004; Sole et al., 2008; Pritchard et al., 2009; Thomas et al., 2009). Marine-terminating outlet glaciers have also shown increases in total ice discharge (Rignot et al., 2004; Howat et al., 2007) and velocity (Rignot and Kanagaratnam, 2006; Joughin et al., 2010), with velocity speedups recently extending to the northwest (Khan et al., 2010). By the end of this century, Greenland's contribution to global sea level rise may total  $\sim 17\text{--}54$  cm (Pfeffer et al., 2008), and perhaps reach an annual rate  $\sim 0.7\text{--}0.8$  mm yr<sup>-1</sup> (Fettweis et al., 2008).

While ice discharge is the primary form of mass loss for most marine-terminating outlet glaciers (Mernild et al., 2010a), meltwater runoff possibly contributes more than half the total mass loss for the ice sheet as a whole (van den Broeke et al., 2009). Mass loss estimates using Gravity Recovery and Climate Experiment (GRACE) gravity data also require knowledge of meltwater runoff, but must currently use modeled estimates rather than direct observations (Velicogna et al., 2005). Increased meltwater production has also been linked to ice velocity increases in fast moving outlet glaciers (Joughin et al., 1996; Shepherd et al., 2009; Andersen et al., 2010), as well as seasonal speedups of the broader, slower moving ice sheet (Zwally et al., 2002; Joughin et al., 2008a; van de Wal et al., 2008; Palmer et al., 2011). Meltwater can be transported to the bed through moulins and possibly well-developed englacial drainage networks (Catania and Neumann, 2010), with large inputs of meltwater from supraglacial lakes decreasing basal friction and increasing short-term ice velocities (Box and Ski, 2007; Das et al., 2008; Schoof, 2010). While meltwater processes are less important to marine-terminating glaciers than are destabilized calving fronts (Thomas et al., 2003; Amundson et al., 2008; Joughin et al., 2008b) and enhanced ice-bottom melting from warm ocean waters (Holland et al., 2008; Howat et al., 2008; Rignot and Steffen, 2008; Straneo et al., 2010; Motyka et al., 2011; Yin et al., 2011), dynamic changes on land-terminating ice are attributed to bedrock lubrication from increased meltwater (Zwally et al., 2002; Sundal et al., 2009; Bartholomew et al., 2010).

## Hydrologic controls on coastal suspended sediment plumes

V. W. Chu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A prime obstacle to quantifying and incorporating runoff processes into models of ice sheet dynamics is a scarcity of direct observations of meltwater exiting the ice sheet, both in rivers draining the ice sheet and from beneath marine-terminating glaciers (e.g., Rignot and Steffen, 2008). Therefore, the amount of meltwater that truly reaches the ocean (rather than refreezing or being retained by the ice sheet) is presently unknown. Meltwater production on the ice sheet surface can be modeled from climate data (Box et al., 2006; Fettweis, 2007; Ettema et al., 2009; Robinson et al., 2010), or observed using remote sensing (Abdalati and Steffen, 1997; Smith et al., 2003; Mote, 2007; Tedesco, 2007; Hall et al., 2008). However, its release from the ice sheet edge to the ocean remains largely unstudied, consisting of a handful of modeling efforts (Bøggild et al., 1999; Lewis and Smith, 2009; Mernild et al., 2010b, 2011) and site-specific field studies (Rasch et al., 2000; Stott and Grove, 2001; Chu et al., 2009; Mernild and Hasholt, 2009; McGrath et al., 2010).

Buoyant sediment plumes that develop in fjords downstream of outlet glaciers and rivers offer a link between ice sheet hydrology and the ocean that can plausibly be observed using satellite remote sensing (Chu et al., 2009; McGrath et al., 2010). Sediment is produced by abrasion as ice moves over underlying bedrock and subsequently transported by meltwater, with sediment output affected by glaciological variables like glacier size, sliding speed, ice flux, and meltwater production, as well as erosional susceptibility of the bedrock (Hallet et al., 1996). Plumes are formed when sediment-rich freshwater runoff from the ice sheet enters the fjord – either directly, for marine-terminating glaciers, or via rivers, for land-terminating glaciers – and floats over denser saline marine water. As meltwater enters the fjord, a buoyant plume typically develops provided sediment concentrations do not exceed  $\sim 40\,000\text{ mg l}^{-1}$  (Mulder and Syvitski, 1995). These features are readily observed in satellite imagery, allowing remote estimation of water quality characteristics including suspended sediment concentration (SSC) (e.g., Curran and Novo, 1988; Doxaran et al., 2002; Hu et al., 2004; Miller and McKee, 2004). The area and length of buoyant plumes have also been measured as a proxy for hydrologic outflows from the land surface to ocean (e.g., Thomas and

Weatherbee, 2006; Halverson and Pawlowicz, 2008; Lihan et al., 2008; Chu et al., 2009; McGrath et al., 2010).

In the upper fjord environment where rivers first enter the coastal zone, plume spreading and mixing are driven predominantly by the kinetic energy of river discharge (Syvitski et al., 1985), but plume characteristics are still controlled by a complex combination of factors both on land and after entering the fjord. Sediment-rich meltwater from land-terminating outlet glaciers may encounter lakes, outwash plains, or braided river valleys, all of which can act as traps or sources for sediment (Busskamp and Hasholt, 1996; Hasholt, 1996). For marine-terminating outlet glaciers, sediment export to the ocean is dominated by the distinctly different mechanisms of iceberg rafting and/or en- and sub-glacially transported meltwater runoff (Andrews et al., 1994). In both environments, as plumes move farther downstream, sediment transport and settling rates are further influenced by tides (Castaing and Allen, 1981; Dowdeswell and Cromack, 1991; Bowers et al., 1998; Halverson and Pawlowicz, 2008), wind (Stumpf et al., 1993; Whitney and Garvine, 2005), and sea ice (Hasholt, 1996).

Here, buoyant sediment plumes that develop in upper fjord environments immediately downstream (~15–20 km, with a maximum of 50 km) of outlet glaciers and rivers that drain the Greenland ice sheet are mapped and analyzed using optical satellite imagery, to identify the distribution and temporal characteristics of sediment and meltwater release to coastal waters. Of particular interest is how well meltwater production on the ice sheet accounts for observed spatial and temporal variations in SSC, and to what extent outlet glacier environments complicate this relationship. SSC is used instead of plume area or length (Chu et al., 2009; McGrath et al., 2010) in order to expand the method beyond a river mouth. Optical images from the Moderate Resolution Imaging Spectroradiometer (MODIS) satellite are used from 2000–2009 to sample buoyant plume SSC in ~230 fjords with near-daily temporal resolution at 500 m spatial resolution. These observations are then compared with a proxy for ice sheet surface melting (Polar MM5 modeled positive degree-days, PDD), routed through potential drainage basins derived from ice surface and bedrock topography (Lewis and Smith, 2009), as

## Hydrologic controls on coastal suspended sediment plumes

V. W. Chu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



well as geomorphic characteristics. The end result is a synoptic, ten-year year analysis of spatiotemporal plume behavior around Greenland and a first assessment of some important controls on their distribution and development.

## 2 Data and methods

5 To explore controls on sediment plume development, we considered (1) daily ice sheet surface melt using modeled PDD, routed into the fjords following potential drainage basins; (2) near-daily fjord SSC from calibrated MODIS satellite imagery aggregated into 100 km coastal gridcells; and (3) outlet glacier environments. Each of these steps is described in the following Sects. 2.1–2.3 next.

### 10 2.1 Ice sheet surface melt

A key driver of sediment plume behavior explored here is ice sheet hydrology as represented by production of meltwater on the ice sheet surface. The fifth generation Polar Mesoscale Model (PMM5) provides a gridded 24 km resolution output of 3-hourly temperatures across the ice sheet surface from 2000–2009 (Box et al., 2006). Data  
15 were provided in a polar stereographic projection and a mask was applied to extract temperature data over the ice sheet. From these data, time series of daily positive degree-days were extracted by averaging the three-hourly temperatures greater than 0°C for each day. PDD is a traditional measure of melt intensity based on relating the cumulative depth of ice and snow melt to the sum of positive air temperatures over a  
20 specified time interval, usually a day. It is widely used because of its simplicity and has outperformed more sophisticated energy balance models for use in catchment-scale hydrologic modeling (Ohmura, 2001; Hock, 2003). Here, positive degree days are used in its un-cumulative form.

As a proxy for meltwater volume produced within each hydrologic drainage basin,  
25 the aforementioned PDD data were totaled over topographically determined-basins

## Hydrologic controls on coastal suspended sediment plumes

V. W. Chu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and assumed to drain only to corresponding ice sheet outlet glaciers and rivers at the ice sheet edge.

The drainage basins, each unique to a 100 km coastal gridcell (for fjord sediment detection, described in Sect. 2.2.3) were defined using a previously derived vector dataset of ice sheet drainage basins based on potentiometric flow networks (Lewis and Smith, 2009), modeled from a combination of bedrock topography and surface topography by assuming hydrostatic pressure conditions and no conduit flow within the ice sheet. Basins were aggregated as necessary to correspond to each 100 km coastal gridcell, with final drainage basin area,  $B$ , and ice edge length,  $l$ , defined as the total length of the ice sheet edge bounded within each drainage basin. Melt area,  $A_{\text{PDD}}$ , was calculated for each drainage basin by totaling the number of 24 km pixels with a daily PDD greater than  $0^{\circ}\text{C}$ . The fraction of each drainage basin experiencing active melting,  $F_{\text{PDD}}$ , was defined as melt area divided by drainage basin size ( $F_{\text{PDD}} = \frac{A_{\text{PDD}}}{B}$ ). Melt penetration distance,  $D_{\text{PDD}} = \frac{A_{\text{PDD}}}{l}$ , represented the average inland distance from the ice edge that experienced surface melting.

## 2.2 Remote sensing of sediment plumes

Methodology for MODIS remote sensing of sediment plumes is shown in the Fig. 1 flowchart. SSC was estimated by (1) classifying 10 years of available daily MODIS imagery into ice-free “open water” (ranging from clear water to sediment-rich water) areas in the fjords, (2) aggregating the 500 m data into 100 km gridcells to attain a high frequency temporal sampling, and (3) transform MODIS reflectance into SSC using an empirical relationship developed from field water samples.

### 2.2.1 MODIS 500 m satellite imagery and quality

The MODIS instrument on NASA’s Terra satellite acquired daily coverage over Greenland from 2000–2010 with seven bands in the visible and infrared spectra at 500 m spatial resolution and two bands at 250 m resolution. Time series of MODIS Level 2

## Hydrologic controls on coastal suspended sediment plumes

V. W. Chu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



500 m surface reflectance product (MOD09) (Vermote et al., 2002), atmospherically corrected for gases, aerosols, and thin cirrus clouds, was used for the melt season (1 May–30 September) each year. These Level 2 data are aggregated into a daily product and available as tiles in a sinusoidal grid projection. Seven MODIS tiles were needed to cover all of Greenland (Fig. 2). MODIS data are freely available and were downloaded from the NASA Warehouse Inventory Search Tool (<https://wist.echo.nasa.gov/api/>). Note that while MODIS data were available for 2010, PMM5 PDD data were only available until 2009. Therefore, MODIS data were processed and displayed for 2010 but not included in 10-year averages for comparison with PDD.

Only high-quality “clear-sky” MODIS pixels were used from each daily MODIS image. High-quality clear-sky images were defined as having: (1) a near-nadir view with adequate solar illumination (i.e., satellite overpass between 13:00 and 17:00 UTC), (2) minimal cloud cover, and (3) minimal atmospheric interference. Atmospheric interference and cloud cover were determined using MODIS 1 km resolution Quality Assurance (QA) datasets that provided quality flags for each band, including cloud state and aerosol quantity.

### 2.2.2 Classification and validation of “open water”

“Open water” pixels were defined as high quality MODIS pixels ranging from clear water to sediment-rich water, free of both clouds and ice, and distinguished using reflectance thresholds. MODIS band 1 (620–670 nm), band 2 (841–876 nm), band 3 (459–479 nm), band 4 (545–565 nm), and band 6 (1628–1652 nm) were used with thresholds in a simplified classification scheme to mask out land, ice (including land-fast ice, sea ice, and calving icebergs), and clouds. Particular difficulty in distinguishing sediment-rich water from melting ice was due to similar spectral responses in the seven available MODIS bands. Land and river pixels were identified primarily by a lower reflectance in band 2 than band 1. To distinguish clouds and ice, both of which show high reflectance in the visible bands, band 6 was used. Clouds were identified with high band 6 reflectance > band 1 reflectance. Remaining pixels with lower band 6 values

## Hydrologic controls on coastal suspended sediment plumes

V. W. Chu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



were classified into ice (and melting ice) and open water. Ice was distinguished with band 2 reflectance greater than  $0.5 \times$  band 1 reflectance. Finally, open water (OW) was then produced as a range of clear water to sediment-rich water free of clouds and ice.

The “open water” pixel classification was verified manually using 10 ASTER Level 2 15 m surface reflectance (AST 07) images and 27 Landsat TM/ETM+ 30 m images (Fig. 2), taking advantage of their higher spatial resolutions to between sediment-rich water and melting sea ice and icebergs from calving glaciers. Melting ice proved difficult to discern in the MODIS data, both spatially owing to the low resolution and spectrally with just seven available bands at 500 m resolution. Scenes for both sensors were selected to be during the summer melt season where plumes could be expected as well as in areas representing contrasting outlet-water interface types and different spatial locations along the coast of Greenland.

Data density and sediment persistence were calculated for each 500 m MODIS pixel from the classified open water data to characterize the frequency of data recovery and high sediment concentration. Data density,  $DD_{500\text{ m}}$ , was defined as the percentage of non-ice, non-cloud open water days over the entire period, and ranges from 0 (no open water) to 100 % (open water detected every day),  $DD_{500\text{ m}} = \frac{OW}{\text{total available data}}$ . While open water is the primary classification used in further analysis, a sediment persistence metric,  $F_{SSC}$ , was used to distinguish high SSC from low SSC to understand the temporal aspects of highly concentrated sediment lingering in the fjords. The threshold of band 1  $> 0.12$  developed in Chu et al. (2009) was used for identifying the highest sediment concentrations, designated the “plume” as opposed to the “brackish plume” in the sediment-rich Kangerlussuaq Fjord, so the same threshold used all around Greenland should identify only the highest SSCs and provide a conservative estimate of persistence. Sediment persistence was defined as the fraction of high-SSC days ( $OW_{SSC}$ ) out of OW ( $F_{SSC} = \frac{OW_{SSC}}{OW}$ ).

## Hydrologic controls on coastal suspended sediment plumes

V. W. Chu et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

### 2.2.3 MODIS spatial sampling and aggregation

Twenty-eight hundred regions of interest (ROIs) covering 7500 km<sup>2</sup> over 230 fjords were manually delineated to enable multiple MODIS sampling of all fjords directly draining the ice sheet via land-terminating and marine-terminating glaciers (Fig. 3). This restriction of analysis to fjords immediately draining the ice sheet reduced sampling of plumes triggered by melting snow packs, coastal erosion, and other sedimentary processes not necessarily triggered by ice sheet meltwater runoff. ROIs were typically digitized within ~15–20 km and not more than 50 km of river mouths and outlet glacier termini, rather than further down-fjord or in the open ocean.

To reduce the loss of open water data from clouds and obtain more precise temporal sampling, the 500 m native-resolution MODIS data were aggregated into 100 km × 100 km coastal gridcells to yield the final dataset for all further analyses (Fig. 3). First, a 100 km fishnet was overlaid onto the seven mosaicked MODIS tiles to summarize the SSC data within the ROIs. MODIS tiles were reprojected from the original sinusoidal projection to match the PMM5 model output polar sinusoidal projection. For each 100 km gridcell,  $R_{\text{peak}}$  was determined from the population of ROI pixels within the gridcell.  $R_{\text{peak}}$  was defined as the median of the top 20 OW MODIS band 1 reflectance values (based on empirical model described in Sect. 2.2.4) to avoid biases from ROI placement or number of ROIs per 100 km gridcell. Furthermore, to help mitigate the effects of data loss from cloud and ice interference, a 7-day moving interval was applied over the raw data in conjunction with the spatial resampling, effectively allowing the resulting daily value to derive from a sample population anywhere within the 100 km gridcell ROIs anywhere from three days before to three days after the day of interest. While this assumes that a sample from a partially cloudy 100 km gridcell is equivalent to a cloud-free box,  $R_{\text{peak}}$  represents the average plume state within a week, allowing a week for the best pixels to be sampled.

## Hydrologic controls on coastal suspended sediment plumes

V. W. Chu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 2.2.4 Calibration/validation of SSC

The final data product of daily  $R_{\text{peak}}$  aggregated into 100 km coastal gridcells was transformed into SSC using an empirical relationship between remotely sensed reflectance and in situ measurements of SSC.

Field samples of SSC were necessary to understand plume characteristics, how SSC relates to the presence of freshwater, and how varying levels of sediment affect the spectral reflectance of the water. In situ water quality data were collected 3 June 2008 in Kangerlussuaq Fjord, southwest Greenland, with surface measurements of SSC, salinity, spectral reflectance, optical depth, and temperature collected every 1 km along a 22 km transect as described in Chu et al. (2009). Additional surface water samples from a marine-terminating fjord ~80 km north of Illulisat were collected 4 July 2007 and culled for the points overlapping with digitized ROIs near the coast where buoyant sediment plumes were found, yielding four locations around 69.79° N, 50.53° W (Fig. 2). These sites provided laboratory measurements of SSC but did not include in situ spectral reflectance. These additional measurements supplemented the more extensive Kangerlussuaq Fjord dataset by providing in situ SSCs from an environment dominated by marine-terminating glaciers. An empirical model relating SSC to MODIS reflectance (Fig. 4) was constructed using all available field samples and simultaneous MODIS band 1 (620–670 nm) reflectance as per Chu et al. (2009), yielding a new revised model of:

$$R(\text{band 1}) = 3.02\ln(\text{SSC}) + 1.12R^2 = 0.86$$

with  $R(\text{band 1})$  as the reflectance (%) for MODIS band 1 (620–670 nm) and SSC measured in  $\text{mg l}^{-1}$ . The model shows reflectance very sensitive to lower SSCs but saturation at high SSCs beyond  $100 \text{ mg l}^{-1}$ .

A new data density measurement,  $\text{DD}_{100 \text{ km}}$ , was calculated from the aggregated data, and a threshold of  $\text{DD}_{100 \text{ km}} > 45\%$  was applied to produce the final gridcells used for further analysis.

TCD

5, 2365–2407, 2011

### Hydrologic controls on coastal suspended sediment plumes

V. W. Chu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## 2.3 Outlet glacier environment

Outlet glacier environment provides insight into the physical mechanisms by which sediment is dispersed from glacier outlets to fjords. While sediment transport in rivers (e.g., Hasholt, 1996; Rasch et al., 2000; Knudsen et al., 2007; Russell, 2007; Hasholt and Mernild, 2008) and sediment deposition in fjords (e.g., Syvitski et al., 1996; Reeh et al., 1999; Mugford and Dowdeswell, 2010) have previously been studied in Greenland, the active sediment plumes themselves are less studied (Chu et al., 2009; Lund-Hansen et al., 2010; McGrath et al., 2010). Though glacial erosion is responsible for some of the largest sediment yields to the ocean (Gurnell et al., 1996; Hallet et al., 1996), this paper focuses specifically on the fine glacial sediments transported by meltwater and remaining in suspension in coastal waters, rather than total sediment flux and/or deposition processes.

Outlet glacier type (marine- or land-terminating) was used to determine whether sediment concentrations differed based on the type of outlet meltwater source. Lewis and Smith (2009) provide georeferenced locations of all confirmed glacier meltwater outlets (i.e., land-terminating glaciers ending in rivers or lakes or marine-terminating glaciers with the presence of a sediment plume) and all unconfirmed glacier meltwater outlets (i.e., marine-terminating glaciers with no visible plume). These three categories of outlet types were further generalized into marine-terminating and land-terminating glacier outlets, and only those that led into a corresponding 100 km coastal gridcell fjord area were counted and summarized within the gridcell. Marine-terminating outlets ranged from those with visible plumes and minimal iceberg activity to those heavily calving icebergs forming a “sikussak” complex of fused icebergs and sea ice attached to the glacier terminus (Syvitski, 1996). The ASTER/ETM+ remote sensing validation process also proved useful for identifying gridcells dominated by fjords with heavily calving marine-terminating glaciers and no visible plume. Land-terminating outlets release meltwater at the edge through proglacial lakes or rivers, and only those that eventually transport meltwater to the fjord through rivers or floodplains were included here. The

### Hydrologic controls on coastal suspended sediment plumes

V. W. Chu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



number of land-terminating outlets ( $N_L$ ) and number of marine-terminating outlets ( $N_M$ ) for each drainage basin were determined using the outlets in Lewis and Smith (2009). The counts were normalized by the total number of outlets to compute fractions of land-terminating glaciers ( $F_L = \frac{N_L}{N_L+N_M}$ ) and marine-terminating glaciers ( $F_M = \frac{N_M}{N_L+N_M}$ ).

### 3 Results

MODIS-derived estimates of coastal fjord SSC ranging from  $\sim 0.7$  to  $1925 \text{ mg l}^{-1}$  were retrieved all around the ice sheet except in northern Greenland where persistent sea ice precludes detection of open water (Fig. 5, Table 1). Though the maximum SSC value from all gridcells over 10 years is extrapolated from the empirical model relating MODIS reflectance to SSC beyond field data values, it is within the range of SSC values observed in other studies (Hasholt, 1996). PMM5-derived PDD totals for ice sheet hydrologic basins draining into 100 km coastal gridcells, range from 0 to  $165^\circ \text{C per day}$ . Highest intensity is found along the ice sheet edge, decreasing exponentially farther inland with increasing elevations (Fig. 5, Table 1). Table 1 displays 10-year mean values for each parameter and each coastal gridcell/drainage basin pair. Basin mean melt area ( $A_{\text{PDD}}$ ) ranges from  $32\text{--}16\,277 \text{ km}^2$ , with mean melt penetration distance ( $D_{\text{PDD}}$ ) ranging from  $0.3\text{--}112 \text{ km}$  inland. Drainage basin size ( $B$ ) ranges from  $553\text{--}142\,175 \text{ km}^2$  with ice edge lengths ( $I$ ) varying between 35 and 344 km. Adequate MODIS data density is found for 47 out of 83 coastal 100 km gridcells. The raw, native-resolution 500 m data show that on average a Greenland coastal ROI pixel experiences at most 26 % of all days between 1 June and 30 September classified as open water (on average 14 %), that is, a water pixel without cloud or ice interference, from a total of 1531 days over the ten years ( $\text{DD}_{500 \text{ m}}$ ). On average the coastal gridcells contain 5 marine-terminating glacier outlets ( $M$ ) and 4 land-terminating glacier outlets ( $L$ ). Spatially and temporally aggregated 100 km gridcells for SSC show an improved data density, with a minimum threshold of  $\text{DD}_{100 \text{ km}} > 45\%$  yielding the 47 gridcells for further analysis and a mean  $\text{DD}_{100 \text{ km}}$  of 75 %. 10-year mean sediment persistence ( $F_{\text{SSC}}$ ) averages 0.11 for the

## Hydrologic controls on coastal suspended sediment plumes

V. W. Chu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



entire ice sheet, meaning gridcells show high-SSC values for one-tenth of the open water days detected on average.

### 3.1 Regional characteristics: mean SSC, mean PDD, data availability, and outlet glacier environment

The aforementioned characteristics of Greenland as a whole mask strong regional differences around the edge of the ice sheet. In Table 1, the 100 km coastal gridcells are further aggregated into six regions (Fig. 3) based on climatic regions of Ohmura and Reeh (1991), typical values of observed SSCs, and dominant outlet glacier types. As these regions are considerably different from Ohmura and Reeh, a new naming convention is used, i.e., Northwest Region, West Region, Southwest Region, Southeast Region, East Region, and Northeast Region. Their Humoldt region is the Northwest region, their Jakobshavn region is split into the West and Southwest regions, their Godthab and Julianehab regions are merged into the Southwest region, their Angmagssalik region is the Southeast Region, their Scoresbysund region is the East regions, and their Tunu region is the Northeast region.

The Northwest Region (Fig. 3) consists of four coastal gridcells (numbered 1–4) with the smallest average  $B$  ( $4214 \text{ km}^2$ , Table 1) and exhibits a moderately low 10-year average SSC ( $55 \pm 63 \text{ mg l}^{-1}$ ), except for one uniquely high SSC gridcell (Gridcell 2), and a moderately low average PDD ( $1.3 \pm 0.5 \text{ }^\circ\text{C}$ ). Gridcell 2, off the coast west of Humboldt Glacier, has a mean SSC of  $162 \pm 4 \text{ mg l}^{-1}$ , above average not only for the region but for the entire ice sheet as well and a high number of land-terminating glaciers ( $N_L = 10$ ). In contrast, Gridcell 4 directly to the south captures a high number of marine-terminating glaciers ( $N_M = 14$ ) with a large swath of glaciers near to the coast, generating one of the lowest mean SSC values ( $4 \text{ mg l}^{-1}$ ). This region has a high  $DD_{100 \text{ km}}$  (86 %) and a high  $F_{\text{SSC}}$  (0.11), meaning high concentrations of sediment are detected for one-tenth of the melt season, with many protected fjords and a relatively high average  $N_L$ .

The West Region (Fig. 3) has eight coastal gridcells (numbered 5–12), including several large marine-terminating glaciers including Jakobshavn Isbrae (Gridcell 12),

## Hydrologic controls on coastal suspended sediment plumes

V. W. Chu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Store Glacier (Gridcell 11), and Rink Glacier (Gridcell 9). The West Region has a mean SSC of  $57 \pm 47 \text{ mg l}^{-1}$  (Table 1) similar to the Northwest Region, but a higher mean PDD ( $5.9 \pm 3.8^\circ\text{C}$ ). This region contains high velocity marine-terminating glaciers that transition from mostly glaciers near the coast to longer glaciers with floating tongues or are near-floating (Thomas et al., 2009) farther south. This region has the largest average  $B$  ( $49\,131 \text{ km}^2$ ) due to the Jakobshavn basin, the largest one of all the gridcells at a size of  $99\,210 \text{ km}^2$ . A moderate  $DD_{100\text{km}}$  (77%) and a moderately low  $F_{\text{SSC}}$  (0.07) reflect the mix of outlet and fjord types, with a low average  $N_L$  (2) and a moderate average  $N_M$  (5), but the highest fraction of marine-terminating glaciers ( $F_M = 0.77$ ).

The Southwest Region (Fig. 3) is made up of eleven coastal gridcells (numbered 13–23) and has the highest average values in most parameters: highest mean SSC ( $262 \pm 168 \text{ mg l}^{-1}$ , Table 1), highest mean PDD ( $12.5 \pm 8.2^\circ\text{C}$ ), highest average  $A_{\text{melt}}$  ( $6002 \text{ km}^2$ ), highest average  $F_{\text{SSC}}$  (0.21), highest average  $N_L$  as well as  $F_L$  (0.81), and the highest  $DD_{100\text{km}}$  (89%). In particular, Gridcell 14, which encompasses Russell Glacier and the downstream Kangerlussuaq Fjord, shows the highest mean SSC ( $555 \pm 422 \text{ mg l}^{-1}$ ) and by far the highest mean PDD ( $30.8 \pm 34.3^\circ\text{C}$ ). This region encompasses the southwest with its gently sloping coast and its warmer climate giving way to the geomorphological uniqueness of this region. The extensive coastal land area affords every gridcell higher  $F_L$  than  $F_M$  and many large braided rivers to disperse the sediment into long, protected fjords. Going farther south in the region with less land area shows decreased mean SSC, decreased mean PDD, and decreased  $F_{\text{SSC}}$ .

The Southeast Region (Fig. 3) has eleven coastal gridcells (numbered 24–34), encompassing the same latitude range as the Southwest Region, and shows a marked difference from the Southwest with very low averages SSC ( $27 \pm 29 \text{ mg l}^{-1}$ , Table 1), PDD ( $1.1 \pm 0.7^\circ\text{C}$ ),  $F_{\text{SSC}}$  (0.06). A moderate average  $DD_{100\text{km}}$  (73%) despite loss of OW from icebergs reflects the removal of those gridcells that did not meet the data density threshold, particularly those encompassing the heavily calving Kangerdlugssuaq and Helheim Glaciers. Persistent sikussaks at the ends of these glaciers extend several kilometers and prevent any detection of OW. These glaciers also have very large

## Hydrologic controls on coastal suspended sediment plumes

V. W. Chu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

drainage basins, so their removal is indicated by a fairly small average  $B$  (9142 km<sup>2</sup>). In contrast with the southwest, the southeast has the highest average  $N_M$  (11, also highest  $F_M = 0.90$ ) and lowest average  $N_L$  (1, also lowest  $F_L = 0.1$ ). Most glaciers are near to the coast with steeper slopes and fairly low melting.

The East Region (Fig. 3) consists of six coastal gridcells (numbered 35–40) around the Scoresby Sund area and shows the lowest mean SSC ( $22 \pm 11$  mg l<sup>-1</sup>, Table 1) and the lowest mean PDD ( $0.9 \pm 0.4$  °C). Average  $F_{SSC}$  is also lowest of all regions (0.05), and two gridcells (Gridcells 35 and 36) do not show any sediment persistence, indicating that only very low concentrations of sediment exist in those areas. Average DD<sub>100 km</sub> is one of the lowest (61%), and outlets are fairly abundant in both marine-terminating and land-terminating types draining into protected fjords with  $F_M$  and  $F_L$  both 0.5. The presence of a large land area as well as some islands allows for smaller glaciers and snowpatches not connected to the ice sheet to produce sediment, so ROIs were placed to avoid those areas as much as possible.

The Northeast Region (Fig. 3) has seven coastal gridcells (numbered 41–47). This region has the lowest average DD<sub>100 km</sub> (58%), and the northernmost Gridcell 47 has the lowest individual gridcell DD<sub>100 km</sub> (48%). The low DD<sub>100 km</sub> of open water pixels free of ice is problematic in the northeast as well as in the northern areas; open water detection is prevented by the persistence of sea ice and/or iceberg calving. This region has the lowest mean PDD ( $0.8 \pm 0.7$  °C), but an intermediate mean SSC ( $63 \pm 41$  mg l<sup>-1</sup>). Both land-terminating and marine-terminating outlets are fairly low in number, with slightly more  $N_L$  (4) compared  $F_M$  (3).

## 3.2 Hydrologic controls on fjord SSC: spatial, interannual, seasonal, and high frequency

### 3.2.1 Spatial variability

A high spatial correlation between 2000–2009 PDD and SSC confirms that more ice sheet melting leads to more suspended sediment being mobilized by meltwater.

## Hydrologic controls on coastal suspended sediment plumes

V. W. Chu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Figure 5b shows a strong correlation ( $R = 0.70$ ,  $p < 0.001$ ) between the 10-year mean PDD and 10-year mean SSC moving counterclockwise around the ice sheet from the northwest to the northeast. Mean intensities of PDD and SSC for individual gridcells illustrate results from the grouped regions. The Northwest and West Regions show a slightly decoupled intensity between PDD and SSC, with Gridcell 2 in the Northwest experiencing a low average PDD but high average SSC, and Gridcells 5–8 in the West revealing higher average PDDs with lower average SSCs. The Southwest Region is distinct in high values of both, with a few gridcells (Gridcells 16–19) associated with high plume SSC's despite low ice sheet PDD. A distinct drop in both PDD and SSC denotes the movement from the western half of Greenland to the eastern half, which is characterized by overall lower average PDD and SSC intensities and less distinct jumps in both datasets. Furthermore, PDD is a significant driver of SSC in high PDD areas but not in low PDD areas. Splitting the data in half with 24 high PDD gridcells and 23 low PDD gridcells, the high PDD portion show that PDD is strongly correlated with SSC ( $R = 0.61$ ,  $p = 0.002$ ), while the low PDD data are not correlated ( $R = -0.16$ ,  $p = 0.47$ ).

### 3.2.2 Interannual variability

Interannual variations between ice sheet PDD and plume SSC track each other for the northern regions (Northwest, West, East, and Northeast), but not for the southern regions (Southwest and Southeastw) or for Greenland as a whole (Fig. 6). Nine gridcells that do show strong relationships are concentrated in the northern parts of Greenland; and for the most part the southern regions show slight anti-correlation interannually (Fig. 7d, Table 1). Gridcell 13 in the Southwest Region containing solely Kangerlussuaq Fjord where a strong interannual correlation was previously found between ice sheet melt area and sediment plume area (Chu et al., 2009) also displays this slight anti-correlation, similar to most gridcells in the Southwest Region. This region highly influences the overall interannual relationship averaged Greenland owing to high values of both PDD and SSC.

## Hydrologic controls on coastal suspended sediment plumes

V. W. Chu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



### 3.2.3 Seasonal variability

Average seasonal climatologies of ice sheet PDD and downstream fjord SSC indicate coinciding seasonal cycles, with sediment plume onset coincident or commencing after surface melt onset (Fig. 8). Seasonal climatologies of PDD and SSC are produced by averaging across the same day-of-year (d.o.y.) over 2000–2009. Cross-correlation between these two seasonal cycles is highest for the West and Southwest Regions ( $R = 0.81$ ,  $p < 0.001$ ,  $R = 0.86$ ,  $p < 0.001$ , respectively, Figs. 7e, 8), especially Grid-cell 14 in the Southwest ( $R = 0.92$ ,  $p < 0.001$ , Table 1). Differences between regions include PDD and SSC intensity and seasonal length of both datasets, with the Southwest Region (in particular Gridcells 17–19) exhibiting highest SSCs and highest PDDs, and longer periods of activity and persistent sediment suspension towards the end of the melt season. Plume decline occurs during PDD decline for all regions, but the Southwest indicates high-SSCs persisting during the decline in melt in contrast to Chu et al. (2009), that found apparent sediment exhaustion with sediment plume areas decreasing prior to declines in melt area. Plume onset generally follows melt onset broadly in the western regions (Northwest, West, and Southwest), but the eastern regions (Southeast, East, and Northeast) show delayed plume onset as well as early melt onset. The Southeast Region is unique in that SSC is detected throughout the beginning of the melt season at low levels ( $\sim 1.62\text{--}2.76 \text{ mg l}^{-1}$ ) indicating no arrival of sediment-rich meltwaters until June. The East Region has very low SSC ( $\sim 0.83\text{--}0.97 \text{ mg l}^{-1}$ ) through the melt onset period, but much data are lost to the presence of sea ice until the end of May; the Northeast Region farther north shows even greater data loss with persistent sea ice until June.

### 3.2.4 High frequency variability

High frequency time series from 2000–2009 of ice sheet PDD and plume SSC show high short term variability (Fig. 9). SSC values from 2010 are also shown (recall PMM5 PDD data were not available after 2009). The eastern regions (Southeast, East, and

## Hydrologic controls on coastal suspended sediment plumes

V. W. Chu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Northeast), all with much lower intensities of PDD, show a more variable melt season onset than the western regions. Overall, near-daily data generated by spatial and temporal aggregation do not correlate well and the temporal relationship between PDD and SSC matches better in average seasonal climatologies.

### 3.3 Land-terminating vs. marine-terminating glacier outlets

Plume characteristics vary with outlet glacier environment, here categorized into two broad types: land-terminating and marine-terminating. The spatial density of land-terminating outlet glaciers ( $N_L$ ) is positively correlated with averaged 2000–2009 SSC around the ice sheet ( $R = 0.32$ ,  $p = 0.03$ ), while the number of marine-terminating outlet glaciers ( $N_M$ ) shows a negative correlation ( $R = -0.42$ ,  $p = 0.003$ ) (Fig. 7a). Normalizing the number of land-terminating and marine-terminating outlets by total number of outlets for each gridcell shows that the  $F_L$  is more correlated with SSC than  $N_L$  ( $R = 0.58$ ,  $p < 0.001$ ). Outlet types for the most part cluster in regions, with the Southwest showing highest  $F_L$  (0.81) and the Southeast showing highest  $F_M$  (0.91). These overall trends reverse in northern regions, with relatively high  $F_L$  (0.59) in the Northeast and high  $F_M$  (0.60) in the Northwest. The erosional susceptibility of the underlying bedrock, characterized by the geology of the ice-free area, shows mostly latitudinal variation dominated by Archaean and early Proterozoic crystalline basement rocks (Geological Survey of Denmark and Greenland (GEUS), 2003). Softer rock types might be more easily eroded and produce more sediment, yet areas with highest SSCs are made up of hard gneisses that are less likely to erode.

### 3.4 Other factors

$F_{SSC}$ , a measure of how long high SSC plumes persist, is closely related to mean SSC around Greenland ( $R = 0.72$ ,  $p < 0.001$ ) and is highest in the southwest with highs also in the northwest and northeast (Fig. 7b). This high spatial correlation between  $F_{SSC}$  and mean SSC reflects that  $F_{SSC}$  is calculated using high-SSC values, but still

## Hydrologic controls on coastal suspended sediment plumes

V. W. Chu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



highlights areas with a persistence of high SSC days. The Southwest Region, again containing Gridcell 13 and the Kangerlussuaq Fjord, shows that protected fjords and many land-terminating outlets in the southwest allow sediment to persist longer toward the end of the season.

DD<sub>100 km</sub> has a positive spatial correlation ( $R = 0.42$ ,  $p = 0.003$ ) with SSC, highlighting areas in the southeast that often lose data due to iceberg calving yet still show low mean SSC (Fig. 7c). While DD<sub>100 km</sub> is a measure of data retrieval, it does represent some physical processes that prevent suspended sediment detection and partially relates to factors such as predominance of marine-terminating outlet glaciers that heavily calve icebergs. This relationship represents another environmental factor affecting SSC detection and intensity; data density reflects both the outlet glacier environment and the climatic regime of the region.

#### 4 Discussion and conclusion

MODIS-derived plume suspended sediment concentration (SSC) successfully maps average plume distribution around Greenland, which is controlled by both ice sheet hydrology and outlet glacier type. Spatial variations in 10-year average positive degree days (PDDs), a proxy for meltwater generation on the ice sheet surface, is the most significant driver of spatial variations in 10-year mean SSC, confirming that higher ice sheet melting produces higher SSC in surrounding coastal waters. Furthermore, the corresponding seasonal development of PDD and SSC track each other rather well, with broadly coincident timings of melt onset and plume detection (Fig. 8). However, the Southeast and East Regions show delayed plume onset, lagging melt onset by ~3–4 weeks (a lag is also seen in the Northeast Region, but is due to delayed open water detection from sea ice), whereas the western regions (Northwest, West, and Southwest) do not. In the Southeast and East Regions, despite delayed plume onset, plume growth coincides with the rising limb of PDD, but not in the Northeast. The Southwest Region reveals highly correlated seasonal climatologies of SSC and PDD (as averaged

## Hydrologic controls on coastal suspended sediment plumes

V. W. Chu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



over 2000–2009, Fig. 8), but also a unique lingering of high SSC persisting ~1 month during the waning of the melt season. This is notably different from the results of Chu et al. (2009), which suggest sediment exhaustion in Kangerlussuaq Fjord. This difference may be due to use of aggregated data, but is more likely due the use of SSC rather than plume area. For example, in an elongate, protected fjord environment such as Kangerlussuaq Fjord, sediment may remain suspended but plume area may contract, possibly owing to circulation. This suggests that while ice sheet meltwater runoff is a dominant control of regional plume SSC development, fjord geometry in addition to outlet glacier types are also a factor.

Buoyant plumes most readily detected downstream of rivers draining land-terminating glaciers, owing to high SSC and minimal obstruction by calving ice. Marine-terminating glaciers, in contrast, produce lower SSC (2–100 mg l<sup>-1</sup> range) and are often obstructed by calf and/or sea ice. Greenland's proglacial rivers, like other proglacial systems, are characterized by extremely high suspended sediment loads (e.g., 5–22 000 mg l<sup>-1</sup>, Hasholt, 1996). Therefore, the spatial distribution of 10-year mean SSC reveals highest concentrations (~200–500 mg l<sup>-1</sup>) in the Southwest Region, where  $F_L$  (fraction of land-terminating glacier outlets) is highest. This differs from other studies that project highest total sediment export in areas with major calving glaciers such as the northeast and southeast (Hasholt, 1996). One reason for this contrast is that the present study identifies freshwater signals represented by fine sediments suspended in ocean surface waters, rather than total depth-integrated sediment export as measured in terrestrial rivers. Also, while other studies present SSC from rivers draining the ice sheet, here SSC is measured from fjord plumes, showing lower average values due to plume spreading and mixing with marine waters. Thus, the MODIS-derived SSCs presented here are useful for detecting fine sediments associated with freshwater release, but not total sediment export, are sensitive to outlet glacier type, and are lower than corresponding SSC values from terrestrial samples of meltwater runoff from the ice sheet.

## Hydrologic controls on coastal suspended sediment plumes

V. W. Chu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



While Chu et al. (2009) found a strong interannual relationship between ice sheet surface melt extent and plume area, the present study finds no corresponding coherence between ice sheet PDD and plume SSC. Indeed, Gridcell 13, representing the Kangerlussuaq Fjord studied by Chu et al. (2009) and McGrath et al. (2010), has a fairly anti-correlated interannual relationship between the two variables (Fig. 6b). This contrast stems solely from the use of SSCs rather than plume area, as the two melt datasets are highly correlated ( $R = 0.96$ ,  $p < 0.001$ ). One reason for this may be the extrapolation of SSCs using a non-linear empirical model, which also does not encompass the full range of sediment-rich water reflectances. Future studies should provide more field samples from a wider array of plume states, particularly at the high-SSC range, to further refine the model. However, while plume dimensions are good indices of ice sheet hydrologic variations, their use is limited to unique environments like Kangerlussuaq Fjord, that develop large plumes at river mouths with consistent absence of calf and/or sea ice. The present approach, using SSC instead of plume dimensions, allows study of plumes in other coastal environments including marine-terminating outlet glaciers where calving ice confounds clear measurement of plume area or length.

Heavily calving glaciers (e.g., Helheim, Kangerdlugssuaq, Jakobshavn) and northern fjords perennially blocked by sea ice are a challenge to open-water detection. While sediment from marine-terminating glacier bottom melting should rise to fjord surface waters owing to high buoyance of the meltwater plume (Mugford and Dowdeswell, 2011), heavily calving glaciers confound remote sensing of open water reflectance, particularly with 500 m MODIS data. Furthermore, iceberg melting is a dominant mechanism of sediment dispersal from calving marine-terminating glaciers (Andrews et al., 1994), and can complicate plume SSC retrieval because those sediments do not originate from terrestrial runoff sources. However, this sediment melt-out process typically occurs beneath the surface (Azetsu-Scott and Syvitski, 1999), unobservable with remote sensing. Therefore, in fjords with moderate iceberg interference, any detection of a surface sediment plume likely reflects input of ice sheet meltwater runoff rather

## Hydrologic controls on coastal suspended sediment plumes

V. W. Chu et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

than sediment from icebergs. While the northern part of Greenland is of interest due to extensive bottom melting (rather than iceberg calving as the primary ablation mechanism, Reeh et al., 1999), data scarcity from persistent sea ice precludes adequate SSC recovery from MODIS reflectance products. Therefore, higher spatial resolution sensors are required to study plume characteristics in areas of pervasive iceberg and sea ice cover.

At high temporal frequencies (near-daily, Fig. 9) ice sheet PDD and plume SSC are generally uncoupled, suggesting that spatio-temporal aggregation is not effective for resolving the well-known temporal resolution limitations of MODIS in narrow fjord environments (Chu et al., 2009; McGrath et al., 2009). This lack of correlation at near-daily time scales is perhaps unsurprising, owing to the very large 100 km aggregations required to produce near-daily time series. Also, other physical factors besides terrestrial runoff influence plume dynamics, including tides and wind (e.g., Dowdeswell and Cro-mack, 1991; Whitney and Garvine, 2005; Halverson and Pawlowicz, 2008). Detecting short-term (days to ~1 week) variations in terrestrial runoff from sediment plumes remains challenging from a satellite remote sensing approach.

Regardless of these limitations, this study provides a first synoptic assessment of remotely-sensed buoyant coastal sediment plumes around Greenland, and links them to outflows of ice sheet runoff from land- and marine-terminating outlet glaciers. Meltwater production on the ice sheet surface, as approximated by PDD, is the most significant driver of spatial variations in suspended sediment concentrations of coastal ocean waters. On average, land-terminating outlet glaciers produce higher plume SSCs than marine-terminating outlet glaciers, although MODIS retrievals from the latter are often obstructed by calf ice. Despite known complexities in plume formation and development processes, remotely sensed sediment plumes appear to supply viable evidence of meltwater release. As such, their detection and monitoring from space represents one of the few ways to observe hydrologic release of meltwater from the Greenland ice sheet to the global ocean over broad spatial and temporal scales.

## Hydrologic controls on coastal suspended sediment plumes

V. W. Chu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



*Acknowledgements.* This work was supported by the NASA Cryospheric Sciences Program grant NNG05GN89G and the NASA Graduate Student Researchers Program fellowship NNX09AM25H, and the MODIS imagery was also provided by NASA. We thank R. Carlos and S. Jimenez for help in MODIS downloading, and K. Savage for help in MODIS processing and classification validation.

## References

- Abdalati, W. and Steffen, K.: Snowmelt on the Greenland ice sheet as derived from passive microwave satellite data, *J. Climate*, 10, 165–175, 1997.
- Allison, I., Alley, R. B., Fricker, H. A., Thomas, R. H., and Warner, R. C.: Ice sheet mass balance and sea level, *Antarctic Science*, 21, 413–426, doi:10.1017/s0954102009990137, 2009.
- Amundson, J. M., Truffer, M., Luthi, M. P., Fahnstock, M., West, M., and Motyka, R. J.: Glacier, fjord, and seismic response to recent large calving events, Jakobshavn Isbrae, Greenland, *Geophys. Res. Lett.*, 35, L22501, doi:10.1029/2008gl035281, 2008.
- Andersen, M. L., Larsen, T. B., Nettles, M., Elosegui, P., van As, D., Hamilton, G. S., Stearns, L. A., Davis, J. L., Ahlstrom, A. P., de Juan, J., Ekstrom, G., Stenseng, L., Khan, S. A., Forberg, R., and Dahl-Jensen, D.: Spatial and temporal melt variability at Helheim Glacier, East Greenland, and its effect on ice dynamics, *J. Geophys. Res.-Earth Surface*, 115, F04041, doi:10.1029/2010jf001760, 2010.
- Andrews, J. T., Milliman, J. D., Jennings, A. E., Rynes, N., and Dwyer, J.: Sediment thicknesses and Holocene glacial marine sedimentation-rates in 3 East Greenland fjords (ca. 68-degrees N), *J. Geology*, 102, 669–683, 1994.
- Bartholomew, I., Nienow, P., Mair, D., Hubbard, A., King, M. A., and Sole, A.: Seasonal evolution of subglacial drainage and acceleration in a Greenland outlet glacier, *Nature Geosci.*, 3, 408–411, doi:10.1038/ngeo863, 2010.
- Bhattacharya, I., Jezek, K. C., Wang, L., and Liu, H. X.: Surface melt area variability of the Greenland ice sheet: 1979–2008, *Geophys. Res. Lett.*, 36, L20502, doi:10.1029/2009gl039798, 2009.
- Bøggild, C. E., Knudby, C. J., Knudsen, M. B., and Starzer, W.: Snowmelt and runoff modelling of an Arctic hydrological basin in west Greenland, *Hydrol. Processes*, 13, 1989–2002, 1999.
- Bowers, D. G., Boudjelas, S., and Marker, G. E. L.: The distribution of fine suspended sedi-

## Hydrologic controls on coastal suspended sediment plumes

V. W. Chu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Hydrologic controls on coastal suspended sediment plumes

V. W. Chu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- ments in the surface waters of the Irish Sea and its relation to tidal stirring, *Int. J. Remote Sens.*, 19, 2789–2805, 1998.
- Box, J. E. and Ski, K.: Remote sounding of Greenland supraglacial melt lakes: implications for subglacial hydraulics, *J. Glaciol.*, 53, 257–265, 2007.
- 5 Box, J. E., Bromwich, D. H., Veenhuis, B. A., Bai, L. S., Stroeve, J. C., Rogers, J. C., Steffen, K., Haran, T., and Wang, S. H.: Greenland ice sheet surface mass balance variability (1988–2004) from calibrated polar MM5 output, *J. Climate*, 19, 2783–2800, 2006.
- Box, J. E., Yang, L., Bromwich, D. H., and Bai, L. S.: Greenland Ice Sheet Surface Air Temperature Variability: 1840–2007, *J. Climate*, 22, 4029–4049, doi:10.1175/2009jcli2816.1, 2009.
- 10 Burgess, E. W., Forster, R. R., Box, J. E., Mosley-Thompson, E., Bromwich, D. H., Bales, R. C., and Smith, L. C.: A spatially calibrated model of annual accumulation rate on the Greenland Ice Sheet (1958–2007), *J. Geophys. Res.-Earth Surface*, 115, F02004, doi:10.1029/2009jf001293, 2010.
- Busskamp, R. and Hasholt, B.: Coarse bed load transport in a glacial valley, Sermilik, South East Greenland, *Zeitschrift Fur Geomorphologie*, 40, 349–358, 1996.
- 15 Castaing, P. and Allen, G. P.: Mechanisms controlling seaward escape of suspended sediment from the Gironde – a macrotidal estuary in France, *Marine Geology*, 40, 101–118, 1981.
- Catania, G. A. and Neumann, T. A.: Persistent englacial drainage features in the Greenland Ice Sheet, *Geophys. Res. Lett.*, 37, L02501, doi:10.1029/2009gl041108, 2010.
- 20 Chen, J. L., Wilson, C. R., and Tapley, B. D.: Satellite gravity measurements confirm accelerated melting of Greenland ice sheet, *Science*, 313, 1958–1960, doi:10.1126/science.1129007, 2006.
- Chu, V. W., Smith, L. C., Rennermalm, A. K., Forster, R. R., Box, J. E., and Reeh, N.: Sediment plume response to surface melting and supraglacial lake drainages on the Greenland ice sheet, *J. Glaciol.*, 55, 1072–1082, 2009.
- 25 Curran, P. J. and Novo, E. M. M.: The relationship between suspended sediment concentration and remotely sensed spectral radiance – A review, *J. Coastal Res.*, 4, 351–368, 1988.
- Das, S. B., Joughin, I., Behn, M. D., Howat, I. M., King, M. A., Lizarralde, D., and Bhatia, M. P.: Fracture propagation to the base of the Greenland Ice Sheet during supraglacial lake drainage, *Science*, 320, 778–781, 2008.
- 30 Dowdeswell, J. A. and Cromack, M.: Behavior of a glacier-derived suspended sediment plume in a small Arctic inlet, *J. Geology*, 99, 111–123, 1991.
- Doxaran, D., Froidefond, J. M., Lavender, S., and Castaing, P.: Spectral signature of highly

## Hydrologic controls on coastal suspended sediment plumes

V. W. Chu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

turbid waters – Application with SPOT data to quantify suspended particulate matter concentrations, *Remote Sens. Environ.*, 81, 149–161, 2002.

Ettema, J., van den Broeke, M. R., van Meijgaard, E., van de Berg, W. J., Bamber, J. L., Box, J. E., and Bales, R. C.: Higher surface mass balance of the Greenland ice sheet revealed by high-resolution climate modeling, *Geophys. Res. Lett.*, 36, L12501, doi:10.1029/2009gl038110, 2009.

Fettweis, X.: Reconstruction of the 1979–2006 Greenland ice sheet surface mass balance using the regional climate model MAR, *The Cryosphere*, 1, 21–40, doi:10.5194/tc-1-21-2007, 2007.

10 Fettweis, X., Hanna, E., Gallée, H., Huybrechts, P., and Ericum, M.: Estimation of the Greenland ice sheet surface mass balance for the 20th and 21st centuries, *The Cryosphere*, 2, 117–129, doi:10.5194/tc-2-117-2008, 2008.

Gurnell, A., Hannah, D., and Lawler, D.: Suspended sediment yield from glacier basins, in: *Erosion and Sediment Yield: Global and Regional Perspectives*, edited by: Walling, D. E. and Webb, B. W., IAHS, 97–104, 1996.

Hall, D. K., Williams, R. S., Luthcke, S. B., and Digirolamo, N. E.: Greenland ice sheet surface temperature, melt and mass loss: 2000–06, *J. Glaciol.*, 53, 81–93, 2008.

Hallet, B., Hunter, L., and Bogen, J.: Rates of erosion and sediment evacuation by glaciers: A review of field data and their implications, *Global Planet. Change*, 12, 213–235, 1996.

20 Halverson, M. J. and Pawlowicz, R.: Estuarine forcing of a river plume by river flow and tides, *J. Geophys. Res.-Oceans*, 113, C09033, doi:10.1029/2008jc004844, 2008.

Hanna, E., Huybrechts, P., Steffen, K., Cappelen, J., Huff, R., Shuman, C., Irvine-Fynn, T., Wise, S., and Griffiths, M.: Increased runoff from melt from the Greenland Ice Sheet: A response to global warming, *J. Climate*, 21, 331–341, doi:10.1175/2007jcli1964.1, 2008.

25 Hasholt, B.: Sediment transport in Greenland, *Erosion and Sediment Yield: Global and Regional Perspectives (Proceedings of the Exeter Symposium)*, July 1996, 105–114, 1996.

Hasholt, B. and Mernild, S. H.: Hydrology, sediment transport and water resources of Ammassalik Island, SE Greenland, *Geografisk Tidsskrift-Danish J. Geography*, 108, 73–95, 2008.

Hock, R.: Temperature index melt modelling in mountain areas, *J. Hydrol.*, 282, 104–115, doi:10.1016/s0022-1694(03)00257-9, 2003.

30 Holland, D. M., Thomas, R. H., De Young, B., Ribergaard, M. H., and Lyberth, B.: Acceleration of Jakobshavn Isbrae triggered by warm subsurface ocean waters, *Nature Geosci.*, 1, 659–664, doi:10.1038/ngeo316, 2008.

## Hydrologic controls on coastal suspended sediment plumes

V. W. Chu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Howat, I. M., Joughin, I., and Scambos, T. A.: Rapid changes in ice discharge from Greenland outlet glaciers, *Science*, 315, 1559–1561, 2007.
- Howat, I. M., Joughin, I., Fahnestock, M., Smith, B. E., and Scambos, T. A.: Synchronous retreat and acceleration of southeast Greenland outlet glaciers 2000–06: ice dynamics and coupling to climate, *J. Glaciol.*, 54, 646–660, 2008.
- Hu, C. M., Chen, Z. Q., Clayton, T. D., Swarzenski, P., Brock, J. C., and Muller-Karger, F. E.: Assessment of estuarine water-quality indicators using MODIS medium-resolution bands: Initial results from Tampa Bay, FL, *Remote Sensing of Environment*, 93, 423–441, doi:10.1016/j.rse.2004.08.007, 2004.
- Joughin, I., Tulaczyk, S., Fahnestock, M., and Kwok, R.: A mini-surge on the Ryder Glacier, Greenland, observed by satellite radar interferometry, *Science*, 274, 228–230, 1996.
- Joughin, I., Das, S. B., King, M. A., Smith, B. E., Howat, I. M., and Moon, T.: Seasonal speedup along the western flank of the Greenland Ice Sheet, *Science*, 320, 781–783, 2008a.
- Joughin, I., Howat, I., Alley, R. B., Ekstrom, G., Fahnestock, M., Moon, T., Nettles, M., Truffer, M., and Tsai, V. C.: Ice-front variation and tidewater behavior on Helheim and Kangerdlugssuaq Glaciers, Greenland, *J. Geophys. Res.-Earth Surface*, 113, F01004, doi:10.1029/2007JF000837, 2008b.
- Joughin, I., Smith, B. E., Howat, I. M., Scambos, T., and Moon, T.: Greenland flow variability from ice-sheet-wide velocity mapping, *J. Glaciol.*, 56, 415–430, 2010.
- Khan, S. A., Wahr, J., Bevis, M., Velicogna, I., and Kendrick, E.: Spread of ice mass loss into northwest Greenland observed by GRACE and GPS, *Geophys. Res. Lett.*, 37, L06501, doi:10.1029/2010gl042460, 2010.
- Knudsen, N. T., Yde, J. C., and Gasser, G.: Suspended sediment transport in glacial meltwater during the initial quiescent phase after a major surge event at Kuannersuit Glacier, Greenland, *Geografisk Tidsskrift-Danish J. Geography*, 107, 1–7, 2007.
- Krabill, W., Hanna, E., Huybrechts, P., Abdalati, W., Cappelen, J., Csatho, B., Frederick, E., Manizade, S., Martin, C., Sonntag, J., Swift, R., Thomas, R., and Yungel, J.: Greenland Ice Sheet: Increased coastal thinning, *Geophys. Res. Lett.*, 31, L24402, doi:10.1029/2004GL021533, 2004.
- Lewis, S. M. and Smith, L. C.: Hydrologic drainage of the Greenland Ice Sheet, *Hydrol. Processes*, 23, 2004–2011, doi:10.1002/hyp.7343, 2009.
- Lihan, T., Saitoh, S. I., Iida, T., Hirawake, T., and Iida, K.: Satellite-measured temporal and spatial variability of the Tokachi River plume, *Estuarine Coastal and Shelf Science*, 78, 237–

---

**Hydrologic controls  
on coastal  
suspended sediment  
plumes**

---

V. W. Chu et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

249, doi:10.1016/j.ecss.2007.12.001, 2008.

Lund-Hansen, L. C., Andersen, T. J., Nielsen, M. H., and Pejrup, M.: Suspended Matter, Chl-a, CDOM, Grain Sizes, and Optical Properties in the Arctic Fjord-Type Estuary, Kangerlussuaq, West Greenland During Summer, *Estuaries and Coasts*, 33, 1442–1451, doi:10.1007/s12237-010-9300-7, 2010.

Luthcke, S. B., Zwally, H. J., Abdalati, W., Rowlands, D. D., Ray, R. D., Nerem, R. S., Lemoine, F. G., McCarthy, J. J., and Chinn, D. S.: Recent Greenland ice mass loss by drainage system from satellite gravity observations, *Science*, 314, 1286–1289, 2006.

McGrath, D., Steffen, K., Overeem, I., Mernild, S. H., Hasholt, B., and van den Broeke, M.: Sediment plumes as a proxy for local ice-sheet runoff in Kangerlussuaq Fjord, West Greenland, *J. Glaciol.*, 56, 813–821, 2010.

Mernild, S. H. and Hasholt, B.: Observed runoff, jokulhlaups and suspended sediment load from the Greenland ice sheet at Kangerlussuaq, West Greenland, 2007 and 2008, *J. Glaciol.*, 55, 855–858, 2009.

Mernild, S. H., Howat, I. M., Ahn, Y., Liston, G. E., Steffen, K., Jakobsen, B. H., Hasholt, B., Fog, B., and van As, D.: Freshwater flux to Sermilik Fjord, SE Greenland, *The Cryosphere*, 4, 453–465, doi:10.5194/tc-4-453-2010, 2010a.

Mernild, S. H., Liston, G. E., Steffen, K., van den Broeke, M., and Hasholt, B.: Runoff and mass-balance simulations from the Greenland Ice Sheet at Kangerlussuaq (Søndre Strømfjord) in a 30-year perspective, 1979–2008, *The Cryosphere*, 4, 231–242, doi:10.5194/tc-4-231-2010, 2010b.

Mernild, S. H., Liston, G. E., Hiemstra, C. A., Christensen, J. H., Stendel, M., and Hasholt, B.: Surface Mass Balance and Runoff Modeling Using HIRHAM4 RCM at Kangerlussuaq (Søndre Strømfjord), West Greenland, 1950–2080, *J. Climate*, 24, 609–623, doi:10.1175/2010jcli3560.1, 2011.

Miller, R. L. and McKee, B. A.: Using MODIS Terra 250 m imagery to map concentrations of total suspended matter in coastal waters, *Remote Sens. Environ.*, 93, 259–266, 2004.

Mote, T. L.: Greenland surface melt trends 1973–2007: Evidence of a large increase in 2007, *Geophys. Res. Lett.*, 34, L22507, doi:10.1029/2007GL031976, 2007.

Motyka, R. J., Truffer, M., Fahnestock, M., Mortensen, J., Rysgaard, S., and Howat, I.: Submarine melting of the 1985 Jakobshavn Isbrae floating tongue and the triggering of the current retreat, *J. Geophys. Res.-Earth Surface*, 116, F01007, doi:10.1029/2009jf001632, 2011.

Mugford, R. I. and Dowdeswell, J. A.: Modeling iceberg-rafted sedimentation in

## Hydrologic controls on coastal suspended sediment plumes

V. W. Chu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

high-latitude fjord environments, *J. Geophys. Res.-Earth Surface*, 115, F03024, doi:10.1029/2009jf001564, 2010.

Mulder, T. and Syvitski, J. P. M.: Turbidity currents generated at river mouths during exceptional discharges to the world oceans, *J. Geology*, 103, 285–299, 1995.

5 Ohmura, A.: Physical basis for the temperature-based melt-index method, *J. Appl. Meteorol.*, 40, 753–761, 2001.

Ohmura, A. and Reeh, N.: New precipitation and accumulation maps for Greenland, *J. Glaciol.*, 37, 140–148, 1991.

10 Palmer, S., Shepherd, A., Nienow, P., and Joughin, I.: Seasonal speedup of the Greenland Ice Sheet linked to routing of surface water, *Earth Planet. Sci. Lett.*, 302, 423–428, doi:10.1016/j.epsl.2010.12.037, 2011.

Pfeffer, W. T., Harper, J. T., and O’Neel, S.: Kinematic constraints on glacier contributions to 21st-century sea-level rise, *Science*, 321, 1340–1343, doi:10.1126/science.1159099, 2008.

15 Pritchard, H. D., Arthern, R. J., Vaughan, D. G., and Edwards, L. A.: Extensive dynamic thinning on the margins of the Greenland and Antarctic ice sheets, *Nature*, 461, 971–975, doi:10.1038/nature08471, 2009.

Rasch, M., Elberling, B., Jakobsen, B. H., and Hasholt, B.: High-resolution measurements of water discharge, sediment and solute transport in the river Zackenbergelven, Northeast Greenland, *Arctic Antarctic and Alpine Research*, 32, 336–345, 2000.

20 Reeh, N., Mayer, C., Miller, H., Thomsen, H. H., and Weidick, A.: Present and past climate control on fjord glaciations in Greenland: Implications for IRD-deposition in the sea, *Geophys. Res. Lett.*, 26, 1039–1042, 1999.

Rignot, E. and Kanagaratnam, P.: Changes in the velocity structure of the Greenland ice sheet, *Science*, 311, 986–990, 2006.

25 Rignot, E. and Steffen, K.: Channelized bottom melting and stability of floating ice shelves, *Geophys. Res. Lett.*, 35, L02503, doi:10.1029/2007gl031765, 2008.

Rignot, E., Braaten, D., Gogineni, S. P., Krabill, W. B., and McConnell, J. R.: Rapid ice discharge from southeast Greenland glaciers, *Geophys. Res. Lett.*, 31, L10401, doi:10.1029/2004GL019474, 2004.

30 Rignot, E., Box, J. E., Burgess, E., and Hanna, E.: Mass balance of the Greenland ice sheet from 1958 to 2007, *Geophys. Res. Lett.*, 35, L20502, doi:10.1029/2008gl035417, 2008.

Robinson, A., Calov, R., and Ganopolski, A.: An efficient regional energy-moisture balance model for simulation of the Greenland Ice Sheet response to climate change, *The*

---

## Hydrologic controls on coastal suspended sediment plumes

V. W. Chu et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Cryosphere, 4, 129–144, doi:10.5194/tc-4-129-2010, 2010.

Russell, A. J.: Controls on the sedimentology of an ice-contact jokulhlaup-dominated delta, Kangerlussuaq, west Greenland, *Sedimentary Geology*, 193, 131–148, doi:10.1016/j.sedgeo.2006.01.007, 2007.

5 Schoof, C.: Ice-sheet acceleration driven by melt supply variability, *Nature*, 468, 803–806, doi:10.1038/nature09618, 2010.

Shepherd, A., Hubbard, A., Nienow, P., King, M., McMillan, M., and Joughin, I.: Greenland ice sheet motion coupled with daily melting in late summer, *Geophys. Res. Lett.*, 36, L01501, doi:10.1029/2008gl035758, 2009.

10 Smith, L. C., Sheng, Y. W., Forster, R. R., Steffen, K., Frey, K. E., and Alsdorf, D. E.: Melting of small Arctic ice caps observed from ERS scatterometer time series, *Geophys. Res. Lett.*, 30, CRY2, doi:10.1029/2003gl017641, 2003.

Sole, A., Payne, T., Bamber, J., Nienow, P., and Krabill, W.: Testing hypotheses of the cause of peripheral thinning of the Greenland Ice Sheet: is land-terminating ice thinning at anomalously high rates?, *The Cryosphere*, 2, 205–218, doi:10.5194/tc-2-205-2008, 2008.

Stott, T. A. and Grove, J. R.: Short-term discharge and suspended sediment fluctuations in the proglacial Skeldal River, north-east Greenland, *Hydrol. Processes*, 15, 407–423, 2001.

15 Straneo, F., Hamilton, G. S., Sutherland, D. A., Stearns, L. A., Davidson, F., Hammill, M. O., Stenson, G. B., and Rosing-Asvid, A.: Rapid circulation of warm subtropical waters in a major glacial fjord in East Greenland, *Nature Geoscience*, 3, 182–186, doi:10.1038/ngeo764, 2010.

Stumpf, R. P., Gelfenbaum, G., and Pennock, J. R.: Wind and tidal forcing of a buoyant plume, Mobile Bay, Alabama, *Cont. Shelf Res.*, 13, 1281–1301, 1993.

25 Sundal, A. V., Shepherd, A., Nienow, P., Hanna, E., Palmer, S., and Huybrechts, P.: Evolution of supra-glacial lakes across the Greenland Ice Sheet, *Remote Sens. Environ.*, 113, 2164–2171, doi:10.1016/j.rse.2009.05.018, 2009.

Syviatski, J. P. M., Asprey, K. W., Clattenburg, D. A., and Hodge, G. D.: The prodelta environment of a fjord – suspended particle dynamics, *Sedimentology*, 32, 83–107, 1985.

30 Syviatski, J. P. M., Andrews, J. T., and Dowdeswell, J. A.: Sediment deposition in an iceberg-dominated glacial marine environment, East Greenland: Basin fill implications, *Global Planet. Change*, 12, 251–270, 1996.

Tedesco, M.: Snowmelt detection over the Greenland ice sheet from SSM/I brightness temperature daily variations, *Geophys. Res. Lett.*, 34, L02504, doi:10.1029/2006gl028466, 2007.

Tedesco, M., Serreze, M., and Fettweis, X.: Diagnosing the extreme surface melt event over southwestern Greenland in 2007, *The Cryosphere*, 2, 159–166, doi:10.5194/tc-2-159-2008, 2008.

Thomas, A. C. and Weatherbee, R. A.: Satellite-measured temporal variability of the Columbia River plume, *Remote Sens. Environ.*, 100, 167–178, doi:10.1016/j.rse.2005.10.018, 2006.

Thomas, R. H., Abdalati, W., Frederick, E., Krabill, W. B., Manizade, S., and Steffen, K.: Investigation of surface melting and dynamic thinning on Jakobshavn Isbrae, Greenland, *J. Glaciol.*, 49, 231–239, 2003.

Thomas, R., Frederick, E., Krabill, W., Manizade, S., and Martin, C.: Recent changes on Greenland outlet glaciers, *J. Glaciol.*, 55, 147–162, 2009.

van den Broeke, M., Smeets, P., Ettema, J., van der Veen, C., van de Wal, R., and Oerlemans, J.: Partitioning of melt energy and meltwater fluxes in the ablation zone of the west Greenland ice sheet, *The Cryosphere*, 2, 179–189, doi:10.5194/tc-2-179-2008, 2008.

van den Broeke, M., Bamber, J., Ettema, J., Rignot, E., Schrama, E., van de Berg, W. J., van Meijgaard, E., Velicogna, I., and Wouters, B.: Partitioning Recent Greenland Mass Loss, *Science*, 326, 984–986, doi:10.1126/science.1178176, 2009.

van de Wal, R. S. W., Boot, W., van den Broeke, M. R., Smeets, C., Reijmer, C. H., Donker, J. J. A., and Oerlemans, J.: Large and rapid melt-induced velocity changes in the ablation zone of the Greenland Ice Sheet, *Science*, 321, 111–113, doi:10.1126/science.1158540, 2008.

Velicogna, I.: Increasing rates of ice mass loss from the Greenland and Antarctic ice sheets revealed by GRACE, *Geophys. Res. Lett.*, 36, L19503, doi:10.1029/2009gl040222, 2009.

Velicogna, I., Wahr, J., Hanna, E., and Huybrechts, P.: Short term mass variability in Greenland, from GRACE, *Geophys. Res. Lett.*, 32, L05501, doi:10.1029/2004GL021948, 2005.

Vermote, E. F., El Saleous, N. Z., and Justice, C. O.: Atmospheric correction of MODIS data in the visible to middle infrared: first results, *Remote Sens. Environ.*, 83, 97–111, 2002.

Whitney, M. M. and Garvine, R. W.: Wind influence on a coastal buoyant outflow, *J. Geophys. Res.-Oceans*, 110, C03014, doi:10.1029/2003jc002261, 2005.

Wouters, B., Chambers, D., and Schrama, E. J. O.: GRACE observes small-scale mass loss in Greenland, *Geophys. Res. Lett.*, 35, L20501, doi:10.1029/2008gl034816, 2008.

Yin, J., Overpeck, J. T., Griffies, S. M., Hu, A., Russell, J. L., and Stouffer, R. J.: Different magnitudes of projected subsurface ocean warming around Greenland and Antarctica, *Nature Geoscience*, 4, 524, doi:10.1038/ngeo1189, 2011.

Zwally, H. J., Abdalati, W., Herring, T., Larson, K., Saba, J., and Steffen, K.: Surface melt-

## Hydrologic controls on coastal suspended sediment plumes

V. W. Chu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- induced acceleration of Greenland ice-sheet flow, *Science*, 297, 218–222, 2002.
- Zwally, H. J., Li, J., Brenner, A. C., Beckley, M., Cornejo, H. G., Dimarzio, J., Giovinetto, M. B., Neumann, T. A., Robbins, J., Saba, J. L., Yi, D. H., and Wang, W. L.: Greenland ice sheet mass balance: distribution of increased mass loss with climate warming; 2003–07 versus 1992–2002, *J. Glaciol.*, 57, 88–102, 2011.

5

TCO

5, 2365–2407, 2011

## Hydrologic controls on coastal suspended sediment plumes

V. W. Chu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Hydrologic controls on coastal suspended sediment plumes**

V. W. Chu et al.

**Table 1.** All parameters for each of the 47 gridcells, which are grouped and averaged into six regions. 10-year mean SSC and 10-year mean PDD are shown with standard deviations and ranges (minimum PDD is 0 °C for every gridcell), and region averages are shown with standard deviations of gridcell averages within the region. *B* is drainage basin size, *l* is ice edge length, *A*<sub>PDD</sub> is melt area within each drainage basin, *F*<sub>PDD</sub> is percent of drainage basin actively melting, *D*<sub>PDD</sub> is melt penetration distance, *N*<sub>M</sub> is number of marine-terminating glacier outlets, *F*<sub>M</sub> is the fraction of marine-terminating outlets, *N*<sub>L</sub> is number of land-terminating glacier outlets, *F*<sub>L</sub> is the fraction of land-terminating outlets, *DD*<sub>500m</sub> is average data density of open water from raw 500 m data, *DD*<sub>100km</sub> is data density of open water from aggregated 100 km gridcells, *F*<sub>SSC</sub> is sediment persistence (fraction of high-SSC days from total open water days), *R* (interannual) is the correlation between interannual variations of SSC and PDD over 2000–2009, and *R* (seasonal) is the correlation between 10-year mean seasonal cycles.

Gridcell	10-yr Mean SSC (mg l <sup>-1</sup> )	Max SSC (mg l <sup>-1</sup> )	Min SSC (mg l <sup>-1</sup> )	10-yr Mean PDD (°C)	Max PDD (°C)	<i>B</i> (km <sup>2</sup> )	<i>l</i> (km)	<i>A</i> <sub>PDD</sub> (km <sup>2</sup> )	<i>F</i> <sub>PDD</sub>	<i>D</i> <sub>PDD</sub> (km)	<i>N</i> <sub>L</sub>	<i>F</i> <sub>L</sub>	<i>N</i> <sub>M</sub>	<i>F</i> <sub>M</sub>	<i>DD</i> <sub>500m</sub> (%)	<i>DD</i> <sub>100km</sub> (%)	<i>F</i> <sub>SSC</sub> (%)	<i>R</i> (Interann.)	<i>R</i> (Seasonal)
1	40 ± 277	723	0.8	1.23 ± 3.78	14.13	2.847	141	830	0.29	5.9	3	0.33	6	0.67	22.2%	92.2%	0.14	0.61	0.48
2	162 ± 4	1,784	1.0	2.23 ± 0.03	27.69	6.587	212	1,720	0.26	8.1	10	0.71	4	0.29	14.4%	85.6%	0.16	0.26	0.75
3	13 ± 7.5	377	0.8	0.78 ± 1.20	7.41	711	73	310	0.44	4.2	5	0.50	5	0.50	20.4%	89.4%	0.07	0.02	0.63
4	4 ± 3.5	65	0.9	1.10 ± 1.31	12.38	6711	189	1041	0.16	5.5	1	0.07	14	0.93	8.4%	75.1%	0.09	-0.18	0.46
Northwest	55 ± 63	737	0.9	1.34 ± 0.54	15.40	4214	154	975	0.29	5.9	5	0.40	7	0.60	16.4%	85.6%	0.11	0.18	0.58
5	8 ± 27	226	0.7	5.09 ± 0.07	42.46	55652	207	4531	0.08	21.9	0	0.00	6	1.00	5.4%	57.7%	0.09	0.17	0.54
6	10 ± 16	112	1.0	3.81 ± 4.92	26.73	36968	212	3176	0.09	14.9	0	0.00	5	1.00	11.9%	67.3%	0.04	0.33	0.74
7	42 ± 95	686	0.9	5.29 ± 7.05	44.24	46506	245	8645	0.08	17.8	3	0.27	8	0.73	13.7%	81.6%	0.06	0.58	0.87
8	38 ± 58	677	0.8	7.86 ± 9.82	59.84	41373	121	4721	0.11	39.0	4	0.44	5	0.56	11.8%	84.3%	0.13	0.83	0.89
9	18 ± 26	215	1.0	1.50 ± 2.97	23.43	30978	148	1763	0.06	11.9	3	0.50	3	0.50	19.8%	73.7%	0.02	0.42	0.81
10	140 ± 290	1925	1.1	2.30 ± 3.61	29.15	10166	203	2089	0.21	10.3	0	0.00	4	1.00	19.3%	78.8%	0.04	0.83	0.55
11	109 ± 227	1553	0.9	7.74 ± 9.42	49.37	72190	170	5495	0.08	32.3	1	0.17	5	0.83	16.1%	91.4%	0.05	0.32	0.76
12	93 ± 220	1616	0.9	13.25 ± 15.49	88.72	99210	224	9863	0.10	44.1	3	0.43	4	0.57	16.5%	79.9%	0.10	0.46	0.08
West	57 ± 47	876	0.9	5.85 ± 3.82	45.74	49131	186	4410	0.10	24.0	2	0.23	5	0.77	14.3%	76.9%	0.07	0.49	0.65
13	112 ± 119	1149	0.8	23.12 ± 24.21	113.25	39964	258	10551	0.26	40.9	13	0.87	2	0.13	22.6%	94.4%	0.19	-0.09	0.62
14	555 ± 422	1875	1.7	30.79 ± 34.27	164.93	55891	146	16,277	0.29	111.9	8	1.00	0	0.00	25.9%	78.4%	0.44	-0.08	0.92
15	238 ± 209	1,568	1.1	10.76 ± 12.86	65.63	21152	111	6725	0.32	60.7	3	1.00	0	0.00	16.9%	89.5%	0.33	-0.09	0.65
16	207 ± 187	939	1.9	3.93 ± 4.38	20.56	1200	39	1200	1.00	30.9	2	1.00	0	0.00	23.9%	91.3%	0.33	-0.23	0.61
17	312 ± 405	1887	0.9	11.10 ± 12.83	82.69	33462	135	7803	0.23	58.0	5	0.71	2	0.29	17.3%	81.6%	0.20	0.19	0.81
18	552 ± 481	1887	1.0	9.64 ± 10.41	57.70	8603	135	4314	0.50	31.8	3	0.75	1	0.25	14.4%	84.9%	0.21	-0.25	0.82
19	397 ± 345	1772	0.8	13.59 ± 13.85	72.97	12796	159	4926	0.38	31.0	6	1.00	0	0.00	14.1%	91.3%	0.17	-0.01	0.35
20	129 ± 213	1820	0.8	16.55 ± 16.11	86.57	21386	245	6272	0.29	25.6	3	0.50	3	0.50	9.9%	87.1%	0.09	-0.21	0.81
21	61 ± 76	443	0.8	5.53 ± 4.86	23.93	3134	131	1384	0.44	10.6	7	0.88	1	0.13	15.9%	93.5%	0.13	-0.10	0.52
22	250 ± 351	1925	1.0	12.19 ± 13.09	74.05	15793	338	6219	0.39	18.4	12	0.57	9	0.43	15.0%	94.4%	0.12	-0.45	0.31
23	72 ± 115	623	0.7	0.96 ± 0.96	6.13	553	35	349	0.63	9.8	4	0.67	2	0.33	7.2%	92.9%	0.12	-0.26	0.85
Southwest	262 ± 168	1444	1.0	12.54 ± 8.16	69.85	19449	157	6002	0.43	39.0	6	0.81	2	0.19	16.7%	89.0%	0.21	-0.14	0.66
24	6 ± 13	133	1.1	0.30 ± 0.45	3.12	3052	74	176	0.06	2.4	1	0.09	10	0.91	5.6%	61.5%	0.01	0.12	0.42
25	4 ± 6	62	1.2	1.84 ± 3.10	27.41	5344	106	0.29	0.41	0.20	12	0.00	12	0.00	7.8%	98.9%	0.01	0.04	0.28
26	26 ± 62	704	0.8	1.05 ± 2.10	23.29	11994	227	891	0.07	3.9	3	0.18	14	0.82	5.6%	99.9%	0.01	-0.39	0.46
27	91 ± 184	1838	0.9	0.27 ± 0.60	5.88	4808	125	227	0.05	1.8	0	0.00	4	1.00	9.0%	68.5%	0.10	-0.32	0.32
28	5 ± 9	373	0.9	1.27 ± 2.35	14.78	11014	344	963	0.09	2.8	3	0.13	20	0.87	8.3%	89.5%	0.04	-0.23	0.10
29	3 ± 5	49	1.0	1.04 ± 1.83	11.73	13047	184	1026	0.08	5.6	0	0.00	15	1.00	5.0%	82.0%	0.01	-0.37	0.09
30	3 ± 3	22	0.9	2.14 ± 3.42	25.77	22121	318	2590	0.12	8.2	2	0.15	11	0.85	7.7%	78.4%	0.08	-0.01	0.64
31	2 ± 1	11	0.9	0.30 ± 0.42	2.41	753	110	219	0.29	2.0	1	0.25	3	0.75	10.3%	81.0%	0.02	0.00	0.63
32	44 ± 90	679	0.9	1.84 ± 2.85	17.58	8391	250	1542	0.18	6.2	1	0.05	19	0.95	7.1%	67.1%	0.12	0.20	0.40
33	64 ± 91	687	1.3	1.78 ± 2.97	25.66	13117	302	1472	0.11	4.9	3	0.30	7	0.82	7.2%	56.8%	0.16	0.05	0.30
34	49 ± 103	806	0.9	0.56 ± 1.12	8.53	6917	197	635	0.09	3.2	0	0.00	8	1.00	9.0%	50.9%	0.13	0.43	0.09

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Hydrologic controls on coastal suspended sediment plumes**

V. W. Chu et al.

**Table 1.** Continued.

Gridcell	10-yr Mean SSC (mg l <sup>-1</sup> )	Max SSC (mg l <sup>-1</sup> )	Min SSC (mg l <sup>-1</sup> )	10-yr Mean PDD (°C)	Max PDD (°C)	<i>B</i> (km <sup>2</sup> )	<i>l</i> (km)	<i>A</i> <sub>POD</sub> (km <sup>2</sup> )	<i>F</i> <sub>PDD</sub>	<i>D</i> <sub>PDD</sub> (km)	<i>N</i> <sub>L</sub>	<i>F</i> <sub>L</sub>	<i>N</i> <sub>M</sub>	<i>F</i> <sub>M</sub>	DD <sub>500m</sub> (%)	DD <sub>100km</sub> (%)	<i>F</i> <sub>SSC</sub> (%)	<i>R</i> (Interann.)	<i>R</i> (Seasonal)
Southeast	27 ± 29	489	1.0	1.13 ± 0.68	15.10	9142	202	994	0.12	4.9	1	0.10	11	0.90	7.5%	73.1%	0.06	-0.04	0.34
35	11 ± 17	108	1.0	1.51 ± 2.72	21.38	6235	126	1111	0.18	8.8	3	0.50	3	0.50	10.8%	56.6%	0.00	0.41	0.33
36	4 ± 3	22	0.7	0.83 ± 1.28	8.03	1464	62	363	0.25	5.9	1	0.25	3	0.75	10.2%	62.6%	0.00	-0.15	0.66
37	23 ± 31	211	0.8	0.67 ± 0.99	7.96	1884	117	318	0.17	2.7	2	0.33	4	0.67	13.4%	53.9%	0.07	0.62	0.54
38	26 ± 28	149	1.1	1.34 ± 2.88	20.75	29747	283	1462	0.05	5.2	9	0.64	5	0.36	12.2%	65.6%	0.04	0.68	0.44
39	35 ± 37	185	0.7	0.42 ± 0.95	8.66	7563	170	302	0.04	1.8	7	0.58	5	0.42	20.9%	65.1%	0.08	0.17	0.55
40	31 ± 44	350	0.8	0.45 ± 1.01	9.36	44 694	310	490	0.01	1.6	6	0.67	3	0.33	6.8%	63.4%	0.09	-0.11	0.72
East	22 ± 11	171	0.8	0.87 ± 0.42	12.69	15264	178	674	0.12	4.3	5	0.50	4	0.50	12.4%	61.2%	0.05	0.27	0.54
41	17 ± 35	333	0.9	0.91 ± 2.19	28.13	27 790	242	841	0.03	3.5	6	0.60	4	0.40	19.0%	65.4%	0.04	-0.25	0.66
42	104 ± 202	1579	0.9	0.47 ± 1.03	9.89	10232	178	501	0.05	2.8	3	0.60	2	0.40	22.9%	65.6%	0.08	0.46	0.57
43	84 ± 97	603	1.0	1.13 ± 2.49	19.23	17 528	170	1273	0.07	7.5	6	0.86	1	0.14	20.7%	65.6%	0.10	0.18	0.76
44	13 ± 35	362	0.9	0.05 ± 0.15	1.10	1165	96	32	0.03	0.3	1	0.50	1	0.50	8.9%	51.0%	0.06	-0.20	0.18
45	21 ± 31	260	0.9	0.41 ± 0.71	4.80	2790	138	231	0.08	1.7	5	0.45	6	0.55	13.0%	58.3%	0.16	0.75	0.79
46	115 ± 141	1497	1.0	2.29 ± 5.56	56.98	142 175	164	2675	0.02	16.3	2	0.50	2	0.50	14.1%	53.7%	0.27	0.32	0.30
47	91 ± 3	396	1.4	0.50 ± 4.15	10.37	8043	180	437	0.05	2.4	3	0.60	2	0.40	17.9%	45.4%	0.11	0.61	0.49
Northeast	63 ± 41	719	1.0	0.82 ± 0.68	18.64	29 960	167	856	0.05	4.9	4	0.59	3	0.41	16.7%	57.9%	0.12	0.27	0.54
Gridcell average	94 ± 121	793	0.9	4.54 ± 6.37	33.38	21 823	176	2685	0.19	16.2	4	0.44	5	0.56	13.5%	74.7%	0.11	0.13	0.54
Avg gridcell min	2	11	0.7	0.05	1.10	553	35	32	0.01	0.3	0	0	0	0	5.0%	45.4%	0	-0.45	0.08
Avg gridcell max	555	1925	1.9	30.79	164.93	142 175	344	16 277	1.00	111.9	13	1.00	20	1.00	25.9%	99.9%	0.44	0.83	0.92

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back


Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






**Fig. 1.** Methodology flowchart of MODIS remote sensing of sediment plumes. Daily 500 m MODIS reflectance data were classified to identify open water pixels ranging from clear water to sediment-rich water, spatially validated using higher-resolution ASTER and Landsat TM/ETM+. Resulting 500 m open water classification produced measures of sediment persistence and data density. To increase temporal sampling, data were aggregated into 100 km coastal gridcells, whose value was represented by  $R_{peak}$  within the gridcell and a 7-day interval.  $R_{peak}$  was derived from fjord ROIs that met a data density threshold. SSC was extracted for each 100 km coastal gridcell using an empirical model relating MODIS band 1 reflectance ( $R_{peak}$ ) and SSC. New measures of sediment persistence and data density along with average SSC were calculated from these data.

## Hydrologic controls on coastal suspended sediment plumes

V. W. Chu et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	






**Fig. 2.** MODIS mosaic of Greenland (RGB = bands 1, 4, 3) with Landsat TM/ETM+ (RGB = bands 3, 2, 1) and ASTER (RGB = bands 2, 1, 1) image locations used for classification validation. Remote sensing analysis included all fjords directly draining the ice sheet through land-terminating glaciers (and downstream rivers) or from marine-terminating glaciers where plumes may form. Landsat and ASTER images show examples of different outlets and resulting plumes, including: (1) fjords fed by land-terminating glaciers, with large sediment plumes forming after transport through a river floodplain, (2) fjords with marine-terminating glaciers, few icebergs and a visible plume, and (3) fjords with marine-terminating glaciers heavily calving ice and no visible plume.

## Hydrologic controls on coastal suspended sediment plumes

V. W. Chu et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
⏪	⏩
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	





**Fig. 3.** Coastal 100 km gridcells and their corresponding ice sheet drainage basins are overlaid onto an image showing data density ( $DD_{500m}$ ) for 10 years of MODIS 500 m data, representing the number of open water days classified for each pixel out of a total of 1531 available and ranging from 0–40.6%. Regions of interest (ROIs, red circles) are manually drawn to restrict open water data to fjords draining the ice sheet. Gridcells are grouped into six regions for summary analysis: Northwest (NW), West (W), Southwest (SW), Southeast (SE), East (E), and Northeast (NE). The north/northeast and a swath along the southeast contain very low frequencies of detected open water days due to the presence of sea ice and glaciers heavily calving icebergs.

**Hydrologic controls on coastal suspended sediment plumes**

V. W. Chu et al.


Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
⏪	⏩
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	





## Hydrologic controls on coastal suspended sediment plumes

V. W. Chu et al.



**Fig. 5.** (a) Map of 10-year mean ice sheet PDD (blue) and fjord plume SSC (red circles). (b) Spatial variation of 10-year mean PDD (grey line) and SSC (black line), starting in the northwest and going counterclockwise towards the northeast ( $R = 0.70$ ,  $p < 0.001$ ). (c) Maps of mean yearly SSC (red circles) and PDD (blue) for 2000–2009.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version


Interactive Discussion

## Hydrologic controls on coastal suspended sediment plumes

V. W. Chu et al.



**Fig. 6.** Interannual variations of ice sheet PDD and plume SSC for the entire ice sheet and for each of the six regions from 2000–2009. The West and Northeast regions track each other rather well, while the Southwest and Southeast regions are fairly anti-correlated.



**Fig. 7.** Maps of parameters and analyses for each of the 47 gridcells. **(a)** Glacier outlet type shows the number and proportion of outlets in each gridcell that is land-terminating ( $N_L$ ) or marine-terminating ( $N_M$ ), with the size of the circle representing the total number of outlets reaching the fjord. **(b)** Sediment persistence ( $F_{SSC}$ ) shows the fraction of open water days with sediment-rich water detected. **(c)** Data density ( $DD_{100km}$ ) shows the fraction of total days with open water detected (free of ice and clouds) for the aggregated 100 km data. **(d)** Correlation between interannual averages of PDD and SSC over 2000–2009. **(e)** Correlation between mean seasonal cycles of PDD and SSC.

## Hydrologic controls on coastal suspended sediment plumes

V. W. Chu et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶

Back Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## Hydrologic controls on coastal suspended sediment plumes

V. W. Chu et al.




**Fig. 8.** Mean seasonal development of ice sheet PDD and downstream fjord SSC for each of the six regions. A seasonal climatology is calculated by taking day-of-year averages over 2000–2009. Seasonal cycles broadly coincide with plume onset coincident with or after surface melt onset, though the East and Northeast regions show difficulty detecting onset due to presence of sea ice. High SSCs persist in the Southwest region during the waning of the melt season.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

## Hydrologic controls on coastal suspended sediment plumes

V. W. Chu et al.



**Fig. 9.** Examples of high frequency time series of SSC and PDD for one gridcell in each of the six regions from 2000–2009. SSC is displayed for 2010 but not used in analysis due to PDD only being available up to 2009.