

The Cryosphere Discuss., 5, 721–753, 2011
www.the-cryosphere-discuss.net/5/721/2011/
doi:10.5194/tcd-5-721-2011
© Author(s) 2011. CC Attribution 3.0 License.



This discussion paper is/has been under review for the journal The Cryosphere (TC).
Please refer to the corresponding final paper in TC if available.

Temperature variability and thermal offset in steep alpine rock and ice faces

A. Hasler, S. Gruber, and W. Haeberli

Glaciology, Geomorphodynamics and Geochronology, Department of Geography,
University of Zurich, Switzerland

Received: 8 February 2011 – Accepted: 15 February 2011 – Published: 1 March 2011

Correspondence to: A. Hasler (andreas.hasler@geo.uzh.ch)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper

TCD

5, 721–753, 2011

Temperature variability in steep alpine rock and ice faces

A. Hasler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Temperature variability in steep alpine rock and ice faces

A. Hasler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

(near-) surface temperatures measurements exist (Gruber et al., 2004; Pogliotti et al., 2008; Allen et al., 2009; PERMOS, 2010; Wegmann et al., 1998; Coutard and Fran-
 cou, 1989; Matsuoka, 2008; Matsuoka and Sakai, 1999); (b) only few boreholes for
 temperature measurements in steep bedrock permafrost exist in the European Alps
 (PERMOS, 2010; Noetzli et al., 2010; Wegmann, 1998); (c) no empirical study on the
 temperatures of steep ice faces is known to the authors. One use of the surface tem-
 perature measurements is the validation of distributed surface energy balance models
 to extrapolate rock face temperatures in space and time and to assess permafrost
 distribution (Gruber et al., 2004; Noetzli et al., 2007). An other one is the long-term
 observation of these temperatures as a proxy for the permafrost conditions in steep
 bedrock (PERMOS, 2010).

In this study we address the question how representative surface temperature mea-
 surements are and whether systematic deviations (thermal offsets) between the mean
 annual rock/ground surface temperature (MAGST) and the permafrost temperature be-
 low exist. We investigate the variability of rock temperatures and thermal offsets and
 their dependence on surface and near-surface characteristics. As a special case of sur-
 face characteristics we additionally investigated the thermal condition in a thin (<10 m)
 ice cover of a steep rock face, which usually are called *ice faces* and that indicate
 underlying permafrost. We analyse mean annual temperatures and thermal offsets
 derived from 17 shallow temperature profiles in bedrock, rock clefts and ice at Mat-
 terhorn and Jungfrauoch (Swiss Alps). The approach to analyse the dependency of
 subsurface temperatures on the different characteristics is descriptive-heuristic, hence
 we quantitatively describe differences (Sect. 4) and try to understand this observed
 variation in terms of the driving processes (Sect. 5).

A recent study on the small-scale variability of mean annual ground surface tem-
 peratures (MAGST) in gently mountain slopes, found a variability of 0.16–2.5 °C within
 10 m × 10 m footprints (Gubler et al., 2011). With the rough micro-topography typical
 for steep fractured bedrock, we expect MAGST-variabilities at the upper end of this
 range. The variation of ground temperatures with shallow depths below surface is

Temperature variability in steep alpine rock and ice faces

A. Hasler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

usually described with the thermal offset (TO), which is defined as the difference between the temperature below the active layer at the permafrost table (TTOP) and the MAGST (Burn and Smith, 1988). This effect is well-known in arctic soils, and Gruber and Haeberli (2007) proposed three possible sources of TOs making its importance also likely in steep fractured bedrock: (1) variable thermal conductivity due to saturation and phase changes of pore water (thermal diode effect of rock); (2) changes of the heat transport across clefts as a consequence of freeze/thaw/runoff of cleft ice (thermal diode effect of clefts); (3) ventilation effects within loose block cover on less steep parts of rock faces. All these processes are expected to reduce temperatures at depth compared with MAGST. For practical reasons we use the measured offset between top and bottom in profiles as indicator of TO. In fractured bedrock the strict definition of TO as TTOP–MAGST is impractical anyway, because of: (a) highly variable MAGST; (b) processes that cause offsets that are not limited to the active layer; and (c) highly variable active layer thickness. Hence we call the temperature difference between near-surface and shallow depth of the profile *thermal offset* even though the lower thermistors are often above the permafrost table and the true offset between MAGST and permafrost temperature is expected to be larger in some cases. However, the values for offsets and temperature variability given in this study have an exemplary character and indicate possible ranges because many degrees of freedom exist in the possible variations of controlling parameters.

2 Measurement setup

2.1 Field sites

In this study, distributed temperature measurements from two permafrost field sites in the Swiss Alps – Matterhorn and Jungfrauoch – are analysed. The sites are located at similar elevation and in comparable topographic situations but differ concerning their geological structure and near-surface characteristics. In proximity of both sites rock

falls of small to medium magnitude ($\approx 1000\text{--}150\,000\text{ m}^3$) occurred within the last century. The Matterhorn is part of the main divide of the western Alps that marks the Swiss-Italian border. The Matterhorn field site (mh) is at the north-east ridge called Hörnligrat at an elevation of 3450 m a.s.l. and comprises both sides of the ridge with main orientations southeast and north (Fig. 1). The bottoms of both rock faces are glaciated, on the south-eastern side by a large plateau causing strong reflection of solar radiation. Jungfrauoch (3500 m a.s.l.) is a mostly glaciated saddle of the northern Alpine range dividing the northern Pre-alps from the glaciated Aletsch basin. The “Sphinx” is an exposed rock ridge in the saddle with diverse tourist and research facilities. The measurement locations are on the northern and southern side of the Sphinx (Fig. 1).

The mean annual air temperature (MAAT, average 1961–1990) is approximately $-6.7\text{ }^\circ\text{C}$ at the Matterhorn field site and $-7.3\text{ }^\circ\text{C}$ at the Jungfrauoch (Hiebl et al., 2009) and currently subject to an accelerating warming trend (Beniston, 2005). Except for some occasional rainfall in summer, all precipitation falls as snow, hence liquid water is mainly supplied by snow melt. Due to the location at the northernmost high-alpine ridge with corresponding orographic cloud formation, the Jungfrauoch receives less annual solar radiation and more precipitation than Matterhorn-Hörnligrat. The southern rock faces at both field sites experience extreme solar radiation due to reflection from the glaciers underneath, making strong daily cycles with positive rock surface temperatures common in clear-sky conditions during all seasons.

The structure at the two field sites differs mainly with respect to fracturation: although metamorphic crystalline rocks prevail at both sites, the frequency and aperture of clefts is significantly different. At Jungfrauoch 5–20 clefts per meter and apertures of 0.5 mm to 3 cm are typical (Fig. 3) while at Matterhorn clefts are less frequent ($0.5\text{--}5\text{ cl m}^{-1}$) but have larger typical apertures (3–30 cm) (Fig. 4). This difference affects the thermal properties of these rock masses, because the thermal parameters of the inter-joint rock mass are overprinted by the geometric setting of the discontinuities: changes in water content and phase state within these discontinuities will influence the overall

Temperature variability in steep alpine rock and ice faces

A. Hasler et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

thermal conductivity in a near-surface layer more than what may be expected from the laboratory-derived thermal parameters of intact rock samples. The described difference in the cleft characteristics but in a similar topographic situation was an important motivation for the selection of the two complementary field sites.

2.2 Instrumentation

At both field sites, wireless sensor networks (WSNs) that record environmental parameters and transmit the data to an Internet server were installed. The conception and setup of these WSNs are described in detail by Beutel et al. (2009) and Hasler et al. (2008). Beside geotechnical and hydrological parameters, temperature measurements with totally 100 temperature sensing elements (YSI 44006 NTC-thermistors) where recorded with high temporal resolution since July 2008 at Matterhorn and since February 2009 at Jungfrauoch. Several differing sensors can be attached to one network node of the WSN, which is then termed *sensor node*, while the expression *base station* is used for the central node that transmits the data off the mountain. Sensor nodes are labelled with abbreviations of the field site mh for Matterhorn and jj for Jungfrauoch) and a number for the location (Fig. 1). Custom-built sensor rods measure the temperature and electrical resistance of the rock at four depths (0.1, 0.35, 0.6 and 0.85 m) in a 0.9 m deep boreholes, which are perpendicular to the surface (Hasler et al., 2008). Similarly, thermistor chains and thermistor–moisture chains measure four to eight temperatures within clefts or in ice faces. For the clefts, the precise physical context of the measured value is more complicated than for the other cases, because the temperature at the sensing element is influenced by the temperature of the air and the rock surface within the cleft or even by ice or percolating water. The measured temperatures within a profile are labelled T_1 – T_4/T_8 with increasing depth (e.g. $T_1 = 0.1$ m and $T_4 = 0.85$ m depth for all sensor rods; cf. Table 1). The depth of these measurements is not exactly defined for all sensors and depends on the installation at each location (see Sect. 2.3). In addition to these multiplex sensors, rock surface temperatures (T_s) are measured with individual thermistors placed 2 cm below the surface in

Temperature variability in steep alpine rock and ice faces

A. Hasler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



small inclined borings minimizing disturbance from solar radiation on the cables. Two sensor rods (jj04 and jj09) were not considered for this study due to malfunction.

2.3 Description of the measurement locations

Nine sensor nodes at Matterhorn and eight sensor nodes at Jungfrauoch perform cleft, rock, or ice temperature measurements relevant for this study (Table 1). At Matterhorn the sensors mh01, mh02, mh05, and mh10 lie on the southeast side of the ridge and are exposed to intense solar irradiation, while mh07 and mh12 are on the ridge and the other sensors are in the shaded north face (Fig. 1). At Jungfrauoch the locations jj01–jj04 are on the southwest slope and the other locations including the two ice boreholes are on the northern side of the Sphinx (Fig. 1). The aspect and slope indicated in Table 1 is the orientation of the surface 1–2 m around the sensor. Furthermore, Table 1 contains remarks on special features such as the micro-topographic situation, snow cover, wetness or fracturation. The depths of thermistors (perpendicular to the surface) is indicated for all sensor rods and for cleft temperatures where it is clearly defined. Otherwise, an estimated depth-range and the corresponding number of thermistors is denoted (Table 1). The depths of the thermistor chains in the ice face are not constant with time as the face accumulates small quantities of ice (about $0.05\text{--}0.3\text{ m a}^{-1}$), however, the distance between sensors remains constant. Based on the installation depth of 0.3 m and the evolution of the amplitudes of the uppermost sensor, we estimate its average depth during the measured period as 0.7 m from the surface (Table 1). The location of the surface temperature measurements are in similar local orientations at 0.1–1 m distance from the boreholes of the sensor rods or from the clefts.

The rock temperature measurements at Matterhorn aim to record the thermal conditions in snow-free and compact rock as a reference for the cleft temperature measurements and comparison to RST-measurements in other areas. Therefore near-vertical bedrock of the three main aspects that persist at this field site (NW, NE, SE) was instrumented with sensor rods. For the location mh10 and mh12, however, no sufficiently large compact rock mass could be found and clefts in proximity of the boreholes exist

Temperature variability in steep alpine rock and ice faces

A. Hasler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(Table 1). At Jungfrauoch, the locations of the rock temperature measurements are selected to cover gradients in surface and near surface conditions. For the two main aspects (N, S) different locations with respect to slope angle (snow retention), microtopography (water availability; only at S) and fracturation where selected (Table 1). The two sensors that failed are the snow-covered one in the north face (jj09) and the sensor in unfractured rock at the south side (jj04). As a consequence, the effect of snow cover in the northern face and difference caused by fracturation for the south side could not be assessed at this field site.

3 Data processing and quality

The raw data series contain invalid measurements or data gaps and the sampling interval of two minutes is slightly irregular. This demands a processing prior to the calculation of mean annual temperatures and thermal offsets within the profiles. First, invalid data is filtered and the remaining data is aggregated to regular intervals. After this, data gaps are filled. As these processing steps but also the characteristics and timing of the data acquisition introduce uncertainty into the computation of mean annual temperature an uncertainty analysis concludes this section.

3.1 Data validation, filtering and accuracy

Each multiplex sensor measures stable reference resistors during each cycle of temperature measurements. Deviations in these reference values correlate with invalid temperature measurements if the source of error is within the data acquisition system and not at the sensing element itself. A threshold is applied on the reference values to filter invalid measurements from the raw data before averaging to ten-minute aggregates. This down-sampling does not lead to a significant loss of information because of the inertia of ground temperatures. Due to this down-sampling the resulting time series are usually continuous because invalid data is rare (with the exception of mh01). A

Temperature variability in steep alpine rock and ice faces

A. Hasler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



time series for more than a year are available and no gap filling was needed while in most of the other datasets 5–10% of the data was missing. The effect of the gap filling on MAT was evaluated by introducing the same gaps into the complete time series; this showed that an approximation of the true MAT to better than ± 0.1 °C was achieved with gap filling compared to ± 1 °C if gaps contain no values. Sensors mh02 and mh12 contain larger gaps and therefore introduce a larger uncertainty into the MAT estimate (Table 2).

3.3 Uncertainty analysis of mean annual temperatures and thermal offsets

Three main sources of uncertainty (Table 2) affect our estimate of the MAT: (a) systematic measurement errors (U_{meas}), (b) data gaps (U_{gap}), and (c) the period for which the mean is calculated (U_{time} , $U_{\text{time-to}}$). U_{meas} is given by the measurement accuracy (Sect. 3.1) because the bias from the measurement is systematic over the whole time series and is not significantly reduced by the averaging. For U_{gap} the values are estimated dependent on quantity of missing data in the averaging window (Sect. 3.2) but lower values are chosen in case of the ice temperatures due to smooth time series and correspondingly better performance of the gap-filling algorithm. MAT calculations are influenced by the start and end date of the averaging window on the long term (inter-annual variation of MAT) but also on the short term (seasonal) if the temperature time series show strong weekly variations. Figure 6 shows the temperature time series and the seasonal variation of the MAT for the sensor rod at mh10 (rock). This variation is considered as uncertainty U_{time} for the comparison of the MATs. This is because the variation of the MAT is not correlated between locations. The MAT values calculated for 1 October 2009 to 1 October 2010 (for all sensors except jj01, mh04 and mh12) are approximately in the middle of this variation range (cf. Fig. 6). The variation of the MAT is, however, influenced by data gaps, hence for three sensors the part of the time series with large gaps is excluded from the estimation of U_{time} that is performed by the difference of 2.5 and 97.5% quantiles (Table 2). As the running annual means of the temperatures at different depth but at the same location are correlated (Fig. 6),

Temperature variability in steep alpine rock and ice faces

A. Hasler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the thermal offset (TO) varies less over time. For that reason $U_{\text{time_to}}$ is estimated as a measure of the uncertainty in thermal offset calculation, introduced by timing of the averaging interval, which is in most cases smaller than U_{time} (Table 2).

The total uncertainties of the MAT (U_{mat}) and the TO (U_{to}) are calculated by quadratic addition of the uncertainties according to the addition of spreads:

$$U_{\text{mat}} = \sqrt{U_{\text{meas}}^2 + U_{\text{gap}}^2} = U_{\text{time}}^2 \quad (1)$$

$$U_{\text{to}} = \sqrt{2 \cdot U_{\text{meas}}^2 + U_{\text{gap}}^2 + U_{\text{time_to}}^2} \quad (2)$$

Contrary to Eq. (1), the U_{meas} term is multiplied by a factor of two in Eq. (2) because the independent uncertainties of two temperature measurements contribute to U_{to} . U_{gap} of two measurements in the same profile are correlated and therefore their single consideration is a worst case. However this influence is negligible in most cases anyway Table 2. The resulting uncertainties that are relevant for the interpretation of the MAT and the TO are listed in Table 2.

4 Results

4.1 Mean annual temperatures (MATs)

Figure 7 gives an overview of the MATs of the clefts, rock (MAGT) and ice ordered by location and type (colors). The representation as profiles with z being the distance from surface does not show the real distance between the sensors and lateral offsets in the thermistor position are masked in case of the cleft and ice temperatures. The MAT values from the north-oriented locations cluster around -6°C and are slightly warmer ($0.5\text{--}1.5^\circ\text{C}$) than the MAAT (Fig. 7). Remarkable is the exact match in mean annual rock/ground temperature (MAGT) of mh11 and jj05, which are both north oriented in intact steep rock (Fig. 7; for a better differentiation of these values see also Fig. 8).

Temperature variability in steep alpine rock and ice faces

A. Hasler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Temperature variability in steep alpine rock and ice faces

A. Hasler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The mean temperatures at the surface (MAGST) at the more sun-exposed locations are 1–8 °C higher than the shaded ones and the same is true for the near-surface cleft temperatures (MAT of T_1) (Fig. 7). The difference in MAGT between sunny and shaded locations is more pronounced at Matterhorn than at Jungfrauoch. This is because the south face MAGT at Matterhorn (mh10) is 3–4 °C higher for T_1 (0.1 m) and 2 °C higher for T_4 (0.85 m) than the ones at Jungfrauoch (jj01, jj02 and jj03) (Fig. 7).

Cleft MATs of the east-oriented locations at Matterhorn are significantly lower than the MAGT at locations with comparable orientation: the cleft at mh07 is 4 °C colder at the top and 3 °C colder at depth than the rock at mh12 which is only a few meters above in the same face; the two clefts mh02 and mh05 are 2–3 °C colder than mh12 although they face more toward south; at depth even the radiation exposed profile mh01 is colder than mh12 (Fig. 7). The MAT-profiles from the ice faces start around –5.5 °C near the surface and show a constant positive temperature gradient with depth of approximately +0.2 °C m⁻¹ (Fig. 7). The near-surface MAT in the ice is 0.2–0.8 °C warmer than in the rock face just above the ice face, hence the difference found is very marginal in this case.

4.2 Thermal offsets (TOs)

To quantify the temperature difference between the near-surface and greater depth, the MAT of T_4 is subtracted from the one of T_1 or for the ice face from T_8 and T_1 respectively, hence, positive differences indicate a warming with depth (Fig. 8). The rock surface temperature measurements (T_s) are not considered for this calculation to avoid a mix of rock and cleft temperatures and to keep sensor rod measurements with and without T_s comparable. For mh02 T_6 is taken instead of T_4 because the MAT of the later one is missing. As stated in Sect. 1, we call this temperature difference between near-surface and depth *thermal offset* (TO), using this term in a more general way than other publications. The difference in depth between the two thermistors that are used for the TO calculation is only constant for the rock temperatures (Fig. 7), hence, the

TO-values in Fig. 8 are directly comparable for the rock measurements but smaller or larger depth range needs to be considered for the cleft and ice temperatures.

To ease interpretation of the MAT and TO in Fig. 8, the local orientation (main aspect) of the measurement locations is indicated at the top. The location labels given in the middle of the figure help to read other attributes from Table 1. At the bottom, the location type is denoted, which corresponds to the colours of the bars from the thermal offset. The error bars in the TO-graph show U_{to} (Table 2) and indicate the significance of thermal offsets values.

In total, seven TOs are negative, four are positive and six lie within the uncertainty range (Fig. 8). More than half (4) of the clearly negative TOs are detected within the clefts, the two most positive TOs consist of the ice face measurements. The locations with warmest surface temperatures (mh01, mh10 and mh12) have most negative TOs and are all located at Matterhorn (2 in rock, 1 in cleft). From the rock temperatures at Jungfraujoch, only one sensor shows negative and one positive TO, whereas all the other sensors have no significant TO. This is in contrast to the Matterhorn data where seven out of nine cleft and rock sensors show a significant TO (Fig. 8). A further regularity is, that all sensors with a slight or significant positive TO are relatively flat and accumulate often a snow cover (cf. Table 1).

4.3 Seasonal temperature variation and inter-annual variability of MAT

To reveal some processes that are responsible for TOs and variations in MAT, a qualitative analysis of the time series from the measured data, the MATs and the smoothed temperature difference ($\Delta T = T_4 - T_1$) is presented in three examples. The data from jj05 serves as a reference for a rock temperature profile that has no significant TO (Fig. 9): the 30-days running mean of ΔT has similar negative and positive amplitudes ($\pm 2^\circ\text{C}$) and results in a TO close to zero if averaged over a year. This is also shown with the overlapping MATs that at the same time indicates the small seasonal variation (compare with U_{time} in Table 2). In Fig. 10 two examples of time series are presented to illustrate, which periods of the year are responsible for the thermal offsets and what

Temperature variability in steep alpine rock and ice faces

A. Hasler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



explains large variations of the MATs and TOs between different years: the time series of the cleft mh01 shows large seasonal variations and very large daily amplitudes in spring and summer that are not symmetrical with the temperatures at depth and cause a negative ΔT from March to November for both years (Fig. 10). Similar seasonal patterns are found at all sensors with large negative TOs (mh07, mh10, mh12). In contrast, at jj01 (rock) positive temperatures and large daily amplitudes at T_1 are limited to the snow free period in summer and the winter temperatures are smoothed by the snow cover (Fig. 10). Because the snow-free periods differ between 2009 and 2010 and the temperatures at depth are buffered by thawing ground ice (zero curtains), the summer ΔT varies strongly and the TO changes from positive to negative values (Fig. 10).

5 Discussion

The general near-surface rock temperature pattern with a MAGT being slightly higher than the MAAT in shaded rock faces and several °C higher at radiation-exposed locations corresponds to other studies and reports from steep high-alpine bedrock (Coutard and Francou, 1989; Gruber et al., 2004; PERMOS, 2010). However, the temperatures are 2–3 °C lower than the ones of Gruber et al. (2004) for this elevation in the Swiss Alps. The lower MAGTs at Jungfrauoch compared to Matterhorn may be explained partly by less direct solar irradiation due to more cloud cover determined by the more western orientation of the sensors with more convective cloudiness in the afternoon and the climatic situation (orographic clouds at the northern divide). The data from the defect sensor jj04 (T_3 and T_4 have sufficient data to calculate annual means), however suggests that MAGT in the range of -0.5 °C occur at the south slope of Jungfrauoch as well. Hence, we assume other factors such as snow retention (jj01, jj02), cooling by melt water (jj02) and local shading (jj03, jj02) due to the micro-topographic situation as mainly responsible for the lower near-surface MAGTs at the Jungfrauoch south face (Table 1; Fig. 3). The same cooling effect by local snow cover and more shading due

Temperature variability in steep alpine rock and ice faces

A. Hasler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



to the concave micro-topography may be responsible for the lower cleft MATs at mh02 and mh05 in comparison with the near-surface MAT of mh01 that has the same orientation. This net cooling effect of the snow cover is in contrast to the net warming effect on more gentle slopes where thick snow cover causes a preponderance of “warming” by winter insulation over the “cooling” by increased albedo and latent heat consumption (Keller and Gubler, 1993). In steep slopes at high elevation the thinner snow cover and summer snowfalls could result in a reverse effect. This is supported by the data from jj01 showing that the surface remains snow-covered in the period with most intense solar irradiation (June and July) and that winter cooling indicated by upward heat fluxes ($\Delta T = +4^\circ\text{C}$; larger than e.g. at jj05) is not prevented (Fig. 10).

The comparably colder cleft temperatures at depth (Sect. 4.1) at locations without snow cover (mh01, mh07) need an alternative explanation (even though a part ($\approx 0.5^\circ\text{C}$) of the cooling with depth at mh01 may originate from lateral heat fluxes through the ridge). The large negative TOs of these clefts and the contrast to the rock surface temperature at mh01 point to strong non-conductive effects responsible for this cooling. Air ventilation is a likely source of cooling at depth because irradiation is reduced in open clefts and the temperature in the lower cleft approximates air temperature depending on the intensity of the sensible heat exchange similar to effects in coarse debris layers (Harris and Pedersen, 1998; Hanson and Hoelzle, 2004). A second cooling effect may be the latent energy consumption by the melt of snow that is deposited in larger clefts (cf. Fig. 4). However, this process is only active if cleft temperatures are at 0°C . The negative TO of $0.5\text{--}1.5^\circ\text{C}$ measured in rock (mh10, mh12, jj06) is well explained by the cooling within the clefts because all three boreholes are in proximity to open clefts (Table 1). Changes in thermal conductivity due to phase change of cleft and pore water (in case of jj06 the borehole crosses two clefts) could be an additional source of a negative TO at mh03 and mh06 (Gruber and Haerberli, 2007; Pogliotti et al., 2008). The seasonal pattern of ΔT (Sect. 4.3) fits best to the ventilation hypothesis for clefts because: (a) the outward heat flux in winter would reduce ΔT (all sensors); (b) radiation can not directly affect the upper most thermistor

Temperature variability in steep alpine rock and ice faces

A. Hasler et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

gradient ($dT/dz < 0.02 \text{ } ^\circ\text{C m}^{-1}$). The lateral heat flux may be, however, ten times larger ($ht = 0.3 \text{ W m}^{-2}$) due to the warm southern side and the heat introduced by the infrastructure within the ridge (Wegmann, 1998) induce a temperature gradient within the ice face in the order of $0.2 \text{ } ^\circ\text{C m}^{-1}$.

6 Conclusions and perspectives

The thermal conditions of steep bedrock permafrost and ice faces were studied based on 17 shallow temperature profiles. On the basis of two-year time series from two field sites in the Swiss Alps, we calculated the mean annual temperatures (MAT) and their offsets (TO) within the profiles and analyzed them with respect to their micro-topographic situation, surface and near-surface characteristics. The main findings are:

- Differences in MAT and TO are highly significant with respect to the uncertainty introduced by measurement errors, data gaps and temporal variations.
- When using MAGST as an indication for the permafrost temperature in mountain faces, one needs to account for thermal offset, similar to arctic lowland areas.
- The ice face investigated in this study has similar MAT as the rock beside and no clear evidence for TOs by latent heat release from percolation effects was found.
- Snow cover likely reduces MAGST ($2\text{--}3 \text{ } ^\circ\text{C}$) of mid-steep ($45\text{--}70 \text{ } ^\circ\text{C}$) locations in radiation-exposed faces at high elevation because it often persists for the period with most intense radiation (June).
- Ventilation effect of clefts causes negative TOs and lower temperatures at depth ($\approx 1.5 \text{ } ^\circ\text{C}$) for strongly fractured near-vertical bedrock at radiation-exposed locations.
- Other processes such as thermal diode effects and local shading may support colder MAGTs but could not be quantified with the available data.

Temperature variability in steep alpine rock and ice faces

A. Hasler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Local warming within clefts by heat advection of percolating water shows minor effects on MAT, however, it should be considered in respect of rock stability.
- Summarizing the previous statements we postulate that radiation-exposed steep rock faces with intermitted snow patches and/or large fractures are up to 3°C colder at depth than expected from MAGST at snow-free locations.

The lowering of rock temperatures in rock faces should be considered for the estimation of permafrost occurrence, which may in fact extend to lower elevations by several hundred meters in the south and east sector than expected so far. Corresponding effects could be parameterized by the use of surface and near-surface characteristics that affect snow retention and ventilation. However the following limitations should be considered if these findings are extrapolated: (a) the two effects should be considered to be complementary rather than cumulative, because snow reduces the efficiency of the ventilation; (b) the ventilation effects depends on cleft aperture and frequency, hence near-surface characteristics need to provide an information on this aspect; and (c) the effect of snow cover could change with elevation due to a changed duration of the snow-free period. To estimate the latter, the presented data may be used to calibrate the snow cover in a physically oriented permafrost model: first the model should be validated with the snow-free sensors before the data from less steep locations with snow are used to calibrate the duration and thickness of the snow cover. The measured gradients in the near-surface layer can serve as a direct estimate of the heat flux through the snow cover if temperatures at the surface are below zero. As long as no further analysis and model-based spatial extrapolation of these findings is performed, we suggest to include the up to 3°C lower temperatures in radiation-exposed rock faces in the uncertainty indications of MAGT estimates.

Temperature variability in steep alpine rock and ice faces

A. Hasler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Acknowledgements. We would like to thank the PermaSense team, J. Beutel, R. Lim, M. Yücel, I. Talzi, T. Gsell, M. Keller, L. Thiele and C. Tschudin, who made these challenging measurements possible. The presented research was funded within the project PermaSense by the Swiss Federal Office for the Environment (FOEN) and the Swiss National Foundation (SNF) NCCR-MICS. The fieldwork at Jungfrauoch was supported by the International Foundation High Altitude Research Stations Jungfrauoch and Gornergrat (HFSJG) and benefited from the friendly support of the Jungfrau Railways and their staff.

References

- Allen, S. K., Gruber, S., and Owens, I. F.: Exploring steep bedrock permafrost and its relationship with recent slope failures in the Southern Alps of New Zealand, *Permafrost Periglac.*, 20, 345–356, 2009.
- Beniston, M.: Mountain climates and climatic change: An overview of processes focusing on the European Alps, *Pure Appl. Geophys.*, 162, 1587–1606, 2005.
- Beutel, J., Gruber, S., Hasler, A., Lim, R., Meier, A., Plessl, C., Talzi, I., Thiele, L., Tschudin, C., Woehrle, M., and Yucel, M.: PermaDAQ: A scientific instrument for precision sensing and data recovery in environmental extremes, in: *Proceedings of the 2009 International Conference on Information Processing in Sensor Networks*, IEEE Computer Society, 265–276, 2009.
- Burn, C. R. and Smith, C. A. S.: Observations of the “thermal offset” in near-surface mean annual ground temperatures at several sites near Mayo, Yukon Territory, Canada. *Arctic*, 41, 99-104, 1988.
- Coutard, J. P. and Francou, B.: Rock temperature measurements in two alpine environments: implications for frost shattering, *Arctic Alpine Res.*, 21, 399–416, 1989.
- Gruber, S. and Haeberli, W.: Permafrost in steep bedrock slopes and its temperature-related destabilization following climate change, *J. Geophys. Res.*, 112, F02S18, doi:10.1029/2006JF000547 2007.
- Gruber, S., Hoelzle, M., and Haeberli, W.: Rock-wall temperatures in the Alps: modelling their topographic distribution and regional differences, *Permafrost Periglac.*, 15, 299–307, 2004.

Temperature variability in steep alpine rock and ice faces

A. Hasler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Temperature variability in steep alpine rock and ice faces

A. Hasler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Gubler, S., Fiddes, J., Gruber, S., and Keller, M.: Scale-dependent measurement and analysis of ground surface temperature variability in alpine terrain, *The Cryosphere Discuss.*, 5, 307–338, doi:10.5194/tcd-5-307-2011, 2011.
- 5 Haeberli, W., Wegmann, M., and Vonder Muehll, D.: Slope stability problems related to glacier shrinkage and permafrost degradation in the Alps, *Ecologiae Geologicae Helvetiae*, 90, 407–414, 1997.
- Hanson, S. and Hoelzle, M.: The thermal regime of the active layer at the Murtel rock glacier based on data from 2002, *Permafrost Periglac.*, 15, 273–282, 2004.
- Harris, S. A. and Pedersen, D. E.: Thermal regimes beneath coarse blocky materials, *Permafrost Periglac.*, 9, 107–120, 1998.
- 10 Hasler, A., Gruber, S., Font, M., and Dubois, A.: Advective heat transport in frozen rock clefts – conceptual model, laboratory experiments and numerical simulation, *Permafrost Periglac. Process.*, in review, 2011.
- Hasler, A., Talzi, I., Beutel, J., Tschudin, C., and Gruber, S.: Wireless sensor networks in permafrost research: concept, requirements, implementation, and challenges, in: *Proceeding of the Ninth International Conference on Permafrost*, Fairbanks, Alaska, USA, 669–674, 2008.
- Hiebl, J., Auer, I., Böhm, R., Schöner, W., Maugeri, M., Lentini, G., Spinoni, J., Brunetti, M., Nanni, T., and Perec Tadi, M.: A high-resolution 1961–1990 monthly temperature climatology for the greater Alpine region, *Meteorol. Z.*, 18, 507–530, 2009.
- 20 Hoelzle, M., Darms, G., and Suter, S.: Evidence of accelerated englacial warming in the Monte Rosa area, Switzerland/Italy, *The Cryosphere Discuss.*, 4, 2277–2305, doi:10.5194/tcd-4-2277-2010, 2010.
- Keller, F. and Gubler, H.: Interaction between snow cover and high mountain permafrost, Murtel-Corvatsch, Swiss Alps, in: *Proceedings of the Sixth International Conference on Permafrost*, 332–337, 1993.
- 25 Luethi, M. P. and Funk, M.: Modelling heat flow in a cold, high-altitude glacier: interpretation of measurements from Colle Gnifetti, Swiss Alps, *J. Glaciol.*, 47, 314–324, 2001.
- Matsuoka, N.: Frost weathering and rockwall erosion in the southeastern Swiss Alps: Long-term (1994–2006) observations, *Geomorphology*, 99, 353–368, 2008.
- 30 Matsuoka, N. and Sakai, H.: Rockfall activity from an alpine cliff during thawing periods, *Geomorphology*, 28, 309–328, 1999.
- Noetzi, J., Gruber, S., Kohl, T., Salzmann, N., and Haeberli, W.: Three-dimensional distribution and evolution of permafrost temperatures in idealized high-mountain topography, *J. Geophys.*

Temperature variability in steep alpine rock and ice faces

A. Hasler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Res., 112, F02S13, doi:10.1029/2006JF000545, 2007.

Noetzli, J., Gruber, S., and Poschinger, A.: Modellierung und Messung von Permafrosttemperaturen im Gipfelgrat der Zugspitze, Deutschland, *Geographica Helvetica*, 65, 2, 113–123, 2010.

5 PERMOS: Permafrost in Switzerland 2006/2007 and 2007/2008, edited by: Noetzli, J. and Vonder Muehll, D., *Glaciological Report (Permafrost) No. 8/9 of the Cryospheric Commission of the Swiss Academy of Sciences*, 2010.

Pogliotti, P., Cremonese, E., Morra di Cella, U., Gruber, S., and Giardino, M.: Thermal diffusivity variability in alpine permafrost rock walls, in: *Proceeding of the Ninth International Conference on Permafrost*, Fairbanks, Alaska, USA, 1427–1432, 2008.

10 Wegmann, M.: *Frostdynamik in hochalpinen Felswänden am Beispiel der Region Jungfraujoch – Aletsch*, *Mitteilungen der Versuchsanstalt für Wasserbau, Hydrologie und Glaziologie der ETH Zürich*, 161, 1998.

15 Wegmann, M., Gudmundsson, G. H., and Haeberli, W.: Permafrost changes in rock walls and the retreat of Alpine glaciers: a thermal modelling approach, *Permafrost Periglac.*, 9, 23–33, 1998.

Temperature variability in steep alpine rock and ice faces

A. Hasler et al.

Table 1. Type and orientation of measurement locations with depth of the thermistors.

location	type	aspect / °	slope / °	characteristics	depths of T_s (*), T_1 , T_2 , ... /m
mh01	cleft	95 (E)	75	intense solar radiation	0.02*, 0.1, 0.4, 0.7, 0.5
mh02	cleft	80 (E)	50	corner, often snow, wet	0.1, 0.3, 0.4–0.8 [6]
mh03	cleft	350 (N)	65	lower part snow	0.02*, 0.1, 0.4, 0.6–0.8 [5]
mh04	cleft	320 (N)	70	gully, often snow	0.05, 0.2, 0.2–0.5 [4]
mh05	cleft	90 (E)	60	small corner, often snow	0.1, 0.8, 1.8, 1.5
mh07	cleft	50 (E)	90	large ventilated cleft	0.1, 1, 2, 3
mh10	rock	140 (S)	90	int. solar rad., cleft at 1 m	0.02*, 0.1, 0.35, 0.6, 0.85
mh11	rock	340 (N)	70	occasionally snow, no clefts	0.02*, 0.1, 0.35, 0.6, 0.85
mh12	rock	45 (E)	85	snow free, clefts beside	0.02*, 0.1, 0.35, 0.6, 0.85
jj01	rock	215 (S)	30	often snow, wet	0.1, 0.35, 0.6, 0.85
jj02	rock	220 (S)	50	gully, often snow, wet	0.1, 0.35, 0.6, 0.85
jj03	rock	190 (S)	80	shaded, dry, small clefts	0.1, 0.35, 0.6, 0.85
jj05	rock	330 (N)	85	no macro clefts	0.1, 0.35, 0.6, 0.85
jj06	rock	335 (N)	75	large clefts at 0.15 and 0.4 m	0.1, 0.35, 0.6, 0.85
jj07	rock	330 (N)	75	limestone! occasional. snow	0.1, 0.35, 0.6, 0.85
jj08	ice	340 (N)	45	firn, 7 m from rock wall	0.7, 1, 1.4, 2.1, 2.8, 3.5, 4.2, 4.9
jj10	ice	330 (N)	45	firn, below single cliff	0.7, 1, 1.4, 2.1, 2.8, 3.5, 4.2, 4.9

* rock surface temperature (T_s) measured beside cleft or sensor rod.

[X] number in brackets indicates number of thermistors in the given depth range without exact depth information.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Temperature variability in steep alpine rock and ice faces

A. Hasler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 2. Uncertainties of the MAT and 563 the TO calculation.

location	U_{meas} °C	U_{gap} °C	U_{time} °C	$U_{\text{time_to}}$ °C	U_{mat} °C	U_{to} °C
mh01	0.2	0.1	0.3	0.2	0.37	0.36
mh02	0.2	0.2	0.35	0.25	0.45	0.43
mh03	0.2	0	0.4	0.3	0.45	0.41
mh04	0.2	0	0.35	0.25	0.40	0.38
mh05	0.2	0	0.35	0.05	0.40	0.29
mh07	0.2	0	0.25*	0.05*	0.32	0.29
mh10	0.2	0	0.4	0.2	0.45	0.35
mh11	0.2	0	0.2*	0.1*	0.28	0.30
mh12	0.2	0.5	0.5	0.3	0.73	0.65
jj01	0.2	0.1	0.5	0.35	0.55	0.46
jj02	0.2	0.1	0.1	0.05	0.24	0.30
jj03	0.3	0.1	0.3	0.2	0.44	0.48
jj05	0.2	0.1	0.15	0.05	0.27	0.30
jj06	0.2	0.1	0.3	0.25	0.37	0.39
jj07	0.2	0.1	0.3	0.2	0.37	0.36
jj08	0.2	0.05	0.1*	0.05*	0.23	0.29
jj10	0.2	0.05	0.05*	0.02*	0.21	0.29

U_{meas} , U_{gap} and U_{time} ($U_{\text{time_to}}$) are the uncertainties introduced by the measurements, the gaps and the chosen time window. Values indicate the confidence interval on a 95% level.

* Only data after July 2009 was considered for the estimation because of large bias by gaps prior to this date.




Fig. 1. Overview of the Matterhorn field site at Hörnligrat. The circles with numbers indicate the sensor locations. Note the thin snow cover in the Matterhorn east face (left picture taken in November 2009).

Temperature variability in steep alpine rock and ice faces

A. Hasler et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
⏪	⏩
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	

Temperature variability in steep alpine rock and ice faces

A. Hasler et al.




Fig. 2. Overview of the Jungfrauoch field site around the Sphinx observatory. The circles with numbers indicate the sensor locations. All pictures taken in October 2006.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Fig. 3. Close-up of sensors in densely fractured rock at the south side of Sphinx, Jungfrauoch. The picture is taken in April 2007 after a periode with intense irradiation.

**Temperature
variability in steep
alpine rock and ice
faces**

A. Hasler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





Fig. 4. Fractures with large spacing and opening at Matterhorn Hörnligrat (picture from November 2010). The instrument in the center is a crackmeter, not considered in this article.

**Temperature
variability in steep
alpine rock and ice
faces**

A. Hasler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Temperature
variability in steep
alpine rock and ice
faces**

A. Hasler et al.




Fig. 5. Example of gap-filling with original values (red) and the resulting dataset after gap filling (blue). Note that the data in this graph (mh01; T_1) is a worst case concerning gap frequency.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 6. Time series from July 2008 to the end of 2010 of the rock temperature measurements (top) at mh10 with interpolated values in data gaps (grey bars) and corresponding running annual means (bottom) that are represented in the center of the averaging window. The black dots indicate this averaging window for one MAT with the quadrat showing the point in time of its representation. For most sensors this averaging window was chosen to minimize data gaps.

Temperature variability in steep alpine rock and ice faces

A. Hasler et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



Temperature variability in steep alpine rock and ice faces

A. Hasler et al.




Fig. 7. Mean annual temperature (MAT) profiles for clefts (light blue), rock (grey) and ice (dark blue) with depth z measured perpendicular to the surface. Solid symbols are shaded locations (north); hollow symbols are more exposed to solar radiation (south and east). Note that the uppermost MAT of mh01 to mh03 is a rock surface temperature (Table 1).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Temperature variability in steep alpine rock and ice faces

A. Hasler et al.




Fig. 8. Mean annual temperature (MAT) and thermal offset or temperature difference (TO) between cleft top and within cleft (light blue); in shallow rock boreholes (grey) and in ice (dark blue). The black error bars show the uncertainties of the thermal offset estimates on a 95% confidence level. The letters at the top indicate the aspect of the locations (E = east, N = north, S = south). For other attributes see (Table 1).

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion







Fig. 9. Time series of rock temperatures at jj05 ($T_0 = 0^\circ\text{C}$) measured every 10' (top) and the temperature difference $\Delta T = T_4 - T_1$ averaged over 30 days (bottom). Additionally the running MATs are plotted in lines similar to Fig. 6 (bottom).

Temperature variability in steep alpine rock and ice faces

A. Hasler et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



Temperature variability in steep alpine rock and ice faces

A. Hasler et al.




Fig. 10. Time series of mh01 (left) and jj01 (right) with measured temperatures (top) and the temperature difference ΔT averaged over 30 days as well as MATs (bottom).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion