

# Stable isotope and gas properties of two ice wedges from Cape Mamontov Klyk, Laptev Sea, Northern Siberia

T. Boereboom<sup>1</sup>, D. Samyn<sup>1,\*</sup>, H. Meyer<sup>2</sup> and J-L. Tison<sup>1</sup>

[1]{Laboratoire de Glaciologie, Université Libre de Bruxelles, Belgium}

[2]{Alfred Wegener Institute for Polar and Marine Research, Research Unit Potsdam, Potsdam, Germany}

[\*]{now at: Dpt of Mechanical Engineering, Nagaoka University of Technology, Nagaoka, Japan}

Correspondence to: T. Boereboom (thierry.boereboom@ulb.ac.be)

## Abstract

This paper presents and discusses the texture, fabric, water stable isotopes ( $\delta^{18}\text{O}$ ,  $\delta\text{D}$ ) and gas properties (total gas content,  $\text{O}_2$ ,  $\text{N}_2$ ,  $\text{Ar}$ ,  $\text{CO}_2$ , and  $\text{CH}_4$  mixing ratios) of two climatically contrasted (Holocene vs. Pleistocene) ice wedges (IW-26 and IW-28) from Cape Mamontov Klyk, Laptev Sea, in Northern Siberia. The two ice wedges display contrasting structures: one being of relatively “clean” ice and the other showing clean ice at its centre as well as debris-rich ice on both sides (referred to as “ice-sand wedge”). Our multiparametric approach allows discriminating between three different ice facies with specific signatures, suggesting different climatic and environmental conditions of formation and various intensities and nature of biological activity. More specifically, crystallography, total gas content and gas composition reveal variable levels of meltwater infiltration and contrasting contributions from anaerobic and aerobic conditions to the biological signatures. Stable isotope data are shown to be valid for discussing changes in paleoenvironmental conditions and in the temporal variation of the different moisture sources for the snow feeding into the ice wedges infillings. Our data set also give further support to the previous assumption that the ice wedge IW-28 was formed in Pleistocene and the ice wedge IW-26 in Holocene times. This study sheds more light on the conditions of ice wedge growth under changing environmental conditions.

## 1 Introduction

Ice wedges are a common geomorphological feature of permafrost areas in the Northern Hemisphere and widespread in Alaska, Canada and Siberia. The analysis of ice wedges properties is therefore commonly used to provide proxy-data for paleoclimatic and paleoenvironmental reconstructions. First, the isotopic signature ( $\delta^{18}\text{O}$  or  $\delta\text{D}$ ) of the ice is thought to reflect the isotopic characteristics of the winter precipitation which are a function of the temperature at the site (Dereviagin et al., 2002; Lauriol et al., 1995; Meyer et al., 2002a, 2002b; Raffi et al., 2004; Vaikmäe, 1989; Vasil'chuk and Vasil'chuk, 1998). As an example, recent ice wedges in Northeast Siberia show  $\delta^{18}\text{O}/\delta\text{D}$  values between -25‰/-190‰ and -20‰/-140‰ (Opel et al., 2011), while younger Holocene ice wedges values range between -26‰/-195‰ and -22‰/-175‰ and older Holocene ice wedges are -26‰/-202‰ to -24‰/-190‰ (Wetterich et al., 2008). The Pleistocene isotopic signature in the same area clearly differs by decreases of approximately 5‰ and 35‰ in  $\delta^{18}\text{O}$  and  $\delta\text{D}$ , respectively (Popp et al., 2006) showing  $\delta^{18}\text{O}/\delta\text{D}$  values between -33‰/-250‰ and -31‰/-245‰ (Wetterich et al., 2008) with minima from the Last Glacial Maximum (-35‰/-290‰) as documented in a recent overview from Wetterich et al. (2011).

Co-isotopic approaches, use the deuterium excess ( $d$ ) as a tracer for changes in the atmospheric source. For example, Meyer et al. (2010b), who identified the Younger Dryas cold event in Northern Alaskan ice wedges, also brought to the light a rapid reorganization of the Arctic atmospheric circulation following the opening of the Bering Strait, with  $d$ -transitions from about 6 ‰ to about 9 ‰, as also observed in the North GRIP deep ice core.

Other ice wedge properties can be useful to understand and reconstruct the environmental condition prevailing during ice wedge genesis. St-Jean et al. (2011) recently used ice crystallography, bubble content, gas composition and stable O-H isotopes of ground ice to identify contrasting genetic processes between ice wedges formed during the Late Pleistocene period under cold and dry climatic conditions and ice wedges formed during the Holocene in warmer and wetter conditions. Ice wedges formation primarily results from cyclic frost-cracking and crack infilling by ice (e.g. Mackay, 1974). The cracking results from the thermal contraction of the soil the winter (Lachenbruch, 1962) and the origin of the ice infilling can be attributed to several sources (or mixture of sources) such as: a) unmodified winter snow, b) hoarfrost accretion during winter and c) snow melt infiltration in spring, subsequently refreezing in the crack to form “ice veins” (e.g. Lauriol et al., 1995; Opel et al., 2011; St-Jean

et al., 2011). In their work, St-Jean et al. (2011) contrast a Pleistocene ice wedge in which the ice filling rather results from dry snow compaction or hoarfrost accretion and an Holocene ice wedge in which liquid water infiltration and refreezing dominates.

Ice wedges incorporate variable amounts of sediment depending on the sediment, snow and/or water availability at a given time. Frost cracks can be completely filled by blown sediments in very dry conditions forming a “sand wedge” (e.g. Berg and Black, 1966) or can be filled-in by successive ice veins and sediments layers as described in Meyer et al. (2002a) or in Romanovsky (1976). This latter type of structure will be here referred to as an “ice-sand wedge” (also termed “composite wedge”).

In this paper, we propose a multi-parametric approach including stable isotope composition, total gas content, gas composition, as well as textures and fabrics to study two specific ice wedges from the Laptev Sea region. These two ice wedges were sampled from a cliff in the framework of the project “Process studies of permafrost dynamics in the Laptev Sea”, which aimed to address the paleoenvironmental history of the Cape Mamontov Klyk area (Bobrov et al., 2009; Müller et al., 2009; Schirrmeister et al., 2008). These specific ice wedges were chosen for this study due to their contrasting ages (Holocene vs. Pleistocene) as indicated by their stable isotope signature and stratigraphic position in a dated regional sequence. The aim of the study is two-fold: a) to improve our understanding of the imprint of the contrasted Holocene/Pleistocene climatic and environmental conditions on the ice wedge properties and infilling processes and b) to interpret the specific deuterium excess signature of the two ice wedges in terms of potential changes in atmospheric moisture sources.

## **2 Study area and ice wedge description**

The two studied ice wedges are part of a coastal bluff on the Laptev Sea coast, about 300 km west from the Lena Delta (73°36' N; 117°10' E, Fig. 1). This area belongs to the subarctic tundra in a region of continuous permafrost, with a thickness of 400-600 m (Yershov, 1989). The active layer shows a maximum thaw depth of 0.2-0.5 m in July. The area is dominated by a continental arctic climate with long severe winters and short cold summers.

The cliff from which the ice wedges originate is subdivided into four different sedimentological units (A to D, from old to young; Fig. 2a) and described in detail by Schirrmeister et al. (2008). From the bottom of the cliff to its top, we distinguish a lower sand unit (unit A) that consists of yellowish-grey, irregularly-laminated fine-grained sand, lacking

1 visible plant remains. Grass roots in the uppermost horizon of this unit were dated to 44.5 ky  
2 (Schirrmeister et al., 2008). Above, unit B consists of an alternation of four cryoturbated peat-  
3 rich horizons and irregularly-laminated, dark-grey silty to fine-grained sandy interbeds. Unit  
4 C is known as “Ice Complex”, a type of ice-rich permafrost deposit of Late Pleistocene age  
5 widespread in Arctic Siberia attributed at this site to the Sartan period (Schirrmeister et al.,  
6 2008). The Ice Complex deposits are composed of alternating mineral-rich (greyish) and  
7 organic-rich (brownish) sediment layers. The upper unit (unit D) consist of peat soils of  
8 Holocene age originating from fillings of small thermokarst or polygonal ponds developed on  
9 top of the Ice Complex or from Holocene thermoerosional or fluvial deposits.

10 Ice wedge 26 (IW-26, Fig. 2b) was sampled within unit C in the upper part of the cliff, at 18.6  
11 m above sea level, and is therefore embedded within late Pleistocene Ice Complex sediments  
12 (Schirrmeister et al., 2008). IW-26 is about 1.6 m in width and about 2.5 m in height are  
13 visible. Sampling was performed across the full ice-wedge width, approximately 1.2 m below  
14 the surface. The ice samples consisted of foliated grey to white, and in parts yellowish, ice  
15 with small (millimetric) gas bubbles. The wedge ice shows a low content of mineral particles  
16 and organic matter (fine-disperse individual particles or thin layers following the foliation).

17 The second ice wedge studied in this paper, IW-28 (Fig. 2c), has been attributed to unit A in  
18 the lower part of the cliff and was sampled at 1 m above sea level. It developed in sandy  
19 sediments. IW-28 is up to 5 m wide in its upper part at the top of the cliff. The ice wedge is  
20 laterally associated with an ice-sand wedge (ISW-28), a portion of which was also sampled  
21 (Fig. 2c, left of sampling box). In the central section, the wedge ice has a white and milky  
22 appearance, and small millimetric gas bubbles are frequent (Fig. 3d). The mineral content and  
23 the organic matter contents are both low, with particles dispersion similar to that in IW-26 and  
24 foliation layers that are 1-3 mm wide. For the ice-sand wedge portion, the ice is mixed with a  
25 significant amount of fine-grained sediments alternating with clean ice layers.

### 27 **3 Sampling and analytical methods**

28 The ice wedges were sampled as ice blocks using a chain saw, in horizontal transects covering  
29 the entire ice-wedge width and partly including the ice-sand wedge section from now on  
30 referred to as ISW-28 (Fig. 2c). Samples used in this paper for fabric, texture, gas and stable  
31 isotope analyses are derived from horizontal sections ( $\pm 1.5$  cm thickness) in the top part of the  
32 blocks extracted from the ice wedge.

Horizontal thin sections were prepared using a traditional biological microtome (Leitz 1400) for the clean ice and a diamond-wire saw (Well 6234) for the debris-laden ice, following the standard procedures from Langway (1958) and Tison (1994), respectively.

Crystal-size determination was performed using two different techniques. The first was the mean linear intercept method developed by Pickering (1976). The number of grain boundaries ( $\bar{N}$ ) crossed by a random linear traverse of length ( $\bar{L}$ ) across the thin section was averaged over many traverses. The mean grain diameter ( $\bar{d}$ ) was estimated using the following equation:  $\bar{d} = \bar{N}/\bar{L}$ . The second method, proposed by Jacka (1984), estimates the mean diameter by counting the number of entire crystals in a known area and assuming a circular cross-section following the equation  $\bar{d} = \sqrt{4A/\pi N}$ , where  $A$  is the area of thin section studied and  $N$  is the crystal count.

Stable water isotopes were measured at high resolution (approximately every centimeter) with a Finnigan MAT Delta-S mass spectrometer at the Alfred Wegener Institute, Research Unit Potsdam using equilibration techniques. Hydrogen and oxygen isotope ratios are presented as per mil difference relative to V-SMOW (‰, Vienna Standard Mean Ocean Water), with internal  $1\sigma$  errors better than 0.8‰ and 0.1‰ for  $\delta D$  and  $\delta^{18}O$ , respectively (Meyer et al., 2000).

Gas inclusions entrapped in the ice were analysed for their total gas volume (27 samples) and gas composition ( $CO_2$ ,  $O_2$ ,  $N_2$ , Ar, and  $CH_4$ ). Gas composition ( $CH_4$  = 24 samples, Ar = 7 samples, other gases = 30 samples) was measured by gas chromatography (Interscience Trace GC) using an FID detector for  $CO_2$  and  $CH_4$  and a TCD detector for  $O_2$ , Ar and  $N_2$ . Between 20 to 35 grams of sample were used for each measurement. For  $CO_2$ ,  $O_2$ , Ar and  $N_2$ , we used the dry-extraction technique described in Raynaud et al. (1982) and Barnola et al. (1983). For  $CH_4$ , we used the melting-refreezing procedure described in Raynaud et al. (1988) and Blunier et al. (1993). The levels of precision of the measurements are 2.5% for  $CO_2$ , 0.4% for  $O_2$  and  $N_2$ , 2% for Ar and 3% for  $CH_4$ . The total gas content (27 samples) was determined using a Toepler pump and a melting-refreezing extraction technique described in Martinerie et al. (1994). The precision of the measurements is  $\leq 5\%$ .

The limited amounts of ice provided for the gases and fabrics analyses did not yield enough material to perform a full granulometric analysis of the sediment inclusions, which is nevertheless beyond the scope of this paper. At a later stage of the interpretation, however, the

residual sediments collected after the gas analyses have been treated by HCl (1 mol.l<sup>-1</sup>) and H<sub>2</sub>O<sub>2</sub> (30%) for qualitative detection of calcium carbonate or organic matter, respectively.

## **4 Results**

### **4.1. Stable <sup>18</sup>O and <sup>2</sup>H isotopes**

Ice wedge IW-26 shows  $\delta^{18}\text{O}$  values ranging between -22.6‰ and -25.8‰ and  $\delta\text{D}$  values between -170‰ and -191‰, respectively (Fig. 4a). The IW-26 samples plot on a slope of 6.63 in the D-<sup>18</sup>O diagram (Fig. 4c, grey squares). The isotopic composition of ice wedge IW-28 is also shown in Fig. 4a and a clear distinction has to be made between the two different facies of the ice wedge: the ice-sand wedge (ISW-28) and the ice wedge itself (IW-28). These differ both in terms of deuterium-excess (Fig. 4b) and co-isotopic relationship (Fig. 4c). The ice-sand wedge ISW-28 displays  $\delta^{18}\text{O}$  values between -29‰ and -30.9‰ and  $\delta\text{D}$  values between -230‰ and -246‰. The samples show a low variability in  $d$  ( $1\sigma = \pm 0.6\text{‰}$ ) with a mean value of 1.4‰, and they are aligned on a slope of 8.03 in the co-isotopic plot. For the ice wedge part IW-28,  $\delta^{18}\text{O}$  and  $\delta\text{D}$  range between -29.4‰ and -31.9‰, and between -229‰ and -247‰, respectively. The  $d$  also shows a low variability ( $1\sigma = \pm 0.7\text{‰}$ ) with a mean value of 7.5‰ and a slope of 7.44 is measured in the co-isotopic diagram. The transition has  $\delta^{18}\text{O}$  values between -29.5‰ and -29.9‰, a  $\delta\text{D}$  between -223‰ and -237‰, and shows variable  $d$  values intermediate between the other two parts (Fig. 4c).

### **4.2. Ice texture and fabrics**

Figure 3 shows thin section photographs that are typical of the three main facies encountered in the two studied ice wedges. In IW-26, the texture is homogeneous, lacking any significant elongation (Fig. 3a). The crystal diameter is ranging between 0.40 and 0.60 cm (10 to 30 mm<sup>2</sup>, Fig. 5) and c-axis orientations are concentrated in the horizontal plane with preferred orientation perpendicular to the foliation azimuth (Fig. 6a). The ISW-28 texture shows, within a matrix of relatively large equigranular grains, several monocrystalline layers of elongated crystals with a well-developed “ribbon structure”, and crystal growth being geometrically associated to the fine debris layers that surround them, oriented parallel to the (subvertical) foliation plane (Fig. 4b). On Fig. 5, crystal diameter is ranging between 0.2 and 0.4 cm (3 to 10 mm<sup>2</sup>). In IW-28, the central part of the ice wedge shows an equigranular texture (Fig. 4c),

with crystals smaller than in ISW-28 (0.1 to 0.2 cm - 1-3 mm<sup>2</sup>, Fig. 5). Both ISW-28 (Fig. 6b) and IW-28 (Fig. 6c) show c-axes orientation patterns similar to IW-26, although the preferred orientation perpendicular to the foliation azimuth is less obvious for ISW-28 given the limited number of observations (due to larger grains).

### 4.3. Gas properties

The contrast between the three facies (IW-26, ISW-28 and IW-28) is also evident from their gas properties (Fig. 7). Table 1 summarizes minimum, mean and maximum values observed for the three facies. Total gas contents in our ice wedges are lower than in ice resulting from simple snow compaction as in ice sheets (see ranges in Table 1) and the total gas content of IW-28 is higher than those of ISW-28 and IW-26. For CO<sub>2</sub>, all ice wedge mixing ratios are clearly higher than the atmospheric values, the highest values being observed in IW-26 (mean = 62 000 ppmV) and ISW-28 showing lower values (mean = 3 000 ppmV) than IW-28 (mean = 25 000 ppmV). On the contrary, CH<sub>4</sub> mixing ratios show their lowest values in IW-26 with values in the range of atmospheric mixing ratio (mean = 1 ppmV), intermediate values for IW-28 (mean = 8 ppmV) and the highest values in ISW-28 (mean = 55 ppmV). Oxygen shows consistently lower values than the atmosphere (of the order of 10%) and nitrogen is slightly higher, balancing other constituents.

### 4.4. Sediments properties

Residual sediments collected from four ice samples have been treated with HCl and H<sub>2</sub>O<sub>2</sub> (2 samples from ISW-28 and 2 samples from IW-28). The ISW-28 samples have strongly reacted with H<sub>2</sub>O<sub>2</sub> while the IW-28 samples only showed a very limited reaction. We therefore argue that there is organic matter in the ISW-28 sediment inclusions, but only limited amounts in IW-28. On the other hand, the HCl treatment did not show detectable bubbling activity under the binocular for any of the 4 sediment samples treated, supporting that carbonate contents are negligible within the sediment enclosed in our ice wedges.

## 5 Discussion

In this section, we will first summarize and interpret ice texture and fabrics, co-isotopic signature and gas properties in terms of the infilling process of the studied ice wedges (see

also Table 1). We will then attempt to link these processes to the regional paleo-climatic and paleo-environmental setting of the study area.

### **5.1. Ice wedge infilling processes**

#### **IW-26**

IW-26 is characterized by relatively large grains (Table 1), suggesting either older crystals or conditions favouring recrystallization at higher temperature. The equigranular texture implies no preferred direction of growth (as would be the case for the freezing of a water reservoir) and no significant deformation process. The horizontal girdle of the c-axes with a localized maximum could be explained in several ways: a) gravity settling of individual snow grain in a narrow vertical crack; b) preferential growth along c-axis (better conduction) in a horizontal temperature gradient (perpendicular to crack sides); c) effect of the deformation in wedge during cyclic contractions or d) a combined version between these processes. The spherical bubbles (as opposed to tubular shapes expected in the case of water freezing) also imply no directional freezing, comforting a snow-water mixture hypothesis rather than water infilling for the ice wedge genesis. The total gas content is low in comparison with a dry firn densification process (between one tenth and one third of the value of a dry firnification process) suggesting a liquid water contribution to the ice formation or a process expulsing the gases, eventually along the frost crack if the latter is only partially filled or if the infilling is still permeable.

CO<sub>2</sub> mixing ratios in IW-26 are well above the present-day values (Fig. 7, row 2, left and Table 1). This indicates that the ice did not form through simple low temperature pressure-driven dry firnification process of snow enclosing the atmospheric gas composition. Previous studies of ice forming close to a bedrock environment have shown that this is very often the case, mainly because of either the presence of liquid water and/or significant biological activity within the enclosed organic matter fraction (Souchez et al., 1995a). Theoretically, the high CO<sub>2</sub> solubility in water could raise the CO<sub>2</sub> mixing ratio in the dissolved phase up to about 20 000 ppmV (at 0°C and 1 atmosphere, with limited effect of pressure and impurities). Freezing of a CO<sub>2</sub> saturated freshwater reservoir in a closed system could therefore display such high CO<sub>2</sub> values. There are however a number of other potential ways to further increase the CO<sub>2</sub> mixing ratio in our ice wedge ice such as diffusion-driven gas fractionation at the freezing front, CO<sub>2</sub> degassing on calcium carbonate precipitation, thermal trapping or biological respiration. Rejection of impurities at the interface by ice growing in a water



1 reservoir will set up a strong diffusion gradient in the adjacent liquid boundary layer. This  
2 could lead to a further increase of the mixing ratio of the least diffusive gas in the solution, as  
3 for instance CO<sub>2</sub> (Killawee et al., 1998). However, theoretically, this effect cannot increase  
4 the CO<sub>2</sub> concentration by a factor higher than the ratio of the CO<sub>2</sub> diffusion coefficient  
5 (DCO<sub>2</sub>) to the air diffusion coefficient (D<sub>air</sub>) in water; i.e. by a factor of about 1.6 (using  
6 typical values for DCO<sub>2</sub>, DN<sub>2</sub>, DO<sub>2</sub>). This would increase the CO<sub>2</sub> mixing ratio to about 30  
7 000 ppmV, provided that the freezing process can be assimilated in this case to ice growing in  
8 a liquid interface. Killawee et al. (1998) have shown that unidirectional freezing of a  
9 supersaturated CaCO<sub>3</sub> solution leads to a strong CO<sub>2</sub> enrichment of the refrozen water due to  
10 the CO<sub>2</sub> degassing resulting from the carbonate precipitation. The lack of reaction of our ice  
11 wedge sediments to an HCl “attack” however does not support the existence of significant  
12 amounts of CaCO<sub>3</sub>. Finally, biological respiration is the most likely active process since  
13 bacteria and microbial communities have been identified in permafrost ice and in ice wedges  
14 (Gilichinsky et al., 2007; Katayama et al., 2007; Steven et al., 2008).

15 The considerations above suggest that the CO<sub>2</sub> mixing ratios of IW-26 result from a  
16 combination of purely physical processes (solubility) and biological respiration processes.  
17 Isotopic measurements of the <sup>13</sup>C of CO<sub>2</sub> or of the <sup>18</sup>O of the O<sub>2</sub> in our samples would further  
18 support this assessment (Cardyn et al., 2007; Souchez et al., 2006), but these data are  
19 currently not available. Interestingly though, the range of CO<sub>2</sub> values in the IW-26 ice wedge  
20 is similar to that observed in the basal ice of the GRIP ice core (up to 130 000 ppmV), where  
21 δ<sup>18</sup>O<sub>air</sub> values as low as -39‰ clearly indicate the occurrence of bacterial respiration processes  
22 (Souchez et al., 2006). Tison et al. (1998) and Souchez et al. (1995b), among others,  
23 demonstrated that the GRIP basal ice developed under climatically mild pre-ice sheet  
24 periglacial conditions. The latter are thus potentially similar to those conditions existing  
25 during the formation of our ice wedges. Another approach would be to test for the  
26 conservation of the sum of O<sub>2</sub> and CO<sub>2</sub> concentration, given the stoichiometry of the  
27 respiration reaction, as suggested by Souchez et al. (1995a). However this approach is not  
28 valid in our case since a liquid water phase is present allowing most of the carbon dioxide  
29 produced by the respiration process to dissolve as HCO<sub>3</sub><sup>-</sup> into the solution, at least at normal  
30 pH levels (Zeebe and Wolf-Gladrow, 2001).

31 CH<sub>4</sub> mixing ratios are rather low (Fig. 7, row 3, left - Table 1), in the range of Pleistocene –  
32 present day atmospheric values and we therefore assume no significant contribution from a  
33 substrate source or bacterial production.

The  $\delta^{18}\text{O}/\delta\text{D}$  values in IW-26 are in the range of Late glacial to Holocene ice wedges (Fig. 6b in Wetterich et al., 2011). The slope (6.63) of the linear regression line in the co-isotopic diagram of Fig. 3c is however quite different from the known values of the Local Meteoric Water line in the region (slope of 7.6 at Tiksi Station, H. Meyer, unpublished data). Paleoclimatic interpretation of the stable isotopes signature in ice wedges is usually challenged by the possibility that water phase changes during evaporation/sublimation or melting/refreezing may affect the isotopic composition through fractionation, and therefore potentially alter the original paleoclimatic signature of the ice wedge. Refreezing processes will shift the  $\delta$  values towards less negative values (e.g., max. +3‰ for  $\delta^{18}\text{O}$  and +21‰ for  $\delta\text{D}$  in a single refreezing process) and will also decrease the slope of the  $\delta^{18}\text{O}$  -  $\delta\text{D}$  co-isotopic relation (Souchez and Lorrain 1991). This is described by the “freezing slope”, which is expressed in the more general case of open systems as:

$$S = \frac{\alpha-1}{\beta-1} * \frac{1000+\delta_i\text{D}}{1000+\delta_i^{18}\text{O}} \quad (1)$$

where  $\alpha = 1.0208$  for deuterium

$$\beta = 1.003 \text{ for } ^{18}\text{O}$$

$\delta_i$  = isotopic composition of the initial liquid

This concept has been used by St. Jean et al. (2011) to associate lower co-isotopic slopes to melting-refreezing processes in Holocene ice wedges. It is however important to underline that a “freezing slope” as such will only develop if the scale of the freezing process (size of the water reservoir) is considerably larger than the resolution of the isotopic sampling. In the case of ice wedges, typical crack-widths are of the order of a few centimeters (Harris et al., 1988), and annual “ice veins” are generally observed to be between 1 mm and 1 cm in width. It is therefore quite unlikely to develop a “freezing slope” from a several meters wide ice wedge, using a stable isotopes data set at a resolution of a few centimeters (about 1 cm in our case). This is probably also one of the reasons why  $\delta^{18}\text{O}/\delta\text{D}$  signatures of ice wedges might have paleoclimatic significance. Small-scale refreezing or evaporation/sublimation processes, if present, are however likely candidates to induce localized small-scale variability towards lower slopes, which might explain the observed lower co-isotopic slope and the lowering of the correlation coefficient ( $r^2 = 0.94$ ) observed in the case of IW-26.

To summarize, our IW-26 analyses enable us to conclude that ice-vein formation in this ice wedge results from refreezing of a water-snow mix, with a large proportion of snow, no

directional freezing, although a slight influence of the horizontal temperature gradient on the depositional c-axes orientations might exist. The co-isotopic slope is probably affected by small scale freezing dispersion of the samples. Relatively low gas content and a mixed signature of solubility and respiration processes are also demonstrated for CO<sub>2</sub>. These properties are consistent with an ice wedge developing under warm and wet Holocene conditions.

## **ISW-28**

ISW-28 shows medium grains size suggesting less recrystallization and/or younger age than IW-26 (which is actually precluded by the relative stratigraphical position of the two ice wedges, as discussed before). The alternation of equigranular and rectangular crystals (ribbon layer) elongated parallel to the crack side is the main characteristic of the ice-sand wedge part. This type of structure has, to our best knowledge, never been described in ice wedge and resembles the “stratified facies” found in basal glacier ice (e.g. Tsijiore Nouve, Switzerland) or ice resulting from film water refreezing on subglacial cavity floors. In our case we assume interface freezing on the cold surface of the frost crack walls of a thin running water film originating from summer melting of the surface lips of the partially open (unfilled) crack (as opposed to a “top-down” or “side-center” refreezing of an “in situ” bulk water reservoir). Sediments at layer boundaries suggest entrainment with the melt water from the surface, rejection from the growing ice and, finally, freeze-in during “shut-down” phases of the water supply. The horizontal girdle drawn by the c-axes distribution does not show a localized maximum, but, as mentioned before, the number of sample is relatively low. The observed high small-scale variability of total gas content is probably due to the alternation of layer with bubbles (episodes of wet snow metamorphism) and layers without (ribbon layer formed through water film freezing). CO<sub>2</sub> is higher than atmospheric values (solubility effect) but well below solubility limit of 20 000 ppm, therefore suggesting an “open system” freezing with gas rejection to atmosphere. CH<sub>4</sub> mixing ratios are much higher than the atmospheric value, suggesting methane production in bubbles or in soil, in close vicinity to the bubbles before their occlusion in the ice vein. Note that the sediments in the ice-sand wedge part are rich in organic matter and provide an adequate substrate for methanogenic bacteria.

Methane production implies an anaerobic environment for the methanogenic archaea to work efficiently (e.g. Wagner and Liebner, 2009). Although ice wedges can be a source of methane (e.g. Meyer et al., 2010a), the range of observed methane mixing ratios seems to be negligible compared to other methane sources studied for the carbon budgets in permafrost area (IPCC,

2007). Apart from a few values close to atmospheric in ISW-28, the O<sub>2</sub> mixing ratio fluctuates around 10% (Fig. 7, row 4), being only half the atmospheric O<sub>2</sub> value. It is hard to tell if this value is the gaseous composition of the local environment (in the crack into the frozen soil) before the bubbles “close-off” or if this value is a result of biogeochemical post-occlusion processes. Whichever way, it does not imply a fully anaerobic environment. However, for O<sub>2</sub> measurements only ±35 g of samples were taken, each containing a large number of millimetric bubbles. As already suggested for the GRIP basal ice (where both high CO<sub>2</sub> and high CH<sub>4</sub> values were measured in the same samples) and in other environments like soils and marine sediments (Souchez et al., 1995b), the mean O<sub>2</sub> concentration might well reflect a situation where some bubbles are depleted in oxygen and others have concentrations close to atmospheric levels. The bubbles can therefore provide contrasted closed system micro-environments in some cases favourable to aerobic respiration (high CO<sub>2</sub>, low CH<sub>4</sub>) and in others to bacterially mediated methanogenesis (high CH<sub>4</sub>, low CO<sub>2</sub>).

The δ<sup>18</sup>O and δD values match well with Pleistocene stadial signature in the studied area (Fig. 6c in Wetterich et al., 2011). The slope in the co-isotopic diagram (8.03) could be indicative for the variability of the autumn – winter snow input to the crack during the integrated period of the ice-sand wedge history. Snow samples in this region do not receive additional secondary moisture (because open water bodies are frozen in winter) and therefore are generally situated close to the GMWL (Meyer et al., 2002a, 2002b) with slopes in the co-isotopic diagram close to 8.

ISW-28 ice veins properties suggest a mixed origin altering episodes of consolidation of a “water-snow” mix (granular) and of events of running water film refreezing at the sides of the open frost crack. The fact that the crack is not always filled with snow might indicate periods favouring debris incorporation either through an increased input of particulate contribution from the surface or because of a thinner snow cover (lower precipitations). It is worth noting that a globally colder-drier period (as suggested by the low precipitation) does not necessarily mean less melting in the period when frost cracks are open, if debris availability allows strong summer albedo effects. The biological activity in ISW-28 is characterized by low to moderate aerobic respiration but high methanogenic activity related to availability of organic matter and lower than atmospheric oxygen levels. These conditions obviously existed during the early stages of IW-28 development (ISW-28).

## **IW-28**

1 IW-28 shows the smallest grain size. This is potentially the combined result of a lower  
2 Arrhenius growth rate at lower temperature and less liquid water availability (which would  
3 otherwise enhance crystal growth rate). Clearly these factors took over the “time factor”  
4 (older ice) in the recrystallization process. The equigranular ice does not show preferred  
5 direction of freezing and the horizontal girdle shows one localized maximum sub-  
6 perpendicular to the foliation, as for the IW-26. As in IW-26 also, the spherical bubbles  
7 indicate no directional freezing and favour a consolidation process of a snow-water mixture  
8 for the origin of the ice veins. However, the highest total gas content with values reaching  
9 about 50% of those observed in dry firnification), suggest less liquid water contribution than  
10 in IW-26. This might be linked to the small-scale hydrodynamical processes at play as the  
11 meltwater percolates into the snow infilling. Colder temperatures in the surface permafrost  
12 from the previous winter would favour rapid freezing in the surface layer and limit water  
13 percolation further down-crack. Detailed vertical gas content profiles would certainly be  
14 useful in documenting these processes further.

15 CO<sub>2</sub> mixing ratios contrast with those observed in the ISW-28 section and are closer to those  
16 observed in ice wedge IW-26, showing the potential combination of pure physical solubility  
17 processes with biological respiration, although the latter is less developed than in IW-26. CH<sub>4</sub>  
18 mixing ratio, largely higher than the atmospheric values, are also intermediate between those  
19 of IW-26 and ISW-28.  $\delta^{18}\text{O}$  and  $\delta\text{D}$  values are similarly low as compared to ISW-28 and  
20 might suggest a slightly colder period for IW-28.

21 IW-28 should therefore be formed from a water-snow mix, at relatively low temperature than  
22 IW-26, with a large proportion of snow, no directional freezing and less sediments input than  
23 in ISW-28. The latter results in moderate CH<sub>4</sub> mixing ratios, although these are higher than  
24 those measured in the Holocene ice wedge IW-26. Fabrics are still dominated by horizontal c-  
25 axes including a single maximum roughly perpendicular to the crack sides, suggesting gravity  
26 settling control or contraction deformation effects rather than recrystallization growth, given  
27 the isotropy of the bubbles and crystal shapes.

28 Our interpretation of ice wedge filling process and ice vein formation contrasts with St-Jean et  
29 al. (2011) which suggest that the Holocene ice wedges result from the freezing of liquid water  
30 and that the Pleistocene ice wedges result rather from snow densification and/or hoarfrost  
31 accretion. Our study suggests variable water-snow mix as source of the ice with higher water  
32 contribution during the Holocene period. Coherently with Lacelle et al. (2011) and St-Jean et

al. (2011) data, however, our  $\delta(\text{N}_2/\text{Ar})$  and  $\delta(\text{O}_2/\text{Ar})$  results (Fig. 8) indicate an increasing influence of biological processes with the increasing amount of water in the water-snow mix. Interestingly, the proximity of the debris source could result in a larger proportion of melting as compared to the snow matrix, and enhanced methanogenesis even though the climate is colder. The latter is supported by a study of Meyer et al. (2010a) in Northern Alaska, where an ice wedge component formed during the colder Younger Dryas displays higher methane concentrations as compared to an ice wedge section formed during the warmer Allerød. Clearly, environmental, physical and biological controls need to be considered when discussing ice wedge properties.

## 5.2. Links to the regional paleoclimatic interpretation

Results from texture, fabric and gas analyses allow to differentiate between three facies within the two studied ice wedges (IW-26, ISW-28 and IW-28), which can also be discussed in terms of the regional paleoclimatic interpretation from available stable isotopes data.

The IW-26 displays least negative  $\delta\text{D}$  and  $\delta^{18}\text{O}$  values, largest crystal sizes, lowest total gas contents and is embedded in the Ice Complex structure of Pleistocene age. However, its isotopic composition is coherent with Holocene age ice wedges ( $\delta^{18}\text{O}$  around -24 to -25‰; Meyer et al., 2000, 2002a, 2002b; Dereviagin et al., 2002; Schirrmeister et al., 2002; Popp et al., 2006; Meyer et al., 2010b). A period of mild temperatures would also increase the rate of biological respiration processes and the availability of organic matter.

The lower isotopic composition of IW-28 suggests a Pleistocene origin (compare Fig. 6b of the recent regional overview in Wetterich et al. (2011)). The situation is however certainly more complex for the ISW-28/IW-28 case as suggested by the strong contrast in their  $d$  values, while showing in both cases low internal variability (mean  $d$ = 1.4‰ for ISW-28 and mean  $d$ =7.5‰ for IW-28) for similar  $\delta\text{D}$  ranges (Fig. 4). Also worth noting is the sharp monotonic transition in  $d$  values at the interface between ISW-28 and IW-28.

There is a number of ways to explain low  $d$  in precipitations (and in the ice that evolves from it), that either belong to the “water cycle processes” or to the “*in-situ* post-deposition processes”. Extensive work has been done dealing with the concept and use of  $d$  in co-isotopic studies, especially in the field of deep ice-core interpretation and the relation to temperature changes at precipitation sites and oceanic sources (Craig, 1961; Dansgaard, 1964; Merlivat and Jouzel, 1979; Yurtsever and Gat, 1981; Jouzel et al., 1982, 2005a, 2005b; Jouzel

and Merlivat, 1984; Johnsen et al., 1989; Masson-Delmotte et al., 2005; Stenni et al., 2001, 2004; Vimeux et al., 2002, 1999). It is commonly admitted that, although humidity and wind speed at the source also play a role, in most cases, the  $\delta^{18}\text{O}_{\text{ice}}$  and  $\delta\text{D}_{\text{ice}}$  primarily depends on the temperature at the site ( $T_{\text{site}}$ ) and to a lesser extent on the temperature at the source ( $T_{\text{source}}$ ). The reverse is true for the  $d$  in the ice. The latter is indeed mainly driven by kinetic isotopic effects during evaporation at the oceanic source, and these will be enhanced for higher temperatures and lower relative humidity, increasing the  $d$  value (Masson-Delmotte et al., 2005; Merlivat and Jouzel, 1979). It should however be kept in mind that at equivalent source temperatures, lower winds would also increase the relative humidity at the source and therefore decrease the  $d$ -excess. As shown in the deep ice core records from Greenland, large topographic and atmospheric circulation changes during the last glacial period (the presence of ice sheets of variable size, larger but fluctuating extension of the sea ice) are responsible for an antiphase relationship between  $\delta$  values ( $T_{\text{site}}$ ) and  $d$  values ( $T_{\text{source}}$ ). Indeed, during mild events, when Greenland was warmer (less negative  $\delta$ 's) and sea ice less extent, the moisture source was closer (colder) and therefore the  $d$  lower than during more severe phases. This results in the samples from these milder periods (less negative  $\delta\text{D}$  and  $\delta^{18}\text{O}$ ) aligning, in a  $\delta\text{D}$ - $\delta^{18}\text{O}$  diagram, on a line with slope close to 8, but with the  $d$ -intercept at +4‰ instead of +8‰ (e.g. Fig 7 in Johnsen et al., 1989).

Today, the main source of the winter precipitation in our study area (west of  $\sim 140^\circ\text{E}$ ) is probably located in the Northern Atlantic (Kuznetsova, 1998; Rinke et al., 1999). However, as discussed by Meyer et al. (2002b) and Wetterich et al. (2011), Pleistocene moisture sources for the same area less straightforward to locate. Two hypotheses were discussed (Table 2): (1) refers to a permanent Atlantic source, with a latitudinal variation similar to that suggested by Johnsen et al. (1989), as described above for Greenland. In that scenario, a contrasting  $d$  signature would represent a temporal shift (within the Pleistocene, e.g. interstadial/stadial) from a milder (Table 2, col. 2) glacial period (higher  $T_{\text{site}}$ ) with a higher relative humidity (i.e. lower  $T_{\text{source}}$  and/or lower winds, and therefore lower  $d$  values in ISW-28) to a colder glacial period (lower  $T_{\text{site}}$ ) with a lower relative humidity (i.e. higher  $T_{\text{source}}$ , and/or higher winds, and therefore higher  $d$  value in IW-28). The amount of precipitation, however, should be lower during cold stadial periods and higher during interstadials (as for the difference between glacials and interglacials in deep ice cores and in permafrost). Alternatively, (2) the combination of the full development of the Eurasian Ice Sheet (the easternmost extent of which reached the western part of Taymyr Peninsula) and of an extended sea ice cover in the

Barents and Kara seas and the Northern Atlantic (stadial, Table 2, col. 5), might have considerably reduced (deflected or even blocked Meyer et al., 2002a) the penetration of Atlantic air masses to the Laptev Sea region. In that case, a stronger influence of the North Pacific winter moisture source, with low  $d$  values and low precipitation (e.g. Clark and Fritz, 1997), could have existed. Note however that the occurrence of this specific North Pacific moisture low  $d$  isotopic signature east of our study site is still somewhat controversial in ground ice (Schwamborn et al., 2006). During interstadials (Table 2, col. 4), on the contrary, a distinct Northern Atlantic source could have been re-established, with potentially higher  $d$  values (more “southerly” sources) and higher precipitation (Lacelle et al., 2011). We will discuss below these two alternatives, in the light of the other ice properties.

Climatic seasonal variations (temperature, precipitation, etc.) are also able to disturb  $d$  (Johnsen et al., 1989; Meyer et al., 2002b) but ice wedges are known to be built from the winter snow precipitation mainly and the spring meltwater derived from it. Post-depositional “*in situ*” phase changes could also be responsible for lowering the  $d$  values of the samples (Meyer et al., 2002b). Isotopic fractionation during ice wedge consolidation was discounted by Michel (1982) based upon the argument that freezing is generally too fast to allow for it. However, conditions might be different for the formation of the “ribbon facies” in the ice-sand wedge for example. Figure 9 shows the effect of equilibrium fractionation during hypothetical 10% refreezing (grey squares) of water resulting from the melting of each of the samples from IW-28 (black dots). The mean isotopic composition of the 10% refrozen fraction was computed using the formulation in Souchez and Lorrain (1991) as in Eq. (2), with a frozen fraction ( $K$ )=0.10.

$$\bar{\delta}_s = 10 * (1000 + \delta_0) * [(1.1 - K)^\alpha - (1 - K)^\alpha] - 1000 \quad (2)$$

where  $\bar{\delta}_s$  =  $\delta$  value of refrozen samples

$\delta_0$  =  $\delta$  value of water (melted IW-28 ice sample, no fractionation on melting (Moser and Stichler, 1980; Souchez and Lorrain, 1991))

$K$  = frozen fraction

$\alpha$  = fractionation coefficient (1.0208 for deuterium and 1.003 for  $^{18}\text{O}$ )

The refrozen IW-28 samples indeed reproduce the  $d$  of the ISW-28 samples and lie on the same co-isotopic line. They are however clearly offset towards less negative  $\delta$  values, and more negative  $\delta$  values than observed in IW-28 would be required to explain the observed



ISW-28  $\delta$ -range. This is not precluded, given the existence of some more negative values from the Sartan period reported by e.g. Wetterich et al. (2011, Fig. 6) in other ice wedges of the area. Experimental sublimation/evaporation processes of a snow pack have also been shown to shift the remaining reservoir towards less negative isotopic values with a slope that is less than 8 (Souchez and Lorrain, 1991 and references within). However, although they could reproduce the observed  $d$  values in ISW-28, both of these post-depositional processes would need to be occurring at a rigorous equivalent rate (e.g., 10% refreezing, fixed evaporation percentage) for all samples to plot on a co-isotopic line as well defined as the one observed for the ISW-28 samples, with a slope close to 8. In fact, the very good alignment of all the ISW-28 samples on a line with a slope of 8.03 ( $r^2=0.98$ , with a fairly low range of the  $\delta$  values) is an argument to favour a “classical” Rayleigh-type “source to sink” fractionation history.

This leaves us therefore with the first hypothesis of a major shift in the moisture source (higher  $d$ ), as the ice wedge growth switches from ISW-28 to IW-28, across the transition. Do the other ice properties allow us to decipher between the two alternative moisture source changes, with opposite conclusions in terms of the  $d$  signature? The answer is not straightforward. Indeed, ISW-28 has very low  $d$ -values (Table 2, col. 6 and Fig. 4) which would either correspond to an interstadial period in hypothesis 1 (Atlantic source, little sea ice cover - Table 2, col. 2) or to a stadial in hypothesis 2 (North Pacific source - Table 2, col. 5). The latter option is coherent with the deduced lower relative precipitation amounts from the observation of the ice-vein properties at ISW-28. However, in that case, the “in-situ” temperature should be lower at ISW-28 than at IW-28 (which would then be typical of re-establishment of an Atlantic source with higher  $d$ -values and higher precipitation). Alternatively, higher  $T_{\text{site}}$  for ISW-28 fit to an interstadial in hypothesis 1 (change in position of Atlantic source), but then higher precipitation amounts should be expected (Table 2, col. 2). Note however that the  $\delta$ -contrast is low between ISW-28 and IW-28, and that it should also be strong if hypothesis 1 was valid, with much higher values for ISW-28, provided that the mechanism is similar to the one presented for the Greenland Ice Sheet “stadial-interstadial” alternation by Johnsen et al., 1989. The latter assumption is hypothetical. Indeed, precipitation and temperature regimes on the downwind side of major ice-sheets might have been quite different to those existing in its centre parts (e.g. foehn effects which would locally increase temperatures and lower precipitations). One could also argue that the “switch” from North Pacific to Atlantic sources in hypothesis 2 could also be primarily linked to changes in

1 the Greenland and Eurasian Ice Sheets geometries, which might in turn not be directly linked  
2 to contrasted global temperature regimes but to other mechanisms such as ice-ocean  
3 interactions.

4 In the ISW-28 ice-sand wedge the dominance of debris input over snow filling, suggests a  
5 period of limited precipitation in accordance with the North Pacific winter moisture source  
6 with a low  $d$  value. However, the thin sections suggest the presence of meltwater and of  
7 melting-refreezing processes in an open-system, sandwiched with refrozen snow layers of  
8 limited extension (also supported by the total gas content measurements). Intuitively, these  
9 would be preferably associated to warmer periods, but they could also simply result from  
10 albedo effects in this debris-rich environment.

## 12 **6 Conclusions**

13 Detailed crystallographic, gas content and composition, and water isotope analyses of the IW-  
14 26 and IW-28 ice wedges allowed us to shed more light on the processes involved in ice  
15 wedge growth. Ice wedge IW-26 has developed in a mild Holocene environment. Higher  
16 temperatures and higher meltwater infiltration resulted in enhanced recrystallization, larger  
17 crystal sizes, lower total gas content and favourable conditions for biological respiration,  
18 inducing relatively high CO<sub>2</sub> levels. Ice wedge ISW-28/IW-28 developed in a relatively mild  
19 Pleistocene environment, during which a large change in atmospheric sources occurred, as  
20 documented by a sharp monotonous transition in deuterium excess from stable values at ca.  
21 1.4‰ in ISW-28 (early growth) to stable values at ca. 7.5‰ in IW-28, that cannot be  
22 explained through post-depositional processes. This transition occurred without significant  
23 changes in the site temperature as indicated by constantly low  $\delta^{18}\text{O}$  and  $\delta\text{D}$  values. It is  
24 however difficult to select potential mechanisms behind the changes in atmospheric sources.  
25 The transition is however concurrent with major changes in the ice wedge morphology. The  
26 ice-sand wedge ISW-28 properties show a smaller contribution of refrozen “snow-water mix”  
27 in the ice veins, with regular episodes of film water open-system refreezing on the crack  
28 walls, associated with important losses of total gas content and abundance of surface-derived  
29 debris inclusions forming the “ribbon facies”. This heterogeneous medium, rich in organic  
30 matter, might have favoured the anaerobic micro-environmental conditions necessary to  
31 explain the maximum CH<sub>4</sub> levels in that unit. Ice veins texture in IW-28 resemble more the

1 refrozen “snow-water mix” of IW-26, but with smaller crystal sizes and larger total gas  
2 content, as would be expected from colder conditions.

3 The changes in  $d$  between ISW-28 and IW-28 are most likely linked with changes in the  
4 moisture source region from Atlantic to Pacific or changes within an Atlantic source region.  
5 The combined observed low  $d$  values and limited snow contribution to the ice vein matrix,  
6 suggesting low precipitation, in ISW-28 is consistent with a North Pacific atmospheric source  
7 for the local precipitation at the time of ISW-28 formation. The length and persistence of this  
8 low  $d$  event is however unknown since it has not been observed at any other ice wedge study  
9 site west of 130°E. The significant contrast in ice veins metamorphic processes between ISW-  
10 28 and IW-28, suggesting a larger contribution of melting-refreezing processes in the former,  
11 could result from slightly milder conditions at ISW-28. However, the close proximity of the  
12 soil-sediment source has certainly fostered melting through albedo feedbacks.

13 Finally, comparing our methane mixing ratios in IW-26 and IW-28 reveals that substrate  
14 availability is as important as the temperature regime in controlling production rates in the  
15 permafrost environment.

## 17 **Acknowledgements**

18 We thank all participants of the Russian-German expedition “Lena Anabar 2003”. Sampling  
19 and analytical work in the AWI Potsdam laboratories was supported by Antje Eulenburg,  
20 Eileen Nebel, Ute Bastian and Lutz Schönicke. This study is part of the Russian-German  
21 cooperative scientific project “System Laptev Sea”.

## References

- Barnola, J. M., Raynaud, D., Neftel, A. and Oeschger, H.: Comparison of CO<sub>2</sub> measurements by two laboratories on air from bubbles in polar ice, *Nature*, 303(5916), 410–413, 1983.
- Berg, T. E. and Black, R. F.: Preliminary measurements of growth of nonsorted polygons, Victoria Land, Antarctica, :In *Antarctic Soils and Soil -Forming Processes*, American Geophysical Union Antarctic Research Series, (1418), 61–108, 1966.
- Black, R. F.: Permafrost - A review, *Bulletin of the Geological Society of America*, 65, 839–856, 1954.
- Black, R. F.: Les coins de glace et le gel permanent dans le Nord de l'Alaska, *Annales de Géographie*, 72(391), 257–271, 1963.
- Black, R. F.: Features indicative of permafrost, *Annual Reviews Inc. Provided by the NASA Astrophysics Data System*, 75–94, 1976.
- Blunier, T., Chappellaz, J. A., Schwander, J., Barnola, J.-M., Despert, T., Stauffer, B. and Raynaud, D.: Atmospheric methane record from a Greenland ice core over the last 1000 years, *Geophysical Research Letters*, 20(20), 2219–2222, 1993.
- Bobrov, A. A., Muller, S., Chizhikova, N. A., Schirrmeister, L. and Andreev, A. A.: Testate amoebae in late quaternary sediments of the Cape Mamontov Klyk (Yakutia), *Biol. Bull*, 36(4), 363–372, 2009.
- Brouchkov, A. and Fukuda, M.: Preliminary Measurements on Methane Content in Permafrost, Central Yakutia, and some Experimental Data, *Permafrost and Periglacial Processes*, 13, 187–197, 2002.
- Cardyn, R., Clark, I. D., Lacelle, D., Lauriol, B., Zdanowicz, C. and Calmels, F.: Molar gas ratios of air entrapped in ice: A new tool to determine the origin of relict massive ground ice bodies in permafrost, *Quaternary Research*, 68(2), 239–248, 2007.
- Clark, I. and Fritz, P.: *Environmental isotopes in hydrogeology*, CRC Press (Lewis Publishers), 1997.
- Corte, A. E.: Relationship between four ground patterns, structure of the active layer, and type and distribution of ice in the permafrost, *U.S. Army Cold Regions Research and Engineering Laboratory*, 1962.
- Craig, H.: Isotopic variations in meteoric waters, *Science*, 133(3465), 1702–1703, 1961.
- Dansgaard, W.: Stable isotopes in precipitation, *Tellus*, 16(4), 436–468, 1964.
- Dereviagin, A. Y., Meyer, H., Chizhov, A. B., Hubberten, H.-W. and Simonov, E. F.: New data on the isotopic composition and evolution of modern ice wedges in the Laptev Sea region, *Polarforschung*, 70, 27–35, 2002.
- Gell, W. A.: *Underground Ice in Permafrost, Mackenzie Delta-Tuktoaktuk*, N.W.T, 1976.
- Gilichinsky, D. A., Nolte, E., Basilyan, A. E., Beer, J., Blinov, A. V., Lazarev, V. E., Kholodov, A. L., Meyer, H., Nikolskiy, P. A., Schirrmeister, L. and Tumskey, V. E.: Dating of syngenetic ice wedges in permafrost with <sup>36</sup>Cl, *Quaternary Science Reviews*, 26(11–12), 1547–1556, doi:10.1016/j.quascirev.2007.04.004, 2007.
- IPCC: *Climate change 2007: The physical Science Basis*, Cambridge University Press., 2007.
- Jacka, T. H.: Laboratory studies on relationships between ice crystal size and flow rate, *Cold*

- 1 Regions Science and Technology, 10(1), 31–42, 1984.
- 2 Johnsen, S. J., Dansgaard, W. and White, J. W. C.: The origin of Arctic precipitation under  
3 present and glacial conditions, *Tellus*, 41B, 452–468, 1989.
- 4 Jouzel, J., Masson-Delmotte, V., Stievenard, M., Landais, A., Vimeux, F., J., S. J.,  
5 Sveinbjornsdottir, A. and White, J. W. C.: Rapid deuterium-excess changes in Greenland ice  
6 cores : a link between the ocean and the atmosphere, *C. R. Géoscience*, 337, 957–969, 2005a.
- 7 Jouzel, J. and Merlivat, L.: Deuterium and oxygen 18 in precipitation: modeling of the  
8 isotopic effects during snow formation, *Journal of Geophysical Research*, 89, 11749–11757,  
9 1984.
- 10 Jouzel, J., Merlivat, L. and Lorius, C.: Deuterium excess in an East Antarctic ice core  
11 suggests higher relative humidity at the oceanic surface during the last glacial maximum,  
12 *Nature*, 299, 688–691, 1982.
- 13 Jouzel, J., Stievenard, M., Johnsen, S. J., Landais, A., Masson-Delmotte, V.,  
14 Sveinbjornsdottir, A., Vimeux, F., von Grafenstein, U. and White, J. W. C.: The GRIP  
15 deuterium-excess record, *Quaternary Science Reviews*, 26(1-2), 1–17, 2005b.
- 16 Katayama, T., Tanaka, M., Moriizumi, J., Nakamura, T., Brouchkov, A., Douglas, T. A.,  
17 Fukuda, M., Tomita, F. and Asano, K.: Phylogenetic analysis of bacteria preserved in a  
18 permafrost ice wedge for 25,000 years, *Appl. Environ. Microbiol.*, 73(7), 2360–2363, 2007.
- 19 Killawee, J. A., Fairchild, I. J., Tison, J.-L., Janssens, L. and Lorrain, R.: Segregation of  
20 solutes and gases in experimental freezing of dilute solutions: Implications for natural glacial  
21 systems, *Geochimica et Cosmochimica Acta*, 62(23/24), 3637–3655, 1998.
- 22 Knight, P. G.: *Glaciers*, Stanley Thornes Ltd., 1999.
- 23 Kuznetsova, L. P.: Atmospheric moisture content and transfer over the territory of the former  
24 USSR, In *Second International Workshop on Energy and Water Cycle in GAMESiberia*,  
25 1997, OhataT, HiyamaT (eds). Research Report of IHAS. Institute for Hydrospheric-  
26 Atmospheric Sciences, Nagoya University: Nagoya, Japan;, 145–151, 1998.
- 27 Lacelle, D., Radtke, K., Clark, I. D., Fisher, D., Lauriol, B., Utting, N. and Whyte, L. G.:  
28 Geomicrobiology and occluded O<sub>2</sub>–CO<sub>2</sub>–Ar gas analyses provide evidence of microbial  
29 respiration in ancient terrestrial ground ice, *Earth and Planetary Science Letters*, 306(1–2),  
30 46–54, doi:10.1016/j.epsl.2011.03.023, 2011.
- 31 Lachenbruch, A. H.: Mechanics of thermal contraction cracks and ice wedge polygons in  
32 permafrost, *Geological Society of America Special Papers*, 70, 1–69, 1962.
- 33 Langway, C. C. J.: Ice fabrics and the universal stage, *SIPRE Technical Report*, 62, 1958.
- 34 Lauriol, B., Duchesne, C. and Clark, I. D.: Systématique du remplissage en Eau des Fentes de  
35 Gel: les Résultats d’une étude Oxygène-18 et Deutérium, *Permafrost and Periglacial*  
36 *Processes*, 6, 47–55, 1995.
- 37 Lawson, W.: Environmental Conditions, Ice Facies and Glacier Behaviour, in *Glacier Science*  
38 *and Environmental Change*, edited by P. G. Knight, pp. 319–328, Blackwell., 2007.
- 39 Mackay, J. R.: Ice-Wedge Cracks, Garry Island, Northwest Territories, *Can. J. Earth Sci.*,  
40 11(10), 1366–1383, doi:10.1139/e74-133, 1974.
- 41 Martinerie, P., Lipenkov, V. Y., Raynaud, D., Chappellaz, J., Barkov, N. I. and Lorius, C.: Air  
42 content paleo record in the Vostok ice core (Antarctica): A mixed record of climatic and  
43 glaciological parameters, *J. Geophys. Res.*, 99(D5), 10565–10576, 1994.

- 1 Masson-Delmotte, V., Jouzel, J., Landais, A., Stievenard, M., Johnsen, S. J., White, J. W. C.,  
2 Werner, M., Sveinbjornsdottir, A. and Fuhrer, K.: GRIP Deuterium Excess Reveals Rapid and  
3 Orbital-Scale Changes in Greenland Moisture Origin, *Science*, 309(5731), 118–121, 2005.
- 4 Merlivat, L. and Jouzel, J.: Global climatic interpretation of the deuterium-oxygen 18  
5 relationship for precipitation., *Journal of Geophysical Research*, 84, 5029–5033, 1979.
- 6 Meyer, H., Dereviagin, A., Siegert, C., Schirrmeister, L. and Hubberten, H.-W.: Palaeoclimate  
7 Reconstruction on Big Lyakhovsky Island, North Siberia—Hydrogen and Oxygen Isotopes in  
8 Ice Wedges, *Permafrost and Periglacial Processes*, 13, 91–105, 2002a.
- 9 Meyer, H., Dereviagin, A. Y., Siegert, C. and Hubberten, H.-W.: Paleoclimate studies on  
10 Bykovsky Peninsula, North Siberia - hydrogen and oxygen isotopes in ground ice,  
11 *Polarforschung*, 70, 37–51, 2002b.
- 12 Meyer, H., Schirrmeister, L., Andreev, A., Wagner, D., Hubberten, H.-W., Yoshikawa, K.,  
13 Bobrov, A., Wetterich, S., Opel, T., Kandiano, E. and Brown, J.: Lateglacial and Holocene  
14 isotopic and environmental history of northern coastal Alaska - Results from a buried ice-  
15 wedge system at Barrow, *Quaternary Science Reviews*, 29(27-28), 3720–3735, doi:doi:  
16 10.1016/j.quascirev.2010.08.005, 2010a.
- 17 Meyer, H., Schirrmeister, L., Yoshikawa, K., Opel, T., Wetterich, S., Hubberten, H.-W. and  
18 Brown, J.: Permafrost evidence for severe winter cooling during the Younger Dryas in  
19 northern Alaska, *Geophys. Res. Lett.*, 37(3), L03501, doi:10.1029/2009GL041013, 2010b.
- 20 Meyer, H., Schönicke, L., Wand, U., Hubberten, H.-W. and Friedrichsen, H.: Isotope studies  
21 of hydrogen and oxygen in ground? ice experiences with the equilibration technique,  
22 *Isotopes in environmental and health studies*, 36, 133–149, 2000.
- 23 Michel, F. A.: Isotope investigations of permafrost waters in northern Canada, University of  
24 Waterloo, Ontario, Canada, Dept. of Earth Sciences., 1982.
- 25 Moser, H. and Stichler, W.: Environmental isotopes in ice and snow, in *Handbook of*  
26 *Environmental Isotope Geochemistry, the terrestrial environment*, 1A, edited by P. Fritz and J.  
27 Fontes, pp. 141–178, Elsevier, Amsterdam., 1980.
- 28 Müller, S., Bobrov, A. A., Schirrmeister, L., Andreev, A. A. and Tarasov, P. E.: Testate  
29 amoebae record from the Laptev Sea coast and its implication for the reconstruction of Late  
30 Pleistocene and Holocene environments in the Arctic Siberia, *Palaeogeography,*  
31 *Palaeoclimatology, Palaeoecology*, 271(3-4), 301–315, 2009.
- 32 Opel, T., Dereviagin, A. Y., Meyer, H., Schirrmeister, L. and Wetterich, S.: Palaeoclimatic  
33 information from stable water isotopes of Holocene ice wedges on the Dmitrii Laptev Strait,  
34 northeast Siberia, Russia, *Permafrost Periglac. Process.*, 22(1), 84–100, doi:10.1002/ppp.667,  
35 2011.
- 36 Paterson, W. S. B.: *The physics of glaciers*, third edition, Butterworth-Heinemann, Oxford.,  
37 1994.
- 38 Pickering, F. B.: *The basis of Quantificative Metallography.*, Metals and Metallurgy Trust.,  
39 1976.
- 40 Popp, S., Diekmann, B., Meyer, H., Siegert, C., Syromyatnikov, I. and Hubberten, H.-W.:  
41 Palaeoclimate Signals as Inferred from Stable-isotope Composition of Ground Ice in the  
42 Verkhoyansk Foreland, Central Yakutia, *Permafrost and Periglacial Processes*, 17, 119–132,  
43 2006.
- 44 Raffi, R., Stenni, B., Flora, O., Polesello, S. and Camusso, M.: Growth processes of an inland

- 1 Antarctic ice wedge, Mesa Range, northern Victoria Land, *Annals of Glaciology*, 39(1), 379–  
2 385, doi:10.3189/172756404781814195, 2004.
- 3 Raynaud, D., Chapellaz, J., Barnola, J. M., Korotkevich, Y. and Lorius, C.: Climatic and CH<sub>4</sub>  
4 cycle implications of glacial-interglacial CH<sub>4</sub> change in the Vostok ice core, *Nature*,  
5 333(6174), 655–657, 1988.
- 6 Raynaud, D., Delmas, D., Ascencio, J. M. and Legrand, M.: Gas extraction from polar ice  
7 cores: a critical issue for studying the evolution of atmospheric CO<sub>2</sub> and ice-sheet surface  
8 elevation, *Annals of Glaciology*, 3, 265–268, 1982.
- 9 Rinke, A., Dethloff, K., Spekat, A., Enke, W. and Hesselbjerg Christensen, J.: High resolution  
10 climate simulations over the Arctic, *Polar Research*; Vol 18, No 2 (1999): Special issue:  
11 Proceedings of the International Symposium on Polar Aspects of Global Change [online]  
12 Available from: <http://www.polarresearch.net/index.php/polar/article/view/6567>, 1999.
- 13 Romanosky, N. N.: The scheme of correlation of polygonal wedge structures, *Biuletyn*  
14 *Peryglacjalny*, (26), 287–294, 1976.
- 15 Samyn, D., Fitzsimons, S. and Lorrain, R.: Strain-induced phase changes within cold basal ice  
16 from Taylor Glacier, Antarctica, indicated by textural and gas analyses, *Journal of*  
17 *Glaciology*, 51(175), 611–619, 2005.
- 18 Schirrmeister, L., Grosse, G., Kunitsky, V., Magens, D., Meyer, H., Dereviagin, A.,  
19 Kuznetsova, T., Andreev, A., Babi, O., Kienast, F., Grigoriev, M., et al.: Periglacial  
20 landscape evolution and environmental changes of Arctic lowland areas for the last 60,000  
21 years (Western Laptev Sea coast, Cape Mamontov Klyk), *Polar Research*, 27(2), 249–272,  
22 2008.
- 23 Schirrmeister, L., Siegert, C., Kuznetsova, T., Kuzmina, S., Andreev, A., Kienast, F., Meyer,  
24 H. and Bobrov, A.: Paleoenvironmental and paleoclimatic records from permafrost deposits in  
25 the Arctic region of Northern Siberia, *Quaternary International*, 89, 97–118, 2002.
- 26 Schwamborn, G., Meyer, H., Fedorov, G., Schirrmeister, L. and Hubberten, H.-W.: Ground  
27 ice and slope sediments archiving late Quaternary paleoenvironment and paleoclimate signals  
28 at the margins of El'gygytgyn Impact Crater, NE Siberia, *Quaternary Research*, 66(2), 259–  
29 272, doi:10.1016/j.yqres.2006.06.007, 2006.
- 30 Souchez, R., Janssens, L., Lemmens, M. and Stauffer, B.: Very low oxygen concentration in  
31 basal ice from Summit, Central Greenland, *Geophysical Research Letters*, 22(15), 2001–2004,  
32 1995a.
- 33 Souchez, R., Jouzel, J., Landais, A., Chapellaz, J., Lorrain, R. and Tison, J.-L.: Gas isotopes  
34 in ice reveal a vegetated central Greenland during ice sheet invasion, *Geophysical Research*  
35 *Letters*, 33, L24503, doi:10.1029/2006GL028424, 2006.
- 36 Souchez, R., Lemmens, M. and Chapellaz, J.: Flow-induced mixing in the GRIP basal ice  
37 deduced from the CO<sub>2</sub> and CH<sub>4</sub> records, *Geophysical Research Letters*, 22(1), 41–44, 1995b.
- 38 Souchez, R. and Lorrain, R.: *Ice Composition and Glacier Dynamics*, Springer-Verlag., 1991.
- 39 Stenni, B., Jouzel, J., Masson-Delmotte, V., Röthlisberger, R., Castellano, E., Cattani, O.,  
40 Falourd, S., Johnsen, S. J., Longinelli, A., Sachs, J. P., Selmo, E., et al.: A late-glacial high-  
41 resolution site and source temperature record derived from the EPICA Dome C isotope  
42 records (East Antarctica), *Earth and Planetary Science Letters*, 217(1-2), 183–195, 2004.
- 43 Stenni, B., Masson-Delmotte, V., Johnsen, S., Jouzel, J., Longinelli, A., Monnin, E.,  
44 Röthlisberger, R. and Selmo, E.: An Oceanic Cold Reversal During the Last Deglaciation,

- 1 Science, 293(5537), 2074–2077, 2001.
- 2 Steven, B., Pollard, W. H., Greer, C. W. and Whyte, L. G.: Microbial diversity and activity  
3 through a permafrost/ground ice core profile from the Canadian high Arctic, *Environmental*  
4 *Microbiology*, 10(12), 3388–3403, doi:10.1111/j.1462-2920.2008.01746.x, 2008.
- 5 St-Jean, M., Lauriol, B., Clark, I. D., Lacelle, D. and Zdanowicz, C.: Investigation of ice-  
6 wedge infilling processes using stable oxygen and hydrogen isotopes, crystallography and  
7 occluded gases (O<sub>2</sub>, N<sub>2</sub>, Ar), *Permafrost Periglac. Process.*, 22(1), 49–64,  
8 doi:10.1002/ppp.680, 2011.
- 9 Tison, J.-L.: Diamond -wire saw cutting technique for investigating textures and fabrics of  
10 debris-laden ice and brittle ice, *J. Glaciol.*, 40(135), 410 – 414, 1994.
- 11 Tison, J.-L., Souchez, R., Wolff, E. W., Moore, J. C., Legrand, M. R. and de Angelis, M.: Is a  
12 periglacial biota responsible for enhanced dielectric response in basal ice from the Greenland  
13 Ice Core Project ice core?, *Journal of Geophysical Research*, 1998, 103(D15), 18885–18894,  
14 1998.
- 15 Vaikmäe, R.: Oxygen isotopes in permafrost and ground ice - a new tool for paleoclimatic  
16 investigations, *Leipzig.*, 1989.
- 17 Vasil'chuk, Y. K. and Vasil'chuk, A. C.: Oxygen-isotope and C<sup>14</sup> data associated with Late  
18 Pleistocene syngenetic ice-wedges in mountains of Magadan region, Siberia, *Permafrost and*  
19 *Periglacial Processes*, 9(2), 177–183, 1998.
- 20 Vimeux, F., Cuffey, K. and Jouzel, J.: New insights into Southern Hemisphere temperature  
21 changes from Vostok ice cores using deuterium excess correction over the Last 420,000  
22 Years, *Earth and Planetary Science Letters*, 203(3-4), 829–843, 2002.
- 23 Vimeux, F., Masson-Delmotte, V., Jouzel, J., Stiévenard, M. and Petit, J. R.: Glacial-  
24 interglacial changes in ocean surface conditions in the southern Hemisphere, *Nature*, 398,  
25 410–413, 1999.
- 26 Wagner, D. and Liebner, S.: Global warming and carbon dynamics in permafrost soils:  
27 Methane production and oxidation, in *Permafrost Soils*, edited by R. Margesin, pp. 219–236,  
28 Springer Berlin Heidelberg., 2009.
- 29 Wetterich, S., Kuzmina, S., Andreev, A. A., Kienast, F., Meyer, H., Schirrmeister, L.,  
30 Kuznetsova, T. and Sierralta, M.: Palaeoenvironmental dynamics inferred from late  
31 Quaternary permafrost deposits on Kurungnakh Island, Lena Delta, Northeast Siberia, Russia,  
32 *Quaternary Science Reviews*, 27(15-16), 1523–1540, 2008.
- 33 Wetterich, S., Rudaya, N., Tumskey, V., Andreev, A. A., Opel, T., Schirrmeister, L. and  
34 Meyer, H.: Last Glacial Maximum records in permafrost of the East Siberian Arctic,  
35 *Quaternary Science Reviews*, 30(21–22), 3139–3151, doi:10.1016/j.quascirev.2011.07.020,  
36 2011.
- 37 Yershov, E. D.: *Geokryologija SSSR (Geocryology of the USSR)*, Nedra, Moscow., 1989.
- 38 Yurtsever, Y. and Gat, J. R.: Atmospheric waters, in *Stable Isotope Hydrology: Deuterium*  
39 *and Oxygen in Water Cycle*, edited by T. Rep., pp. 103–145, IAEA, Vienna., 1981.
- 40 Zeebe, R. E. and Wolf-Gladrow, D.: CO<sub>2</sub> in Seawater: Equilibrium, Kinetics, Isotopes, 65,  
41 Elsevier., 2001.
- 42



1 Table 1. Summary of results and comparison between IW-26, ISW-28 and IW-28.. Values  
2 separated with / are the minimum and the maximum, (number) corresponds to mean values.

	IW-26	ISW-28	IW-28
Crystal area range (mm <sup>2</sup> )	10.18 – 27.34	4.15 – 9.62	0.95 – 4.15
Crystal shape	equigranular	granular/rectangular elongated vertically sediments at grain boundary	equigranular
c-axes	1 max in horizontal girdle	horizontal girdle	1 max in horizontal girdle
Bubble size	~1mm	~1mm	~1mm
Bubble shape	spherical	spherical	spherical
Total gas content (ml/kg) (Meteoric ice =75-140*)	10/36 (homogeneous) (27)	11/42 (heterogeneous) (25)	27/50 (homogeneous) (45)
N <sub>2</sub> (%)	77/86 (82)	78/89 (85)	82/89 (86)
O <sub>2</sub> (%)	5/13 (10)	10/21 (15)	10/13 (12)
Ar (%)	(1.16)	(1.03)	(0.93)
CO <sub>2</sub> (ppmV)	35 000/110 000 (62 000)	860/7000 (3000)	10 000/48 000 (25 000)
CH <sub>4</sub> (ppmV)	0.5/1.6 (1)	41/72 (55)	7/10 (8)
δ <sup>18</sup> O (‰)	-30.6/-22.6 (-24.9)	-30.9/-28.9 (-29.9)	-31.9/-29.4 (-31)
δD (‰)	- 230/- 170 (-188)	-246/-230 (-237)	-247/-229 (-240)
δ <sup>18</sup> O/δD slope (‰)	6.63 (r <sup>2</sup> =0.94)	8.03 (r <sup>2</sup> =0.98)	7.44 (r <sup>2</sup> =0.98)

\*Martinerie et al. (1992)


1 Table 2: Comparison of relative values of  $\delta$ ,  $d$  and precipitation amount for the two sections  
2 of ice wedge ISW-28/IW-28 with expected values for the two atmospheric moisture sources  
3 hypotheses discussed in the text.

4

(1)	(2)	(3)	(4)	(5)	(6)	(7)
variables	Hyp. 1: Atlantic source		Hyp. 2: Atlantic vs. North Pacific		ISW-28	IW-28
	low sea ice Interstadial	high sea ice Stadial	Interstadial	Stadial		
$\delta$ -values (T° site)	high	low	high	low	low	lower
$d$ -values (T° source)	<b>low</b>	high	high	<b>low</b>	<b>low</b>	high
Precipitation (distance controlled)	high	low	high	low	low	high

5

1




2

3

4 Figure 1: Location of the Cape Mamontov Klyk sampling site in Arctic Siberia. The studied  
5 ice wedges were exposed on a coastal cliff on the Laptev Sea coast.

1



2

3 Figure 2: a) General stratigraphic scheme of the main section of Mamontov Klyk (adapted  
 4 from Schirrmeister et al., 2008) with positions and general settings b) and c) of ice wedges 26  
 5 (IW-26) and 28 (IW-28) respectively. The sampling areas are outlined with white boxes, on  
 6 approximately 160 cm and 190 cm for respectively IW-26 and IW-28. Rectangular  
 7 parallelepipeds have been removed from the cliff with chain-saw of which the upper part (1.5  
 8 cm thickness) has been used in this work. Note that IW-28 was sampled in both the ice-sand  
 9 wedge (ISW-28, left of box) and in the ice wedge itself (IW-28, right of box).

10






Figure 3: Thin section photographs of the textural properties of the ice wedges. a) IW-26, b) ISW-28 (ice-sand wedge), the left part of ice wedge 28 with debris sub vertical layers (note the crystal elongation along the foliation), c) IW-28, clean ice in the central part of ice wedge 28 and d) thick section photograph of ice from IW-28 in transmitted light allowing bubble content observation.

1



2

3 Figure 4: Isotopic data for the two ice wedges: a)  $\delta^{18}\text{O}$  and  $\delta\text{D}$  profiles, b)  $d$  and c)  $\delta^{18}\text{O}$ - $\delta\text{D}$   
 4 relationship. Numbers on the bottom x-axis of a) and b) refer to ice sample ordering,  
 5 increasing from left to right, in the outcrop. The equivalent horizontal distance from extreme  
 6 left is shown on the upper x-axis.



1

2 Figure 5: Crystals size measurements using the two techniques described in the “sampling and  
 3 analytical methods” section.

4




Figure 6: Schmidt equal-area diagrams for the c-axes of horizontal thin sections. a) a total of six thin sections from IW-26 ( $n = 199$ ) showing horizontal girdle with 1 maximum, b) a total of two thin sections from ISW-28 ( $n = 70$ ) showing horizontal girdle, c) a total of five thin sections from IW-28 ( $n = 150$ ) showing horizontal girdle with 1 maximum.








Figure 7: Gas properties of ice wedges IW-26 (left) and ISW-28/IW-28 (right).



1



2

3 Figure 9:  $d$  (a) and co-isotopic plots (b) of observed data for ISW-28 (open circles) and IW-28  
 4 (black dots) compared to a simulation (grey squares) on the effect of hypothetical equilibrium  
 5 refreezing of 10% of melted fraction in IW-28 sample, using Eq. (1). The refrozen theoretical  
 6 IW-28 samples reproduce the  $d$  of the ISW-28 samples and lie on the same co-isotopic line.  
 7 They are however clearly offset towards less negative  $\delta$  values.