

## ***Interactive comment on “The multiphase physics of sea ice: a review” by E. C. Hunke et al.***

**S. Ackley (Referee)**

stephen.ackley@utsa.edu

Received and published: 7 September 2011

General Comments: Since this Review paper is not the usual presentation of scientific results, it cannot be subjected to the usual criteria for a Journal article, e.g. original work, new results, scientific accuracy, etc. I therefore offer some personal criteria to judge the quality of this review such as: Is the work reviewed a relatively complete summary of the subject area? Does the review provide a source of relevant material that can be used as background for other papers on the specific details of the subject, such as field work or modeling studies? The context of this article is the authors' statement that the review resulted from a Workshop on The Multi-Phase Physics of Sea Ice: Growth, Desalination and Transport Processes held in 2010. As a result, the authors have sacrificed the breadth of sea ice physics, experimental and field data, for instead looking at Mushy Layer Physics and how it explains basically steady state growth of sea ice in a highly controlled laboratory situation, which was an emphasis at

C922

that workshop. It may be useful for understanding the physics of the medium. However, in the 2nd half of the paper the authors review thermohaline descriptions of the sea ice component of global climate models. Since these rely on older physics, and make simplifications for computational efficiency, the relevance of Mushy Layer Physics to at least current numerical modeling is questionable. A better connection between the first half and second half of the paper is therefore sorely missed. I personally therefore did not find this review particularly useful as it is lacking in both completeness and as a source. There are only two figures in the paper (four, if you count the three parts of Figure 2 individually). Comparatively, the chapter on Sea ice Salinity in the book by W.F. Weeks (On Sea Ice, U of Alaska Press 2010) has 43 figures. That chapter is of similar length to this paper and also includes a succinct description of Mushy Layer Physics, including about 5 figures, none of which are included in this paper, despite the emphasis here on that topic. I think a Review titled the Multiphase Physics of Sea ice should include a figure of the Phase Diagram, and a discussion on the calculation (and dependence) of Brine Volume and Brine Salinity on Temperature. As well, Figure 1, after Malmgren 1927, implies that the evolution of the salinity profile is pretty standard. However, while Eicken 1992 is in the Reference List, his classification of the salinity profiles observed in Antarctic sea is not referred to. These indicate that of the order of only one third of the measured profiles in Antarctic sea ice are similar to those shown in Figure 1 (The “C” profile). Three other profile models are found due to the variations of driving forces, including surface flooding and high ocean heat flux, that are found to a different degree in the Antarctic. One premise of the Review is that the evolution of ice salinity makes a large difference in the way that sea ice is modeled, and drives many of the results. However, if the salinity profiles over the majority of the Antarctic ice pack do not conform to the assumed “C” profile, how can modeling results give an accurate representation? Maybe they do, but by not showing that they do, the reader is left with the impression that the 1927 results from Malmgren are the observed state of the art which the modelers are working toward replicating.

In the abstract, the focus of the review is identified as mushy-layer theory (since it

C923

“describes general multiphase materials” and on numerical approaches now being explored to model the multiphase evolution of sea ice. The implication is that these numerical approaches are using mushy layer theory but, with the absence of any model derived figures (figs 1 is Malmgren’s observed profiles and Fig 2 is pictures of brine pockets, and brine tubes observed in the laboratory and field), at least no comparisons are available. The entire review is a narrative that could be described as a continuous annotated bibliography. No information can be extracted from the review itself without using the original references. The first half of the paper, on mushy layer theory is actually a review of a previous paper, Notz and Worster 2009, so the first half is a review of that review? If all you’re going to talk about is a single paper, why go through it again in abbreviated form? Why not just say: “the main processes of salt removal from sea ice are initial rejection at the growing interface and gravity drainage from sea ice, followed by flushing from meltwater in the summer. Except for Flushing, mushy layer theory (Notz and Worster 2009) best explains the underlying physics of these processes. See that paper for details.” This article therefore represents the end product of the workshop and reflects the participation there, rather than the more comprehensive view taken in other reviews. Since these other reviews are recently available in book form, (Dieckmann and Thomas, 2nd Edition Sea Ice Physics Chemistry and Biology, and W.F. Weeks, On Sea Ice), I think an attempt to expand this review to be as useful is perhaps a duplicative effort, so recommend that the paper remain as a TC Discussion paper rather than trying to correct it into an article in The Cryosphere.

Specific Comments:

p. 1950 l.7 The statement that brine can drain from the ice, taking other constituents with it, depends on the ice porosity is a bit inaccurate. The permeability is the correct term, and it depends on structure as well as porosity. The two terms are used interchangeably throughout the paper but they should be distinguished. p.1951 There is a very qualitative description of the variation of brine, salts and ice as a function of temperature with no significant information passed on. This would be a good spot for

C924

the Phase Diagram. (e.g. It is stated that “solid salts start to precipitate at -2.2C” yet no identification of this as Calcium Carbonate with highly significant role in CO<sub>2</sub> exchange. “other salts remain in solution. . .below -50C”. First of all these temperatures are rarely, if ever achieved in natural sea ice, so the temperature range of interest is the NaCl eutectic (down to -21 C) when 99% of the brine is converted to solid salts. The other salt that comes out is Mirabilite, sodium sulfate, at -7 to -8C (if I had the phase diagram I could be more accurate. . .) Above temperatures of -10C are of more interest since the cutoff in permeability occurs (typically) in that range and the highest volume of sea ice will be above -10C for most of its existence. Why give a qualitative description when real values are easily mentioned? p.1952 Mention of the carbon cycle and the precipitation and dissolution of CaCO<sub>3</sub> as a pathway for CO<sub>2</sub> exchange, however It’s quite disconnected from the incomplete description in the previous page where the highest temperature salt precipitated as calcium carbonate is not identified. p. 1952 on iron accumulation, dissolved iron is mentioned. but a more recent paper by Lannuzel et al 2011, shows that particulate iron accumulated outweighs dissolved iron by 23:1. p.1954 Brine salinity is therefore only a function of temperature T. (see Cox and Weeks for empirical functions). In Willy Weeks new book, there is a whole chapter on the Phase Diagram, which is based on empirical data but follows fundamental principles, since Gibbs, of physical chemistry. Yet, it’s not worthy of mention, never mind showing, in this review. Also, it would be helpful to actually give the empirical functions of Cox and Weeks.

P.1954 Brief overview of mushy-layer theory. I think there is bit of overemphasis on Mushy Layer Theory as providing all the answers, especially considering the later sections on numerical modeling which, for the most part, do not include mushy layer theory. Most of the advances using mushy layer theory deal with the case, using laboratory results, corresponding to constant heat flux from above resulting in one dimensional freezing of the ice. This can be used to predict the formation of brine channels, the release of salt from the ice, and the conduction of heat as the ice is freezing. However, the authors mention Feltham 2006 for a derivation of classical parameterizations of

C925

sea-ice properties from mushy layer theory. Is it therefore necessary to use full mushy-layer theory in numerical calculations, or are these parameterizations more efficient to apply computationally? Similarly, I personally think that many of the problems worth looking at are those dealing with temperature cycling, when the ice flops between cold and impermeable to warm and permeable, or in the warming case, when the structure laid down by the freezing process is already there. If the main channel structure is already there, is it necessary to use mushy layer theory to determine the fluid flow paths again? Or can some statistical ensemble of pipes of varying sizes be used with simpler, more efficient, equations? p. 1960 For the sake of discussion, I would like the authors' to comment on the following point. An argument is given (P.1960) that mushy layer theory somehow invalidates the Burton et al sea ice rejection (which uses  $k_{eff}$ , the effective partition coefficient, and does not depend on gravity), since salt rejected would be reincorporated when the mushy layer equations are applied without gravity. Sure, and all those oceanographic parameterizations that use the "eddy diffusivity" (eddies are not diffusive, are they?) are no good either. But, does  $k_{eff}$  actually implicitly include gravity, through the brine densification, so the argument the  $k_{eff}$  is actually the same as  $k$ , (the salt molecular partition coefficient) is analogous to calling the eddy diffusivity the molecular diffusivity? Isn't it what numerical modelers would call a "parameterization"? p. 1960 and on it goes, now we recapitulate the inferiority of the brine diffusion or brine pocket migration argument, but we knew that in 1968 (Untersteiner). p. 1961 This argument on brine expulsion, now refuted by mushy layer theory (Notz and Worster 2009). What isn't mentioned however is the possible role of brine expulsion in pushing brine upward in thin sea ice, possibly accounting for high salinities at the top layer of the sea ice?

p.1964 Approaches. After about a 15 pg exposition of the merits of applying mushy layer theory to sea ice, we are given a short paragraph on how it is applied to direct numerical simulations of individual crystals in the metallurgy fields, divided into sharp and diffuse interface solutions. After these descriptions of the numerical setup, with references, the following statement is made: "The heavy computational burden of ex-

C926

PLICITLY tracking the microstructure makes these techniques unsuitable for modeling a complete sea-ice layer, although they may prove useful in determining appropriate sub-grid scale parameterizations, such as the permeability, for averaged models." The next section then deals with Volume averaged simulations, where individual ice crystals and brine inclusions no longer need to be explicitly modeled, so "the problem becomes much more computationally tractable". Then, simulations specifically for forming sea ice, compared to the lab results of Cox and Weeks 1975 and Wettlaufer 1997a. results in "The ice growth rate and brine drainage rate were comparable to the experimental results, but the simulations lacked the observed delayed onset of drainage." Are the authors implying that the physics is validated, and that the deficiencies are a result of the need for volume averaging for computational efficiency? Maybe volume averaging instead causes some important physics to be overlooked? Is it good enough to use numerically if the onset of drainage is delayed? One dimensional simulations p. 1966 "For computational efficiency, the thermohaline description in the the sea ice component of global climate models is one-dimensional." I think this statement indicates the primary disconnect in the review between the modeling and the theory. The review in the section on Multi-Layer Physics of Sea Ice has dealt nearly exclusively with the Mushy Layer theory, i.e. a medium with convecting chimneys separated by a porous medium. If most of the numerical simulations are dealing with other than Mushy Layer Physics, wouldn't the audience be better served by a description of the physics that is actually used, rather than one that isn't applied?

p. 1966-1970 These descriptions of the various model results would be helped by some figures that show some of the important results. p.1970 "lack sufficient observational data to narrow the range of model parameters, a potential hindrance for further development." (puzzling statement, very little of the observational data that is available has been used or cited, so weakens the case that sufficient observational data doesn't exist) p.1972, figures needed to show the impact of salinity variations on sea ice from Vancoppenolle et al 2009a from their sensitivity simulations. p.1972 Couple of statements that are a little misleading. "only brine convection in the lowermost ,porous sea

C927

ice (Reeburgh, 1984) and the flooding of the surface by seawater (Fritsen et al 1994) seem intense enough to provide the required nutrients to sustain biological growth in the ice. “Fritsen et al actually showed that later freezing in flooded layers drove convection that caused the near-surface intense biological growth. So convection in both the uppermost, (or full thickness convection) and lowermost layers can both sustain biology. The mechanism for surface flooding itself to have high biology is still somewhat ephemeral, e.g. may require some waves, floe surging etc to mix surface seawater into flooded layers. The next statement that “vertical profiles of dissolved macronutrient concentrations and salinity have a similar shape” is also misleading. Several of these studies have shown for example that nutrients in the sea ice do not scale with salinity and are either depleted or enhanced because of biological activity. So similar shape, with the implication that nutrients are passive tracers of salinity, masks the possible true behavior. p.1973, algae, nutrients, trace metals, gases, are incorrectly here identified as passive tracers. Some of them might be, but for the most part none of them really are, e.g. algae stays within the ice while brine is rejected, nutrients are taken up proportional to biology, and CO<sub>2</sub> and DMS gases have radically different profiles from salinities. p.1973 Some success is ascribed to Vancoppenolle et al 2010 and Jeffrey et al 2011 in one dimensional models based on transport equations “containing some of the mushy-layer physics”, again here some display of results would have been helpful. p.1974 “There appear to be potential nonlinear interactions between brine dynamics and biogeochemical sources and sinks” Would this statement and the following couple of sentences have been a much better introduction to this section than the previous couple of pages which instead lead the reader into believing that there is merit in considering biogeochemicals as passive tracers?(See above comments.) Observations p.1975 Mention is made of the time series of ice salinity evolution and the lack of temporally and spatially resolved data from ice core studies. However, as reviewed by Weeks in On Sea ice, studies by Kovacs have determined functions that describe ice thickness and salinity that use a large quantity of observed cores from both the Arctic and Antarctic. As well,” the lack of data to test numerical models that describe

C928

the microstructure of sea ice with high spatial resolution”, given the various parameterizations described here that obviate the “high spatial resolution” for “computational efficiency” seems a bit harsh? While data may be sparse in relation to the widespread areas covered by sea ice, don't the similarities in the data from many areas suggest that high spatial and temporal resolution may not be quite as necessary? Doesn't Malmgren's data from a handful of cores in 1927 look quite similar to the detailed profiles of Nakawo and Sinha in 1981? Doesn't Eicken's (1992) classification of salinity profiles from Antarctic sea ice give some of the generality of this behavior as well? Its generally, however, hard to argue that better and more observations are needed, particularly in a time-series sense, for better comparison to numerical models. But their case is weakened when known observations and classifications are ignored. p.1977 Bit misleading to refer to ESA's Soil Moisture and Ocean Salinity satellite, which, as far as I know, would have no application to determining sea ice salinity and microstructure.

---

Interactive comment on The Cryosphere Discuss., 5, 1949, 2011.

C929