

## ***Interactive comment on “Basal crevasses in Larsen C Ice Shelf and implications for their global abundance” by A. Luckman et al.***

**J. Woodward (Referee)**

john.woodward@northumbria.ac.uk

Received and published: 20 September 2011

This is a well structured, important paper providing insight into basal crevasse formation on Antarctic ice shelves. The paper neatly highlights the importance of basal crevasses and provides a novel method for recognising them from remote sensing imagery. The remote sensing imagery is tied to ground-penetrating radar profiles from the Larsen C Ice Shelf. The GPR profiles are thoroughly convincing and well displayed. Numerical modelling is then applied to predict crevasse penetration heights. I am not best placed to review numerical modelling papers, so refrain from commenting, suffice to say a more complete modelling assessment of crevasse formation for the Ice Shelf would be an obvious next step in developing this work.

I have only limited questions regarding this paper, as it is well written, well presented  
C977

and well researched. Minor points are:

1. I was interested in the comments in lines 9-17 on page 2038, referring to the temporal pattern of crevasse formation. S1 crevasses seem to form every 5 years, with S2 crevasses forming every 14 years. Why is this? What controls the timing of crevasse formation? This could be revisited and commented upon in the discussion.
2. Line 12 of page 2046 indicates other remote sensing images are not shown because of ‘considerations of space’. Please produce this figure and include it for completeness. Space should not be an issue in an interactive online journal.

---

Interactive comment on The Cryosphere Discuss., 5, 2035, 2011.