

This discussion paper is/has been under review for the journal The Cryosphere (TC).
Please refer to the corresponding final paper in TC if available.

Spatial patterns of North Atlantic Oscillation influence on mass balance variability of European Glaciers

B. Marzeion¹ and A. Nesje²

¹Center of Climate and Cryosphere, University of Innsbruck, Innsbruck, Austria

²Bjerknes Centre for Climate Research, and Department of Earth Science, University of Bergen, Bergen, Norway

Received: 18 November 2011 – Accepted: 27 November 2011 – Published: 3 January 2012

Correspondence to: B. Marzeion (ben.marzeion@uibk.ac.at)

Published by Copernicus Publications on behalf of the European Geosciences Union.

TCD

6, 1–35, 2012

NAO influence on mass balance variability

B. Marzeion and A. Nesje

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

We present and validate a set of minimal models of glacier mass balance variability. The most skillful model is then applied to reconstruct 7735 individual time series of mass balance variability for all glaciers in the European Alps and Scandinavia. Subsequently, we investigate the influence of atmospheric variability associated with the North Atlantic Oscillation (NAO) on the glaciers' mass balances.

We find a spatial coherence in the glaciers' sensitivity to NAO forcing which is caused by regionally similar mechanisms relating the NAO forcing to the mass balance: In Southwestern Scandinavia, winter precipitation causes a correlation of mass balances with the NAO. In Northern Scandinavia, temperature anomalies outside the core winter season cause an anti-correlation between NAO and mass balances. In the Western Alps, both temperature and winter precipitation anomalies lead to a weak anti-correlation of mass balances with the NAO, while in the Eastern Alps, the influences of winter precipitation and temperature anomalies tend to cancel each other, and only on the southern side a slight anti-correlation of mass balances with the NAO prevails.

1 Introduction

The North Atlantic Oscillation (NAO) is the most prominent mode of atmospheric variability over the North Atlantic Ocean and Northwestern Europe (Visbeck et al., 2001; Wanner et al., 2001; Hurrell et al., 2003). One measure of the state of the NAO is the NAO index, calculated as the difference in sea level pressure between Stykkisholmur, Iceland, and Ponta Delgada, Azores (Hurrell, 1995b). The impact of NAO variability is most pronounced during the winter months, and strongest over Western Norway (Nesje et al., 2000). During the positive phase, i.e. stronger than normal pressure gradient between Iceland and Azores, stronger than normal westerlies, above normal precipitation and mild temperatures prevail across the Eastern North Atlantic and Northwestern Europe. During pronounced negative phases of the NAO, opposite patterns

NAO influence on mass balance variability

B. Marzeion and A. Nesje

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



of temperature and precipitation anomalies are observed (see Hurrell, 1995a; Wanner et al., 2001; Hurrell et al., 2003, for an overview over the NAO and its influence on European climate).

By influencing precipitation and temperature anomalies, the NAO also exerts control over glacier mass balances. Reichert et al. (2001) find that for Nigardsbreen, an eastern outlet glacier from the Jostedalsgreen ice cap, the impact of winter precipitation overrules the impact of positive temperature anomalies, leading to a correlation of mass balance and NAO. Nesje et al. (2000) show that the control of the NAO over the glacier mass balance in Scandinavia gradually decreases with increasing continentality (i.e., towards the east).

In the European Alps, the relation between climate anomalies originating from the Atlantic and glacier mass balance is more complex, as the Alps are situated in the transition zone between Northern Europe, experiencing warm and wet conditions during positive phases of the NAO, and Southern Europe, experiencing drier than normal conditions, and only small temperature deviations (Wanner et al., 2001). The influence of the NAO in the Alps is therefore generally less pronounced than in Scandinavia (Six et al., 2001), and more ambiguous (Scherrer et al., 2004). Huss et al. (2010) find a rather strong influence of the (longer term) Atlantic Multidecadal Oscillation on mass balances of glaciers in the Swiss Alps. Using a coupled general circulation model to drive a glacier mass balance model, Reichert et al. (2001) find that the mass balance of Rhonegletscher in the Western Alps is more sensitive to precipitation than temperature anomalies, resulting in an anticorrelation of the mass balance with the NAO. However, Steiner et al. (2008) show that winter precipitation is less important for length variations of Alpine glaciers than for Norwegian glaciers. Moreover, Marzeion et al. (2011) present a reconstruction of mass balance anomalies of glaciers in the European Alps based on a simple model derived from monthly values of temperature and precipitation and a set of mass balance measurements needed both to train and to validate the mass balance model, and they find that glaciers (especially in the Central and Eastern) Alps are more sensitive to temperature anomalies than to precipitation anomalies.

NAO influence on mass balance variability

B. Marzeion and A. Nesje

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



NAO influence on mass balance variability

B. Marzeion and A. Nesje

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



As input, Marzeion et al. (2011) use the HISTALP dataset (Auer et al., 2007)¹, which provides monthly precipitation sums and 2 m temperatures on a 5 × 5 minute grid of the greater alpine region, covering the years 1780 to 2008 (temperature) and 1801 to 2003 (precipitation), and area-integrated mass balances obtained from Cogley (2009)². Here, we first transfer their modeling approach from the HISTALP data to CRU TS 3.0 data (Mitchell and Jones, 2005)³ in Sect. 2.1. As in Marzeion et al. (2011), in the following we refer to this model as the *individually trained model*, and to the model using the mean parameters of the individually trained model the *mean model* (see below for a detailed description). In Sect. 2.2, we present another model of the glacier mass balance, which is based solely on climatological values of precipitation and temperature data as input, and does not depend on measured mass balances for training, but only for validation. In the following, we refer to this model as the *climatologically derived model*. A detailed validation of the models is presented together in Sect. 2.3. We then apply the models to all European glaciers contained in the extended format of the World Glacier Inventory (WGI-XF) data base (Cogley, 2005)⁴, and perform a correlation-based analysis of the response of the glaciers' mass balances to NAO-related variability in temperature and precipitation (see Fig. 1 for an overview of the spatial distribution of all glaciers considered in this study). The results are shown in Sect. 3, and we discuss them and conclude in Sect. 4.

¹available at <http://www.zamg.ac.at/histalp>

²available at <http://people.trentu.ca/~gcogley/glaciology/>

³available at <http://badc.nerc.ac.uk/>

⁴available at <http://people.trentu.ca/~gcogley/glaciology/>

2 Glacier mass balance models

2.1 Individually trained and mean model

The area-integrated annual mass balance MB of a glacier can be understood as the sum of the monthly precipitation onto the glaciers surface and monthly run off from the glacier. In Marzeion et al. (2011), the modeled mass balance MB_{model} is therefore derived by introducing two parameters, a and μ , such that

$$MB_{\text{model}} = \sum_{i=1}^{12} (aP_i - \mu(\max(0, T_i))) \quad (1)$$

where i indicates the month, P_i is the monthly solid precipitation onto the glacier surface (in their of case obtained from HISTALP data, which includes specific information on the solid fraction of total precipitation), T_i is the monthly mean air temperature at the location of the glacier terminus (in their case obtained from HISTALP data, corrected with a temperature lapse rate for any differences between the height of the glacier terminus and the height of the HISTALP data point). In this model, a can be understood as a parameter representing effects of a precipitation lapse rate, aeolian transport of snow, and avalanching, and μ can be understood as a parameter relating monthly mean temperatures at the glacier terminus to production of glacier melt. For each individual glacier with N existing annual mass balance measurements MB_{measured} , the optimal parameters a_{opt} and μ_{opt} can be estimated by requiring that the mean square error

$$mse_{\text{model}} = \frac{1}{N} \sum_{k=1}^N (MB_{k,\text{measured}} - MB_{k,\text{model}})^2, \quad (2)$$

where k is the year, is minimal. From this procedure, individual parameters estimates can be derived for each glacier with existing mass balance measurements (individually trained model). If a mass balance measurements exist for a number of glaciers within

NAO influence on mass balance variability

B. Marzeion and A. Nesje

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the region of interest, it is possible to construct the mean model by employing the mean of the parameters obtained from the individually trained models to reconstruct the mass balance of another glacier in the region of interest. While for a detailed discussion of the concept of the individually trained and the mean model we refer to Marzeion et al. (2011), the transfer of the model from the HISALP to the CRU TS 3.0 data as the driving data set necessitates the estimation of the solid fraction of precipitation onto the glacier surface, and a new validation of model performance and robustness. This is achieved by a cross validation procedure presented in Sect. 2.3.

2.2 Climatologically derived model

Kaser et al. (2010) develop a model of the monthly climatological glacier mass balance, based on monthly climatological precipitation and temperature at the glacier location, in order to estimate the potential of seasonally delayed runoff to water availability in large river basins. For a glacier in equilibrium with monthly climatological temperatures and precipitation, the annual mass gain from precipitation $\sum_{i=1}^{12} P_{i,\text{clim}}$, where $P_{i,\text{clim}}$ is the climatological monthly solid precipitation integrated over the surface of the glacier, has to equal the annual mass loss by melting of ice, $\sum_{i=1}^{12} R_{i,\text{clim}}$, where $R_{i,\text{clim}}$ is the climatological monthly melt from the glacier. The determination of $R_{i,\text{clim}}$ requires the consideration of energy budget of the glacier (see e.g., Kuhn, 1987; Oerlemans, 2000; Mölg and Hardy, 2004), but air temperature is a reasonable proxy for the energy available to the glacier for producing melt (Sicart et al., 2008; Ohmura, 2001). We therefore assume $R_{i,\text{clim}} = -\mu_{\text{clim}}(\max(0, T_{i,\text{clim}} - T_{\text{melt}}))$, where $T_{i,\text{clim}}$ is the monthly climatological air temperature at the glacier terminus, and T_{melt} is the monthly mean air temperature above which melt at the glacier terminus occurs. T_{melt} does not necessarily have to be 0°C , since the diurnal cycle and intra-monthly variability may lead to above-freezing temperatures even if the monthly mean is below 0°C , but it turns out that the model performance is both quite insensitive to the exact value of T_{melt} , and highest for $T_{\text{melt}} = 0^\circ\text{C}$ (see Marzeion et al. (2011) for a more detailed discussion), so that we assume $T_{\text{melt}} = 0^\circ\text{C}$ in the following.

NAO influence on mass balance variability

B. Marzeion and A. Nesje

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Then, if we assume that a glacier in its current extent is in equilibrium with monthly climatological temperatures and precipitation (see Sect. 4 for a discussion of the implications of this assumption), the annual climatological mass balance has to fulfill

$$MB_{\text{clim}} = \sum_{i=1}^{12} (P_{i,\text{clim}} + R_{i,\text{clim}}) = 0, \quad (3)$$

5 which leads directly to an estimate of μ_{clim} (see Eq. 6).

The variable annual mass balance of a glacier can then be estimated as

$$MB_{\text{model}} = \sum_{i=1}^{12} (P_i - \mu_{\text{clim}}(\max(0, T_i))). \quad (4)$$

where P_i is the monthly solid precipitation onto the glacier surface, and T_i is the monthly mean air temperature at the location of the glacier terminus.

10 The CRU CL 2.0 data set (New et al., 2002)⁵ includes $10' \times 10'$ resolution fields of monthly mean climatological temperature and total precipitation, while the CRU TS 3.0 data set (Mitchell and Jones, 2005) provides time series of monthly temperature and total precipitation on a $30' \times 30'$ grid. It is therefore possible to estimate vertical gradients of temperature and precipitation on the scale of the CRU TS 3.0 grid based
 15 on the CRU CL 2.0 climatology, if the topography around the location of interest is rough, i.e. if there is spread in the elevation of the CRU CL 2.0 grid surrounding the location of interest, and if the horizontal gradients in the area of interest are small compared to the vertical gradients. The first condition is fulfilled for all the the locations of interest in this study, since the glaciers are located in mountainous terrain. We
 20 assume that the second condition is only fulfilled for temperature, since especially in Scandinavia, the horizontal gradients in precipitation are large (see Sect. 4 for a more detailed discussion). We therefore estimate vertical lapse rates of temperature for all glaciers in the WGI-XF data set by regressing temperature in 3×3 CRU CL 2.0 grid

⁵available at <http://www.cru.uea.ac.uk/cru/data/hrg/tmc/>

NAO influence on mass balance variability

B. Marzeion and A. Nesje

| | |
|--------------------------|--------------|
| Title Page | |
| Abstract | Introduction |
| Conclusions | References |
| Tables | Figures |
| ◀ | ▶ |
| ◀ | ▶ |
| Back | Close |
| Full Screen / Esc | |
| Printer-friendly Version | |
| Interactive Discussion | |



points around the location of the glacier onto elevation. Figure 2, panel a, shows the result. The correlation between temperature and elevation is very high (typically > 0.95) and above the 95 % confidence interval for all glaciers. For precipitation, we assume a lapse rate of 2% precipitation increase every 100 m altitude increase. This lapse rate is on the lower side of the values typically gained from determining the lapse rates based on the CRU CL 2.0 climatology in those areas where the horizontal gradients in the climatology are small.

In order to estimate μ_{clim} for each glacier, we first determine the monthly climatological temperature $T_{i,\text{clim}}$ at the glacier terminus as

$$T_{i,\text{clim}} = T_{i,\text{CRU}_{\text{clim}}} + \gamma_T (z_{\text{terminus}} - z_{\text{CRU}}), \quad (5)$$

where $T_{i,\text{CRU}_{\text{clim}}}$ is the monthly climatological temperature of the CRU TS 3.0 data set at the location of the glacier⁶, γ_T is the temperature lapse rate at the location of the glacier determined as described above, z_{terminus} is the altitude of the glacier terminus, and z_{CRU} is the altitude of the CRU TS 3.0 grid point.

We then estimate the fraction of solid precipitation by assuming that if the temperature at terminus elevation is below 2°C, solid precipitation equals total precipitation. If the temperature at the highest elevation of the glacier (obtained using γ_T) is above 2°C, there is no solid precipitation. Within this range, we interpolate the solid fraction linearly. Note that the choice of 2°C as the threshold temperature is arbitrary, but yields the best model performance in the cross validation.

$$\frac{\sum_{i=1}^{12} P_{i,\text{clim}}}{\sum_{i=1}^{12} \max(0, T_{i,\text{clim}})} = \mu_{\text{clim}}, \quad (6)$$

where $P_{i,\text{clim}}$ is the monthly climatological solid precipitation estimate based on the CRU

⁶It would be possible to use the CRU CL 2.0 data set here. But for reasons of consistency when we apply the climatologically derived model in the variable climate, we use the monthly climatology derived from CRU TS 3.0 here.

NAO influence on mass balance variability

B. Marzeion and A. Nesje

| | |
|--------------------------|--------------|
| Title Page | |
| Abstract | Introduction |
| Conclusions | References |
| Tables | Figures |
| ◀ | ▶ |
| ◀ | ▶ |
| Back | Close |
| Full Screen / Esc | |
| Printer-friendly Version | |
| Interactive Discussion | |



TS 3.0 data set at the location of the glacier, gives the estimate of μ_{clim} . Figure 2, panel b, shows μ_{clim} for each of the glaciers considered in this study.

2.3 Model validation

The individually trained, and the mean model, depend on the measured mass balance values of the glaciers. In order to validate these models using the measured mass balances, we perform a cross validation of these two models (Michaelson, 1987, and Hofer et al., 2010, give a detailed description of cross validation applications in climate). The climatologically derived model is built independent of the measured mass balances, and can therefore directly be validated with the measurements.

2.3.1 Cross validation of the individually trained model

As in Marzeion et al. (2011), for each glacier in the region of interest with existing mass balance measurements, we first construct a time series of modeled annual mass balance MB_{model} , where each annual value is independent of the measured value of that year. This is achieved by performing the model optimization required for estimating the model parameters a_{opt} and μ_{opt} N times, where N is the number of mass balance measurements, leaving out of the optimization the years $k \pm t_{\text{lag}}$, where k is the year of the measurement, and t_{lag} is the length of autocorrelation in the measured mass balance time series. This yields the model parameters $a_{k,\text{cross}}$ and $\mu_{k,\text{cross}}$, which are then used to determine $\text{MB}_{k,\text{model,cross}}$. Figure 3 shows the standard deviation of $a_{k,\text{cross}}$ and $\mu_{k,\text{cross}}$ for all glacier with existing mass balance measurements, as a function of the number of mass balance measurements existing for that glacier. It becomes apparent that the robustness of the estimates of the model parameters increases with the number of mass balance measurements, and that a minimum number of measurements is necessary to achieve reliable estimates of $a_{k,\text{cross}}$ and $\mu_{k,\text{cross}}$. For this reason, we reject all glaciers with fewer than 12 years of mass balance measurements from all further analysis.

NAO influence on mass balance variability

B. Marzeion and A. Nesje

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Figure 4, panel d, shows the modeled versus measured mass balances (normalized) of the individually trained model for all glacier with more than 12 measured mass balances, panel a shows the correlation coefficient for each glacier. We also calculate the bias of the model, and the skill score

$$5 \quad SS = 1 - \frac{mse_{\text{model}}}{mse_{\text{ref}}}, \quad (7)$$

where mse_{model} is the mean square error of MB_{model} , and mse_{ref} is the mean square error of a reference model (in this case, we use the mean of the measured mass balances as the reference model). Figure 5 shows that the variability of the modeled time series is smaller than that of the measured mass balance time series. But still, there is considerable skill in the modeled time series (mean skill score 0.51), and the correlation between modeled and measured mass balances is reasonably high (mean value of 0.67). The results of the cross validation of the individually trained model are summarized in Table 1 (see Marzeion et al. (2011) for a more detailed description and discussion of the cross validation procedure).

15 We finally determine a_{opt} and μ_{opt} as the mean values of $a_{k,\text{cross}}$ and $\mu_{k,\text{cross}}$, and apply them in Eq. (1) to obtain the time series MB_{model} of the individually trained model over the entire period of data available in the CRU TS 3.0 data set. Figure 6, panel a, shows the result for an exemplary glacier, Ålfotbreen⁷ in Western Norway (black line), together with the measured mass balances (green line).

20 2.3.2 Cross validation of the mean model

In order to cross validate the mean model, we construct time series of MB_{model} for each glacier with more than 12 existing mass balance measurements, independent of the mass balance measurements of that glacier. This is achieved by determining

⁷Ålfotbreen was chosen as an example, because with respect to glaciers in Western Norway, it exhibits a typical relation to the NAO, and it has a long history of mass balance measurements.

NAO influence on mass balance variability

B. Marzeion and A. Nesje

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



$a_{\text{mean,cross}}$ and $\mu_{\text{mean,cross}}$ as the mean of a_{opt} and μ_{opt} of all the glaciers in the region, except for the glacier where the cross validation is being carried out.

Marzeion et al. (2011) show that if the Alps are considered as one region, the mean model yields reasonable results. Here, we define 5 different regions for determining the mean parameters: Northern Scandinavia, Southern Scandinavia, Scandinavia (i.e., Northern and Southern Scandinavia combined), the Alps, and Europe as one region containing all glaciers considered here. The outlines of the different regions are shown in Fig. 1. The results of the cross validation of the mean model are summarized in Table 1. If measured by correlation between modeled and measured mass balances, the performance of the mean model is higher than that of the individually trained model, independent of how the region is defined. The mean model works best for the Alps (mean correlation of 0.74), but the performance suffers only little when applied over all of Europe (mean correlation 0.3). The fact that the correlation improves for the mean model compared to the individually trained model can be understood as the result of a vastly increased data basis of the mean model compared to the individually trained model. The mean model also slightly improves the representation of variability of the mass balances (Fig. 5), but it exhibits a considerable bias (see Table 1). As a result, the skill score becomes negligible (except for the Alps). This implies that the model's results cannot be used in applications where absolute values of the mass balance are needed. But for correlation-based analyses, which do not depend on bias and the amplitude of the variability, the model results are applicable.

Figure 4, panel e, shows the modeled versus measured mass balances (normalized) of the mean model for all glacier with more than 12 measured mass balances (the mean being applied over all of Europe), panel b shows the correlation coefficient for each glacier.

We finally determine a_{mean} and μ_{mean} as the mean values of a_{opt} and μ_{opt} , and apply them in Eq. (1) to obtain the time series MB_{model} of the mean model over the entire period of data available in the CRU TS 3.0 data set. Figure 6, panel b, shows the result for Ålfotbreen (black line), together with the measured mass balances (green line).

NAO influence on mass balance variability

B. Marzeion and A. Nesje

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.3.3 Validation of the climatologically derived model

Since the climatologically derived model is built independent of mass balance measurements, it can be validated directly using the measured mass balances of glaciers where available. This would be possible independent of the number of mass balance measurements, but in order to be able to directly compare the climatologically derived model with the individually trained and the mean model, we here show the validation based only on those glaciers for which more than 12 measured mass balances are available. Figure 4, panel f, shows the modeled versus measured mass balances (normalized) of the climatologically derived model, panel c shows the correlation coefficient for each glacier. From Fig. 5 it is apparent that a weakness of the climatologically derived model is an underestimation of the variability of the mass balances. But the summary of the validation (see Table 2) shows that the climatologically derived model shows an improvement over the individually trained and mean models in terms of correlation, and over the mean model in terms of skill score.

Figure 6, panel c, shows the result of the climatologically derived model for Ålfotbreen (black line), together with the measured mass balances (green line).

3 Results

The (cross) validation of the three different models has shown that for a correlation-based analysis, the climatologically derived model provides the most reliable results. In the following, we therefore discard of the individually trained and mean models, and only apply the climatologically derived model.

3.1 Decomposition of the forcing fields

In order to bring out the influence of the NAO on the glacier mass balances in Europe, we first determine T_{glacier} as the temperature at the glacier terminus using Eq. (5), and

NAO influence on mass balance variability

B. Marzeion and A. Nesje

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



P_{glacier} as the estimated solid precipitation onto the glacier surface from CRU TS 3.0, and then decompose T_{glacier} and P_{glacier}

$$T_{\text{glacier}} = T_{\text{glacier}_{\text{clim}}} + T_{\text{glacier}_{\text{NAO}}} + T'_{\text{glacier}}, \quad (8)$$

$$P_{\text{glacier}} = P_{\text{glacier}_{\text{clim}}} + P_{\text{glacier}_{\text{NAO}}} + P'_{\text{glacier}}, \quad (9)$$

where the subscript $_{\text{clim}}$ denotes the monthly mean climatological values, the subscript $_{\text{NAO}}$ denotes monthly anomalies associated with NAO variability, and the prime denotes all other anomalies. In order to determine $T_{\text{glacier}_{\text{NAO}}}$ and $P_{\text{glacier}_{\text{NAO}}}$, we perform a regression of T_{CRU} and P_{CRU} on the monthly NAO index (Hurrell, 1995b)⁸, individually for each month of the year. Generally speaking, the NAO influence is strongest in winter, with positive NAO anomalies leading to above-average temperatures all over Europe, while the precipitation anomalies tend to be positive in Northern Europe (with exceptions in the northwestern part of Northern Norway), and negative around the Mediterranean (Fig. 1, see Hurrell et al., 2003, for a detailed discussion of the impacts of the NAO on European climate).

Figure 7 shows exemplary results of the decomposition for the location of Ålfotbreen, for the hydrological years 1995 and 1996 (vertical, light blue bar in Fig. 6). The comparison of panel a with panels d and g shows that while the NAO variability does not show a strong seasonality, its impacts on temperature and precipitation are only evident during winter.

3.2 Constructing the mass balance time series

For each glacier we then construct a total of 5 time series of mass balance variability:

1. $MB_{\text{clim}} = P_{\text{glacier}_{\text{clim}}} - \mu_{\text{clim}}(\max(0, T_{\text{glacier}_{\text{clim}}}))$ (10)

is the monthly, climatological mass balance of the glacier, where $\sum_{i=1}^{12} MB_{i, \text{clim}} = 0$, i indicating the months of the hydrological year, follows directly from Eq. (7).

⁸available at <http://www.cgd.ucar.edu/cas/jhurrell/indices.html>

NAO influence on mass balance variability

B. Marzeion and A. Nesje

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



NAO influence on mass balance variability

B. Marzeion and A. Nesje

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



MB_{clim} is shown in Fig. 7, panel h.

$$2. MB_{anom} = [P_{glacier} - \mu_{clim}(\max(0, T_{glacier}))] - MB_{clim} \quad (11)$$

is the monthly deviation of the mass balance from the climatological mass balance, shown in Fig. 7, panel i.

$$5. 3. MB_{NAO} = \left[\left(P_{glacier_{NAO}} + P_{glacier_{clim}} \right) \right. \quad (12)$$

$$\left. - \left(\mu_{clim}(\max(0, T_{glacier_{NAO}} + T_{glacier_{clim}})) \right) \right] - MB_{clim}$$

is the monthly deviation of the mass balance from the climatological mass balance associated with NAO variability, shown in Fig. 7, panel j.

$$4. MB(T_{NAO}, P_{clim}) = \left[P_{glacier_{clim}} - \left(\mu_{clim}(\max(0, T_{glacier_{NAO}} + T_{glacier_{clim}})) \right) \right] - MB_{clim} \quad (13)$$

is the monthly deviation of the mass balance from the climatological mass balance associated with only NAO temperature variability, shown in Fig. 7, panel k.

$$5. MB(T_{clim}, P_{NAO}) = \left[\left(P_{glacier_{NAO}} + P_{glacier_{clim}} \right) - \quad (14)$$

$$\left(\mu_{clim}(\max(0, T_{glacier_{clim}})) \right) \right] - MB_{clim}$$

is the monthly deviation of the mass balance from the climatological mass balance associated with only NAO precipitation variability, shown in Fig. 7, panel l.

3.3 The spatial patterns of NAO influence on mass balance variability

Figure 8 shows the correlation between monthly NAO index, and monthly modeled mass balance anomalies MB_{anom} .⁹ The strongest relationship between NAO and mass balance becomes apparent in Western Norway (confirming the results of Nesje et al. (2000) and Reichert et al. (2001), who investigated single glaciers in these regions). In Northern Scandinavia, and in the Alps, the mass balances are anti-correlated with the NAO, and in the Alps the connection is generally weaker. In the case of the Alps the anti-correlation is significant only in the western part (here, Reichert et al. (2001) also find an anti-correlation for Rhonegletscher, but they find the NAO influence to be stronger), and south of the main ridge. Panel b reveals that there is hardly any relation between the correlation and the altitude of the glacier terminus within each region.

As a measure of how strong the NAO influence is on the mass balance, we show correlations between MB_{anom} and MB_{NAO} in Fig. 9. As is to be expected from Fig. 8, there is a minimum of NAO-driven mass balance variability in mid-Scandinavia, the influence of the NAO is strongest in Western Norway and for the glaciers in the very north of Scandinavia. Within the Alps, the NAO influence is generally smaller, but interestingly, slightly grows towards the east.

Since there is a distinct seasonality in the connection between NAO and temperature and precipitation anomalies, it is instructive to look into the seasonality of the mass balances' response to NAO forcing. Figure 10, panels a and b show the correlation between monthly NAO index, and monthly modeled mass balance anomalies MB_{anom} (i.e., the same as Fig. 8), but only during the winter months December, January, and February, when the influence of the NAO on precipitation and temperature in Europe is strongest. Panels c and d show the same, but for the remainder of the year.

Generally speaking, the connection between the winter mass balance and winter NAO is – perhaps not surprisingly – stronger than the connection over the entire year.

⁹Note in this and the subsequent figures that the glaciers with existing mass balances, where the model validation takes place, are well distributed in the whole sample of WGI-XF glaciers.

But the distinction between winter and rest of the year also allows for an insight into the mechanism of NAO influence. During winter, even a strong positive NAO, with corresponding warm temperature anomalies does not lift air temperatures at the glaciers' termini above freezing. The winter signal is therefore predominantly a precipitation signal, and it becomes apparent that the anti correlation between NAO and mass balance in Northern Scandinavia is caused predominantly by warmer summer temperatures. In Western Norway, the relation is more ambiguous: both winter and summer signal contribute to the positive correlation, only in the northern part of Southern Norway the warm summer temperatures' influence is apparent.

In the Alps, a clear distinction is possible between the western and eastern part: In the west, negative winter accumulation and warmer summer temperatures contribute to the anti correlation with the NAO alike. In the eastern part, however, during the winter a positive anomaly becomes apparent. This is the region that does not show a significant correlation with the NAO when the entire year is considered. The reason is apparent in panel c: the positive signal caused by winter accumulation is canceled by a negative signal during the rest of the year, when warmer temperatures prevail. Since the positive precipitation signal of the NAO is limited to the northern boundary of the Alps (see also Fig. 1), the negative signal shows through in the Southern Alps when the whole year is considered.

Another approach to understanding how the relative influences of temperature and precipitation set up the entire signal is followed in Fig. 11, showing the correlation between MB_{anom} and $MB(T_{NAO}, P_{clim})$ (panels a and b), and between MB_{anom} and $MB(T_{clim}, P_{NAO})$ (panels c and d). The mass balance anomalies created by NAO-related temperature anomalies correlate with the full mass balance anomalies everywhere, but the correlations insignificant (and even negative for a few glaciers) in Western Norway. I.e., a warm NAO signal contributes negative mass balance anomalies everywhere, but in Western Norway, the positive winter mass balance anomalies take the lead.

NAO influence on mass balance variability

B. Marzeion and A. Nesje

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

4 Discussion

One obvious difference between the climatologically derived and the individually trained (and mean) models is that we do not estimate the increase of precipitation with altitude (or aeolian transport and avalanching, i.e. the parameter a in the individually trained and mean models) in the climatologically derived model individually. One strong benefit of the climatologically derived model is the independence on any mass balance measurements, and the resulting applicability in regions that have fewer mass balance measurements than are necessary to obtain robust parameter estimates for the individually trained (and mean) models. A feasible way of obtaining an estimate of precipitation increase with altitude would be the determination of a vertical precipitation gradient from the CRU CL 2.0 data set, analogous to the determination of the temperature lapse rates. However, as mentioned above, this approach requires that the vertical gradients in the data set are large compared to the horizontal gradients. For precipitation, this is not always the case: especially in Scandinavia, the maximum of precipitation is rather close to the coast, and not over the highest elevation. Calculating precipitation lapse rates analogous to the temperature lapse rates therefore results in negative vertical gradients (i.e., less precipitation with increasing altitude) over large parts of Scandinavia, which is not realistic. For temperature, the problem of mixing horizontal with vertical gradients is less significant, as there is a strong dependence of temperature on altitude by default, and the horizontal temperature gradients are small compared to this default vertical gradient. We tested the validity of this approach: Prescribing a constant vertical lapse rate in temperature decreases the model's performance (independent of the value of the prescribed lapse rate). If we apply a variable (derived analogous to temperature) precipitation lapse rate, the performance slightly suffers in large parts of Scandinavia, and is hardly affected at all in the Alps.

The main deficit of the climatologically derived model compared to the individually trained model is the relatively strong bias (the same is true for the mean model), leading to the weak skill scores. While the bias does not affect our correlation-based analysis,

NAO influence on mass balance variability

B. Marzeion and A. Nesje

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



5 it does prevent the model from being applied to reconstruct absolute values of mass balance. The construction of the climatologically derived model implies that over the entire CRU TS 3.0 period, the mean mass balance is very close to zero (it is not exactly zero, since the interannual variability in temperature and precipitation is not evenly distributed over the months of the year). The bias is therefore a measure of how far away the glaciers are from equilibrium with the climatological forcing.

10 It is questionable whether the slight advantage of the climatologically derived model over the mean model is real, or an artifact of the different validation procedures: The dependence of the mean model on measured mass balances necessitates a cross validation, which implies that for the validation, information has to be withheld from the model, leading to a slight (and principally not determinable) decrease of the model's measured performance.

15 In comparison with the results from Marzeion et al. (2011), the performance of the mean model driven by CRU data is weaker than when driven with HISTALP data. This is to be expected, as the spatial resolution of the HISTALP data is six times that of the CRU TS 3.0 data. In this respect, we find the decrease of performance remarkably small (skill score in the Alps drops from 0.34 for the HISTALP data to 0.22 for the CRU TS 3.0 data, and correlation drops from 0.82 for the HISTALP data to 0.74 for the CRU TS 3.0 data).

20 5 Conclusions

25 Qualitatively, our results closely resemble the findings of Nesje et al. (2000) and Reichert et al. (2001), who investigate the impact of NAO variability on selected glaciers in Scandinavia and the Alps. While the correlations values we find are less pronounced, we are able to show that there is a regional coherence in the mechanisms and impact of NAO variability on glacier mass balance, which allows for the classification of all European glaciers into four regions:

NAO influence on mass balance variability

B. Marzeion and A. Nesje

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



NAO influence on mass balance variability

B. Marzeion and A. Nesje

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1. In Southwestern Scandinavia, the variability of winter precipitation leads to a correlation between glacier mass balances and NAO, even though temperature anomalies work against this (in line with the findings of Reichert et al., 2001). There is a gradient within the region, with the correlation getting weaker (and even negative) towards the Northeast (in line with the results of Nesje et al., 2000).
2. In Northern Scandinavia, temperature anomalies cause an anti-correlation between mass balance and NAO. The anti-correlation gets stronger towards the North.
3. In the Western Alps, winter precipitation and temperature anomalies cause a (in our case, weak) anti-correlation between mass balances and NAO (in line with the results of Reichert et al., 2001).
4. In the Eastern Alps, along the northern boundary of the Alps winter precipitation works towards a weak positive correlation between NAO and mass balances, but this effect is balanced by the effect of temperature. In the more central and southern parts, the lack of a precipitation signal leads to a weak anti-correlation between mass balances and NAO.

Acknowledgements. We wish to thank Graham Cogley for providing the WGI-XF and global mass balance datasets, and Regine Hock for discussions on the model set up. This is publication no. A383 from the Bjerknes Centre for Climate Research.

References

- Auer, I., Böhm, R., Jurkovic, A., Lipa, W., Orlik, A., Potzmann, R., Schöner, W., Ungersböck, M., Matulla, C., Briffa, K., Jones, P., Efthymiadis, D., Brunetti, M., Nanni, T., Maugeri, M., Mercalli, L., Mestre, O., Moisselin, J.-M., Begert, M., Müller-Westermeier, G., Kveton, V., Bochnicek, O., Stastny, P., Lapin, M., Szalai, S., Szentimrey, T., Cegnar, T., Dolinar, M., Gajic-Capka, M., Zaninovic, K., Majstorovic, Z., and Nieplova, E.: HISTALP—historical instrumental climatological surface time series of the Greater Alpine Region, *Int. J. Climatol.*, 27, 17–46, doi:10.1002/joc.1377, 2007. 4
- Cogley, J. G.: Extended Format for the World Glacier Inventory, Tech. Rep. Trent Technical Note 2005-1, Department of Geography, Trent University, Peterborough, Ontario, Canada, revised September 2008, 2005. 4
- Cogley, J. G.: Geodetic and direct mass-balance measurements: comparison and joint analysis, *Ann. Glaciol.*, 50, 96–100, doi:10.3189/172756409787769744, 2009. 4
- Hock, R.: Temperature index melt modelling in mountain areas, *J. Hydrol.*, 282, 104–115, doi:10.1016/S0022-1694(03)00257-9, 2003.
- Hofer, M., Mölg, T., Marzeion, B., and Kaser, G.: Empirical-statistical downscaling of reanalysis data to high-resolution air temperature and specific humidity above a glacier surface (Cordillera Blanca, Peru), *J. Geophys. Res.*, 115, D12120, doi:10.1029/2009JD012556, 2010. 9
- Hurrell, J.: Decadal trends in the North Atlantic Oscillation: regional temperatures and precipitation, *Science*, 269, 676–679, 1995a. 3
- Hurrell, J. W.: NAO Index Data provided by the Climate Analysis Section, NCAR, Boulder, USA, available at: <http://www.cgd.ucar.edu/cas/jhurrell/indices.html>, 1995b (last access: 28 December 2011). 2, 13
- Hurrell, J. W., Kushnir, Y., Ottersen, G., and Visbeck, M., eds.: The North Atlantic oscillation: climatic significance and environmental impact, Geophysical Monograph, American Geophysical Union, Washington, DC, USA, 2003. 2, 3, 13
- Huss, M., Hock, R., Bauder, A., and Funk, M.: 100-year mass changes in the Swiss Alps linked to the Atlantic Multidecadal Oscillation, *Geophys. Res. Lett.*, 37, L10501, doi:10.1029/2010GL042616, 2010. 3
- Kaser, G., Grosshauser, M., and Marzeion, B.: Contribution potential of glaciers to water availability in different climate regimes, *P. Natl. Acad. Sci. USA*, 107, 20223–20227,

NAO influence on mass balance variability

B. Marzeion and A. Nesje

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**NAO influence on
mass balance
variability**B. Marzeion and A. Nesje

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

doi:10.1073/pnas.1008162107, 2010. 6

Kuhn, M.: Micro-meteorological conditions for snow melt, *J. Glaciol.*, 33, 24–26, 1987. 6

Marzeion, B., Hofer, M., Jarosch, A. H., Kaser, G., and Mölg, T.: A minimal model for re-
constructing interannual mass balance variability of glaciers in the European Alps, *The*
5 *Cryosphere Discuss.*, 5, 2799–2839, doi:10.5194/tcd-5-2799-2011, 2011. 3, 4, 5, 6, 9, 10,
11, 18

Michaelsen, J.: Cross-validation in statistical climate forecast models, *J. Appl. Meteorol.*, 26,
1589–1600, 1987. 9

Mitchell, T. D. and Jones, P. D.: An improved method of constructing a database of monthly
10 climate observations and associated high-resolution grids, *Int. J. Climatol.*, 25, 693–712,
doi:10.1002/joc.1181, 2005. 4, 7

Mölg, T. and Hardy, D. R.: Ablation and associated energy balance of a horizontal glacier
surface on Kilimanjaro, *J. Geophys. Res.*, 109, D16104, doi:10.1029/2003JD004338, 2004.
6

15 Nesje, A., Lie, Ø., and Dahl, S.: Is the North Atlantic Oscillation reflected in Scandinavian
glacier mass balance records?, *J. Quaternary Sci.*, 15, 587–601, 2000. 2, 3, 15, 18, 19

New, M., Lister, D., Hulme, M., and Makin, I.: A high-resolution data set of surface climate over
global land areas, *Clim. Res.*, 21, 1–25, 2002. 7

Oerlemans, J.: Analysis of a 3 year meteorological record from the ablation zone of
20 Morteratschgletscher, Switzerland: energy and mass balance, *J. Glaciol.*, 46, 571–579,
doi:10.3189/172756500781832657, 2000. 6

Ohmura, A.: Physical basis for the temperature-based melt-index method, *J. Appl. Meteorol.*,
40, 753–761, 2001. 6

Reichert, B. K., Bengtsson, L., and Oerlemans, J.: Midlatitude forcing mechanisms for glacier
25 mass balance investigated using general circulation models, *J. Climate*, 14, 3767–3784,
2001. 3, 15, 18, 19

Scherrer, S., Appenzeller, C., and Laternser, M.: Trends in Swiss Alpine snow days:
the role of local- and large-scale climate variability, *Geophys. Res. Lett.*, 31, L13215,
doi:10.1029/2004GL020255, 2004. 3

30 Scart, J., Hock, R., and Six, D.: Glacier melt, air temperature, and energy balance in different
climates: the Bolivian Tropics, the French Alps, and Northern Sweden, *J. Geophys. Res.*,
113, D24113, doi:10.1029/2008JD010406, 2008. 6

Six, D., Reynaud, L., and Letréguilly, A.: Bilans de masse des glaciers alpins et scandinaves,

leurs relations avec l'oscillation du climat de l'Atlantique nord: Alpine and Scandinavian glaciers mass balances, their relations with the North Atlantic Oscillation, C.R. Acad. Sci. II A, 333, 693–698, 2001. 3

Steiner, D., Pauling, A., Nussbaumer, S., Nesje, A., Luterbacher, J., Wanner, H., and Zumbühl, H.: Sensitivity of European glaciers to precipitation and temperature – two case studies, Climatic Change, 90, 413–441, 2008. 3

Visbeck, M. H., Hurrell, J. W., Polvani, L., and Cullen, H. M.: The North Atlantic Oscillation: past, present, and future, P. Natl. Acad. Sci. USA, 98, 12876–12877, 2001. 2

Wanner, H., Brönnimann, S., Casty, C., Gyalistras, D., Luterbacher, J., Schmutz, C., Stephenson, D. B., and Xoplaki, E.: North Atlantic Oscillation – concepts and studies, Surv. Geophys., 22, 321–381, 2001. 2, 3

TCD

6, 1–35, 2012

NAO influence on mass balance variability

B. Marzeion and A. Nesje

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



NAO influence on mass balance variability

B. Marzeion and A. Nesje

Table 1. Summary of parameters and results of the individually trained and the mean model. All numbers given are the mean values obtained during the cross validation procedure, with their standard deviation. Negative skill scores imply zero skill, and have therefore been replaced with zero for the calculation of the mean and standard deviation of the skill scores.

| | No. Glaciers | No. MB | a_{opt} | μ_{opt} | r | SS | Bias [mm] |
|---------------------------------------|--------------|--------|-----------------|--------------|-----------------|-----------------|-----------------|
| individually trained | 71 | 1579 | 2.10 ± 1.39 | 71 ± 150 | 0.50 ± 0.45 | 0.46 ± 0.24 | -124 ± 1484 |
| individually trained (no. MB > 12) | 39 | 1339 | 2.10 ± 1.54 | 107 ± 51 | 0.67 ± 0.15 | 0.51 ± 0.14 | 0 ± 19 |
| Northern Scandinavia (no. MB > 12) | 7 | 208 | 4.10 ± 1.77 | 116 ± 60 | 0.73 ± 0.10 | 0.09 ± 0.15 | -543 ± 1714 |
| Southern Scandinavia (no. MB > 12) | 10 | 372 | 2.95 ± 0.78 | 76 ± 26 | 0.72 ± 0.15 | 0.12 ± 0.25 | -544 ± 2627 |
| Scandinavia (no. MB > 12) | 17 | 580 | 3.42 ± 1.36 | 93 ± 46 | 0.73 ± 0.13 | 0.14 ± 0.24 | -587 ± 2375 |
| Alps (no. MB > 12) | 22 | 759 | 1.08 ± 0.59 | 118 ± 52 | 0.74 ± 0.11 | 0.22 ± 0.31 | -506 ± 2393 |
| all (no. MB > 12) | 39 | 1339 | 2.10 ± 1.54 | 107 ± 51 | 0.73 ± 0.14 | 0.07 ± 0.16 | -258 ± 3039 |

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



NAO influence on mass balance variability

B. Marzeion and A. Nesje

Table 2. Summary of the results of the climatologically derived model. All numbers given are the mean values with their standard deviation.

| | r | SS | Bias [mm] |
|----------------|-----------------|-----------------|----------------|
| N. Scandinavia | 0.75 ± 0.09 | 0.11 ± 0.10 | 35 ± 579 |
| S. Scandinavia | 0.77 ± 0.07 | 0.24 ± 0.17 | -237 ± 413 |
| Scandinavia | 0.76 ± 0.08 | 0.19 ± 0.16 | -125 ± 491 |
| Alps | 0.74 ± 0.11 | 0.31 ± 0.18 | 62 ± 284 |
| all | 0.75 ± 0.10 | 0.26 ± 0.18 | -19 ± 394 |

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 1. Maps of regression coefficients of monthly total precipitation (**a**) and monthly mean temperature (**b**) on the NAO index (shading), values where the correlation between NAO index and precipitation/temperature is below the 99% confidence interval have been omitted. Shown are the regression coefficients of the month with the maximum NAO effect, which are generally winter months. Gray background shading: Topography of the CRU TS 3.0 data. Black markers: locations of the glaciers with more than 12 mass balance measurements. Green dots: locations of the glaciers contained in the WGI-XF data base. The boxes indicate the locations of the regions referred to in the text, figures, and tables.

NAO influence on mass balance variability

B. Marzeion and A. Nesje

| | |
|--------------------------|--------------|
| Title Page | |
| Abstract | Introduction |
| Conclusions | References |
| Tables | Figures |
| ◀ | ▶ |
| ◀ | ▶ |
| Back | Close |
| Full Screen / Esc | |
| Printer-friendly Version | |
| Interactive Discussion | |






Fig. 2. Maps of the model parameters of the climatologically derived model: **(a)** temperature lapse rates. Values where the correlation between temperature and elevation is below the 95% confidence interval have been omitted. **(b)** μ_{clim} . Markers with black circle indicate the results of the glaciers with more than 12 existing measurements of annual mass balance. Gray background shading: Topography of the CRU TS 3.0 data.

NAO influence on mass balance variability

B. Marzeion and A. Nesje

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 3. Standard deviation of $a_{k,\text{cross}}$ **(a)** and $\mu_{k,\text{cross}}$ **(b)** as a function of the number of mass balance measurements. Black dots are the values of glaciers accepted into the final set, gray dots are those of the glaciers rejected from the final set. The gray vertical line indicates the minimum number of mass balance measurements necessary to obtain robust results for $a_{\text{optimized}}$ and $\mu_{\text{optimized}}$.

NAO influence on mass balance variability

B. Marzeion and A. Nesje

| | |
|--------------------------|--------------|
| Title Page | |
| Abstract | Introduction |
| Conclusions | References |
| Tables | Figures |
| ◀ | ▶ |
| ◀ | ▶ |
| Back | Close |
| Full Screen / Esc | |
| Printer-friendly Version | |
| Interactive Discussion | |



NAO influence on mass balance variability

B. Marzeion and A. Nesje




Fig. 4. Upper row: correlation coefficients of the individually trained **(a)**, mean **(b)**, and climatologically derived model **(c)** for all glaciers with more than 12 measured annual mass balances. See Tables 1 (for a and b) and 2 (for c) for the mean values and standard deviation. Gray background shading: Topography of the CRU TS 3.0 data. Lower row: Modeled versus measured normalized annual mass balances of the individually trained model **(d)** and the mean model **(e)** obtained by cross validation, and of the climatologically derived model **(f)**, for all glaciers with more than 12 measured mass balances.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


**NAO influence on
mass balance
variability**

B. Marzeion and A. Nesje




Fig. 5. Standard deviation of the annual mass balance plotted against the area of the glaciers. Green dots show MB_{measured} , black dots show the values of the individually trained model, blue dots show the results of the mean model, and red dots show the results of the climatologically derived model. Horizontal lines show the respective means.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



NAO influence on mass balance variability

B. Marzeion and A. Nesje




Fig. 6. Timeseries of the reconstruction for Ålfotbreen. **(a)** green: standardized MB_{measured} , black: standardized MB_{model} , light gray shading: $\pm 2 \cdot \text{rmse}$, dark gray shading: $\pm 1 \cdot \text{rmse}$. **(b)** green: standardized MB_{measured} , black: standardized $MB_{\text{mean model}}$, light gray shading: $\pm 2 \cdot \text{rmse}$, dark gray shading: $\pm 1 \cdot \text{rmse}$. **(c)** green: standardized MB_{measured} , black: standardized $MB_{\text{clim model}}$, light gray shading: $\pm 2 \cdot \text{rmse}$, dark gray shading: $\pm 1 \cdot \text{rmse}$. **(d)** annual temperature anomaly at the location of the glacier. **(e)** annual precipitation anomaly at the location of the glacier, light blue: total precipitation, dark blue: estimated solid precipitation. **(f)** winter mean of the NAO index. The vertical light blue bar indicates the two years show in detail in Fig. 7.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



NAO influence on mass balance variability

B. Marzeion and A. Nesje




Fig. 7. Exemplary detail of the monthly time series of the reconstruction for Ålfotbreen for the hydrological years 1995 and 1996. **(a)** NAO index. **(b)** Temperature climatology, $T_{\text{glacier}_{\text{clim}}}$. **(c)** Temperature anomaly, $T_{\text{glacier}_{\text{NAO}}} + T'_{\text{glacier}}$. **(d)** Temperature anomaly associated with NAO, $T_{\text{glacier}_{\text{NAO}}}$. **(e)** Precipitation climatology, $P_{\text{glacier}_{\text{clim}}}$. **(f)** Precipitation anomaly, $P_{\text{glacier}_{\text{NAO}}} + P'_{\text{glacier}}$. **(g)** Precipitation anomaly associated with NAO, $P_{\text{glacier}_{\text{NAO}}}$. (In panels e-g: light blue: total precipitation, dark blue: estimated solid precipitation) **(h)** Mass balance climatology, MB_{clim} . **(i)** Mass balance anomaly, MB_{anom} . **(j)** Mass balance anomaly associated with NAO variability in temperature and precipitation, MB_{NAO} . **(k)** Mass balance anomaly associated with NAO temperature anomaly, $MB(T_{\text{NAO}} \cdot P_{\text{clim}})$. **(l)** Mass balance anomaly associated with NAO precipitation anomaly, $MB(T_{\text{clim}} \cdot P_{\text{NAO}})$.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 8. Correlation between modeled monthly mass balance anomalies, i.e. MB_{anom} , and monthly NAO index. Markers with black circle **(a)** and circle markers **(b)** indicate the results of the glaciers with more than 12 existing measurements of annual mass balance. Values below the 95 % confidence interval have been omitted. Gray background shading: Topography of the CRU TS 3.0 data. In panel **(b)** black is for glaciers in the Alps, red for glaciers in Southern Scandinavia, blue for glaciers in Northern Scandinavia.

NAO influence on mass balance variability

B. Marzeion and A. Nesje

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 9. Correlation between modeled monthly mass balance anomalies obtained using the full variability in the forcing, and modeled monthly mass balance anomalies obtained using only NAO-related variability in the forcing, i.e. between MB_{anom} and MB_{NAO} . Markers with black circle **(a)** and circle markers **(b)** indicate the results of the glaciers with more than 12 existing measurements of annual mass balance. Values below the 95 % confidence interval have been omitted. Gray background shading: Topography of the CRU TS 3.0 data. In panel **(b)** black is for glaciers in the Alps, red for glaciers in Southern Scandinavia, blue for glaciers in Northern Scandinavia.

NAO influence on mass balance variability

B. Marzeion and A. Nesje

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 10. Mean of the monthly correlations between modeled mass balance anomalies, i.e. MB_{anom} , and NAO index during December, January, and February (**a,b**), and during March to November (**c,d**). Markers with black circle (**a,c**) and circle markers (**b,d**) indicate the results of the glaciers with more than 12 existing measurements of annual mass balance. Values below the 95 % confidence interval have been omitted from the calculation of the mean. Gray background shading: Topography of the CRU TS 3.0 data. In panels (**b**) and (**d**) black is for glaciers in the Alps, red for glaciers in Southern Scandinavia, blue for glaciers in Northern Scandinavia. Note that the correlation values close to zero are the result of calculating the mean of significant, non-zero correlations.

NAO influence on mass balance variability

B. Marzeion and A. Nesje

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



NAO influence on mass balance variability

B. Marzeion and A. Nesje




Fig. 11. Correlation between modeled monthly mass balance anomalies obtained using the full variability in the forcing, and modeled monthly mass balance anomalies obtained using only NAO-related temperature variability and climatological precipitation, i.e. between MB_{anom} and $MB(T_{NAO}, P_{clim})$ (a,b); and correlation between modeled monthly mass balance anomalies obtained using the full variability in the forcing, and modeled monthly mass balance anomalies obtained using only NAO-related precipitation variability and climatological temperature, i.e. between MB_{anom} and $MB(T_{clim}, P_{NAO})$ (c,d). Markers with black circle (a,c) and circle markers (b,d) indicate the results of the glaciers with more than 12 existing measurements of annual mass balance. Values below the 95% confidence interval have been omitted. Gray background shading: Topography of the CRU TS 3.0 data. In panels (b) and (d), black is for glaciers in the Alps, red for glaciers in Southern Scandinavia, blue for glaciers in Northern Scandinavia.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

