

Sea ice inertial
oscillations

F. Gimbert et al.

This discussion paper is/has been under review for the journal The Cryosphere (TC).
Please refer to the corresponding final paper in TC if available.

Sea ice inertial oscillation magnitudes in the Arctic basin

F. Gimbert^{1,3}, D. Marsan², J. Weiss³, N. C. Jourdain⁴, and B. Barnier⁴

¹ISTerre, UMR – CNRS, UMR 5275, Université Joseph Fourier, Grenoble, France

²ISTerre, UMR – CNRS, Université de Savoie, Le Bourget du Lac, France

³LGGE, UMR – CNRS, Université Joseph Fourier, Grenoble, France

⁴LEGI, UMR – CNRS, Université Joseph Fourier, Grenoble, France

Received: 30 April 2012 – Accepted: 26 May 2012 – Published: 18 June 2012

Correspondence to: F. Gimbert (florent.gimbert@ujf-grenoble.fr)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

An original method to quantify the amplitude of inertial motion of oceanic and ice drifters, through the introduction of a non-dimensional parameter M defined from a spectral analysis, is presented. A strong seasonal dependence of the magnitude of sea ice inertial oscillations is revealed, in agreement with the corresponding annual cycles of sea ice extent, concentration, thickness, advection velocity, and deformation rates. The spatial pattern of the magnitude of the sea ice inertial oscillations over the Arctic basin is also in agreement with the sea ice thickness and concentration patterns. This argues for a strong link between the magnitude of inertial motion on one hand, the dissipation of energy through mechanical processes, and the cohesiveness of the cover on the other hand. Finally, a significant pluri-annual evolution towards greater magnitudes of inertial oscillations in recent years, in both summer and winter, is reported, thus concomitant with reduced sea ice thickness, concentration and spatial extent.

1 Introduction

The spectacular evolution of the Arctic sea ice cover over the last few decades is not restricted to the shrinking of the ice extent (Comiso et al., 2008; Stroeve et al., 2008), its thinning (Rothrock et al., 2008; Kwok and Rothrock, 2009), or, consequently, a continued decline of the ice volume (Lindsay et al., 2009). Kinematics is affected as well, and its evolution plays a central role in the changes currently taking place in the Arctic Ocean. As observed from buoy drift data, the sea ice mean speed over the Arctic increased at a rate of 9 % per decade from 1979 to 2007, whereas the mean deformation rate increased by more than 50 % per decade over the same period (Rampal et al., 2009). These two aspects of recent sea ice evolution, i.e. strong decline in terms of ice extent and thickness, and accelerated kinematics, are strongly coupled within the albedo feedback loop. Increasing deformation means increasing fracturing, hence more lead opening and a decreasing albedo (Zhang et al., 2000). As a result, ocean

TCD

6, 2179–2220, 2012

Sea ice inertial oscillations

F. Gimbert et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Sea ice inertial oscillations

F. Gimbert et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



warming, in turn, favours sea ice thinning in summer and delays refreezing in early winter, i.e. strengthens sea ice decline. This thinning should decrease the mechanical strength, therefore allowing even more fracturing, hence larger speed and deformation. A consequence is the acceleration of the export of sea ice through Fram strait with a significant impact on sea ice mass balance (Rampal et al., 2009, 2011; Haas et al., 2008), and ice age (Nghiem et al., 2007). Moreover, sea ice mechanical weakening decreases the likelihood of arch formation along Nares strait, therefore allowing old, thick ice to be exported through this strait (Kwok et al., 2010).

The principle of an intimate link between the ice state (concentration, thickness) in one hand, and the mechanical behaviour of the cover and its strength on the other hand, although rather intuitive, needs however to be quantified more precisely. Beyond the trivial reduction of strength in proportion to the thinning of an ice plate, one can expect a more complex effect of the compactness of the ice cover on its average strength as, e.g., in granular media (Rajchenbach, 2000; Aranson and Tsimring, 2006). This is the combination of these two effects that we aim to explore here. Ideally, one would like to check whether the sea-ice cover responds differently from year to year to the same mechanical forcing. We here propose to tackle this problem by analysing its response, as deduced from ice drifter trajectories, to the specific inertial forcing. The effect of the Coriolis force on geophysical fluids dynamics has been studied for more than a century. Interestingly, the first studies of oceanic inertial oscillations (Ekman, 1905) were prompted by the observations of Nansen, made during the Fram's journey along the Transpolar drift, that sea ice was moving with a 20° – 40° angle to the right of the wind direction (Nansen, 1902). Indeed, as the Coriolis force acts perpendicularly to the particle velocity, it induces a deviation of the trajectory to the right in the Northern Hemisphere. This deviation generates inertial oscillations, characterized by the frequency

$$f_0 = 2 \sin \phi \text{ cycles day}^{-1} \quad (1)$$

where ϕ is the latitude, i.e. close to a semi-diurnal frequency ($2 \text{ cycles day}^{-1}$) in the Arctic.

Within the ice-free ocean, these oscillations are triggered by wind forcing events such as storms or moving fronts (Price, 1983; Gill, 1984; D'Asaro et al., 1995). These events generate inertial oscillations within a few hours, which then decrease progressively with a characteristic time scale of a few days (Park et al., 2009). This damping is essentially due to the internal friction within the Ekman ocean layer, as well as to the radiation of inertial waves toward the thermocline.

On ice-covered oceans, we expect other processes to strengthen the damping of these oscillations, such as friction at the ice/ocean interface or, more importantly, the internal ice stresses resulting from solid mechanical interactions within the ice cover such as fracturing, friction between adjacent floes and shearing of leads, or crushing during convergent deformation and ridge formation, i.e. the "ice internal friction" (Lepparanta, 2004; Colony and Thorndike, 1980). These terms are apparent in the momentum balance of sea ice dynamics:

$$\frac{DU_i}{Dt} + (\Omega \mathbf{k} \wedge \mathbf{U}_i) = \frac{1}{\rho_i} \nabla \cdot \boldsymbol{\sigma} + \frac{1}{\rho_i h_i} (\boldsymbol{\tau}_a + \boldsymbol{\tau}_w) \quad (2)$$

where $D/Dt = \partial/\partial t + \mathbf{U}_i \cdot \nabla$ is the Lagrangian time derivative, ρ_i the ice density, h_i the ice thickness, \mathbf{U}_i the ice velocity vector expressed in Cartesian coordinates, $\boldsymbol{\sigma}$ the internal stress tensor, $\boldsymbol{\tau}_a$ and $\boldsymbol{\tau}_w$ respectively the wind and oceanic "stresses" (forces per unit ice area). The last term on the left hand side of Eq. (2) is the Coriolis force, with $\Omega = 2\pi/24 \text{ rad h}^{-1} = 7.10^{-5} \text{ rads}^{-1}$ and \mathbf{k} aligned along the south to north pole axis.

Here, we neglect the contribution of the sea surface tilt to the momentum balance, which is small compared to the other contributions (Steele et al., 1997). In Eq. (2), the wind forcing $\boldsymbol{\tau}_a$ excites the oscillations, whereas the oceanic drag $\boldsymbol{\tau}_w$ and the internal stress term $\nabla \cdot \boldsymbol{\sigma}$ damp those oscillations (Colony and Thorndike, 1980).

We expect the response of sea ice to the Coriolis force in the frequency domain to be within two following extremes. A buoy moving in free drift, i.e. fixed on an ice floe that

Sea ice inertial oscillations

F. Gimbert et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



drifts according to wind and ocean currents without any mechanical interaction with other ice floes (the term $\nabla \cdot \sigma$ is then negligible compared to the others), is expected, in first order approximation, to follow the oceanic fluid parcel and thus to oscillate in a similar way, although another source of damping of the oscillations might come from the friction between the ice bottom and the ocean surface (Lepparanta et al., 2012). In contrast, on an highly cohesive ice cover experiencing strong internal stresses, the corresponding contribution $\nabla \cdot \sigma$ will dominate the other terms in Eq. (2) so that the oscillations are immediately damped out and thus become undetectable. As shown later in Sect. 3, the amplitude of inertial oscillations also depends on the advection velocity, as intense advection tends to fade inertial loops. Recast in terms of ice mechanical behavior, this plays an opposite to the role of internal mechanical interactions: indeed, a weaker, looser ice cover is expected to drift and advect faster (Rampal et al., 2009), which implies a negative contribution on inertial oscillations. Consequently, larger inertial oscillations will, in average, genuinely measure (actually underestimate) a weakening of the energy dissipation through mechanical processes that take place within the ice cover.

To measure the amplitude of the inertial oscillations of the ice drifters is therefore to estimate the level of mechanical dissipation within the ice cover, and therefore its degree of cohesiveness. As such, its evolution and its spatial pattern can highlight the changes in ice conditions and ice cover mechanical behaviour. Internal ice stress measurements have been performed directly from specific sensors (Richter-Menge and Elder, 1998; Richter-Menge et al., 2002). The seasonal variations of these local stress measurements have shown an increase of ice-motion induced stresses as the winter season progresses and the cohesiveness/compactness of the ice cover develops (Richter-Menge and Elder, 1998). While of major importance to analyze sea ice mechanical behaviour and rheology (Weiss et al., 2007), these measurements are however limited to the local scale and do not carry any information about a possible large-scale, long-term trend of the mechanical state of the cover. Our approach is thus complementary, as it allows a much better sampling, both in time and space.

Sea ice inertial oscillations

F. Gimbert et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Sea ice inertial oscillations

F. Gimbert et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

The idea to relate the amplitude of the inertial oscillations to the degree of consolidation of the ice cover was already formulated by McPhee (1978) as well as Colony and Thorndike (1980). The latter studied the role of the mechanical behaviour of the ice on the coherency of inertial motion between different buoys, an approach different from what is described below as we analyze trajectories individually. More recently, from buoys trajectories near the Antarctic Peninsula, Geiger and Perovich (2008) observed an increase of inertial motion related to the degradation of the ice pack during the spring breakup. Hutchings et al. (2012) also discussed the link between the degree of disconnection of the ice cover, the mechanical dissipation of energy, and the amplitude of inertial motion. These last authors based their discussion on the analysis of strain-rate records obtained from a dense array of buoys in the Weddell sea, i.e. they did not analyze individual trajectories. However, although sea ice inertial oscillations have been studied by several others authors (Colony and Thorndike, 1980; McPhee, 1978; Hunkins, 1967; Lammert et al., 2009), including recently on ice strain-rate records (Kwok et al., 2003), this is the first time that a systematic analysis is performed at the Arctic basin and multidecadal scales.

In this paper, we propose a method to quantify the magnitude of inertial oscillations from lagrangian (buoys) trajectories (Sect. 3). We then apply this methodology to the International Arctic Buoy Program dataset (IABP) covering 30 yr of data in the Arctic Ocean (Sect. 4). As shown below, this analysis is in full agreement with the above expectations: inertial oscillations are very weak or absent in a highly cohesive ice cover, such as in winter within the Central Arctic, but are well developed in summer at the periphery of the basin, i.e. in regions of less concentrated, loose ice. In addition, a significant strengthening (on average) of these oscillations is observed, suggesting a mechanical weakening of the Arctic sea ice cover. This is confirmed in Gimbert et al. (2012), where a simple ocean-sea ice coupled dynamical model explains these seasonal, geographical and pluri-annual variations of inertial oscillation magnitude in term of changes within sea ice internal mechanical properties, through an associate decrease of sea ice internal friction in recent years.

2 Data set

2.1 IABP buoy data set

The sea ice drifting buoy data set, provided by the International Arctic Buoy Program (IABP), consists of approximately 650 buoy tracks recorded from January 1979 to December 2008. These ice drifters are dropped every year at the end of winter, mostly in April, and drift according to the ice motion. Their positions are tracked by GPS receivers or Argos transmitters with a position uncertainty of the order of 100 m and 300 m, respectively. The raw buoys' positions are irregularly sampled through time. Thus, in order to get a regular sampling, the buoy positions provided by the IABP result from a cubic interpolation of the raw positions with a 3 h time re-sampling (see the IABP documentation at <http://iabp.apl.washington.edu/data.html> for further details). This interpolation procedure acts as a low-pass filter, thus most likely reducing the position error around 100 m or below (Lindsay and Stern, 2003). Figure 1 shows all the buoy tracks on a polar stereographic map, where buoy locations are defined as $x\mathbf{e}_1 + y\mathbf{e}_2$, where \mathbf{e}_1 and \mathbf{e}_2 are two orthogonal unit vectors with \mathbf{e}_1 being along the Greenwich meridian and $(x, y) = (0, 0)$ at North Pole.

2.2 Oceanic buoy data set

In order to get an example of the amplitude of inertial oscillations in the absence of sea ice, i.e. when the damping of oscillations is mainly due to the internal friction of the Ekman layer without contribution from internal ice stresses, we consider trajectories of buoys drifting in the North Atlantic, along the Northern Spanish coast (Fig. 2).

This oceanic buoy dataset, provided by the SHOM (Service Hydrographique et Oceanographique de la Marine), consists of 4 buoy tracks (B239, B241, B242, B245). These buoys were deployed between 6 and 11 December 2006 in the framework of the CONGAS project (Le Cann and Serpette, 2009). The buoys' positions were tracked by GPS receivers with a position uncertainty of the order of several tens of meters. The

Sea ice inertial oscillations

F. Gimbert et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Sea ice inertial oscillations

F. Gimbert et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



temporal sampling was regular, equal to 1 h. Thus, the buoys' positions do not result from any re-sampling or re-interpolation. Buoys B239 and B245 operated during 1 yr, whereas buoys B241 and B242 operated only 1 month. For each latitude (ϕ) – longitude (φ) buoy position, we define the orthogonal base ($\mathbf{e}'_1, \mathbf{e}'_2$) of this coordinate system as $\mathbf{x}_{(\phi, \varphi)} = x\mathbf{e}'_1 + y\mathbf{e}'_2$ using a Lambert projection centered in the coordinates $\phi = 43^\circ$ and $\varphi = 6^\circ$.

3 Quantifying the magnitude of inertial oscillations

3.1 Observations

We analyze here several cases of inertial oscillations observed over an ice-free ocean or over sea ice.

Figure 3a shows 1 month of the trajectory of the oceanic buoy B245. Inertial oscillations are noticeable during this time period: being in the Northern Hemisphere, the buoy drift trajectory exhibits cycloids in the clockwise direction. The intermittent character of inertial oscillations is also clearly visible. Periods of strong inertial oscillations, marked by cycloidal loops (red boxes in Fig. 3a) might have been triggered by storms. Following these periods, the oscillations are progressively damped out and the loops nearly disappear (green boxes in Fig. 3a), before a new storm or moving front can trigger new inertial oscillations.

We compute the $u_x(\tilde{x}, \tilde{y}, \tilde{t})$ (along the x axis) and $u_y(\tilde{x}, \tilde{y}, \tilde{t})$ (along the y axis) speeds as follow:

$$\begin{aligned} u_x(\tilde{x}, \tilde{y}, \tilde{t}) &= (x(t + \Delta t) - x(t)) / \Delta t \\ u_y(\tilde{x}, \tilde{y}, \tilde{t}) &= (y(t + \Delta t) - y(t)) / \Delta t \end{aligned} \quad (3)$$

with $\Delta t = 1$ h and where the mid-points \tilde{x} , \tilde{y} and \tilde{t} are defined as $\tilde{x} = (x(t + \Delta t) + x(t)) / 2$, $\tilde{y} = (y(t + \Delta t) + y(t)) / 2$ and $\tilde{t} = ((t + \Delta t) + t) / 2$. The Fourier transform $\hat{U}_b(\omega)$ of the buoy velocities u_x and u_y , for a selected buoy trajectory which starts at time t_0 and ends at

time t_{end} as:

$$\hat{U}_b(\omega) = \frac{1}{N} \sum_{t=t_0}^{t_{\text{end}}-\Delta t} e^{-i\omega t} (u_x(\tilde{x}, \tilde{t}) + i u_y(\tilde{y}, \tilde{t})) \quad (4)$$

where N is the number of velocity samples along the trajectory and $\omega = 2\pi f$. This vectorial Fourier transform distinguishes negative and positive frequencies: peaks at $f < 0$ or $f > 0$ are associated with clockwise, counter-clockwise, displacements, respectively.

Figure 3b shows the Fourier spectrum $|\hat{U}_b(\omega)|$ of the buoy velocities associated to the trajectory plotted on Fig. 3a. The Fourier transform reveals a peak of magnitude 5.2 km day^{-1} at the inertial oscillation frequency ($f_0 \approx -1.35 \text{ cycles day}^{-1}$ using the average latitude $\phi = 1/N \sum_t \phi(t) \approx 42^\circ$ in Eq. (1)). This peak corresponds to the component of the buoy motion associated to the Coriolis force. The peak at $f = 0$ represents the advective component of the buoy's motion. Finally, the two peaks observed at $f = -2 \text{ cycles day}^{-1}$ and $f = 2 \text{ cycles day}^{-1}$ are associated with a semi-diurnal tidal oscillation. Unlike the inertial oscillation, the tidal oscillation does not rotate and the associated peak is therefore observed at positive and negative frequencies.

We now consider IABP ice drifters. A buoy moving in free drift, i.e. fixed on an ice floe that drifts according to wind and ocean currents alone, hence without any mechanical interaction with other ice floes, is expected to oscillate in a way similar to e.g. the oceanic B245 buoy of Fig. 3, subject to similar forcing conditions, although another source of damping of the oscillations might come from the friction between the ice bottom and the ocean surface.

Figure 4a shows an IABP buoy trajectory within the Fram strait during the summer period, where ice is highly fragmented and loosely packed. The clockwise-cycloids are clearly visible. We observe bursts in the inertial oscillation intensity (red rectangles) followed by a period of decaying intensity (green rectangle).

The velocity of the buoy is computed using Eq. (3), with stereographic coordinates x and y . The inertial oscillations are evidenced by a strong peak observed on the

Sea ice inertial oscillations

F. Gimbert et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Sea ice inertial oscillations

F. Gimbert et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



loops indicate that the rotation velocity is larger than the advection velocity. In contrast, no full loops are observed within the green boxes, showing that the rotation velocity is lower than the advection velocity. Lepparanta (2004, p. 157) proposed a simple ratio of cycloidal rotation velocity over the advection velocity to qualitatively estimate the strength of inertial oscillations.

We here propose to evaluate in a quantitative way the time-dependent oscillation magnitude using the Fourier spectrum at the inertial frequency f_0 . Since inertial oscillations are best quantified by means of a spectral analysis, we here perform such an analysis on a sliding time window. This will enable us to define a time-varying quantity M that measures how inertial oscillations evolve with time as a buoy drifts. For a given buoy location defined by the coordinates $(x_{\text{cur}}, y_{\text{cur}}, t_{\text{cur}})$, where “cur” stands for “current”, we define a gaussian window function $g_{\text{cur}}(t)$ centered on t_{cur} with a characteristic duration of the order of several inertial time periods:

$$g_{\text{cur}}(t) = \exp\left(\frac{-(t - t_{\text{cur}})^2}{2(nT_o)^2}\right) \quad (5)$$

where $T_o = -1/f_0$ is the inertial time period and n controls the width of the window. In this analysis, we set $n = 1$, i.e., a time interval around t_{cur} that is long enough to really probe the inertial time scale, and short enough to properly measure rapid time variations in the inertial oscillation magnitude.

Gaps (missing data) are present in the IABP buoy records. Their duration varies from one sample (i.e. a time interval of 6 h between successive positions) to several weeks. Therefore, we introduce in our calculations a selection condition such that no gap of data is allowed for values of t that verify:

$$g_{\text{cur}}(t) > P \quad (6)$$

where we set $P = 0.01$. No M -value is thus computed at time t_{cur} if one or more sampling time t verifying condition (6) is lacking. Moreover, since missing data even occurring far from t_{cur} (i.e. at values of t such that $g_{\text{cur}}(t) < P$) can play a non negligible role

on the Fourier transforms, the gaussian of Eq. (5) is then truncated at t_{lim}^+ and t_{lim}^- in the following way:

$$\begin{aligned} \forall t > t_{\text{lim}}^+ \quad g_{\text{cur}}(t) &= 0 \\ \forall t < t_{\text{lim}}^- \quad g_{\text{cur}}(t) &= 0 \end{aligned} \quad (7)$$

where $g_{\text{cur}}(t_{\text{lim}}^+ > t_{\text{cur}}) = g_{\text{cur}}(t_{\text{lim}}^- < t_{\text{cur}}) = P$. This truncation occurs at ± 1.5 days from t_{cur} .

By the use of condition (6), we can compute 1.3×10^6 M -values from the IABP dataset, which represents 83 % of the total number of observations available.

We then compute the velocity $\mathbf{W}_{\text{cur}}(t)$ as follow:

$$\mathbf{W}_{\text{cur}}(t) = \mathbf{U}_b(t) \times g_{\text{cur}}(t) \quad (8)$$

where $\mathbf{U}_b(t) = u_x(t)\mathbf{e}_1 + u_y(t)\mathbf{e}_2$.

We compute the normalized Fourier spectrum at time t_{cur} of the velocity time series $\mathbf{W}_{\text{cur}}(t)$ at the inertial frequency f_0 as:

$$\hat{W}_{\text{cur}}(f_0) = \Delta t \sum_{t=t_0}^{t_{\text{end}}} e^{-i\omega_0 t} \times (u_x(t) + iu_y(t)) \times g_{\text{cur}}(t) \quad (9)$$

where $\omega_0 = 2\pi f_0$, and t_0 and t_{end} are the starting and ending times of the trajectory. This value is likely to represent the amplitude of the inertial oscillations. We further normalize it in order to define an adimensionalized parameter M that measures the magnitude (still at t_{cur}) of the inertial oscillation:

$$M = \frac{|\hat{W}_{\text{cur}}(f_0)|}{1.27} \times \frac{4}{\pi \overline{W}_{\text{cur}}} \quad (10)$$

Sea ice inertial oscillations

F. Gimbert et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



where

$$\overline{W}_{\text{cur}} = \frac{1}{1.27} \int_{t_0}^{t_{\text{end}}} dt |W_{\text{cur}}(t)|. \quad (11)$$

is the (current, at time t_{cur}) mean of the norm of the drift velocity, and the value 1.27 is equal to $\int_{t_0}^{t_{\text{end}}} g_{\text{cur}}(t) dt$.

In order to figure out how the parameter M defined here varies with the geometry exhibited by the buoys' trajectories, we consider an ideal case of a buoy trajectory led by two (constant in time) velocity components: an advection velocity U and a rotation velocity V , induced by the Coriolis force. The equation of the trajectory described by the buoy under these considerations is:

$$\begin{cases} u_x(t) = U + V \cos \omega_0 t \\ u_y(t) = V \sin \omega_0 t \end{cases} \quad (12)$$

Here, M essentially measures the ratio V/U . Figure 6 illustrates trajectories computed from Eq. (12) for several ratios V/U . For a buoy's trajectory that describes a perfect cycloid, i.e. for $U = V$ (see Fig. 6b), the $4/\pi$ factor of Eq. (10) ensures that $M = 1$. In that ideal case, corresponding to a buoy that behaves in free drift without being submitted to any damping source, M is equal to 1, independently of the value of the advection velocity U considered. In contrast, for the other cases, M increases with decreasing advection velocity U . When considering $U = 0$ or $U \ll V$, i.e. a buoy trajectory that describes a circle, the associated M -value is equal to $\frac{4}{\pi} = 1.27$, which constitutes the upper bound value of M . For example, by taking $V/U = 2$ (Fig. 6a), the trajectory exhibits loops and $M = 1.2$. On the opposite, for $U > V$, values of M are lower than 1, with the limit case of $M \approx 0$ when $U \gg V$, i.e. when the buoy trajectory is similar to a straight line. Intermediate values are illustrated on Fig. 6c, d computed using $V/U = 0.5$, that corresponds to $M = 0.6$, and $V/U = 0.15$, that corresponds to $M = 0.2$, respectively.

Sea ice inertial oscillations

F. Gimbert et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



For $M = 0.2$, the inertial oscillations are difficult to distinguish while a clear cycloid is observed for $M = 0.6$.

3.3 Application to the buoy trajectories

Figure 7 shows the values of M computed using the different trajectories plotted in Sect. 3.1. Large M -values are obtained for the oceanic buoy trajectory plotted on Fig. 3, with an average of 0.56. For the ice drifter of Fig. 4, we also get large M values with an average of 0.8: the $M > 1$ values are all associated with cycloidal loops observed on Fig. 4a (red rectangles), followed by their decay as characterized by a decrease of M (green rectangles). In contrast, for the ice drifter of Fig. 7c, the M values are much lower: the mean M value over the time period is only equal to 0.13, illustrating the relative absence of inertial loops.

As the raw buoys' positions of the IABP ice drifters dataset are irregularly sampled through time, then interpolated (cubic interpolation) and re-sampled at a regular 3 h interval, we investigated the effect of this procedure in the values of M computed for the IABP dataset. To do so, we artificially degraded the oceanic buoy dataset, which samples buoys positions every hour, by randomly removing a given % of raw positions. A cubic interpolation followed by a re-sampling at 3 h is then performed on the degraded dataset, as done for IABP data, and the associated M -values (noted M_D) are re-computed (Fig. 8). The cubic interpolation alone, i.e. without degrading the raw data, has almost no influence on M (Fig. 8a). On Fig. 8b, we can see that the deviation of the M_D time series from the M time series becomes visible only when more than 50 % of the raw buoys positions are missing. For 50% of missing data, the error on individual M values is lower than 5%. More importantly, as we focus in the following on M values averaged over trajectories, the departure of averaged M_D values from the average M value for increasing missing data ratios is shown on Fig. 8c. Average M_D values do not deviate significantly from the average M value for missing data ratios up to 60 %, which corresponds to an average time sampling of about 2:50 h and a time sampling maximum gap of about 6 h. Thus, for sampling times lower than these threshold values,

Sea ice inertial oscillations

F. Gimbert et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



we can consider that the computed M values are not affected by an irregular sampling through time. While not shown here, similar results have been obtained by considering ice drifters initially regularly sampled at 1 h during the TARA field campaign (Gascard et al., 2008). This indicates the robustness of IABP interpolation procedure coupled to our estimation of the amplitude of inertial oscillations.

These examples demonstrate that the parameter M is appropriate to quantify the inertial oscillation magnitude. Small average values of M (say lower than 0.2) correspond to buoy trajectories within a strongly cohesive ice pack, while large values of M (say greater than 0.6) correspond to buoy trajectories that we can consider to be nearly in free drift condition.

4 Analysis of 30 yr of IABP data

In this section, we analyse the spatial and temporal patterns of the inertial oscillation magnitude M , and check whether a significant trend can be observed over the Arctic basin during the 30 yr of the IABP dataset, that would reveal a significant change in ice conditions. As previously explained in Sect. 3.1, inertial oscillations are caused by sudden changes in external forces (strong wind), which then decay due to kinetic energy dissipation within the Ekman layer or friction that takes place at the ice/water interface, or to internal ice stresses. Therefore, more frequent and stronger storms over the years should imply larger M -values on average. Cyclonic activity over the Arctic Ocean shows a maximum during summer that could partly explain an annual cycle of the inertial oscillation magnitude (Serreze and Barrett, 2008) (see below). On the other hand, no significant trend in cyclonic activity has been found over the last 50 yr (Serreze and Barrett, 2008). Similarly, the average wind speed over the Arctic basin, as estimated from the ERA-40 reanalysis dataset, does not show any significant trend over the period 1979–1999 (Rampal et al., 2009).

Sea ice inertial oscillations

F. Gimbert et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



4.1 Seasonal variation

In this section, we compute M for buoys located within the Central Arctic basin, as delimited by the thick black line on Fig. 1, and then investigate intra-annual (monthly) variations by grouping M values into monthly periods, i.e.: one average monthly value M for all the M values occurring in January, whatever the calendar year, etc. The Central Arctic is here delimited by all buoys positions lying further than 150 km away from the coasts and the Fram strait. This way, we skip from the analysis buoys possibly stuck on fast ice.

Figure 9 shows the monthly M value averaged over the 30 yr of record. To estimate the associated error bars, we checked using a bootstrap method (Rampal et al., 2009) that the standard deviation ΔM indeed varies as $\frac{M}{\sqrt{N}}$, where N is the number of M -values used to calculate the monthly mean M . The monthly mean M value reaches a minimum of 0.168 in May and a maximum of 0.294 in September, and exhibits an obvious annual cycle. We define the summer season by the three months of July, August and September and the winter period by the rest of the year. These results are consistent with other observations: sea ice concentration, sea ice thickness and sea ice deformation also describe an annual cycle (Rampal et al., 2009; Rothrock et al., 2008). The annual cycle of ice thickness within the Arctic basin has a maximum on 30 April (Rothrock et al., 2008), in phase with our minimum M -value in May. Being thinner and less concentrated in summer, sea ice is less cohesive and more deformable during this time period. This leads, in average, to a lower damping of the inertial oscillations, and therefore larger M -values. To reinforce this point, it is important to note that, through the evaluation of M , our analysis underestimates the effect of the decrease of mechanical energy dissipation within the ice cover. Indeed, larger advection velocities recorded in summer (Rampal et al., 2009) should lead to an associated decrease of the M values (cf Sect. 3.2), while the opposit, i.e. an increase of the M values in summer, is observed. On the other hand, we cannot exclude that a stronger cyclonic activity

Sea ice inertial oscillations

F. Gimbert et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

during summer (Serreze and Barrett, 2008) could reinforce the annual cycle described by the average M -values by triggering more oscillations.

4.2 Spatial pattern

The question arises as to whether the seasonal changes in the M -values evidenced in the previous section can be associated to a spatial pattern. In this section, we select values of M associated to buoys located within the Central Arctic basin and within the Fram strait (thick black and grey lines on Fig. 1). From these values, we build a seasonal M data set by separating the M values computed in winter, i.e. all the M -values recorded for all the 30 winter seasons, from the M values computed in summer. The seasonal spatial patterns of M for both seasons are plotted on Fig. 10. The spatially averaged M value, denoted $\overline{M}(X_j, Y_j)$, is computed for each grid point j of coordinates (X_j, Y_j) as:

$$\overline{M}(X_j, Y_j) = \frac{1}{\sum_i w_{ij}(x_i, y_i)} \sum_i w_{ij}(x_i, y_i) M_i(x_i, y_i) \quad (13)$$

where the summation is performed over all the buoys positions (x_i, y_i) contained in a circle of radius $L = 400$ km centered on the coordinates (X_j, Y_j) . The weight $w_{ij}(x_i, y_i)$ is defined as

$$w_{ij}(x_i, y_i) = e^{-d^2/2L^2} \quad (14)$$

where $d = \sqrt{(x_i - X_j)^2 + (y_i - Y_j)^2}$ is the distance between the buoy position (x_i, y_i) and the grid point (X_j, Y_j) .

The patterns observed on Fig. 10 are, at least qualitatively, consistent with the observed distribution of the sea ice thickness (Kwok and Rothrock, 2009) and concentration (<http://nsidc.org/data/seaice/index.html>) (Comiso, 1990, updated 2012). In summer, large \overline{M} -values are observed at the edge of the basin in the Beaufort, Chukchi

Sea ice inertial oscillations

F. Gimbert et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Sea ice inertial oscillations

F. Gimbert et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and Laptev seas, i.e. in regions where the multiyear ice cover has been progressively disappearing during the last decade (Maslanik et al., 2011). On the contrary, small values of \overline{M} can be observed along the Canadian coasts, where the average ice thickness is at its maximum (Kwok and Rothrock, 2009). Sea ice behaves more as a strongly cohesive plate in this region, insensitive to the Coriolis force. This zone corresponds to the multi-year ice still remaining nowadays. Large values of \overline{M} are also observed south of Fram strait, which is consistent with what we discussed in Sect. 3.1. In contrast, the winter pattern does not reveal any particular structure within the basin. The values of \overline{M} are small over the whole basin, except in the Barents sea, where the number of observations is too small to allow a firm conclusion. Note, however, unexpected and unexplained slightly larger M -values along the Canadian archipelago in this case, although the difference with the Central Arctic is small. In addition, the Barents sea being characterized by a continental plateau and shallow waters, the tidal oscillation is more marked there, as revealed by a frequency peak at $+2 \text{ cycles day}^{-1}$. This adds also to the inertial peak at $-2 \text{ cycles day}^{-1}$, and could partly explain the large \overline{M} -values in this region.

To further test the link between the state of the sea ice cover and its cohesiveness, as expressed by the amplitude of the inertial oscillations, we perform a correlation analysis between the \overline{M} -values and the open water concentration $1 - \alpha$, where α is the sea ice concentration dataset collected by NSIDC (<http://nsidc.org/data/seaice/index.html>) (Comiso, 1990, updated 2012). This dataset has a spatial resolution of 25 km and consists of ice concentration values sampled every two days from 1979 to 1987 (SMMR), and every day since 1987 (SSM/I). For each value of M , we search for the corresponding value of open water concentration as the closest sample in time and space: a $1 - \alpha$ value is associated to a given \overline{M} -value if we can find, for the same day of record, within one day during the period 1979–1987, a sample that is closer than 25 km. The corresponding correlation coefficient R is equal to $0.245(\pm 0.002)$. This positive correlation is statistically significant as R is more than 120 times greater than the

standard deviation obtained in the null hypothesis at no correlation (numerically computed by randomly reshuffling the \overline{M} and $1 - \alpha$ values). This is consistent with stronger oscillations characteristic of a less compact sea ice cover. However, the coefficient of correlation is small, suggesting that other factors than the ice concentration, such as the ice thickness, control the observed M -values.

In order to check whether this global correlation is, or not, only due to the fact that \overline{M} and $1 - \alpha$ both describe an annual cycle, we group together these values by months. We then compute the cross-correlation coefficient R separately for each month of the year, over the 30 yr of record. While not plotted here, the annual variation of the correlation coefficient is in phase with the annual cycle of M plotted on Fig. 9: the correlation is larger in summer. The correlation coefficient is equal to $0.350(\pm 0.005)$ in August, which means that the \overline{M} -values are spatially correlated with the associated open water concentration values. In winter, the correlation is much less significant, which can be understood by the fact that, during this period, the sea ice concentrations are equal to 1 almost everywhere within the basin and therefore show very little fluctuations.

4.3 Pluri-annual variation

We have shown so far that the M parameter is a useful proxy for the cohesiveness of the cover, and, consequently, the level of internal stresses. We now analyze whether the M -values have changed over the last three decades owing to a change in ice conditions.

4.3.1 M time series

In this section, we use the Central Arctic buoy data set. The temporal sampling of the IABP buoys dataset is heterogeneous, being characterized by variations of data density over the period. Most notably, a larger number of observations is available in the later years, as compared to the early 80s or during the late 90s. To circumvent this problem, the evaluation of the pluri-annual variation of M is done by equally binning

Sea ice inertial oscillations

F. Gimbert et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the M -values in time. The summer and winter datasets are both separately splitted into 10 successive bins, all containing the same number of observations. For each bin, an average value of M , associated to an average value of time, is computed.

Figure 11 shows the time series $M(t)$ between January 1979 and December 2008 for the summer and winter seasons. The error on the average M value is computed using a bootstrap method as explained in Rampal et al. (2009). The bootstrap method is performed individually for each bin and the deviation from the mean ΔM is represented by the errorbars. We see on both plots that the most prominent feature is a significant increase of the inertial oscillation amplitude through time. A linear fit gives a positive trend equal to $1.19 (\pm 0.34) \times 10^{-5} \text{ yr}^{-1}$ (i.e., a 16.5 % increase per decade) for summer and $5.7 (\pm 1.9) \times 10^{-6} \text{ yr}^{-1}$ (i.e., a 11 % increase per decade) for winter. Thus, the increase of inertial oscillations is relatively more marked in summer than in winter.

The use of GPS as the buoy positioning system became more common since the end of the last century. As noted in Sect. 2.1, this system is more accurate than ARGOS positioning. We thus check whether the observed trend on the mean time series $M(t)$ could be a spurious effect caused by reduced noise in recent years. To do so, noised mean time series $M_N(t)$ are built by computing M values on buoy trajectories over which Gaussian noise is added. Because the increase in the average $M(t)$ values is particularly marked from the beginning of the 21 century, Gaussian noise is only added to buoy trajectories recorded during the period 2002–2008. For each IABP position (x_b, y_b) of buoy trajectories recorded from the year 2002 onwards, we define a noised buoy position $(\tilde{x}_b, \tilde{y}_b)$ as $\tilde{x}_b = x_b + \delta x$ and $\tilde{y}_b = y_b + \delta y$ where δx and δy are increments randomly picked from a centered gaussian distribution with a standard deviation σ . Figure 12 shows realizations of mean time series $M_N(t)$ built using $\sigma = 300\text{m}$ and $\sigma = 1000\text{m}$, along with the IABP time series $M(t)$, as on Fig. 11. From the year 2002, the mean time series $M_N(t)$ deviate from $M(t)$, showing that noise on the buoys positions has an influence on the parameter M . However, almost systematically, values of $M_N(t)$ are larger than $M(t)$, showing that noise most generally acts to increase the M values rather than to decrease them.

Sea ice inertial oscillations

F. Gimbert et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Sea ice inertial oscillations

F. Gimbert et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



This can be explained as follows: noise increases the amplitude of the Fourier spectrum at high frequencies. This has two competing consequences; (i) the norm of the drift velocity $\overline{W}_{\text{cur}}$ slightly increases, which tends to decrease M (see Eq. 11) and (ii) the amplitude at the inertial frequency $\hat{W}_{\text{cur}}(f_0)$ is increased, which tends to increase M . The second effect is particularly marked when the inertial motion (so, M) is small, and, most of the time, dominates over the first effect, so that M is increased by noise. Consequently, the evolution observed on Fig. 11 is not the consequence of possible reduced position noise in recent years, as the difference between the periods 1979–2001 and 2002–2008 would have been even larger if the level of noise had remained the same over the entire record. In other words, the observed trend in $M(t)$ is most likely an underestimation of the pluri-annual evolution of the inertial oscillation magnitude.

The same is also true for the annual cycle (Fig. 8): as the effect of noise is stronger for low M -values, winter M -values are, in average, more overestimated than summer M -values.

In addition, Rampal et al. (2009) reported a significant (9 % per decade) increase of sea ice speeds in the Arctic basin over the last three decades. Therefore, as increasing advection velocities act to decrease M (Sect. 3.2), the pluri-annual positive trend on the M -values reported here most likely underestimates the associated decrease of the mechanical energy dissipation within the ice cover in the later years.

4.3.2 Evolution of spatial patterns

The previous section demonstrated a strong evolution of the average inertial oscillation amplitude, although it is more strongly marked in summer. We here analyze the spatial consequences of the observed pluri-annual changes. To do so, we split the whole IABP buoy seasonal data set in two periods. A student' t-test performed on the winter and the summer $M(t)$ -time series at Fig. 11 shows that the most significant changing point occurs after 5 averaged- M values out of the 10. More precisely, splitting the winter time series $\{M_1, M_2, \dots, M_{10}\}$ into two distributions $\{M_1, \dots, M_5\}$ and $\{M_6, \dots, M_{10}\}$, the probability that these two distributions have the same mean is only 0.92 %. We have

verified that this probability is indeed the lowest when splitting the time series just after M_5 . Similarly, we find the same optimal change-point in summer, with an associated probability that the two distributions share the same mean equal to 0.42 %. This leads to define two distinct periods, with Period 1 from 1979 to 2001 and Period 2 from 2002 to 2008, see Fig. 11. We can remark that according to this periodic splitting, Period 1 and Period 2 contain the same number of observations i.e. 5 bins each. As done in Sect. 4.2, the spatial patterns of the M values for the summer and winter seasons associated to each period are shown on Fig. 13. As expected, the changes are stronger in summer. They affect most of the Arctic, but are more pronounced on the siberian side: a drastic increase of the peripheral ice zone area is observed in the later years. Conversely, winter changes are milder north of Canada and Greenland, where the thickest multiyear ice can be found nowadays (Kwok and Rothrock, 2009).

4.3.3 Is the observed evolution from the IABP buoys dataset representative of the whole Arctic basin?

In the two previous sections, care was taken to remove the effect of temporal heterogeneities in the IABP buoys dataset. However, the IABP buoy dataset is spatially heterogeneous: changes in in the spatial sampling by the buoys could affect the overall trend. In order to check whether the results reported in Fig. 11 truly indicate a significant increase of the M -values, we investigate whether this trend could be an artefact of changes in spatial sampling.

Following the procedure described in Rampal et al. (2009) for ice velocities, we formulate a null hypothesis: in this hypothesis, there exists no temporal changes in M -maps over the year, but the effect at spatially sampling these maps in different ways at different periods will cause the M -time series to evolve with time. To construct these M -maps, we consider any buoy position associated to a given M value and recorded at a given year and a given season. For such a position, we compute the mean M_0 value for summer, respectively winter, by considering all the M values recorded in summer, respectively winter, and whatever the year, contained within a circle of radius

Sea ice inertial oscillations

F. Gimbert et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Sea ice inertial oscillations

F. Gimbert et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



$L = 200$ km, using Eq. (13). This mean value is therefore year-independent, and corresponds to our null hypothesis of no inter-annual changes. Then, we calculate our null hypothesis M_0 time series as the mean of the summer, respectively winter M -values, using the same bins as in Fig. 11. Any inter-annual variation observed in the time series $M_0(t)$ could only be explained by changes on the spatial sampling rather than by an actual, global trend: for example, a positive trend would be explained if IABP buoys had a tendency to sample regions associated to large M values more often in the later years as compared to earlier years.

Figure 14 shows the mean time series $M(t)$ and $M_0(t)$ over the whole time period 1979–2008. The trend associated to the null hypothesis is equal to $1.57 (\pm 0.98) \times 10^{-6} \text{ yr}^{-1}$ in summer and $1.28 (\pm 0.51) \times 10^{-6} \text{ yr}^{-1}$ in winter. This means that changes in spatial sampling are responsible for approximately 10% of the observed trend in summer and 20% of the observed trend in winter. We thus conclude that the trend observed in the average M values cannot only be explained by an irregular sampling of the buoys. This increase in M is not an artefact and reveals a genuine increase of the inertial oscillations magnitude over the whole Arctic basin.

5 Conclusions

From the IABP ice drifter trajectories recorded between 1979 and 2008, we analyzed the average amplitude of inertial oscillations over the Arctic sea ice cover. To do so, we defined a non dimensional parameter M that quantifies the magnitude of inertial oscillations relative to advection motion along the trajectories. For a sea ice cover of low concentration, constituted from a loose assembly of floes moving nearly in free drift, inertial motions and M values are large. On the reverse, an highly cohesive sea ice pack, characterized by strong internal stresses, is expected to be associated with low M values. From appropriate averaging of this 30-yr dataset at different space or time scales, we have shown that:

Sea ice inertial oscillations

F. Gimbert et al.

- (i) the M values describe an annual cycle with a minimum reached in May and a maximum in September, in a qualitative agreement with the corresponding annual cycles of sea ice extent (Comiso et al., 2008), concentration, thickness (Rothrock et al., 2008; Kwok and Rothrock, 2009), advection velocity and deformation rates (Rampal et al., 2009).
- (ii) the spatial pattern of M over the Arctic basin is in agreement with the sea ice thickness and concentration patterns. Low M values are observed in Western Arctic, whereas large values are observed within a peripheral zone (Beaufort Sea, Eastern Arctic) and south of Fram strait.
- (iii) a significant increase of average values of M is observed from 1979 to 2008. This increase, although more marked in summer, is observed in both seasons and is associated with the reduction of the thick, multiyear ice zone in recent years.

From the expected link between the magnitude of the oscillations and the degree of consolidation of the ice cover (McPhee, 1978; Colony and Thorndike, 1980; Geiger and Perovich, 2008), we believe that point (iii) is a signature of the mechanical weakening of the Arctic sea ice cover in recent years. However, one may argue from the momentum balance of sea ice, see Eq. (2), that this strengthening of inertial motion might simply and more directly result from the observed thinning of the cover, i.e. a reduction of ice mass per unit area. In addition, this evolution could, to some extent, be the result of a modification of vertical penetration of turbulent momentum within the ocean boundary layer. In Gimbert et al. (2012), we have shown, from a simple ocean-sea ice coupled dynamical model, that these two explanations cannot fully account for the evolution of inertial motion in the Arctic, which actually reveals a genuine mechanical weakening, through an associate decrease of the sea ice internal friction magnitude, of the cover at the basin scale. Such mechanical weakening has (will have) strong consequences in terms of ice drifting speeds, deformation rates, export (Rampal et al., 2009) and therefore on mass balance (Rampal et al., 2011).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Acknowledgements. This study owes much to a suggestion by Stephane Roux. We thank the IABP for the availability of their buoy dataset as well as the SHOM laboratory for having provided us the oceanic buoys trajectories. We also thank Pierre Rampal and Neige Calonne for earlier work on this topic and helpful discussions.



The publication of this article is financed by CNRS-INSU.

References

- Aranson, I. and Tsimring, L.: Patterns and collective behavior in granular media: theoretical concepts, *Rev. Mod. Phys.*, 78, 641–692, 2006. 2181
- Colony, R. and Thorndike, A.: The horizontal coherency of the motion of summer Arctic sea ice, *J. Phys. Oceanogr.*, 10, 1281–1289, doi:10.1175/1520-0485(1980)010<1281:THCOTM>2.0.CO;2, 1980. 2182, 2184, 2202
- Comiso, J.: Bootstrap sea ice concentrations from NIMBUS-7 SMMR and DMSP SSM/I (January 1979 to December 2008), edited by: Maslanik, J. and Stroeve, J., Digital media, NSIDC, Boulder, Colorado, USA, 1990, updated 2012. 2195, 2196
- Comiso, J. C., Parkinson, C. L., Gersten, R., and Stock, L.: Accelerated decline in the Arctic sea ice cover, *Geophys. Res. Lett.*, 35, L01703, doi:200810.1029/2007GL031972, 2008. 2180, 2202
- D'Asaro, E. A., Eriksen, C. C., Levine, M. D., Niiler, P., Paulson, C. A., and Meurs, P. V.: Upper-ocean inertial currents forced by a strong storm. Part I: Data and comparisons with linear theory, *J. Phys. Oceanogr.*, 25, 2909–2936, doi:10.1175/1520-0485(1995)025<2909:UOICFB>2.0.CO;2, 1995. 2182
- Ekman, W.: On the influence of the earth's rotation on ocean-currents., *Ark. Mat. Astron. Fys.*, 2, 1–52, 1905. 2181

Sea ice inertial oscillations

F. Gimbert et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Sea ice inertial oscillations

F. Gimbert et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Gascard, J., Festy, J., le Goff, H., Weber, M., Bruemmer, B., Offermann, M., Doble, M., Wadhams, P., Forsberg, R., Hanson, S., Skourup, H., Gerland, S., Nicolaus, M., Metaxian, J., Grangeon, J., Haapala, J., Rinne, E., Haas, C., Heygster, G., Jakobson, E., Erko, T., Wilkinson, J., Kaleschke, L., Claffey, K., Elder, B., and Bottenheim, J.: Exploring Arctic transpolar drift during dramatic sea ice retreat, *EOS*, 89, 21–22, 2008. 2193
- 5 Geiger, C. and Perovich, D.: Springtime ice motion in the Western Antarctic Peninsula region, *Deep-Sea Res. Pt. II*, 55, 338–350, 2008. 2184, 2202
- Gill, A. E.: On the behavior of inertial waves in the wakes of storms, *J. Phys. Oceanogr.*, 14, 1129–1151, 1984. 2182
- 10 Gimbert, F., Jourdain, N. C., Marsan, D., Weiss, J., and Barnier, B.: Recent mechanical weakening of the Arctic sea ice cover as revealed by larger sea ice inertial oscillations, *J. Geophys. Res.*, 117, C00J12, doi:10.1029/2011JC007633, 2012. 2184, 2202
- Haas, C., Pfaffling, A., Hendricks, S., Rabenstein, L., Etienne, J., and Rigor, I.: Reduced ice thickness in Arctic transpolar drift favors rapid ice retreat, *Geophys. Res. Lett.*, 35, L17501, doi:200810.1029/2008GL034457, 2008. 2181
- 15 Hunkins, K.: Inertial oscillations of fletcher’s ice island (T-3), *J. Geophys. Res.*, 72, 1165–1174, doi:196710.1029/JZ072i004p01165, 1967. 2184
- Hutchings, J., Heil, P., Steer, A., and Hibler III, W. D.: Sub-synoptic scale spatial variability of sea ice deformation in the Western Weddell sea during early summer, *J. Geophys. Res.*, 117, C01002, doi:10.1029/2011JC006961, 2012. 2184
- 20 Kwok, R. and Rothrock, D. A.: Decline in Arctic sea ice thickness from submarine and icesat records: 1958–2008, *Geophys. Res. Lett.*, 36, L15501, doi:10.1029/2009GL039035, 2009. 2180, 2195, 2196, 2200, 2202
- Kwok, R., Cunningham, G. F., and Hibler III, W. D.: Sub-daily sea ice motion and deformation from RADARSAT observations, *Geophys. Res. Lett.*, 30, 2218, 2003. 2184
- 25 Kwok, R., Pedersen, L. T., Gudmandsen, P., and Pang, S. S.: Large sea ice outflow into the nares strait in 2007, *Geophys. Res. Lett.*, 37, L03502, doi:201010.1029/2009GL041872, 2010. 2181
- Lammert, A., Brummer, B., and Kaleschke, L.: Observation of cyclone-induced inertial sea-ice oscillation in fram strait, *Geophys. Res. Lett.*, 36, L10503, doi:10.1029/2009GL037197, 2009. 2184
- 30

Sea ice inertial oscillations

F. Gimbert et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Le Cann, B. and Serpette, A.: Intense warm and saline upper ocean inflow in the southern bay of Biscaya in autumn-winter 2006–2007, *Cont. Shelf Res.*, 29, 1014–1025, doi:10.1016/j.csr.2008.11.015, 2009. 2185
- Lepparanta, M.: The drift of sea ice, Springer, 2004. 2182, 2189
- 5 Lepparanta, M., Oikkonen, A., Shirasawa, K., and Fukamachi, Y.: A treatise on frequency spectrum of drift ice velocity, *Cold Reg. Sci. Technol.*, 76–77, 83–91, doi:10.1016/j.coldregions.2011.12.005, 2012. 2183
- Lindsay, R. W. and Stern, H. L.: The radarsat geophysical processor system: quality of sea ice trajectory and deformation estimates, *J. Atmos. Ocean Tech.*, 20, 1333–1347, 2003. 2185
- 10 Lindsay, R. W., Zhang, J., Schweiger, A., Steele, M., and Stern, H.: Arctic sea ice retreat in 2007 follows thinning trend, *J. Climate*, 22, 165–175, 2009. 2180
- Maslanik, J., Stroeve, J., Fowler, C., and Emery, W.: Distribution and trends in Arctic sea ice age through spring 2011, *Geophys. Res. Lett.*, 38, L13502, doi:10.1029/2011GL047735, 2011. 2196
- 15 McPhee, M.: A simulation of inertial oscillation in drifting pack ice, *Dynam. Atmos. Oceans*, 2, 107–122, doi:10.1016/0377-0265(78)90005-2, 1978. 2184, 2202
- Nansen, F.: Oceanography of the North Polar basin: the Norwegian North Polar Expedition 1893–96, *Scientific Results*, 3, 1902. 2181
- Nghiem, S., Rigor, I., Perovich, D., Clemente-Colón, P., Weatherly, J., and Neumann, G.:
20 Rapid reduction of Arctic perennial sea ice, *Geophys. Res. Lett.*, 34, L19504, doi:10.1029/2007GL031138, 2007. 2181
- Park, J. J., Kim, K., and Schmitt, R. W.: Global distribution of the decay timescale of mixed layer inertial motions observed by satellite-tracked drifters, *J. Geophys. Res.*, 114, C11010, doi:200910.1029/2008JC005216, 2009. 2182
- 25 Price, J. F.: Internal wave wake of a moving storm. Part I. Scales, energy budget and observations, *J. Phys. Oceanogr.*, 13, 949–965, doi:10.1175/1520-0485(1983)013<0949:IWVOAM>2.0.CO;2, 1983. 2182
- Rajchenbach, J.: Granular flows, *Adv. Phys.*, 49, 229–256, 2000. 2181
- Rampal, P., Weiss, J., and Marsan, D.: Positive trend in the mean speed and deformation rate of Arctic sea ice, 1979–2007, *J. Geophys. Res.*, 114, C05013, doi:200910.1029/2008JC005066, 2009. 2180, 2181, 2183, 2193, 2194, 2198, 2199, 2200, 2202
- 30

Sea ice inertial oscillations

F. Gimbert et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Rampal, P., Weiss, J., Dubois, C., and Campin, J.-M.: Ipcc climate models do not capture Arctic sea ice drift acceleration: consequences in terms of projected sea ice thinning and decline, *J. Geophys. Res.*, 116, C00D07, doi:10.1029/2011JC007110, 2011. 2181, 2202
- 5 Richter-Menge, J. A. and Elder, B. C.: Characteristics of pack ice stress in the alaskan Beaufort sea, *J. Geophys. Res.*, 103, 21817–21829, doi:199810.1029/98JC01261, 1998. 2183
- Richter-Menge, J. A., McNutt, S. L., Overland, J. E., and Kwok, R.: Relating Arctic pack ice stress and deformation under winter conditions, *J. Geophys. Res.*, 107, 8040, doi:200210.1029/2000JC000477, 2002. 2183
- 10 Rothrock, D. A., Percival, D. B., and Wensnahan, M.: The decline in Arctic sea-ice thickness: separating the spatial, annual, and interannual variability in a quarter century of submarine data, *J. Geophys. Res.*, 113, C05003, doi:200810.1029/2007JC004252, 2008. 2180, 2194, 2202
- Serreze, M. C. and Barrett, A. P.: The summer cyclone maximum over the Central Arctic Ocean, *J. Climate*, 21, 1048–1065, 2008. 2193, 2195
- 15 Steele, M., Zhang, J., Rothrock, D., and Stern, H.: The force balance of sea ice in a numerical model of the Arctic Ocean, *J. Geophys. Res.*, 102, 21061–21079, doi:199710.1029/97JC01454, 1997. 2182
- Stroeve, J., Serreze, M., Drobot, S., Gearheard, S., Holland, M., Maslanik, J., Meier, W., and Scambos, T.: Arctic sea ice extent plummets in 2007, *Eos Trans. AGU*, 89, 2, 13 pp., doi:200810.1029/2008EO020001, 2008. 2180
- 20 Weiss, J., Schulson, E. M., and Stern, H. L.: Sea ice rheology from in-situ, satellite and laboratory observations: fracture and friction, *EPSL*, 255, 1–8, doi:10.1016/j.epsl.2006.11.033, 2007. 2183
- Zhang, J., Rothrock, D., and Steele, M.: Recent changes in Arctic sea ice: the interplay between ice dynamics and thermodynamics, *J. Climate*, 13, 3099–3114, 2000. 2180
- 25




Fig. 1. Map of the Arctic basin showing the buoy trajectories of the IABP dataset. The positions are sampled every 3h from January 1979 to December 2008 and plotted following a stereographic projection centered on the North Pole. The Laptev sea is poorly covered by this dataset. The thick solid black and grey lines define the Central Arctic basin and the Fram strait region, respectively.

Sea ice inertial oscillations

F. Gimbert et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	






Fig. 2. Map of the Atlantic Ocean showing the oceanic buoy trajectories deployed during the Congas project. The positions are sampled every 1 h and plotted following a Lambert projection.

Sea ice inertial oscillations

F. Gimbert et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	






Fig. 3. (a) 1 month sample of the trajectory of buoy B245 (25 May to 25 June 2007). The trajectory is plotted in latitude-longitude coordinates. Its beginning is marked by the red circle. The red and green boxes outline periods with strong and low cycloidal loop activity, respectively. (b) Fourier spectrum of the buoy velocity. The velocity is computed following Eq. (3) with $\Delta t = 1$ h.

Sea ice inertial oscillations

F. Gimbert et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Sea ice inertial oscillations

F. Gimbert et al.




Fig. 4. (a) 1 month sample of the trajectory of IABP buoy 1897 (30 August to 30 September 1987). The location of the buoy is indicated by the red box in the inset of Fig. 4a. The red boxes show periods when cycloidal loops are best observed, whereas the amplitude of the loops is much lower in the green boxes. (b) Fourier spectrum of the buoy velocity.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Sea ice inertial oscillations

F. Gimbert et al.




Fig. 5. (a) 1 month sample of the trajectory of buoy 12825 (20 April to 21 May 1992). The location of the buoy within the Arctic basin is indicated by the red box in the inset of Fig. 5a. (b) Fourier spectrum of the buoy velocity.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 6. Example of buoy trajectories computed from Eq. (12) using **(a)** $V/U = 2$, giving $M = 1.2$, **(b)** $V/U = 1$, giving $M = 1$, **(c)** $V/U = 0.5$, giving $M = 0.6$ and **(d)** $V/U = 0.15$, giving $M = 0.2$.

Sea ice inertial oscillations

F. Gimbert et al.




Fig. 7. M values for **(a)** the trajectory of the oceanic buoy 255 plotted on Fig. 3a, **(b)** the trajectory of the ice-tethered buoy 1897 plotted on Fig. 4a and **(c)** the trajectory of the buoy 12825 plotted on Fig. 5a. The red and green rectangles respectively correspond to those shown in Figs. 3 and 4.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Sea ice inertial oscillations

F. Gimbert et al.




Fig. 8. (a) M values for the trajectory of the oceanic buoy 255 plotted on Fig. 3 (raw data: blue line, same as Fig. 7a) and the trajectory obtained after a cubic interpolation and a 3 h resampling of the raw positions (red line). (b) M_D time series computed from the degraded datasets of missing data ratio varying between 0 and 60 % (the red line is the same as in a). (c) M_D values averaged over the whole trajectory (1 month) as a function of the missing data ratio. 20 realisations have been done at each given value of missing data ratio. The grey dashed line shows the average raw M value, equal to 0.56.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Sea ice inertial oscillations

F. Gimbert et al.




Fig. 9. Average annual cycle in the *M* values.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Sea ice inertial oscillations

F. Gimbert et al.




Fig. 10. Spatial pattern of inertial oscillation magnitude within the Arctic basin in summer (top) and in winter (bottom). These two fields are computed from the seasonal M dataset. A spatially averaged value of M is computed following Eq. (13) for each point of a 25 km resolution grid. The graphic representation is a linear interpolation of the gridded \bar{M} -values.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Sea ice inertial oscillations

F. Gimbert et al.




Fig. 11. Time series $M(t)$ from January 1979 to December 2008 computed using **(a)** the summer dataset and **(b)** the winter dataset. For each season, the dataset is equally splitted in 10 bins over which the average is computed. One bin corresponds to approximately 3.1×10^4 M -values in summer and 8.0×10^4 M -values in winter. The horizontal lines associated to each datapoint indicate the time period associated to each bin. Vertical lines are errorbars computed from a bootstrap method. Bold lines are linear fits: the trends are $1.19 (\pm 0.34) \times 10^{-5} \text{ yr}^{-1}$ (i.e., 16,5% increase per decade) for summer and $5.7 (\pm 1.9) \times 10^{-6} \text{ yr}^{-1}$ (i.e., 11% increase per decade) for winter.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I◀

▶I

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Sea ice inertial oscillations

F. Gimbert et al.




Fig. 12. Influence of buoy positions' noise on the monthly mean time series $M(t)$ for **(a)** the summer season and **(b)** the winter season. Gaussian noise, with a constant standard deviation σ , has been added to all buoy positions recorded since year 2002. Results for noised mean time series $M_N(t)$ (in yellow and green) for $\sigma = 300\text{m}$ and $\sigma = 1000\text{m}$ are compared with the IABP mean time series $M(t)$ as on Fig. 11.

Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Sea ice inertial oscillations

F. Gimbert et al.




Fig. 13. Spatial repartition of M within the Arctic basin in summer and winter computed using the seasonal dataset of the period 1979–2001 and the seasonal dataset of the period 2002–2008. An average mean value of M is computed following Eq. (13) for each node of a 25 km resolution grid. The smoothing parameter L is equal to 400 km.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Sea ice inertial oscillations

F. Gimbert et al.




Fig. 14. Mean time series $M(t)$ **(a)** in summer (red) and **(b)** in winter (blue), as on Fig. 11 and $M_0(t)$ (black lines), resulting from the null hypothesis. The best linear fit are also plotted, showing that changes in spatial sampling only account for a small fraction of the observed trend.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

