

Net accumulation rates derived from ice core stable isotope records

M. Schwikowski et al.

Net accumulation rates derived from ice core stable isotope records of Pío XI glacier, Southern Patagonia Icefield

M. Schwikowski^{1,2,3}, M. Schläppi^{1,2,3}, P. Santibañez⁴, A. Rivera^{4,5}, and G. Casassa⁴

¹Paul Scherrer Institut (PSI), Villigen, Switzerland

²Department for Chemistry and Biochemistry, University of Bern, Switzerland

³Oeschger Centre for Climate Change Research, University of Bern, Switzerland

⁴Centro de Estudios Científicos (CECS), Valdivia, Chile

⁵Departamento de Geografía, Universidad de Chile, Santiago, Chile

Received: 28 November 2012 – Accepted: 11 December 2012

– Published: 19 December 2012

Correspondence to: M. Schwikowski (margit.schwikowski@psi.ch)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

Pío XI, the largest glacier of the Southern Patagonia Icefield, reached its neoglacial maximum extent in 1994 and is one of the few glaciers in that area which is not retreating. In view of the recent warming it is important to understand glacier responses to climate changes. Due to its remoteness and the harsh conditions in Patagonia, no systematic mass balance studies have been performed. In this study we derived net accumulation rates for the period 2000 to 2006 from a 50 m (33.2 4 m weq) ice core collected in the accumulation area of Pío XI (2600 m a.s.l., 49°16'40'' S, 73°21'14'' W). Borehole temperatures indicate near temperate ice, but the average melt percent is only 16 % ± 14 %. Records of stable isotopes are well preserved and were used for identification of annual layers. Net accumulation rates range from 3.4 to 7.1 water equivalent (m weq) with an average of 5.8 m weq, comparable to precipitation amounts at the Chilean coast, but not as high as expected for the Icefield. Ice core stable isotope data correlate well with upper air temperatures and may be used as temperature proxy.

1 Introduction

The Southern Patagonia Icefield (SPI) is the largest ice body of the Southern Hemisphere outside Antarctica with an area of 13 000 km² and 48 major glaciers (Aniya et al., 1996), many of them calving into fjords and lakes. Since the end of the Little Ice Age (LIA, late 19th Century) most of the Patagonian glaciers have been retreating and thinning, contributing significantly to sea-level rise (Glasser et al., 2011), with only a few in a state of equilibrium or advance (Casassa et al., 2002; Masiokas et al., 2009a,b). Glacier retreat is interpreted primarily as a response to regional atmospheric warming and to a lesser extent to precipitation decrease (Casassa et al., 2002). Among the advancing ones is Pío XI (also called Brügger), the largest glacier of SPI with a length of 65.7 km and 1277 km² total area in 2003. This glacier has the longest record of frontal

Net accumulation rates derived from ice core stable isotope records

M. Schwikowski et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Net accumulation rates derived from ice core stable isotope records

M. Schwikowski et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

variation observations in the SPI and reached its neoglacial maximum extent in 1994, when it destroyed mature 400-yr old trees (Rivera et al., 1997a; Warren and Rivera, 1994; Rivera and Casassa, 1999). Between 1995 and 1999 Pío XI was relatively stable with an overall minor retreat. By 2000 the glacier readvanced again, reaching a position close to the 1994-maximum extent in 2008 (Masiokas et al., 2009b). The anomalous behaviour of Pío XI has been attributed to a surging, although the mechanism is not clear (Rivera and Casassa, 1999; Rivera et al., 1997b). Considering recent warming in Southern Patagonia and the Antarctic Peninsula it is important to understand glacier variations and their responses to climate change (Cook et al., 2005; Rosenblüth et al., 1995; Thomas et al., 2009). A quick response to climate change is expected for glaciers with high accumulation and ablation rates (Oerlemans and Fortuin, 1992). However, changes of equilibrium line altitude (ELA), front, volume, and ice thickness have so far only be deduced from maps, aerial photos, and satellite data. No systematic glacier mass balance or other in-situ data have been obtained yet from Pío XI glacier.

The climate of the SPI is dominated by prevailing intense Westerlies, which are slightly stronger in summer compared to winter (Carrasco et al., 2002). They continuously carry humid air masses from the Pacific Ocean towards the Andes which form an efficient barrier from North to South, resulting in a strong west-east gradient of precipitation with about 7000 mm per year on the Chilean coast and less than 200 mm east of the Andes (Carrasco et al., 2002; Schneider et al., 2003; Villalba et al., 2003). At the Chilean coast more than 300 days of rain per year were observed and mean annual cloudiness exceeds 85 % (Warren and Sugden, 1993). Phenomena like El Niño have minor influence on the Patagonian climate (Warren and Sugden, 1993). Extreme weather conditions were recorded during an ice drilling expedition on Tyn-dall glacier, SPI, with wind speeds of 13.6 ms^{-1} (hourly average, measured over 24 h, 1756 m a.s.l.), snow accumulation of 3.5 m (1.75 m weq) within 24 days from 30 November to 23 December 1999, and visibility of more than 50 m only on 4 out of 24 days (Kohshima et al., 2007).

Precipitation and temperature data required to understand glacier responses are available only from coastal sites or east of SPI. Weather stations show homogenous distribution of precipitation throughout the year (Carrasco et al., 2002) with a decreasing trend for Lago Argentino (period 1940–1990), (Ibarzabal et al., 1996), Puerto Montt (1961–2005) and Coyhaique (1961–2005) (Carrasco et al., 2008). Only for Punta Arenas a slight increase of precipitation is reported for the period 1965–2005 (Carrasco et al., 2008). No longer-term data is available between 48 and 52° S, the latitude of SPI.

Temperature is like precipitation poorly documented in this region. West of SPI typical maritime influence with small seasonal but high annual variations is observed (Enomoto and Nakajima, 1985), whereas east of SPI more continental climate with larger temperature fluctuation during the year prevails (Rosenblüth et al., 1995). Minimum temperatures for the period 1961–2006 show a slight warming south of 47° S and a cooling of 0.04 to 0.18 °C per 10 yr between 40 and 47° S. Most of these tendencies were attributed to the 1976/77 climate shift (Carrasco et al., 2008). For the more recent period 1979–2006, temperature trends over land in southern Chile (38° S–48° S) are weak and show no clear spatial patterns (Falvey and Garreaud, 2009). At higher altitude (850 hPa), NCEP-NCAR reanalysis data reveal a warming trend of 0.5 °C over 40 yr, both in winter and summer, resulting in a shift from snow to rain of ~ 5 % of the precipitation, whereas the total amount has changed little (Rasmussen et al., 2007).

In principle, an accumulation history can be obtained from glacier ice cores, but only few cores with meaningful climatic information have been obtained from Patagonia. This deficiency is due to extremely harsh field conditions with stormy weather and high precipitation rates throughout the year, and to the fact that the low-altitude main plateaus of both the Northern Patagonia Icefield (NPI) and the SPI are strongly influenced by meltwater percolation. Meltwater may cause a loss of climatic signal, which was reported for the lower plateau area of NPI and SPI, e.g. at San Rafael glacier, 1296 m a.s.l. (Yamada, 1987), Nef glacier, 1500 m a.s.l. (Matsuoka and Naruse, 1999), and Tyndall glacier, 1756 m a.s.l. (Shiraiwa et al., 2002). Accumulation rates of several

Net accumulation rates derived from ice core stable isotope recordsM. Schwikowski et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

meters of snow per year were deduced from the preserved part of the stable isotope record, but they are representative only for the year before drilling.

There are a few high-elevated (> 2000 m) potential drilling sites on mountains in the Icefields, and from three of them shallow cores were collected. Records from Perito Moreno glacier (Aristarain and Delmas, 1993), Gorra Blanca Norte (Schwikowski et al., 2006), and San Valentín (Vimeux et al., 2008) showed well preserved chemical and stable isotope signals, less perturbed by melting. Net snow accumulation rates are lower at higher elevated drilling sites, with 1.2 m weq at Perito Moreno (2000 m, 50°38' S, 73°15' W, SPI), 1 m weq at Gorra Blanca Norte (2300 m, 49°8' S, 73°3' W, SPI), and 0.19 m weq at San Valentín (3747 m, 46°35' S, 73°19' W, NPI). All three sites are located east of the ice divide of the Icefields, where lower precipitation rates are expected, but nonetheless, there might be net loss by erosion due to the intense winds (Vimeux et al., 2011).

In this regard, the upper reaches of Pío XI glacier appear to be an excellent candidate to obtain net accumulation information, because it is located at an altitude where little influence of melting is expected, and where wind erosion should not be so critical, since it forms an extended, gently sloping plateau. In August 2006 a 50 m ice core was drilled in the accumulation area (49°16' S, 73°21' W, 2600 m a.s.l.).

In the present study the characteristics of the glacier and the ambient climate are investigated. The potential of the glacier for ice core-based reconstruction of accumulation and temperature is discussed.

2 Sampling site and methods

2.1 Expedition and sampling site

Based on glacier topography, results from a first study in 2001 (Schwikowski et al., 2006), and a reconnaissance flight on 14 April 2006, the northern plateau of Cordón Mariano Moreno (CMM) was selected for ice coring (49°16'40" S, 73°21'14" W, 2600 m

Net accumulation rates derived from ice core stable isotope records

M. Schwikowski et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



a.s.l., Figs. 1 and 2). The site is part of the accumulation area of Pío XI glacier. A possible escape route allowing descending the glacier on skis autonomous from helicopter in case of bad weather conditions was also explored during the reconnaissance flight.

Drilling camp was set up on 19 August 2006 at a location, where a radar survey with a 6 MHz system indicated glacier thicknesses of 170 m. On the northern CMM plateau glacier thickness varied from 70 to more than 300 m. Drilling was performed with an updated version of the electromechanical drill described by Ginot et al. (2002) producing cores with a diameter of 8.2 cm and a maximum core length of 75 cm. The drill was operated in a particularly rigid tent which was developed for harsh field conditions. This tent, manufactured by FS INVENTOR AG Switzerland, consists of an aluminium frame and a plasticized canvas cover. It was easy to set up and resisted the strong winds in Patagonia. After two days of drilling a depth of 50.6 m was reached. At this depth temperate ice containing slight amounts of liquid water was encountered and drilling had to be stopped because of problems with the transport of drilling chips. Borehole temperatures were measured every 10 m with a temperature probe fixed to the drive unit of the ice drill. The temperature probe was equipped with a PT100 sensor incorporated in an aluminium disc which was in contact with the ice. The probe was connected to a transmitter which generated a signal between 4 and 20 mA, where 4 mA corresponded to a temperature of -50°C and 20 mA to a temperature of $+10^{\circ}\text{C}$, respectively (all components from Greisinger Electronic GmbH, Germany, personal communication D. Stampfli). A network of nine stakes was set up around the drilling site, where detailed surface topography was measured with a dual-frequency GPS receiver. Stake heights, which were initially 1 m high at the most, were remeasured after 10 days, at the end of the campaign. Due to heavy snow storms and high wind speeds the drilling camp could not be evacuated before 2 September 2006. The ice core was shipped frozen to the Paul Scherrer Institut.

Net accumulation rates derived from ice core stable isotope records

M. Schwikowski et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.2 Ice core analyses

Processing of the ice core segments was performed in a cold room (-20°C) at the Paul Scherrer Institut. All cores were weighed, length and diameter were measured to determine density, and stratigraphic features were recorded. To avoid any possible contamination a band saw with a stainless steel blade and a Teflon-coated table was used to cut out inner segments of the ice core (Eichler et al., 2000). Sample resolution varied from 3 to 6.7 cm, with an average of 4.7 cm. The melt percent profile was generated with the same 4.7 cm average resolution. Melt features appear bright and bubble-free when the core is backlit. They were quantified by visual evaluation into classes of 0, 25, 50, and 100 %, representing melt layer thicknesses of 0, 1.2, 2.4, 3.5, and 4.7 cm, respectively. Samples were melted at room temperature. Analysis of major ion concentrations and stable isotopes were conducted at Paul Scherrer Institut, Switzerland, whereas biovolume and pollen concentration was determined at Centro de Estudios Científicos, Valdivia, Chile.

The stable isotope ratio $\delta^{18}\text{O}$ was measured by injecting samples in a high temperature (1250°C) reactor, where water is pyrolysed to carbon monoxide (CO) and transported to the mass spectrometer (Finnigan Delta Plus XP) using helium as carrier gas (Gehre et al., 2004). For δD determination the sample was reduced in a chromium reactor prior to mass spectrometric analysis (Finnigan Mat Delta S) (Nelson and Dettman, 2001). Precision was 0.1 ‰ for $\delta^{18}\text{O}$ and 0.5 ‰ for δD .

Ice core samples for biological analysis were cut in 20 cm sections from 0 to 40 m, and 10 cm sections from 40 to 50 m, after removing 0.5 cm of the core surface for eliminating potential contaminants. Sample preparation was conducted according to Santibanez et al. (2008). Microalgae and pollen grains were counted using a fluorescent microscope (OLYMPUS BX-FLA). The total cell number of microalgae and pollen of *Nothofagus* spp. was estimated on each filter by counting the cells along five to seven parallel transects. In the case of *Podocarpaceae*, all pollen grains present were counted.

Net accumulation rates derived from ice core stable isotope records

M. Schwikowski et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3 Results and discussion

3.1 Borehole temperatures and density profile

5 Temperatures measured in the borehole were just slightly below 0 °C, indicating near temperate ice (Fig. 3). Annual mean air temperature at 700 hPa, an altitude roughly corresponding to 3000 m, for the period 2001–2007 was –8.2 °C, ranging from –0.36 °C to –12.44 °C (monthly values, grid 49.5° S, 73.5° W, ERA-Interim). The elevated ice temperature indicates input of latent heat from refreezing of surface melt water formed in summer. Ice lenses from 0.1 to 19 cm thickness, formed by refreezing, are present throughout the core, but alternate with pure firn. The corresponding melt percent averaged over 1 m depth ranges from 0 to 58 % with a mean value of 16 % ± 14 % (Fig. 3).
10 Latent heat released by freezing of one gram of water raises the temperature of 160 g of snow by one degree (Cuffey and Paterson, 2010). A simple estimation based on mean values of the entire core suggests that the latent heat release due to the observed amount of ice lenses would result in warming by 8.7 °C (mean density: 0.66 g cm⁻³, total snow mass per cm⁻²: 3340 g, total ice mass formed by refreezing of melt water: 182 g cm⁻² (refrozen melt water filling the pore space of 0.66 g cm⁻³ dense firn)). The resulting warming is in the order of magnitude needed to explain the difference between the estimated annual mean air temperature and the measured borehole temperatures.

20 The density increases from 0.32 g cm⁻³ at the surface to 0.89 g cm⁻³ at 50.6 m depth with an average value of 0.66 g cm⁻³ (Fig. 3). This indicates that the firn-to-ice transition was probably reached at a depth of 50.6 m, which would explain the observed small amounts of liquid water at this depth. Densities are slightly higher than simulated with a simple empirical model of firn densification in the dry firn zone (Herron and Langway, 1980), especially in the sections where melt percentages exceed 20 % (Fig. 3). Compared to temperate glaciers, the firn-ice transition at Pío XI occurs at greater depths,
25 suggesting that this part of the glacier belongs to the percolation zone using the classification scheme of Cuffey and Paterson, 2010.

Net accumulation rates derived from ice core stable isotope records

M. Schwikowski et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.2 Stable isotope ratios and net accumulation rates

Stable isotope ratios $\delta^{18}\text{O}$ and δD in the Pío XI core fluctuate regularly around a mean value of $-15.1\text{‰} \pm 2.2\text{‰}$ for $\delta^{18}\text{O}$ and $-111.2\text{‰} \pm 18.1\text{‰}$ for δD , respectively (Fig. 4). Amplitudes between maximum and minimum values are similar throughout the core, indicating that the $\delta^{18}\text{O}$ and δD records are not significantly affected by meltwater percolation, consistent with the low average melt percent of 16%. Melting leads in the extreme case to a total loss of signal variability in a depth of a few meters as observed for other ice cores from Patagonia (Matsuoka and Naruse, 1999) or to a reduction of the amplitude in the seasonal stable isotope variation. The latter was reported for ice cores from the Arctic with average melt percent values of 40% (Goto-Azuma et al., 2002) and 55% (Pohjola et al., 2002), significantly exceeding the amount of melt observed in the Pío XI core. The stable isotope ratios $\delta^{18}\text{O}$ and δD in the Pío XI core show significant correlation ($\delta\text{D} = 8.0 \times \delta^{18}\text{O} + 9.2$, $r = 0.98$, $n = 1078$) and the slope and intercept are in good agreement with the Global Meteoric Water Line ($\delta\text{D} = 8 \times \delta^{18}\text{O} + 10$), pointing to a direct marine origin of precipitation and no continental influence. This good agreement is another indicator for a well-preserved stable isotope record, since melting and refreezing cycles cause a decrease of the slope (Zhou et al., 2008). From the Global Network of Isotopes in Precipitation (GNIP) it is known that in the southern temperate zone of South America (south of 40°S), like in the northern, the isotopic composition of precipitation is controlled mainly by changes in temperature, with a maximum of $\delta^{18}\text{O}$ and δD during austral summer and a minimum during winter (Rozanski and Araguas Araguas, 1995). This relationship between stable isotopes in precipitation and air temperature is illustrated in Fig. 5 on the example of the GNIP station Coyhaique ($45^\circ 21' 00''\text{S}$, $72^\circ 4' 12''\text{W}$, 310 m a.s.l.) (IAEA/WMO, 2006) which has the longest record in vicinity of the Southern Patagonia Icefield. Studies of the stable isotope-air temperature relation with a general circulation model (LMDZ4) confirmed this significant positive correlation for Patagonia (Risi et al., 2010). Precipitation in South America is maximal around 50°S , with over 300 days of rainfall per year in

Net accumulation rates derived from ice core stable isotope records

M. Schwikowski et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

some places (Kerr and Sugden, 1994). The amount of precipitation is nearly uniformly distributed throughout the year (Carrasco et al., 2002) as illustrated in Fig. 6 on the example of the coastal stations Cabo Raper (46°48'12" S, 75°38'17" W, 46 m a.s.l.), San Pedro (47°43'0" S, 74°55'0" W, 56 m a.s.l.), and Faro Evangelista (52°24'0" S, 75°6'0" W, 52 m a.s.l.). We therefore assume that the major fluctuations of $\delta^{18}\text{O}$ and δD in the Pío XI ice core do reflect seasonal temperature variations and can be used to identify annual layers as suggested in Fig. 4. Accordingly the ice core covers the period winter 2000 to winter 2006. $\delta^{18}\text{O}$ and δD maxima are attributed to February, generally the month with the highest temperatures.

Average algal biovolume in the Pío XI core is only $43\text{ mm}^3\text{ mL}^{-1}$ (range 0–590 $\text{mm}^3\text{ mL}^{-1}$) which is two to three orders of magnitude lower than values found at other glacier sites (Kohshima et al., 2007; Santibanez et al., 2008; Uetake et al., 2006; Yoshimura et al., 2000). Thus it was impossible to derive seasonal boundaries based on algal biovolume. *Nothofagus* and *Podocarpaceae* pollen concentrations are also low, but nevertheless they show four maxima each, corresponding closely to the summer maxima identified in the $\delta^{18}\text{O}$ and δD records (Fig. 4). Pollen peaks occur generally earlier than the $\delta^{18}\text{O}$ maxima attributed to February, which is reasonable considering that flowering starts during the austral spring (Hechenleitner et al., 2005). The absence of pollen during a particular spring-summer season does not necessarily represent periods without pollen dispersal since the trees do not flower regularly and in some years widespread non-flowering occurs. Thus, the pollen records alone do not allow dating, but they confirm the identification of annual layers based on stable isotopes.

The proposed dating results in high annual net accumulation rates of 3.4 to 7.1 m weq, with an average of 5.8 m weq. Note that a year represents the period 1 February to 31 January. The average net accumulation rate at Pío XI is comparable to precipitation amounts at the Chilean coast, but not as high as the 10 m weq expected for the Icefield (Carrasco et al., 2002). Erosion due to wind drift is a relevant factor that affects snow accumulation, as was detected in the field based on stake observations

Net accumulation rates derived from ice core stable isotope records

M. Schwikowski et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

during the two weeks campaign between 16 August and 2 September 2006 (Fig. 7). The drilling site, located on a high plateau only 300 m from a prominent ridge, is exposed to very high wind, generally from the west, which effectively reduces the snow accumulation. However, this effect is strongest in the proximity of the ridge, where wind is enhanced, and less pronounced at the drilling site. The amount of snow erosion can not be quantified, thus, we consider the obtained net accumulation rates as lower limits. Compared to accumulation rates of 1.2 m weq and 0.36 m weq deduced from Perito Moreno (Aristarain and Delmas, 1993) and 0.36 m weq from San Valentín (Vimeux et al., 2008), two other ice core sites at the eastern margin, the net accumulation rate at Pío XI is much higher. This suggests a strong west-east precipitation gradient within the SPI and/or stronger wind erosion effects at Perito Moreno and San Valentín.

3.3 Relationship between stable isotopes and temperature

For identifying annual layers, the stable isotope records were used, assuming that they reflect regional temperature. In order to check the potential of $\delta^{18}\text{O}$ and δD as temperature proxy, the records were compared to temperature at 700 hPa (approximately 2850 m a.s.l.) from upper air radiosonde data collected at Punta Arenas, 460 km south-east of Pío XI. The depth scale of the ice core was transformed linearly into a time scale starting on 12 December 2000 and ending on 1 September 2006, using the maxima in the stable isotope record, which were assigned to the highest temperature in the Southern Hemisphere summer. This procedure is justified because of the uniform distribution of precipitation throughout the year. All data sets (stable isotope series and air temperature), were averaged to yield three-day means, which represents the lowest resolution in the deepest part of the ice core. Both, $\delta^{18}\text{O}$ and δD data series agree reasonably well with the air temperatures. Pearson correlation coefficients are $r = 0.63$ ($p < 0.001$, $n = 573$) for the individual data and $r = 0.81$ ($p < 0.001$, $n = 19$) for the 29 point moving averages for the correlation between $\delta^{18}\text{O}$ and temperature (Fig. 8). The corresponding $\delta^{18}\text{O}$ /air temperature ratio of $0.54\text{‰}\text{°C}^{-1}$ is in good agreement with ratios from the Antarctic Peninsula ($0.56\text{‰}\text{°C}^{-1}$, Aristarain et al., 1986) and from Argentine Island

Net accumulation rates derived from ice core stable isotope records

M. Schwikowski et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



($0.59 \pm 0.4 \text{‰} \text{°C}^{-1}$, Rozanski et al., 1993). This corroborates that the Pío XI stable isotope composition is mainly controlled by temperature.

It is well known that $\delta^{18}\text{O}$ and δD of precipitation become more negative with increasing altitude (altitude effect). This is attributed to the progressive condensation of atmospheric vapour and rainout of the condensed phase when air masses undergo lifting and adiabatic cooling. This dependence was investigated for $\delta^{18}\text{O}$ values of different ice core records from SPI and NPI and the instrumental stations Punta Arenas, Puerto Montt and Coyhaique which belong to the Global Network of Isotopes in Precipitation (GNIP). For the GNIP data, monthly values from 1995 to 1998 were averaged, because this period was available for all stations (Fig. 9). Assuming a linear relation between altitude and $\delta^{18}\text{O}$ an altitude lapse rate of $0.32 \text{‰} 100 \text{m}^{-1}$ is obtained, comparable to values found in most regions of the world below 5000 m elevation (Poage and Chamberlain, 2001). The $\delta^{18}\text{O}$ average at Pío XI (-15.1‰) agrees well with the stable isotope composition of precipitation at the same altitude at 33°S in Chile (Rozanski and Araguas Araguas, 1995) and thus reflects depletion due to orographic lifting and rainout from the Pacific as the dominant moisture source.

4 Conclusions

The 50 m ice core collected from the accumulation area of Pío XI (2600 m a.s.l., $49^\circ 16' \text{S}$, $73^\circ 21' \text{W}$) is the first core from the Southern Patagonia Icefield, showing well preserved records of stable isotopes throughout. This was expected at the altitude of 2600 m a.s.l. for which annual mean temperatures in the range of -8 to -11°C were estimated. However, borehole temperatures indicated near-temperate ice, contradicting the good preservation of the stable isotope signals. We assume that most meltwater formed at the surface or rain infiltrating the upper firn layers immediately refroze, forming impermeable ice layers observed in the core and increasing the ice temperature due to latent heat release. Meltwater relocation and drainage cannot be

Net accumulation rates derived from ice core stable isotope records

M. Schwikowski et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

totally excluded, but the amount is limited. Otherwise the stable isotope signals would not be preserved.

Annual net accumulation derived from annual layers identified in the profiles of stable isotopes and biological markers ranged from 3.4 to 7.1 m weq. These values are similar to precipitation rates at the Chilean coast, much higher than at other high-elevation sites further to the east, but not as high as expected for the Icefield. This suggests that even at the flat plateau of the Pío XI site influence of snow drift and erosion can not totally be excluded, as was observed during some days in the field. Thus, the derived net accumulation rates present lower limits.

A reasonably good correlation of the stable isotope data with upper air temperatures indicates that the Pío XI stable isotopes are controlled by temperature. They also show the expected altitude effect. Thus, the 50 m Pío XI core contains besides precipitation (net accumulation) a temperature proxy signal (stable isotopes).

Acknowledgements. Special thanks go to the drilling team which did an exceptional job in the field. Precipitation data were kindly provided by Jorge Carrasco, Dirección Meteorológica de Chile. Financial support from National Geographic (Grant No. 7587-04), FONDECYT Centro de Estudios Científicos, Valdivia, Chile (Grant No. 1061269), and the Swiss State Secretariat for Education and Research SER (Grant No. CJRP-1004) is highly acknowledged.

References

- Aniya, M., Sato, H., Naruse, R., Skvarca, P., and Casassa, G.: The use of satellite and airborne imagery to inventory outlet glaciers of the Southern Patagonia Icefield, South America, *Photogramm. Eng. Rem. S.*, 62, 1361–1369, 1996.
- Aristarain, A. J. and Delmas, R. J.: Firn-core study from the Southern Patagonia ice cap, South America, *J. Glaciol.*, 39, 249–254, 1993.
- Aristarain, A. J., Jouzel, J., and Pourchet, M.: Past Antarctic Peninsula climate (1850–1980) deduced from an ice core isotope record, *Climatic Change*, 8, 69–89, 1986.
- Carrasco, J. F., Casassa, G., and Rivera, A.: Meteorological and climatological aspects of the Southern Patagonia icefield, in: *The Patagonian Icefields: a Unique Natural Laboratory for*

Net accumulation rates derived from ice core stable isotope records

M. Schwikowski et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Net accumulation rates derived from ice core stable isotope records

M. Schwikowski et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Environmental and Climate Change Studies, edited by: Casassa, G., Sepulveda, F., and Sinclair, R. M., Kluwer Academic/Plenum Publishers, New York, 29–41, 2002.
- Carrasco, J. F., Osorio, R., and Casassa, G.: Secular trend of the equilibrium-line altitude on the western side of the southern Andes, derived from radiosonde and surface observations, *J. Glaciol.*, 54, 538–550, 2008.
- Casassa, G., Rivera, A., Aniya, M., and Naruse, R.: Current knowledge of the Southern Patagonia icefield, in: *The Patagonian Icefields: a Unique Natural Laboratory for Environmental and Climate Change Studies*, edited by: Casassa, G., Sepulveda, F., and Sinclair, R. M., Kluwer Academic/Plenum Publishers, New York, 67–83, 2002.
- Cook, A. J., Fox, A. J., Vaughan, D. G., and Ferrigno, J. G.: Retreating glacier fronts on the Antarctic Peninsula over the past half-century, *Science*, 308, 541–544, 2005.
- Cuffey, K. M. and Paterson, W. S. B.: *The Physics of Glaciers*, Fourth edition, Elsevier, Burlington, MA, USA, 2010.
- Eichler, A., Schwikowski, M., Gäggeler, H. W., Furrer, V., Synal, H. A., Beer, J., Saurer, M., and Funk, M.: Glaciochemical dating of an ice core from upper Grenzgletscher (4200 m a.s.l.), *J. Glaciol.*, 46, 507–515, 2000.
- Enomoto, H. and Nakajima, C.: Recent climate-fluctuations in Patagonia, in: *Glaciological studies in Patagonia Northern Icefield, 1983–1984*, edited by: Nakajima, C., *Jap. Soc. Snow and Ice*, Nagoya, Japan, 7–14, 1985.
- Falvey, M. and Garreaud, R. D.: Regional cooling in a warming world: recent temperature trends in the southeast Pacific and along the west coast of subtropical South America, *J. Geophys. Res.*, 114, D04102, doi:10.1029/2008jd010519, 2009.
- Gehre, M., Geilmann, H., Richter, J., Werner, R. A., and Brand, W. A.: Continuous flow H-2/H-1 and (18)O/O-16 analysis of water samples with dual inlet precision, *Rapid Commun. Mass Sp.*, 18, 2650–2660, 2004.
- Ginot, P., Stampfli, F., Stampfli, D., Schwikowski, M., and Gäggeler, H.: Felics, a new ice core drilling system for high-altitude glaciers, *Workshop “Ice Drilling Technology 2000”*, *Mem. Natl. Inst. Pol. Res. Spec. Issue*, 56, 38–48, 2002.
- Glasser, N. F., Harrison, S., Jansson, K. N., Anderson, K., and Cowley, A.: Global sea-level contribution from the Patagonian Icefields since the Little Ice Age maximum, *Nat. Geosci.*, 4, 303–307, 2011.
- Goto-Azuma, K., Koerner, R. M., and Fisher, D. A.: An ice-core record over the last two centuries from Penny ice cap, Baffin Island, Canada, *Ann. Glaciol.*, 35, 29–35, 2002.

Net accumulation rates derived from ice core stable isotope records

M. Schwikowski et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Hechenleitner, V., Gardner, M. F., Thomas, P. I., Echeverria, C., Escobar, B., Brownless, P., and Martinez, C.: Plantas amenazadas del centro sur de Chile: distribución, conservación y propagación, Universidad Austral de Chile y Real Jardín Botánico de Edimburgo, 188, 2005.
- Herron, M. M. and Langway Jr., C. C.: Firn densification: an empirical model, *J. Glaciol.*, 25, 373–385, 1980.
- Ibarzabal, T., Donangelo, T., Hoffmann, J. W., and Naruse, R.: Recent climate changes in southern Patagonia, *B. Glacier Res.*, 14, 29–36, 1996.
- Kerr, A. and Sugden, D.: The sensitivity of the South Chilean snowline to climatic-change, *Climatic Change*, 28, 255–272, 1994.
- Kohshima, S., Takeuchi, N., Uetake, J., Shiraiwa, T., Uemura, R., Yoshida, N., Matoba, S., and Godoi, M. A.: Estimation of net accumulation rate at a Patagonian glacier by ice core analyses using snow algae, *Global Planet. Change*, 59, 236–244, 2007.
- Masiokas, M. H., Luckman, B. H., Villalba, R., Delgado, S., Skvarca, P., and Ripalta, A.: Little Ice Age fluctuations of small glaciers in the Monte Fitz Roy and Lago del Desierto areas, south Patagonian Andes, Argentina, *Palaeogeogr. Palaeoclimatol.*, 281, 351–362, 2009a.
- Masiokas, M. H., Rivera, A., Espizua, L. E., Villalba, R., Delgado, S., and Aravena, J. C.: Glacier fluctuations in extratropical South America during the past 1000 years, *Palaeogeogr. Palaeoclimatol.*, 281, 242–268, 2009b.
- Matsuoka, K. and Naruse, R.: Mass balance features derived from a firn core at Hielo Patagónico Norte, South America, *Arct. Antarct. Alp. Res.*, 31, 333–340, 1999.
- Nelson, S. T. and Dettman, D.: Improving hydrogen isotope ratio measurements for on-line chromium reduction systems, *Rapid Commun. Mass Sp.*, 15, 2301–2306, 2001.
- Oerlemans, J. and Fortuin, J. P. F.: Sensitivity of glaciers and small ice caps to greenhouse warming, *Science*, 258, 115–117, 1992.
- Poage, M. A. and Chamberlain, C. P.: Empirical relationships between elevation and the stable isotope composition of precipitation and surface waters: considerations for studies of paleoelevation change, *Am. J. Sci.*, 301, 1–15, 2001.
- Pohjola, V. A., Moore, J. C., Isaksson, E., Jauhiainen, T., van de Wal, R. S. W., Martma, T., Meijer, H. A. J., and Vaikmae, R.: Effect of periodic melting on geochemical and isotopic signals in an ice core from Lomonosovfonna, Svalbard, *J. Geophys. Res.*, 107, 4036, doi:10.1029/2000JD000149, 2002.
- Rasmussen, L. A., Conway, H., and Raymond, C. F.: Influence of upper air conditions on the Patagonia icefields, *Global Planet. Change*, 59, 203–216, 2007.

Net accumulation rates derived from ice core stable isotope recordsM. Schwikowski et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Risi, C., Bony, S., Vimeux, F., and Jouzel, J.: Water-stable isotopes in the LMDZ4 general circulation model: model evaluation for present-day and past climates and applications to climatic interpretations of tropical isotopic records, *J. Geophys. Res.*, 115, D12118, doi:10.1029/2009jd013255, 2010.
- 5 Rivera, A. and Casassa, G.: Volume changes on Pio XI glacier, Patagonia: 1975–1995, *Global Planet. Change*, 22, 233–244, 1999.
- Rivera, A., Aravena, J. C., and Casassa, G.: Recent fluctuations of Glaciar Pio XI, Patagonia: discussion of a glacial surge hypothesis, *Mt. Res. Dev.*, 17, 309–322, 1997a.
- Rivera, A., Lange, H., Aravena, J. C., and Casassa, G.: The 20th Century advance of Glaciar Pío XI, Southern Patagonia icefield, *Ann. Glaciol.*, 24, 66–71, 1997b.
- 10 Rosenblüth, B., Casassa, G., and Fuenzalida, H.: Recent climatic changes in western Patagonia, *B. Glac. Res.*, 13, 127–132, 1995.
- Rozanski, A. and Araguas Araguas, L.: Spatial and temporal variability of stable isotope composition of precipitation over the South American continent, *B. Inst. Fr. Etud. Andin.*, 24, 370–390, 1995.
- 15 Santibanez, P., Kohshima, S., Scheihing, R., Jaramillo, J., Shiraiwa, T., Matoba, S., Kanda, D., Labarca, P., and Casassa, G.: Glacier mass balance interpreted from biological analysis of firn cores in the Chilean lake district, *J. Glaciol.*, 54, 452–462, 2008.
- Schneider, C., Glaser, M., Kilian, R., Santana, A., Butorovic, N., and Casassa, G.: Weather observations across the Southern Andes at 53° S, *Phys. Geogr.*, 24, 97–119, 2003.
- 20 Schwikowski, M., Brütsch, S., Casassa, G., and Rivera, A.: A potential high-elevation ice-core site at Hielo Patagonico Sur, *Ann. Glaciol.*, 43, 8–13, 2006.
- Shiraiwa, T., Kohshima, S., Uemura, R., Yoshida, N., Matoba, S., Uetake, J., and Godoi, M. A.: High net accumulation rates at Campo de Hielo Patagonico Sur, South America, revealed by analysis of a 45.97 m long ice core, *Ann. Glaciol.*, 35, 84–90, 2002.
- 25 Thomas, E. R., Dennis, P. F., Bracegirdle, T. J., and Franzke, C.: Ice core evidence for significant 100-year regional warming on the Antarctic Peninsula, *Geophys. Res. Lett.*, 36, L20704, doi:10.1029/2009GL040104, 2009.
- Uetake, J., Kohshima, S., Nakazawa, F., Suzuk, K., Kohno, M., Kameda, T., Arkhipov, S., and Fujii, Y.: Biological ice-core analysis of Sofiyskiy glacier in the Russian Altai, *Ann. Glaciol.*, 43, 70–78, 2006.
- 30 Villalba, R., Lara, A., Boninsegna, J. A., Masiokas, M., Delgado, S., Aravena, J. C., Roig, F. A., Schmelter, A., Wolodarsky, A., and Ripalta, A.: Large-scale temperature changes across

Net accumulation rates derived from ice core stable isotope records

M. Schwikowski et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

the southern Andes: 20th-century variations in the context of the past 400 years, *Climatic Change*, 59, 177–232, 2003.

Vimeux, F., de Angelis, M., Ginot, P., Magand, O., Casassa, G., Pouyaud, B., Falourd, S., and Johnsen, S.: A promising location in Patagonia for paleoclimate and paleoenvironmental reconstructions revealed by a shallow firn core from Monte San Valentin (Northern Patagonia Icefield, Chile), *J. Geophys. Res.*, 113, D16118, doi:10.1029/2007JD009502, 2008.

Vimeux, F., Maignan, F., Reutenauer, C., and Pouyaud, B.: Evaluation of cloudiness over Monte San Valentin, Northern Patagonia Icefield, from 2000 to 2008 using MODIS satellite images: implications for paleoclimate investigations from ice cores, *J. Glaciol.*, 57, 221–230, doi:10.3189/002214311796405915, 2011.

Warren, C. R. and Rivera, A.: Nonlinear climatic response of calving glaciers – a case-study of Pio-Xi Glacier, Chilean Patagonia, *Rev. Chil. Hist. Nat.*, 67, 385–394, 1994.

Warren, C. R. and Sugden, D. E.: The Patagonian Icefields – a glaciological review, *Arctic Alpine Res.*, 25, 316–331, 1993.

Yamada, T.: Glaciological characteristics revealed by 37.6-m deep core drilled at the accumulation area of San Rafael Glacier, the Northern Patagonia Icefield, *B. Glac. Res.*, 4, 59–67, 1987.

Yoshimura, Y., Kohshima, S., Takeuchi, N., Seko, K., and Fujita, K.: Himalayan ice-core dating with snow algae, *J. Glaciol.*, 46, 335–340, 2000.

Zhou, S. Q., Nakawo, M., Hashimoto, S., and Sakai, A.: The effect of refreezing on the isotopic composition of melting snowpack, *Hydrol. Process.*, 22, 873–882, 2008.




Fig. 1. Left: Map of Southern South America with Northern Patagonia Ice Field (NPI, Campo de Hielo Norte) and location of Pío XI glacier at the Southern Patagonia Ice Field (SPI, Campo de Hielo Sur). The red triangle marks the drilling site. Upper right: Landsat ETM, acquired on 2 April 2003, showing the area of Pío XI glacier with ice front position in 1945 (red line), 2003 (black), and 2005 (orange). Lower right: The northern plateau of Cordón Mariano Moreno including topographic information (contour lines) extracted from SRTM.

Net accumulation rates derived from ice core stable isotope records

M. Schwikowski et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	




Fig. 3. Density profile (blue), temperature (red) in the borehole at Pío XI glacier, and melt percent (black). The melt percent represents averages over 1 m depth. Green dashed lines show densities simulated with an empirical model of firn densification in the dry firn zone (Herron and Langway, 1980), using a firn temperature of -1°C and an annual accumulation rate of 7.1 m weq (left curve) and 3.4 m weq (right curve), respectively.

Net accumulation rates derived from ice core stable isotope records

M. Schwikowski et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Net accumulation rates derived from ice core stable isotope records

M. Schwikowski et al.




Fig. 4. Pío XI ice core stable isotope ratios $\delta^{18}\text{O}$, δD (raw data and 29 point moving average) and pollen records. Proposed attribution of years is indicated with vertical dashed lines.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Net accumulation rates derived from ice core stable isotope records

M. Schwikowski et al.




Fig. 5. Relationship between $\delta^{18}\text{O}$ in precipitation and air temperature at the GNP station Coyhaique ($45^{\circ}21'00''\text{S}$, $72^{\circ}4'12''\text{W}$, 310 m a.s.l.) (IAEA/WMO, 2006).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 6. Monthly averaged precipitation at the coastal sites Cabo Raper (46°48'12" S, 75°38'17" W, 46 m a.s.l., period 1968–2003), San Pedro (47°43'0" S, 74°55'0" W, 56 m a.s.l., period 1968–2003), and Faro Evangelista (52°24'0" S, 75°6'0" W, 52 m a.s.l., period 1899–2003).

Net accumulation rates derived from ice core stable isotope records

M. Schwikowski et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 7. Stake network (solid black circles) deployed around the drilling site (empty circle with x). The numbers at each stake represent snow accumulation in cm day^{-1} . The thick blue line shows the location of a prominent ridge which marks the ice divide between Pío XI Glacier and Viedma Glacier to the North and to the East. The brown lines are contour lines with elevations in m a.s.l., obtained from the SRTM 2000 data. All stakes located lower than 2600 m were lost due to snow burial, showing a snow accumulation larger than 7 cm day^{-1} . At the highest stake, snow accumulation was practically zero, which is due to enhanced wind in the proximity of the ridge. The background image is a Quickbird scene of 26 December 2004, obtained from Google Earth, which shows a crevasse immediately Northeast of the ridge, and some rock outcrops farther downslope.

Net accumulation rates derived from ice core stable isotope records

M. Schwikowski et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



Net accumulation rates derived from ice core stable isotope records

M. Schwikowski et al.




Fig. 8. Pío XI $\delta^{18}\text{O}$ record (thin black line: raw data set adjusted to 100 points per year, thick black line: 29 point moving average) compared to 700 hPa temperature data from Punta Arenas (thin red line: daily raw data set adjusted to 100 points per year, thick red line: 29 point moving average) and to 700 hPa temperature reanalysis data from ERA-Interim (grey dashed line, monthly values, nearest grid point 49.50°S , 73.50°W). For procedure to transform ice core depth scale in time scale see text.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Net accumulation rates derived from ice core stable isotope records

M. Schwikowski et al.




Fig. 9. Mean $\delta^{18}\text{O}$ values as function of elevation compared for different ice cores from SPI and NPI (circles) and nearest GNIP stations (squares). Ice core data from San Valentín, Gorra Blanca, Perito Moreno, Tyndall, Nef, and San Rafael are from (Aristarain and Delmas, 1993; Matsuoka and Naruse, 1999; Schwikowski et al., 2007; Shiraiwa et al., 2002; Vimeux et al., 2008; Yamada, 1987).

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)