

## ***Interactive comment on “Influence of ablation-related processes in the built-up of simulated Northern Hemisphere ice sheets during the last glacial cycle” by S. Charbit et al.***

**R. J. Braithwaite (Referee)**

roger.braithwaite@manchester.ac.uk

Received and published: 14 February 2013

General Comments:

This is a very interesting paper and tackles an important issue for ice sheet modelling.

In the first sentence of his seminal paper, Reeh (1991) says “Models of the dynamics and thermodynamics of ice sheets and glaciers depend on the boundary conditions at the ice-sheet surface involving mass balance and surface temperature”. Cynics will say that Charbit et al (2012) arrive at a very similar conclusion after 22 years of more work by many people. In particular, Charbit et al (2012) find that glacier inception depends critically on the representation of temperature variability for their chosen mass balance

C3014

model (the positive degree-day PDD model) as well as the representation of meltwater refreezing. They do briefly discuss alternative mass balance models, e.g. the (more fundamental?) energy balance approach, although they still prefer the PDD model for modelling over the last glacial cycle.

In this review, I confine myself to commenting on the temperature variability and the refreezing, which lie well within my areas of competence. The other review, by L. Tarasov, is wider ranging.

At several points in their paper (e.g. line 19 on page 4902), Charbit et al (2012) say that the PDD method was first proposed by Reeh (1991). This is untrue and must be corrected. Braithwaite (1977, 1984) was the first to formulate the contribution of random temperature variations to the positive degree-day total (PDD), and to identify the standard deviation of temperature variations (denoted by  $\sigma$  in the present paper) as an important parameter for calculation of PDD. The annual melting was then calculated by the MB1 model (Braithwaite and Thomsen, 1989) as the sum of monthly melt values calculated from mean temperatures for individual months using the Braithwaite (1984) model. The contribution of Reeh (1991) was to estimate these monthly mean temperatures from an annual temperature cycle described by a sine wave and then apply these temperatures to the calculation of annual PDD using the  $\sigma$  parameter. Aside from its applications to ice sheet modelling, the modified PDD approach of Reeh (1991) explains the long-known empirical relation between summer mean temperature and annual accumulation at the glacier ELA (Ohmura et al, 1992 and Braithwaite, 2008).

From the above, I suggest changing “The PDD method has first been proposed by Reeh (1991)” to say “The original PDD method of Braithwaite (1984) was modified by Reeh (1991), and many workers use his modified version”. I think this fairly summarises the situation and gives proper credit to both myself and the Niels Reeh.

Braithwaite (1984) calculated monthly degree-days for standard deviations in the range

C3015

of 1, 2, 3, 4 and 5 deg in his Table 1 but assumed  $\sigma = 4$  deg for the two examples he discussed. From my own later work, I would now prefer the lower part of this range, e.g.  $\sigma = 2$  to 3 deg, but Charbit et al (2012) correctly say that the higher part of the range has been commonly used in the literature. I personally find the suggestion of Fausto et al (2009) quite plausible that  $\sigma$  increases from a relatively low value near the ice sheet margin, with higher temperatures, to a higher value at higher altitudes and lower temperatures.

It is still difficult to describe meltwater refreezing in a physically realistic way that is computationally economic (Reijmer et al. 2012). Reeh (1991) refreezes all melt up to 60% of the annual accumulation and allows runoff for melt in excess of this 60%. He gives no explanation or reference for this 60% figure, which therefore looks very arbitrary and ad hoc, but there is a sound physical basis for this figure. In the late 1980s, Niels Reeh and I discussed the refreezing of meltwater, and Reeh was aware of my MB1 mass balance model (Braithwaite and Thomsen, 1989) where refreezing was estimated using a simple density mixing model, see equation (4) in Braithwaite et al (1994). In this model, snow must be converted to impermeable ice by meltwater refreezing before any runoff is allowed. The ratio of annual melt to annual accumulation at the runoff line is given by the densities of impermeable ice and pre-melt snow. With the values that we assumed for these densities in the late 1980s, the melt/accumulation ratio at the runoff line is 0.6, and Reeh (1991) assumes this value without explanation. After a very strenuous field trip in spring 1992 to measure firn densities in the lower accumulation area of the Greenland ice sheet, Braithwaite et al (1994) refined this ratio to 0.58. Pfeffer et al. (1991) estimate a ratio of 0.7 with a more refined refreezing model but Braithwaite et al (1994) suggest this can be adjusted down to 0.62 if the Braithwaite density values are substituted into the Pfeffer equation. Shumskii (1964, p. 416) suggests a melt/accumulation ratio of only 0.4 at the runoff line but that can be adjusted upwards using the densities from Braithwaite et al (1994). The simple density mixing model (Braithwaite et al, 1994) does not allow any runoff from melting snow, while in real life there may be some runoff from a shallow snow cover, with underlying

C3016

ice, if the snow has been brought to the melting point by latent heat release from refrozen meltwater. The density mixing model may therefore slightly overestimate the amount of refreezing.

I hope the reader will not regard the above comments as too much of a historical digression but the simple point is that the refreezing scheme of Reeh (1991) is more physically based than it looks and may even be more realistic than some later schemes.

Some detailed comments

Title of paper: "build-up" and not "built-up".

Lines 2-3 on page 4899: Please give a standard definition of mass balance.

Lines 22-24 on page 4899: In my papers, I always cite Finsterwalder and Schunk (1887) for stating that melting only occurs under positive temperatures, which is certainly important for the development of the PDD concept, but the word "method" here is going too far.

Lines 26-28 on page 4899: Yes, it is fair to say that Reeh (1991) "proposed a new formulation of the PDD method".

Line 13 on page 4900: Does anyone suggest using a single DDF value for snow and ice? I think the authors mean one DDF value for snow and another DDF value for ice.

Line 19 on page 4902: As said earlier Reeh (1991) did not first propose the PDD method. See remarks above.

Line 12 on page 4902: There is no real reason to assume  $\sigma = 4$  deg.

Line 20 on page 4902: The cited author's name is "Braithwaite" and not "Braithwaithe".

Lines 23-25 on page 4902: The maximum amount of superimposed ice cannot exceed 60% of the annual accumulation and the 60% figure is not "somewhat arbitrary" (see my comments above).

C3017

Lines 6-23 on page 4903: Again, the PDD formulation of Reeh (1991) is not “early”. You should put in a reference to Braithwaite (1995) somewhere in this paragraph as he reviews these the high DDF values and shows that extrapolation of the surface energy balance to lower temperatures will lead to higher values of DDF. When performing the analyses described by Braithwaite (1995) I was more concerned about the possible changes in degree-day factors with rising temperatures, for example under global warming, than with changes at lower temperatures.

Equations (4) and (5) on p. 4905: I am concerned that these inequalities will give “steps” in computed mass balance, although they do partly take account of the temperature dependence of the degree-day factors predicted by the energy balance model (Braithwaite, 1995).

Line 5 on page 4907: There is no real justification for  $\sigma = 4.5$  to  $5.5$  deg.

Sections 3 on page 4907 to section 5 ending on page 4920: I have no comments.

Lines 18-19 on page 4920. The whole purpose of Braithwaite (1995) was to study variations in degree-day factor by extrapolating the energy balance. Although not the immediate purpose of the present paper, I think somebody should do more work in this area. It may be correct that energy balance models are not suitable for on-line forcing of ice sheets over longer time scales but temporal and geographical variations in degree-day factors could be more thoroughly studied by running advanced energy balance models off-line, i.e. by updating the basic approach of Braithwaite (1995) with more sophisticated energy balance models at more sites.

#### References:

Braithwaite, R. J. Air temperature and glacier ablation - a parametric approach. McGill University, Montreal, Canada, 1978.

Braithwaite, R. J. Calculation of degree-days for glacier-climate research. *Zeitschrift für Gletscherkunde und Glazialgeologie* 20, 1-8, 1984.

C3018

Braithwaite, R. J. and Thomsen, H. H. Simulation of runoff from the Greenland ice sheet for planning hydro-electric power, Ilulissat/Jakobshavn, West Greenland. *Ann. Glaciol.* 13, 12-15, 1989.

Braithwaite, R. J., Laternser, M. and Pfeffer, W. T. Variations of near-surface firn density in the lower accumulation area of the Greenland ice sheet, Pâkitsoq, West Greenland. *J. Glaciol.* 40, 136, 477-485, 1994.

Braithwaite, R. J. Positive degree-days factors for ablation on the Greenland ice sheet studied by energy-balance modelling. *J. Glaciol.* 41, 137, 153-160, 1995.

Braithwaite, R. J. Temperature and precipitation climate at the equilibrium-line altitude of glaciers expressed by the degree-day factor for melting snow. *J. Glaciol.* 54, 186, 437-444, 2008.

Charbit, S., Dumas, C., Kageyamam, M., Roche, D. M. and Ritz, C. Influence of ablation-related processes in the build-up of simulated Northern Hemisphere ice sheets during the last glacial cycle. *The Cryosphere Discuss.* 6, 4897-4938, 2012.

Fausto, R. S., Ahlstrom, A. P., van As, D., Johnsen, S. J., Langen, P. L. and Steffen, K. Improving surface boundary conditions with focus on coupling snow densification and meltwater retention in large-scale ice-sheet models of Greenland. *J. Glaciol.* 55, 193, 869-878, 2009.

Finsterwalder, S. and Schunk, H. *Der Suldenferner.* Z. Deutsch. Öster. Alpenver. 18, 72-89, 1887.

Ohmura, A., Kasser, P. and Funk, M. Climate at the equilibrium line of glaciers, *J. Glaciol.* 38, 130, 397-411., 1992.

Pfeffer, W. T., Meier, M. F. and Illangasekare, T. H. Retention of Greenland runoff by refreezing: implications for projected future sea level changes. *J. Geophys. Res.* 96, 22117-22124, 1991.

C3019

Reeh, N. Parameterization of melt rate and surface temperature on the Greenland ice sheet. *Polarforschung* 59, 113-128, 1991.

Reijmer, C. H., van den Broeke, M. R., Fettweis, X, Etterna, J. and Stap, L. B. Refreezing on the Greenland ice sheet: a comparison of parameterizations. *The Cryosphere* 6, 743-763, 2012.

Shumskii, P. A. *Principles of structural glaciology*. New York, Dover Publications, 1964.

---

Interactive comment on *The Cryosphere Discuss.*, 6, 4897, 2012.