

The 2013 Erebus
Glacier tongue
calving event

C. L. Stevens et al.

This discussion paper is/has been under review for the journal The Cryosphere (TC).
Please refer to the corresponding final paper in TC if available.

Brief communication

“The 2013 Erebus Glacier tongue calving event”

C. L. Stevens^{1,2}, P. Sirguey³, G. H. Leonard³, and T. G. Haskell⁴

¹National Institute for Water and Atmospheric Research (NIWA), Greta Point Wellington, New Zealand

²University of Auckland, Dept. Physics, New Zealand

³National School of Surveying, University of Otago, Dunedin, New Zealand

⁴Callaghan Innovation, Gracefield Lower Hutt, New Zealand

Received: 25 March 2013 – Accepted: 30 March 2013 – Published: 25 April 2013

Correspondence to: C. L. Stevens (c.stevens@niwa.cri.nz)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

The Erebus Glacier Tongue, a small floating glacier in southern McMurdo Sound, is one of the best-studied ice tongues in Antarctica. Despite this, its calving on the 27 February 2013 (UTC) was around 10 yr earlier than previously predicted. The calving was likely a result of ocean currents and the absence of fast ice. The subsequent trajectory of the newly-created iceberg supports previous descriptions of the surface ocean circulation in southern McMurdo Sound.

1 Introduction

The Erebus Glacier Tongue (EGT) is a small glacier extending from the western coast of Ross Island into Southern McMurdo Sound. Prior to the events described here the glacier tongue was approximately 12.9 km long (measured from the virtual line connecting the coastline where it intersects with EGT), about 1.5 km wide and was estimated to be 300 m thick at the grounding line. The glacier is protected by the Dellbridge Islands (including Tent and Big Razorback Islands) to the north (Fig. 1). The area is close to Cape Evans, the base for Robert Falcon Scott's 1911–12 expedition, and so was the focus for some of the earliest geophysical observations in Antarctica (Taylor, 1922). Curious features are the glacier's aspect ratio (narrow relative to width) and the striking side-lobes (Holdsworth, 1974, 1982).

On the 27 February 2013 (UTC), and for only the 4th time in the brief recorded history of Antarctica, the Erebus Glacier Tongue calved. It broke off around 1 km west of a fast ice bridge extending south west from Big Razorback Island. In doing so it freed a ~ 3.3 km section of glacier (measured along its longest axis) to form a 3.2 km² penguin-like shaped iceberg (Fig. 1a, b). The reduced length of EGT to its new tip is about 9.9 km.

Previous calving events occurred in 1990, ~ 1942 and 1911. The 1911 calving of the Ross Island Tongue (as it was called then) was reported by Taylor (1922) as having

TCD

7, 1749–1760, 2013

The 2013 Erebus Glacier tongue calving event

C. L. Stevens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



broken off during a southerly blizzard on the night of 1 March 1911. There is some discrepancy in Taylor (1922) as to the position of the new end of the EGT after the calving. The position used here comes from Map I presented at the end of the text in Taylor (1922). The 1942 calving date and position are approximate (Holdsworth, 1982) as there was no occupation of the sound at that time. The 1990 event is the best documented previous calving (Robinson and Haskell, 1990), occurring on 1 March of that year – the same year day as the 1911 calving (Fig. 1c, d). These authors identified ocean swell waves coming from the north in the absence of sea ice as being the primary driver of the calving.

In this brief communication we document the calving and contextualize this using available data on tip position and previous calving events. Furthermore, we use the trajectory of the detached tip interpreted from a sequence of MODIS imagery to examine aspects of circulation in southern McMurdo Sound. This is useful information as the ocean currents in this region are poorly known due to the challenges of high latitude ocean survey work.

2 Calving

The weather at the time of the 2013 calving was clear with moderate wind conditions. The McMurdo Sound weather station reported a maximum wind speed of 12 m s^{-1} over the few days prior to calving (www.ncdc.noaa.gov). A number of remotely sensed data products are available for the area. One hundred and fourteen granules from MODIS sensors aboard both TERRA and AQUA satellites were acquired over the period from year day 57 (26 February) to 80 (21 March) and processed to derive projected false colour composite images (bands 1, 2 and 3) at 250 m spatial resolution (Sirguey et al., 2009). The detached tip of the EGT was positively identified in 38 of these images. Furthermore, higher resolution (15 m) ASTER orthorectified VNIR images were obtained before (12 January 2013) and shortly after the event (27 February 20:25 UTC and 3 March 2013). Finally, Landsat 4 TM images from before and after the 1990 calving

The 2013 Erebus Glacier tongue calving event

C. L. Stevens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



were obtained (29 January 1989; 29 November 1989 and 14 December 1992) to revisit the mapping of the 1990 event in the context of the recent data.

Visual interpretation indicates that, prior to the calving there was loose pack ice in the vicinity but no large objects. A few days prior to calving this loose pack amalgamated in a region to the SE of the EGT tip, but was subsequently advected away leaving open ocean at the time of calving (Fig. 1a, b). The 2013 calving separated along a line running to the south east and so removed more from the southern side of the tongue (Fig. 1b). This was also seen in the 1911 calving, although less exaggerated than 2013, where contemporary maps showed perhaps 1 km more was lost from the south side as compared to the north (Taylor, 1922). Sketches and satellite data indicate that the calving line in the 1990 event was almost north-south (Fig. 1d). The variability in the orientation of this line adds to the ambiguity in tip position by as much as 1.2 km.

The EGT did not split off at the multiyear fast ice bridge from Big Razorback Island, but instead along one of the depression lines running SE–NW (Fig. 2a, b). GPS transects of glacier freeboard from December 2010 indicate that these depressions imply a marked localized thinning of the glacier (Stevens et al., 2013).

Glacier calving can be linked to climate drivers (Frezzotti, 1997). The timing of the calving can be placed in context by considering an update of Holdsworth's (1982) space-time trajectory of the glacier tip (Fig. 2c), which was developed prior to the 1990 calving. Whilst not statistically significant, it is nonetheless noteworthy that the time between calving for the 2013 event was the shortest of the observations so far. Robinson and Haskell (1990) had speculated that, based on previous behavior, it should re-calve sometime in 2020–30. By using the Landsat images pre and post 1990, we estimate that the EGT was flowing at an average rate of 153 m a^{-1} at the time, thus allowing the position of the EGT to be approximated as of 1 March 1990. It appeared that the southwards breaking point of the 2013 calving matched precisely the location of the fracture in 1990, although the EGT extended about 600 m farther at that time.

The comparison with historical estimates suggest that the calving line varies. However, the two events in the satellite era show the southern edge of the calving line at

The 2013 Erebus Glacier tongue calving event

C. L. Stevens et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

least to be consistent. This break point may be related to the lobe features as these appear to form the end points for the depression features in the glacier that subsequently formed the calving line.

As well as confirming the “early” calving of the recent event (i.e. earlier than predicted by Robinson and Haskell 1990 based on previous calving intervals), it also indicates a curious arrest in propagation in the 70 s and 80 s. There is very little change from the tip position identified by Jacobs et al. (1981) and the point at which the glacier calved in 1990. Did the side-lobes keep moving – i.e. was it still flowing and just sloughing off small bergs at the tip? This diagram also suggests a gradual slowing in the propagation speed of the tip over the last century by as much as a factor of nearly two. From the satellite era we estimate that the speed of the tip over the period January 1989 to December 1992 was 153 ma^{-1} . It decreased to 113 ma^{-1} between December 2010 and February 2012. Yet in a macro-sense it needs to be moving at 200 ma^{-1} to grow sufficiently to match records (Fig. 2c).

Additionally, the 2013 calving occurred only two days earlier in the year than previous documented occurrences (neither 1911 nor 1990 were leap years). The apparent consistency of this date calls into question the role of storm-driven ocean waves as they are unlikely to always occur in this small time window – and indeed did not in 2013. A number of mechanisms have been explored for failure including the role of ocean swell and excitation of standing waves in the glacier itself (Squire, 1994).

Tidal and ocean currents have not been a strong focus of earlier examinations of calving. Tidal currents certainly influence floating glaciers (Legresy et al., 2004) and might also be considered to be a driver of failure. Synchronicity of calving with spring tides would be evidence of this. However, the present calving took place a day prior to neap tides (Goring and Pyne, 2003). Conversely the calving in 1990 took place a few days before spring tide.

Several things happen to the EGT as it extends out to the longitude of Tent Island. First, it is exposed to currents and waves heading southward past Tent Island and second, it reaches a bathymetric shoal directly south of Tent Island where the water

The 2013 Erebus Glacier tongue calving event

C. L. Stevens et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

column nearly halves in depth (from around 340 m either side) to as shallow as 190 m. There is likely a feedback whereby the shoaling accelerates the flow.

While tides are not the obvious driver of failure here, the residual circulation flows are significant in the present system. Flow speeds in the region comprise 10 cm s^{-1} regional circulation and tidal speeds on top of this of as much as 15 cm s^{-1} (Stevens et al., 2013). The eastern side of the sound is generally thought to flow southward (Robinson and Williams, 2012). However, observations close to shore indicate a northward flow suggesting there might be flow separation around Tent Island and so flowing back towards the NW (Leonard et al., 2006; Stevens et al., 2013). Such currents were observed in observations near the EGT tip in 2010 (Stevens et al., 2013). This back-eddy may be the reason for the northward bias in the glacier axis as it extends into the sound (although Holdsworth 1982 suggested differential melting may drive this). The effect of the southward current at the tip is exacerbated by the northward inshore flow, thus applying an enhanced moment about a vertical axis to the glacier.

The ASTER image of the newly detached berg (Fig. 1b) also reveals information about the multi-year ice filling the spaces between lobes. The detachment process has removed these ice-bridges and so the lobes are far more exaggerated. This phenomenon was encountered during the experiments described in Stevens et al. (2013) where an upwards-looking side-scan sonar traversing beneath the glacier was expecting to see glacier ice but instead only saw multiyear ice.

3 Iceberg trajectory

Iceberg trajectories are generally dominated by ocean currents, so the trajectory followed by the iceberg provides evidence of surface circulation in southern McMurdo Sound. The sequence of events that followed the calving started with an anticlockwise rotation as the berg is driven by the recirculation south of Tent Island. However, as it drifted farther south it moved out of the influence of this back eddy. At this point the berg virtually stopped and rotated about 180° clockwise. This can be explained if there

The 2013 Erebus Glacier tongue calving event

C. L. Stevens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



is a southward flowing jet located west of Tent Island. This jet will have a faster flowing core so that an object moving across the jet experiences a change in the sign of the moment imparted to the berg (Fig. 1b).

By the end of day 59 (28 February) the berg reached the McMurdo Ice Shelf. The bathymetry is poorly mapped in this area, but what data there are indicate depths in excess of 500 m, so the berg was not grounded (see Fig. 3). It remained adjacent to the ice shelf, moving with the tide over a period of approximately 60 h, reaching speeds of 4 cm s^{-1} . Robinson et al. (2010) found average flows of 18 cm s^{-1} to the SE in this region and tidal speeds up to 30 cm s^{-1} . Clearly the berg was not able to accelerate sufficiently rapidly to reach tidal speeds. However, on day 64, the berg commenced moving to the NW at an average speed exceeding 10 cm s^{-1} , carrying with it sea ice that remained attached to the berg until at least day 80 when the berg was visibly trapped within more forming sea ice. This trajectory matches the integrated distributions of platelet growth which is indicative that ice shelf influenced water flows to the northwest at some point as one moves to the west across McMurdo Sound (Gough, 2012).

On previous calving occasions the iceberg was later found grounded or locked in fast ice to the northwest. In 1911 it moved to Marble Point ($77^\circ 15' \text{ S}$; $164^\circ 20' \text{ E}$) still with a pony chaff depot visible. Similarly, the 1990 berg was later observed near Dunlop Island 23 km north of Marble Point with a sampling station intact and visible. A very similar path was taken by an ice tethered profiler (ITP40 <http://www.who.edu/itp/itp40data.html>) mounted in fast ice that broke away in February 2011 and eventually grounded north of Dunlop Island.

The calving event suggests a number of questions for future research. (i) Analysis of the frequency of calving and implications for change in the region. (ii) Determination of the generation of lobes and importantly their transverse linkages (i.e. why did the highly angled grooves form in the most recently growth period?). (iii) Is there a jet-like surface current moving southward past Tent Island that reduces as one moves to the west and then ultimately turns northward?

TCD

7, 1749–1760, 2013

The 2013 Erebus Glacier tongue calving event

C. L. Stevens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Acknowledgements. The MODIS data used in this study were acquired as part of the NASA's Earth-Sun System Division and archived and distributed by the MODIS Adaptive Processing System (MODAPS). ASTER images were obtained with support from the Global Land Ice Measurement from Space (GLIMS). Bathymetry data is from Davey (2004). Craig Stewart, Pat Langhorne, Paul Woodgate and Ross Venell are thanked for discussions. This work is supported by the New Zealand Royal Society Marsden Fund, NIWA Core-funded research and Antarctica New Zealand.

References

- Davey, F. J.: Ross Sea bathymetry, 1 : 2 000 000, ver. 1.0, Geophys. Map 16, Inst. Of Geol. and Nucl. Sci. Ltd., Lower Hutt, New Zealand, 2004.
- Frezzotti, M.: Ice front fluctuation, iceberg calving flux and mass balance of Victoria Land glaciers, *Antarct. Sci.*, 9, 61–73, 1997.
- Goring, D. G. and Pyne, A.: Observations of sea-level variability in Ross Sea, Antarctica, *New Zeal. J. Mar. Fresh.*, 37, 241–249, 2003.
- Gough, A.: Sea ice near an ice shelf, PhD thesis, University of Otago, Otago, 2012.
- Holdsworth, G.: Erebus Glacier Tongue, McMurdo Sound, Antarctica, *J. Glaciol.*, 13, 27–35, 1974.
- Holdsworth, G.: Dynamics of Erebus Glacier tongue, *Ann. Glaciol.*, 3, 131–137, 1982.
- Jacobs, S. S., Huppert, H. E., Holdsworth, G., and Drewry, D. J.: Thermohaline steps induced by melting of the Erebus Glacier Tongue, *J. Geophys. Res.*, 86, 6547–6555, 1981.
- Legresy, B., Wendt, A., Tabacco, I., Remy, F., and Dietrich, R.: Influence of tides and tidal current on Mertz Glacier, Antarctica, *J. Glaciol.*, 50, 427–435, 2004.
- Leonard, G. H., Purdie, C. R., Langhorne, P. J., Haskell, T. G., Williams, M. J. M., and Frew, R. D.: Observations of platelet ice growth and oceanographic conditions during the winter of 2003 in McMurdo Sound, Antarctica, *J. Geophys. Res.* 111, C04012, doi:10.1029/2005JC002952, 2006.
- Robinson, W. and Haskell, T. G.: Calving of Erebus Glacier tongue, *Nature* 346, 615–616, 1990.

The 2013 Erebus Glacier tongue calving event

C. L. Stevens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Robinson, N. J. and Williams, M. J. M.: Iceberg induced changes to polynya operation and regional oceanography in the southern Ross Sea, Antarctica, from in situ observations, *Antarct. Sci.*, 24, 514–526, 2012.
- Robinson, N. J., Williams, M. J. M., Barrett, P. J., and Pyne, A. R.: Observations of flow and iceocean interaction beneath the McMurdo Ice Shelf, Antarctica, *J. Geophys. Res.*, 115, C03025, doi:10.1029/2008JC005255, 2010.
- Sirguey, P., Mathieu, R., and Arnaud, Y.: Subpixel monitoring of the seasonal snow cover with MODIS at 250 m spatial resolution in the Southern Alps of New Zealand: methodology and accuracy assessment, *Remote Sens. Environ.*, 113, 160–181, doi:10.1016/j.rse.2008.09.008, 2009.
- Squire, V. A., Robinson, W. H., Meylan, M., and Haskell, T. G.: Observations of flexural waves on the Erebus Ice Tongue, McMurdo Sound, Antarctica, and nearby sea ice, *J. Glaciol.*, 40, 377–385, 1994.
- Stevens, C. L., Stewart, C. L., Robinson, N. J., Williams, M. J. M., and Haskell, T. G.: Flow and mixing near a glacier tongue: a pilot study, *Ocean Sci.*, 7, 293–304, doi:10.5194/os-7-293-2011, 2011.
- Stevens, C. L., McPhee, M. G., Forrest, A., Leonard, G. H., Stanton, T., and Haskell, T. G.: The influence of an Antarctic glacier tongue on near-field ocean circulation and mixing, *J. Geophys. Res.*, submitted, 2013.
- Taylor, T. G.: *The Physiography of the McMurdo Sound and Granite Harbour region*, Harrison, London, 1922.

The 2013 Erebus Glacier tongue calving event

C. L. Stevens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 1. ASTER images from **(a)** year day 12 (12 January 2013), **(b)** year day 58 (27 February 2013) showing tip moving south and speculated current shear described in the text. Landsat 4 TM images from 1989 showing **(c)** year day 29 (29 January 1989) and **(d)** year day 349 (14 December 1992). The Erebus Glacier Tongue (EGT), Tent Island (TI), Big Razorback Island (BRI) and Hut Peninsula (HP) are marked in **(a)**.

The 2013 Erebus Glacier tongue calving event

C. L. Stevens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The 2013 Erebus Glacier tongue calving event

C. L. Stevens et al.




Fig. 2. ASTER images of EGT (a) before and (b) after the 2013 calving (zoomed from Fig. 1a, b), and (c) tip space-time diagram. The longitudes of Tent Island and the coast are marked. Squares show longitude at calving (hollow square indicates the date is poorly constrained) whilst circles are tip positions. The 1911–12 data come from Taylor (1922), the 1940s estimates are from Holdsworth (1982), the 1950–1990 data from a mix of USGS aerial photographs and vessel observations (Jacobs et al., 1981). The more recent data come from ASTER satellite images.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


The 2013 Erebus Glacier tongue calving event

C. L. Stevens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 3. EGT tip iceberg trajectory built from a sequence of 38 MODIS and 2 ASTER images from year day 57 (26 February 2013) through to year day 80 (21 March 2013). Estimated time in hours since detachment is indicated at each location. The width of the arrow is indicative of the speed. Note the 180° rotation when passing McMurdo Station (H+10 to H+28).