

Mass balances of  
Mera and Pokalde  
glaciers (Nepal) since  
2007

P. Wagnon et al.

This discussion paper is/has been under review for the journal The Cryosphere (TC).  
Please refer to the corresponding final paper in TC if available.

# Seasonal and annual mass balances of Mera and Pokalde glaciers (Nepal Himalaya) since 2007

P. Wagnon<sup>1,2</sup>, C. Vincent<sup>3</sup>, Y. Arnaud<sup>1</sup>, E. Berthier<sup>4</sup>, E. Vuillermoz<sup>5</sup>, S. Gruber<sup>1</sup>,  
M. Ménégoz<sup>3</sup>, A. Gilbert<sup>3</sup>, M. Dumont<sup>6</sup>, J. M. Shea<sup>2</sup>, D. Stumm<sup>2</sup>, and  
B. K. Pokhrel<sup>7</sup>

<sup>1</sup>IRD/UJF – Grenoble 1/CNRS/G-INP, LTHE UMR 5564, LGGE UMR5183, Grenoble, 38402, France

<sup>2</sup>ICIMOD, GPO Box 3226, Kathmandu, Nepal

<sup>3</sup>UJF – Grenoble1/CNRS, Laboratoire de Glaciologie et Géophysique de l'Environnement (LGGE) UMR5183, Grenoble, 38041, France

<sup>4</sup>LEGOS, CNRS, Université de Toulouse, 14 av. Edouard Belin, 31400 Toulouse, France

<sup>5</sup>Ev-K2-CNR Committee, Via San Bernardino, 145 – 24126 Bergamo, Italy

<sup>6</sup>Météo-France – CNRS, CNRM – GAME UMR 3589, CEN, Grenoble, France

<sup>7</sup>DHM, Department of Hydrology and Meteorology, Babarmahal, Kathmandu, Nepal

Received: 21 June 2013 – Accepted: 28 June 2013 – Published: 5 July 2013

Correspondence to: P. Wagnon (patrick.wagnon@ujf-grenoble.fr)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Abstract

In the Everest region, Nepal, ground-based monitoring programs were started on the debris-free Mera Glacier (27.7° N, 86.9° E; 5.1 km<sup>2</sup>, 6420 to 4940 m.a.s.l.) in 2007 and on the small Pokalde Glacier (27.9° N, 86.8° E; 0.1 km<sup>2</sup>, 5690 to 5430 m.a.s.l., ~ 25 km North of Mera Glacier) in 2009. These glaciers lie on the southern flank of the central Himalaya under the direct influence of the Indian monsoon and receive more than 80 % of their annual precipitation in summer (June to September). Despite a large inter-annual variability with glacier-wide mass balances ranging from  $-0.77 \pm 0.40$  m w.e. in 2011–2012 (Equilibrium-line altitude (ELA) at ~ 6055 m.a.s.l.) to  $+0.46 \pm 0.40$  m w.e. in 2010–2011 (ELA at ~ 5340 m.a.s.l.), Mera Glacier has been shrinking at a moderate mass balance rate of  $-0.10 \pm 0.40$  m w.e. yr<sup>-1</sup> since 2007. Ice fluxes measured at two distinct transverse cross sections at ~ 5350 m.a.s.l. and ~ 5520 m.a.s.l. confirm that the mean state of this glacier over the last one or two decades corresponds to a limited mass loss, in agreement with remotely-sensed region-wide mass balances of the Everest area. Seasonal mass balance measurements show that ablation and accumulation are concomitant in summer which in turn is the key season controlling the annual glacier-wide mass balance. Unexpectedly, ablation occurs at all elevations in winter due to wind erosion and sublimation, with remobilized snow likely being sublimated in the atmosphere. Between 2009 and 2012, the small Pokalde Glacier lost mass more rapidly than Mera Glacier with respective mean glacier-wide mass balances of  $-0.72$  and  $-0.26 \pm 0.40$  m w.e. yr<sup>-1</sup>. Low-elevation glaciers, such as Pokalde Glacier, have been usually preferred for in-situ observations in Nepal and more generally in the Himalayas, which may explain why compilations of ground-based mass balances are biased toward negative values compared with the regional mean under the present-day climate.

## Mass balances of Mera and Pokalde glaciers (Nepal) since 2007

P. Wagnon et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 1 Introduction

Since the erroneous statement in the 2007 Intergovernmental Panel on Climate Change (IPCC) Fourth Assessment Report that Himalayan glaciers could disappear in the coming decades if the Earth keeps warming at the current rate (Cogley et al., 2010), these glaciers have drawn the attention of a growing number of scientists (e.g. Bolch et al., 2012; Kääb et al., 2012). Almost every study stresses that major gaps remain in our knowledge of the behavior of Himalayan glaciers mainly due to “insufficient numbers of in-situ measurements, for which remote sensing only partly substitutes” (Bolch et al., 2012). This paucity of appropriate glacier data is of particular concern while considering that Himalayan glaciers have significant social and economic impacts on densely populated regions through water resources (e.g. Kehrwald et al., 2008; Immerzeel et al., 2010; Kaser et al., 2010; Kääb et al., 2012; Nepal et al., 2013; Schaner et al., 2012; Gardelle et al., 2013) or glacial hazards (e.g. Bajracharya and Mool, 2009).

Karakoram-Himalayan glaciers cover the largest glacierized area on Earth outside the polar regions ( $\sim 40\,800\text{ km}^2$ ; Bolch et al., 2012). They experience contrasting precipitation regimes, with a decreasing influence of the westerlies from west to east, and a simultaneous increasing influence of the Indian and Asian monsoons, allowing two striking gradients to emerge: an east-to-west rainfall gradient along the Himalayan foreland, and a south-to-north gradient across the Himalayan range due to topography and relief (Burbank et al., 2003; Bookhagen and Burbank, 2006, 2010). This complex climate diversity results in a contrasting pattern of glacier change throughout the range, mainly revealed by studies using remote sensing data (e.g. Fujita and Nuimura, 2011; Scherler et al., 2011; Bolch et al., 2012; Kääb et al., 2012; Gardelle et al., 2013; Gardner et al., 2013). Ground-based observations everywhere along the entire range are obviously not possible, but in-situ measurements are eagerly needed to confirm remotely-sensed observations and determine some important variables not measurable from space such as annual and seasonal mass balances, mass balance gradients and snow/ice melt factors for degree-day models.

## Mass balances of Mera and Pokalde glaciers (Nepal) since 2007

P. Wagnon et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Our present study contributes to fill this data gap and presents results from extensive mass balance and dynamics measurements conducted in the Everest region (central Himalaya, Nepal), on Mera Glacier since 2007 (Hinku valley), and on Pokalde Glacier since 2009 (Khumbu valley). Recently, Vincent et al. (2013) provided an exhaustive compilation of all in-situ mass balance series available in western Himalaya (Northern India), showing that these series are short-term ( $< 10$  yr usually), discontinuous and not representative of the mass balance of this large region. In central Himalaya and Nepal especially, the situation is similar even though the Japanese conducted some pioneering surveys in the 1970s on AX010 Glacier ( $\sim 0.4$  km<sup>2</sup>, Shorong Himal,  $\sim 30$  km south-west of Mera Glacier), on Yala Glacier ( $\sim 1.9$  km<sup>2</sup>, Langtang Himal – central Nepal) and on Rikha Samba Glacier ( $\sim 4.6$  km<sup>2</sup>, Lower Mustang – western Nepal) (e.g. Ageta et al., 1980; Ageta and Higuchi, 1984; Fujita and Nuimura, 2011). Fujita et al. (2001a) reported systematic negative glacier-wide mass balances for AX010 Glacier between 1978–1999 ( $-0.6$  m w.e. yr<sup>-1</sup> for 1978–1991;  $-1.0$  m w.e. yr<sup>-1</sup> for 1991–1996 and  $-0.8$  m w.e. yr<sup>-1</sup> for 1996–1999) based on long-term surface-elevation change observations and 4 yr of glaciological annual mass balance observations (1995–1999). Fujita and Nuimura (2011) extended this study until 2010 relying on additional geodetic observations conducted in 2008–2010 and using an energy-mass balance model with downscaled gridded climate datasets. They also included Yala and Rikha Samba glaciers in their study and concluded that glaciers have been experiencing negative area-averaged mass balances for the last three decades (from  $\sim -0.75$  m w.e. yr<sup>-1</sup> for Yala and AX010 glaciers to  $\sim -0.5$  m w.e. yr<sup>-1</sup> for Rikha Samba Glacier located in a drier climate). Kadota et al. (2000) followed by Nuimura et al. (2011) also measured surface lowering of Khumbu Glacier between 1978 and 1995. Even though Japanese glaciologists have intermittently observed the changes of the surface elevation of various Nepalese glaciers by geodetic surveys, only four years of annual mass balances are available on AX010 Glacier (1995–1999), making the 5 yr Mera Glacier series presented here the longest continuous series of measured annual mass balance ever recorded in Nepal. Decadal mass balances measured using the



geodetic method are important but they do not replace seasonal mass balance measurements that are needed to analyze the climatic causes of mass balance changes. Since 2011, annual in-situ mass balance surveys on Yala and Rikha Samba glaciers have been conducted by ICIMOD (International Centre for Integrated Mountain Development, Kathmandu, Nepal) and partner institutes.

Remote sensing studies in Nepal also indicate that glacier mass balances are negative on average. Bolch et al. (2011) reported a mass balance of  $-0.32 \pm 0.08$  m.w.e.yr<sup>-1</sup> between 1970 and 2007 over 62 km<sup>2</sup> of glaciers in the Everest area, while for a larger group of glaciers covering 183 km<sup>2</sup> in the same area, Nuimura et al. (2012) measured a mass balance of  $-0.40 \pm 0.25$  m.w.e.yr<sup>-1</sup> over 1992–2008. Recently, Gardelle et al. (2013) found a region-wide mass balance of  $-0.26 \pm 0.09$  m.w.e.yr<sup>-1</sup> over 2000–2011 for a 1461 km<sup>2</sup> glacierized area, in agreement with the value ( $-0.30 \pm 0.09$  m.w.e.yr<sup>-1</sup>) reported by Kääb et al. (2012) for East Nepal and Bhutan for 2003–2008.

Apart from the geographical description of the glaciers and the study area, the aim of this contribution is to present 5 yr of mass balance, annual surface velocity and thickness measurements of Mera Glacier, and to compare to the seasonal mass balances of Pokalde Glacier. The measurements presented in this study will allow a better assessment and understanding of glacier mass balance in this part of Nepal. Indeed, Mera Glacier is planned to become a benchmark glacier for process-understanding in this part of the Himalayan range, following the basic concepts of international climate-related glacier monitoring (“tier-2” level described in Haeberli et al. (2002) or Paul et al., 2007). This glacier is likely to be included into the *GLACIOCLIM* network whose aim is to select a limited number of glaciers representative of various climates of the world in order to better understand the climate-glacier relationship (<http://www-igge.obs.ujf-grenoble.fr/ServiceObs/index.htm>).

## Mass balances of Mera and Pokalde glaciers (Nepal) since 2007

P. Wagnon et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## 2 Site description

Mera Glacier (27.7° N, 86.9° E, 5.1 km<sup>2</sup>) is a debris-free glacier located in Hinku valley (Dudh Koshi basin, Everest region, central Himalaya), and has been monitored since 2007 (Fig. 1). Mera Peak is popular among the climbers because of its gentle slopes (mean slope of ~ 16°) and its 5 to 7 day access from Lukla airport. From the summit at 6420 m.a.s.l., the glacier flows north and divides into two main branches at 5800 m.a.s.l. The main branch flows north and then west down to its snout at 4940 m.a.s.l. while the second branch is north-east orientated with its lowest elevation at 5260 m.a.s.l. These two branches are referred as Mera and Naulek, respectively (Fig. 1). Naulek is also part of the glacierized complex covering the northern slopes of Naulek Peak located ~ 3.5 km south-east of Mera Peak. Table 1 gives a list of geographical and topographical characteristics of Mera Glacier as well as some basic mass balance information calculated according to mass balance data of this study (see Sect. 5.1). Mera Peak is one of the most southern mountains of the region, and thus occupies a frontal position against the Indian monsoon flux (e.g. Bookhagen and Burbank, 2006).

Pokalde Glacier (27.9° N, 86.8° E; 0.1 km<sup>2</sup>) is located ~ 25 km north of Mera Glacier, in Khumbu valley (Fig. 1). Due to its inner position inside the range, this glacier is efficiently sheltered from the wet summer monsoon influx leading to a drier climate than on Mera Glacier (Bookhagen and Burbank, 2006). This north-orientated glacier flows from 5690 to 5430 m.a.s.l., is accessible from Ev-K2-CNR Pyramid permanent research observatory, and has been monitored since 2009 (Fig. 1).

## 3 Climatic setting

Figure 2 shows the mean annual cycle of monthly precipitation and monthly air temperature (years 2003 to 2012) recorded at four meteorological stations operated by Ev-K2-CNR in Lukla (2660 m.a.s.l.), Namche Bazar (3570 m.a.s.l.), Pheriche (4260 m.a.s.l.) and Pyramid (5035 m.a.s.l.) (Fig. 1). Precipitation sensors at these locations are tipping

TCD

7, 3337–3378, 2013

### Mass balances of Mera and Pokalde glaciers (Nepal) since 2007

P. Wagnon et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



5 buckets usually used for rainfall measurements, and may under-catch part of the solid precipitation. Therefore, precipitation at the two highest stations (Pyramid and Pheriche) is likely underestimated. This data set has gaps (listed in Fig. 2 caption) but is sufficient to describe annual local climate cycles. Both air temperature and precipitation present a pronounced seasonality, with more than 80 % of the annual precipitation falling during summer (June to September). Average monthly temperatures have an annual amplitude of approximately 12 °C at every meteorological station. Both studied glaciers are clearly under the influence of the Indian monsoon system (e.g. Webster et al., 1998; Wang, 2006). During the Indian summer monsoon, moisture originating from the Bay of Bengal moves to the north and northwest and initiates heavy orographic rainfall when colliding with the mountain front of eastern and central Himalaya (e.g. Bookhagen and Burbank, 2006). In winter, the overall monsoon-related circulation weakens allowing the westerly upper-tropospheric synoptic-scale waves, dry over Nepal, to be dominant (e.g. Wang, 2006). The central Himalaya is characterized by a two-step topography triggering two rainfall peaks, the outer peak occurring along the southern Himalayan foothills (~ 900 m a.s.l.) and the inner peak along the southern flanks of the Greater Himalaya (~ 2100 m a.s.l.) (Bookhagen and Burbank, 2006, 2010). The glaciers surveyed in this study are located just beyond this inner rainfall peak, and are submitted to a strong south-to-east negative gradient of precipitation, as illustrated by the decreasing amount of precipitation received by the meteorological stations as they are located higher in elevation and inner in the mountain range (Figs. 2 and 1). Consequently, Mera Glacier with its more southern frontal position is submitted to a wetter climate than Pokalde Glacier.

---

## Mass balances of Mera and Pokalde glaciers (Nepal) since 2007

P. Wagnon et al.

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## 4 Methodology

### 4.1 Mass balance

Since 2007 and 2009, annual mass balance measurements have been carried out on Mera and Pokalde glaciers respectively. Measurements are performed in November or beginning of December each year. Extra stake readings have been performed in April 2009 and April 2013 on Mera Glacier, and every 2 to 5 months on Pokalde Glacier. The direct glaciological method is used for these measurements (Cuffey and Paterson, 2010). In the ablation area, annual mass balance is determined from bamboo stakes inserted up to 10 m deep in the ice. In the mass balance calculations, ice density is assumed to be  $900 \text{ kg m}^{-3}$ , and in the presence of snow, its density is measured in the field. The number of ablation stakes has progressively increased on Mera Glacier from 28 in 2007 to 44 stakes in 2012, and all stakes are located between 5000 and 5550 m a.s.l. Five stakes have been inserted on Pokalde Glacier in 2009 between 5500 and 5600 m a.s.l. (Fig. 1). In the accumulation area above 5600 m a.s.l., 6 cores (5 on Mera Glacier, 1 on Pokalde Glacier) are drilled to measure the annual net accumulation from snow layering (stratigraphy) and density measurements. Recco avalanche reflectors and blue-coloured chalk powder spread out over a  $2 \text{ m}^2$  surface mark an annual horizon. This horizon is then located easily the following year with a Recco detector, and the previous annual surface is identified while drilling. During years of high accumulation, ablation stakes at 5500 m a.s.l. also serve as accumulation stakes.

The annual glacier-wide mass balance  $B_a$  is calculated according to:

$$B_a = \frac{1}{S} \int_S b_a ds \quad (\text{in m w.e.}) \quad (1)$$

where  $b_a$  is the point surface mass balance obtained from the corresponding stake readings or net accumulation measurements and  $S$  is the glacier area. The hypsography of every glacier is derived from digital elevation models (DEMs) derived from

TCD

7, 3337–3378, 2013

## Mass balances of Mera and Pokalde glaciers (Nepal) since 2007

P. Wagnon et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Pleiades stereo-images of 25 November 2012 for Mera Glacier and from SPOT5-HRS stereo-images of 4 January 2011 for Pokalde Glacier (see Sect. 4.4). The glaciers were delineated by visual inspection of the images with the aid of field photographs. Using a single map over the whole measuring period to derive  $B_a$  means that we obtain here the reference-surface glacier-wide balance (Harrison et al., 2005).

On the small Pokalde Glacier, mass balance was obtained for every 10 m altitudinal range using a single linear fit of all available in-situ  $b_a$  measurements (ablation and accumulation). Every 10 m altitudinal range area was then multiplied by its corresponding mass balance, summed over the entire glacier and finally divided by the glacier area  $S$  to get the glacier-wide mass balance (Eq. 1). On the larger Mera Glacier,  $B_a$  was calculated in a similar way but using distinct linear interpolations for the ablation and accumulation zones. In the ablation zone, Mera and Naulek branches were also considered separately because they have fairly distinct mass balance gradients. Figure 3 displays for every surveyed year the regression lines that have been used to derive the glacier-wide mass balance. Each year, the equilibrium-line altitude (ELA) is deduced as the altitude at which the regression line crosses the zero mass balance value.

Accuracy of this overall specific annual mass balance depends on various factors such as the accuracy of stake readings, the accuracy of the density measurements, the density and representativeness of stake or accumulation measurement sites, the accuracy of the map and the method used to assess the mass balance for every altitudinal range. Thibert et al. (2008) conducted a variance analysis applied to all types of errors and obtained an overall uncertainty of  $\pm 0.40$  m w.e. for the glacier-wide mass balance. Although conducted on a glacier in the Alps, the analysis of Thibert et al. (2008) can be generalized to other glaciers because it is based on measurement errors which are similar on every glacier when using the glaciological method. Despite the relatively large error bars, year-to-year relative differences are instructive for climatic purposes because annual values refer to the same map, to the same area-elevation distribution function and to the same measurement network.

## Mass balances of Mera and Pokalde glaciers (Nepal) since 2007

P. Wagnon et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## 4.2 Surface velocity

Annual surface ice velocities have been measured every year in November on Mera Glacier by determining the displacement of 47 ablation and accumulation stakes using a Differential Global Positioning System (DGPS). These measurements have been performed relative to a fixed reference point outside the glacier on firm rocks (Fig. 1). The accuracy in  $x$  (easting),  $y$  (northing) and  $z$  (elevation) of each stake position is estimated at  $\pm 0.2$  m depending mainly on the size of the hole in which the stake has been set-up. Thus the surface ice velocities measured from stake displacements have an accuracy of  $\pm 0.3 \text{ myr}^{-1}$ .

## 4.3 Glacier thickness

Ground penetrating radar measurements have been conducted in November 2009 and November 2012 on Mera Glacier in order to determine ice thickness on 2 transverse cross sections at 5350 and 5520 m a.s.l. as well as close to the top of the glacier at 6350 m a.s.l. The radar system is made up of a transmitter and a receiver, each of which is outfitted with a similar kind of antenna (Hubbard and Glasser, 2005, p. 148). In our study we used the pulse radar transmitter (Icefield Instruments, Canada) based on the Narod transmitter (Narod and Clarke, 1994). The radar survey was carried out at a frequency of 4.2 MHz with antenna length of 10 m. Transmitter and receiver were put in snow sledges, separated by a fixed distance of 20 m and moved together along the profiles with a measurement every 10 m. The positions of the receiver and the transmitter are known through DGPS measurements, with an accuracy of  $\pm 0.1$  m. The speed of electromagnetic wave propagation in ice has been assumed to be  $167 \text{ m}\mu\text{s}^{-1}$  (Hubbard and Glasser, 2005). The field measurements were performed in such a way to obtain reflections from the glacier bed located more or less in the vertical plane with the measurement points at the glacier surface, allowing the determination of the glacier bed in two dimensions. The surface of the bedrock was constructed as an envelope of

# Mass balances of Mera and Pokalde glaciers (Nepal) since 2007

P. Wagnon et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



all ellipse functions, which give all the possible reflection positions between sending and receiving antennae.

#### 4.4 Digital elevation models

A 4 m DEM of Mera Glacier was derived from a stereo-pair of high resolution (0.7 m) Pleiades-1A images acquired 25 November 2012. Pleiades-1A is the first member of a new family of high resolution satellite from the French Space Agency (CNES) and was launched on 17 December 2011 (<http://smc.cnes.fr/PLEIADES/>). To our knowledge, we present here the first DEM derived from Pleiades-1A imagery over a glaciated region. Given the lack of accurate field ground control points (GCPs), GCPs were generated from SPOT5 images and a SPOT5 DEM that were first corrected using DGPS measurements performed in the Khumbu valley.

Our reference dataset is a 40 m DEM and a co-registered 5 m ortho-image computed without GCPs by the French Mapping Institute (IGN, Institut Géographique National) from SPOT5-HRS images acquired 4 January 2011 (Korona et al., 2009; Gardelle et al., 2013). This regional (120 km by 120 km) but relatively coarse dataset is used as reference to ortho-rectify a 2.5 m SPOT5-HRG image (60 km by 60 km) acquired 29 October 2009. A sub-pixel co-registration was achieved using the Cosis-Corr software (Leprince et al., 2007). Approximately 3600 kinematic DGPS measurements, acquired along the trails of the Everest base camp (Khumbu Valley) in April 2011, were overlaid on the SPOT5-HRG image and the image was shifted westward by 10 m so that the trails, clearly visible on the 2.5 m image, matched the location of the DGPS tracks. The 10 m westward shift was also applied to the SPOT5-HRS 40 m DEM. After this horizontal co-registration, the vertical difference between the SPOT5-HRS DEM and the altitude of the kinematic DGPS points was computed. As the difference was very close to zero (mean = 0.7 m, median = 0.1 m; standard deviation = 7.6 m;  $N = 3590$ ), the SPOT5-HRS DEM was not shifted vertically.

The 2.5 m SPOT5-HRG ortho-image and the 40 m SPOT5-HRS DEM, now both co-registered to the DGPS data, were used to identify 25 GCPs for Pleiades raw images.

### Mass balances of Mera and Pokalde glaciers (Nepal) since 2007

P. Wagnon et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





This set of GCPs was then ingested in the ©PCI Geomatica software (version 2013) to compute the DEM. The accuracy of the final 4 m DEM was checked against DGPS measurements performed on Mera Glacier between 20 and 27 November 2012, within a few days of the acquisition of the Pleiades-1A stereo pair. Horizontal and vertical precision of the DGPS measurements on Mera Glacier are estimated to 0.05 m. For a total of 445 points, the mean and median of the elevation difference between the Pleiades DEM and the DGPS are identical, at  $-0.93$  m, and the standard deviation is 1.02 m. The 0.93 m mean vertical shift was added to all altitudes in the Pleiades DEM.

## 5 Results

### 5.1 Annual and cumulative glacier-wide mass balance

The annual glacier-wide mass balances  $B_a$  for Mera and Pokalde glaciers are reported in Table 2, for the periods 2007–2012 and 2009–2012 respectively. Figure 4 displays the annual and cumulative mass balances for these glaciers. On Mera Glacier, the inter-annual variability is large (standard deviation of  $\pm 0.53$  m.w.e.  $\text{yr}^{-1}$ ) with 2 positive (2007–2008 and 2010–2011), 2 negative (2008–2009 and 2011–2012) and a balanced year (2008–2009). Mera Glacier experienced balanced or slightly negative mass balance conditions between 2007 and 2012 (mean  $B_a = -0.10 \pm 0.40$  m.w.e.  $\text{yr}^{-1}$ ) while the mass loss for Pokalde Glacier was more pronounced (mean  $B_a = -0.72 \pm 0.40$  m.w.e.  $\text{yr}^{-1}$  compared to  $-0.26 \pm 0.40$  m.w.e.  $\text{yr}^{-1}$  for Mera Glacier between 2009 and 2012). Over the period 2009–2012, the cumulative mass balance difference between both glaciers is 1.36 m.w.e., with Pokalde Glacier losing mass more rapidly than Mera Glacier. On Mera Glacier the ELA varied from a minimum of 5335 m.a.s.l. in 2010–2011 ( $B_a = 0.46 \pm 0.40$  m.w.e.; AAR = 0.89) to a maximum as high as 6055 m.a.s.l. in 2011–2012 ( $B_a = -0.77 \pm 0.40$  m.w.e.; AAR = 0.10).

## Mass balances of Mera and Pokalde glaciers (Nepal) since 2007

P. Wagnon et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 5.2 Seasonal glacier-wide mass balance

On 23–25 April 2009, stake readings were performed on Mera Glacier from the snout to 5800 m a.s.l. to assess both winter and summer mass balances for the year 2008–2009 (Figs. 4 and 5a). In 2008–2009, the annual mass balance ( $-0.10 \pm 0.40$  m w.e.) reveals that the glacier was close to steady state and some features of winter and summer mass balances can be pointed out. As expected, this Nepalese glacier is a summer-accumulation type glacier (Ageta and Fujita, 1996). In summer 2009, it experienced ablation below 5500 m a.s.l. and significant accumulation above resulting in positive summer mass balance (Fig. 4). Unexpectedly, ablation is non-negligible in winter between November 2008 and April 2009 at every elevation, which leads to negative winter mass balance (Fig. 4). Figure 5a gives the point mass balance as a function of elevation for the year 2008–2009, and also for the winter (6 November 2008–24 April 2009) and the summer (24 April 2009–5 November 2009) seasons. In winter, the glacier lost mass at every elevation, even in its highest parts, due to the combined effect of wind erosion and sublimation. The 2008–2009 glacier-wide winter mass balance is  $-0.20$  m w.e. In summer, the point mass balance as a function of elevation is very similar to the annual mass balance gradient, which means that the annual glacier-wide mass balance is almost exclusively controlled by the summer climatic conditions (Fig. 5a).

Additional stake readings were also performed on 16–21 April 2013, showing that the glacier-wide 2012–2013 winter mass balance is not significantly different from zero (Fig. 4). Indeed point winter mass balance is very close to zero at all elevations (Fig. 5b) confirming that summer is the key season for the annual glacier mass balance. Snow occasionally deposited in winter is systematically blown away after some days, due to strong westerly winds (jet stream) that occur above 5000 m a.s.l. in this region (Wang, 2006). At 6330 m a.s.l., since November 2008, we were never able to locate the blue horizon marker systematically spread over a  $2\text{ m}^2$  area a few tenths of centimetres below the surface, and covered by dense snow in the previous year. However, we were able to recover the Recco reflector tied to a bamboo stake inserted deep enough to

TCD

7, 3337–3378, 2013

### Mass balances of Mera and Pokalde glaciers (Nepal) since 2007

P. Wagnon et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Mass balances of Mera and Pokalde glaciers (Nepal) since 2007

P. Wagnon et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

branches comes from their orientation. Indeed, Naulek has a north-eastern aspect, while Mera has an overall northerly aspect, with some elevations (5400–5500 m a.s.l.) facing north-east and the lowest elevations (below 5350 m a.s.l.) facing west. This variable aspect along Mera branch probably explains the larger scatter of point mass balance data around the regression line on Mera branch than on Naulek branch. In some cases, stakes on the north-east orientated part of Mera branch experience a net mass balance similar to that observed on Naulek branch (see in particular one point mass balance measurement of Mera above the Naulek regression line, in 2008–2009 and 2009–2010; highlighted with a red circle in Fig. 3). The vertical mass balance gradient difference observed between both branches also partly explains why the terminus of Naulek (5260 m a.s.l.) is located at higher elevation than the terminus of Mera (4940 m a.s.l.).

Pokalde Glacier has a north-west aspect but its mass-balance gradient ( $1.51 \text{ m.w.e. (100 m)}^{-1}$ , Table 2) is double to triple that observed on both branches of Mera Glacier. This difference can be explained by the fact that (i) the glacier is much smaller than Mera Glacier, and thus likely experiences enhanced long-wave radiation and heat advection from the surrounding terrain that contribute to enhanced melting (e.g. Olyphant, 1986; Wagnon et al., 2009); (ii) it is located inside the mountain range and thus submitted to a much drier climate than Mera Glacier and (iii) that its vertical mass balance gradient is likely to be more uncertain than that on Mera Glacier due to its narrow elevation range (only 130 m altitude difference between the lowest and the highest stakes). The aridity and size difference between glaciers are also responsible for the altitudinal mass-balance difference observed on stakes, the ablation measured at 5550 m a.s.l. on Pokalde Glacier being equivalent to the ablation 200 m lower on Mera Glacier.

### 5.4 Mean annual surface velocities

Mean annual surface velocity measurements are located on the map of Mera Glacier (Fig. 7). Over the period 2007–2012, no velocity change was detectable, in agreement with balanced conditions over this relatively short time period. Estimated surface



and longitudinal cross sections) carried out at 6350 ma.s.l. show an ice thickness of about 55 m. The accuracy of the calculated ice thickness is determined, in part, by the accuracy of the measurement of the time delays and the antenna spacing. Thanks to clear reflections, interface geometry was generally easy to determine on all profiles.

5 Additional errors may arise because the smooth envelope of the reflection ellipses is only a minimal profile for a narrow, deep valley-shape bed topography, with the result that the ellipse equation will be governed by an arrival from a reflector located toward the side and thus not directly beneath the point of observation. Further errors may be introduced by assuming that all reflection points lie in the plane of the profile rather than an ellipsoid. The overall uncertainty on ice thickness is estimated at  $\pm 15$  m. Given that the uncertainty on ice surface coordinates is low ( $\pm 0.1$  m), the uncertainty on cross section areas mainly arises from the uncertainty on ice thickness and corresponds to  $\pm 6400$  m<sup>2</sup> at 5350 ma.s.l. and  $\pm 9200$  m<sup>2</sup> at 5520 ma.s.l.

## 5.6 Ice fluxes

15 The ice flux  $Q$  (m<sup>3</sup> of ice per year) through the cross sections at 5350 ma.s.l. and 5520 ma.s.l. can be inferred using two distinct methods. First they were calculated using the kinematic method consisting in multiplying the cross sectional area  $S_c$  (m<sup>2</sup>) with the depth-averaged horizontal ice velocity  $U$  (m yr<sup>-1</sup>).

$$Q = US_c \quad (2)$$

20 The depth-averaged horizontal ice velocity was derived from the mean surface ice velocity obtained by averaging the surface velocities available along each cross section (Sect. 5.4 and Fig. 7). Nye (1965) gives ratios of depth-averaged horizontal ice velocity to mean surface ice velocity varying from 0.8 (no sliding) to 1 (maximum sliding). The error range of the ice fluxes obtained with this method combines the error range for the cross section areas and the above-mentioned ratio assumed to vary from 0.8 to 1.

25 We also calculated ice fluxes using annual surface mass balances measured during 2007–2012. Although the dynamic changes are neglected here, this method allows us

# Mass balances of Mera and Pokalde glaciers (Nepal) since 2007

P. Wagnon et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



to estimate the ice fluxes for each section from mass-balance data according to the following equation:

$$Q = \frac{1}{0.9} \sum_z^{z_{\max}} b_i s_i \quad (3)$$

Where  $Q$  is the ice flux (converted into  $\text{m}^3$  of ice per year using an ice density of  $900 \text{ kg m}^{-3}$ , hence the factor  $1/0.9$ ) at a given elevation,  $z$ , and  $b_i$  is the annual mass balance of the altitudinal range  $i$  of map area  $s_i$ . The altitudinal ranges taken into account in the calculation are the areas contributing ice to the cross section and thus located between  $z$  and the altitude of the glacier top  $z_{\max}$ . In case of Mera Glacier, the upper areas only feeding the Mera branch and contributing ice to the two sections have been taken into account. The areas contributing to the two short and steep tongues separating from Mera branch on its left bank at 5600 and 5500 m.a.s.l. (Fig. 7) have been successively discarded for CS\_5520 and CS\_5350. We assume here that on each point of the glacier above this altitude,  $z$ , the surface elevation has remained unchanged from one year to the next. The uncertainties on ice fluxes resulting from surface mass balance are directly derived from the mass-balance uncertainties (see Sect. 4.1) applied to areas contributing ice to each cross section and also from the uncertainties in delineating accurately these areas.

The ice fluxes calculated with both methods at each cross section are given in Table 3. Both methods have been described in details in Azam et al. (2012) and a comparison between methods allowed them to assess the state of balance of Chhota Shigri Glacier (western Himalaya) over the one or two decades prior to their measurements. In case of Mera Glacier, the mean values of ice fluxes at CS\_5350 and CS\_5520 inferred from 5 yr of annual mass-balance data are higher than the respective values obtained with the kinematic method ( $0.51 \times 10^6$  versus  $0.18 \times 10^6 \text{ m}^3 \text{ ice yr}^{-1}$  at 5350 m.a.s.l. and  $0.70 \times 10^6$  versus  $0.38 \times 10^6 \text{ m}^3 \text{ ice yr}^{-1}$  at 5520 m.a.s.l.). The flux difference between both sections ( $0.2 \times 10^6 \text{ m}^3 \text{ ice yr}^{-1}$ ) is the same regardless of the method used here which gives confidence in the results. Moreover, the fact that fluxes obtained with the

Mass balances of  
Mera and Pokalde  
glaciers (Nepal) since  
2007

P. Wagnon et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion







(e.g. Immerzeel et al., 2012). Addressing this issue in Nepal and more generally in the monsoon-influenced Himalaya will not be achieved without implementing snow transport models (e.g. Vionnet et al., 2013) to better understand the redistribution and the sublimation of the snow by the wind at glacier elevations.

## 6.2 Mass-balance gradients

The mass balance gradients reported on Mera Glacier (from 0.48 to 0.85 m.w.e.  $(100\text{ m})^{-1}$  depending on the orientation) are roughly similar to the gradients observed in the Alps or on other mid-latitude glaciers (e.g. Rabatel et al., 2005; Zemp et al., 2009; Shea et al., 2013) or in western Himalaya on Chhota Shigri Glacier (Wagnon et al., 2007). Considered together with the value measured on Pokalde Glacier (1.51 m.w.e.  $(100\text{ m})^{-1}$ ), these gradients from both glaciers of our study are comparable to values reported previously on Yala Glacier (0.81–1.3 m.w.e.  $(100\text{ m})^{-1}$ , 1996) (Fujita et al., 1998); AX010 Glacier (0.81–0.9 m w.e.  $(100\text{ m})^{-1}$ , 1996–1999) (Fujita et al., 2001a) and Rikha Samba Glacier (0.64 m w.e.  $(100\text{ m})^{-1}$ , 1999) (Fujita et al., 2001b). Consequently, the mass-balance gradients of Nepalese glaciers can vary at least threefold, from 0.5 m.w.e.  $(100\text{ m})^{-1}$  to 1.5 m.w.e.  $(100\text{ m})^{-1}$  depending on the location (which drives the local climate and especially the amount of precipitation received by the glacier), the size and the aspect of the glacier.

In the Himalayan region, snow and glacier melt models based on ablation gradient usually use a single value for mass balance gradient (e.g. Racoviteanu et al., 2013) and they could be significantly improved by applying various gradients according to aspect, size, and climatic setting of the glaciers. Mass balance gradients over debris-covered glaciers also need to be investigated, and process-based studies on Changri Nup Glacier, located a few km north-west of Pokalde Glacier, were initiated in 2010 (Lejeune et al., 2013).

### 6.3 Recent mass-balance conditions

Over the period 2007–2012, the glacier-wide mass balance of Mera Glacier was slightly negative at  $-0.10 \pm 0.40$  m.w.e. yr<sup>-1</sup>. Pokalde Glacier lost mass more rapidly than Mera Glacier with a mean glacier-wide mass balance of  $-0.72 \pm 0.40$  m.w.e. yr<sup>-1</sup> compared to  $-0.26 \pm 0.40$  m.w.e. yr<sup>-1</sup> for Mera Glacier over the common 2009–2012 period (Table 2). The very negative mass balance observed on Pokalde Glacier is probably due to its lower accumulation area. Indeed, glaciers located at lower altitudes and usually small tend to have higher melt rates (e.g. Fujita and Nuimura, 2011). Such glaciers are often smaller and easier to access and therefore usually preferred for in-situ observations. Consequently, compilations of ground-based mass balances are probably biased toward negative values compared with the regional mean under the present-day climate (Fujita and Nuimura, 2011; Gardner et al., 2013). Our comparison between Mera and Pokalde glaciers seems to confirm this fact. Low-elevation glaciers like Pokalde Glacier are sometimes reduced to a single ablation zone, with the ELA above their upper reaches like in 2011–2012 (Table 2). These glaciers are no longer in equilibrium with the present climate and would disappear in the near future if climatic conditions similar to 2011–2012 prevail.

Even though the periods do not properly match, these values for Mera Glacier are not statistically different from the values recently reported by Gardelle et al. (2013) between 1999 and 2010 over 1461 km<sup>2</sup> of glacierized area in Everest region ( $-0.26 \pm 0.09$  m.w.e. yr<sup>-1</sup>). Moreover, the ice fluxes obtained by the kinematic method allow us to infer the state of balance of Mera Glacier over the last one or two decades (Azam et al., 2012). As seen in Sect. 5.6, the dynamic behaviour of the glacier is representative for mean decadal mass-balance conditions ranging between  $-0.48$  and  $-0.10$  m.w.e. yr<sup>-1</sup> in agreement again with Gardelle et al. (2013). If conditions observed between 2007 and 2012 prevail in the future, the glacier might accelerate in the coming years as suggested by the fact that cross section ice fluxes inferred from the mean 2007–2012 mass balance data are significantly higher than kinematic ice fluxes. In

the map showing the glacier-elevation changes of this region (Gardelle et al., 2013; Fig. A2), it is noticeable that Mera Glacier, probably due to its more southerly and wetter position, shrinks still less rapidly than glaciers located further north in the mountain range. This striking feature is confirmed by our observations of more negative mass balances for Pokalde Glacier than for Mera Glacier.

Between 2009 and 2012, both glaciers show a similar cumulative centred mass balance (= annual value – mean value over the three years for every glacier) revealing that inter-annual mass balance fluctuations are very similar between both glaciers, regardless of their size and their position inside the mountain range (Fig. 4 and Table 2). This striking consistency, which has been previously observed in other mountain ranges over longer time periods (e.g. Vincent et al., 2005; Huss et al., 2010), indicates that the mass balances of both study glaciers respond to a common regional climatic signal.

#### 6.4 Unclear link between meteorological and mass balance variability

Which climatic signal do these glaciers record? To evaluate the climatic influences on glacier mass balance, we compared annual or seasonal air temperature and precipitation records from Pyramid Observatory (5035 m a.s.l.; Fig. 1) with mass balance data (Table 2). First we had to reconstruct a complete series of monthly temperature and precipitation at Pyramid between November 2007 and October 2012 using data recorded in Khumbu valley. Data gaps in the Pyramid meteorological records were filled by applying a regression equation with the corresponding variable recorded at Pheriche (4260 m a.s.l.) when available or otherwise at Namche Bazar (3570 m a.s.l.). At monthly time-step and over the whole 2003–2012 period, the Pyramid record is well-correlated with the Pheriche and Namche Bazar records for temperature ( $r^2 = 0.95$  and  $0.83$  respectively) and for precipitation ( $r^2 = 0.81$  and  $0.86$  respectively). We were not able to reconstruct precipitation in 2008–2009 because there were too many gaps at all stations. In total only 13 % and 8 % of data have been reconstructed for temperature and precipitation respectively, suggesting that this reconstruction has a limited impact on the analysis.

### Mass balances of Mera and Pokalde glaciers (Nepal) since 2007

P. Wagnon et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



---

**Mass balances of  
Mera and Pokalde  
glaciers (Nepal) since  
2007**P. Wagnon et al.

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

We do not find any clear relationships between meteorological variables and mass balance data, at either seasonal or annual scales (Table 2). At most, it seems that negative mass balances are related to a deficit in precipitation during the monsoon, but this result needs to be confirmed with longer time-series. We attempted to reconstruct these data series at hourly time-step in order to discriminate snow and rain at Pyramid and to extrapolate this to glacier elevations using a temperature lapse rate derived from monsoon data at Pyramid and Pheriche ( $-5.75^{\circ}\text{C km}^{-1}$ ). This analysis shows that some years (2010–2011 and 2011–2012), up to 28 % of the total monsoon precipitation may fall as rain at 5500 m a.s.l. with probably significant impacts on the surface energy balance of glaciers and consequently on mass balance in the ablation area. But again, our reconstructed climatic dataset at hourly time-step was not accurate or long enough to clearly understand the drivers of the large mass-balance inter-annual variability between 2007 and 2012. Since the annual glacier-wide mass balance is mainly controlled by summer ablation and accumulation conditions, this inter-annual variability depends mainly on the Indian monsoon variability. It is likely that moderately negative mass balances of central Himalayan glaciers over the last decade are due to the weakening of the Indian monsoon (Bollasina et al., 2011), leading to higher atmospheric temperatures and lower precipitation (Yao et al., 2012). A thorough climatic analysis including data from the automatic weather station located at 5360 m a.s.l. on Mera Glacier and run since 2011 (Fig. 1) and a comparison with meteorological records available from nearby stations is still required and will be conducted in the future.

In conclusion, our observations on Mera Glacier, combined with findings from recent remote-sensed studies in the Everest area (Bolch et al., 2011; Nuimura et al., 2012; Gardelle et al., 2013), confirm that glaciers in this region are losing mass. Though our observation period is relatively short, the rate of observed specific mass loss is three to four times lower than that observed for glaciers in the European Alps, whose mean mass balance over the last decade is around  $-1 \text{ m.w.e. yr}^{-1}$  (Huss, 2012). Long-term monitoring of Mera and Pokalde glaciers is required to study a poorly-known and seemingly complex relationship between glaciers and climate in eastern Nepal. This

program, planned for the coming years, will include winter and summer mass balance measurements, surveys of thickness changes and energy balance studies on Mera Glacier and on other debris-covered glaciers in the region.

*Acknowledgements.* This work has been supported by the French Service d'Observation GLACIOCLIM and the French National Research Agency through ANR-09-CEP-005-01/PAPRIKA. This study was carried out within the framework of the Ev-K2-CNR Project in collaboration with the Nepal Academy of Science and Technology as foreseen by the Memorandum of Understanding between Nepal and Italy, and thanks to contributions from the Italian National Research Council, the Italian Ministry of Education, University and Research and the Italian Ministry of Foreign Affairs. All people involved in the field or helping for logistical support at the Pyramid, in Kathmandu or in Italy are greatly acknowledged here. We thank J. P. Chazarin, R. Biron and the porters who have been involved in successive field trips, sometimes in harsh conditions. S. Kaspari measured the stakes on 23–25 April 2009, and is greatly acknowledged here. Pleiades-1A images were obtained free of charge in the framework of the “Recette Thématique Utilisateur Pléiades/ORFEO” conducted by the French Space Agency (CNES). The SPOT5 DEM and images were obtained thanks to ISIS proposals #397 and #520. E. Berthier acknowledges support from the CNES through the TOSCA program.



The publication of this article is financed by CNRS-INSU.

## References

- Ageta, Y. and Fujita, K.: Characteristics of mass balance of summer-accumulation type glaciers in the Himalayas and Tibetan Plateau, *Zeitschrift für Gletscherkunde und Glazialgeologie*, 32, 61–65, 1996.
- Ageta, Y. and Higuchi, K.: Estimation of mass balance components of a summer-accumulation type glacier in the Nepal Himalaya, *Geogr. Ann.*, 66, 249–255, 1984.

## Mass balances of Mera and Pokalde glaciers (Nepal) since 2007

P. Wagnon et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Mass balances of Mera and Pokalde glaciers (Nepal) since 2007

P. Wagnon et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Ageta, Y., Ohata, T., Tanaka, Y., Ikegami, K., and Higuchi, K.: Mass balance of Glacier AX010 in Shorong Himal, east Nepal during the summer monsoon season, *Seppyo, J. Jpn. Soc. Snow Ice*, 41, 34–41, 1980.
- Arendt, A., Bolch, T., Cogley, J. G., Gardner, A., Hagen, J.-O., Hock, R., Kaser, G., Pfeffer, W. T., Moholdt, G., Paul, F., Radič, V., Andreassen, L., Bajracharya, S., Beedle, M., Berthier, E., Bhambri, R., Bliss, A., Brown, I., Burgess, E., Burgess, D., Cawkwell, F., Chinn, T., Copland, L., Davies, B., de Angelis, H., Dolgova, E., Filbert, K., Forester, R., Fountain, A., Frey, H., Giffen, B., Glasser, N., Gurney, S., Hagg, W., Hall, D., Haritashya, U. K., Hartmann, G., Helm, C., Herreid, S., Howat, I., Kapustin, G., Khromova, T., Kienholz, C., Koenig, M., Kohler, J., Kriegel, D., Kutuzov, S., Lavrentiev, I., LeBris, R., Lund, J., Manley, W., Mayer, C., Miles, E., Li, X., Menounos, B., Mercer, A., Moelg, N., Mool, P., Nosenko, G., Negrete, A., Nuth, C., Pettersson, R., Racoviteanu, A., Ranzi, R., Rastner, P., Rau, F., Rich, J., Rott, H., Schneider, C., Seliverstov, Y., Sharp, M., Sigurðsson, O., Stokes, C., Wheate, R., Winsvold, S., Wolken, G., Wyatt, F., and Zheltyhina, N.: Randolph Glacier Inventory [v2.0]: A Dataset of Global Glacier Outlines, Global Land Ice Measurements from Space, Boulder Colorado, USA, Digital Media, available at: <http://www.glims.org/RGI/randolph.html>, 2012.
- Azam, M. F., Wagnon, P., Ramanathan, A., Vincent, C., Sharma, P., Arnaud, Y., Linda, A., Pottakkal, J., Chevallier, P., Singh, V. B., and Berthier, E.: From balance to imbalance: a shift in the dynamic behaviour of Chhota Shigri Glacier (Western Himalaya, India), *J. Glaciol.*, 58, 315–324, doi:10.3189/2012JoG11J123, 2012.
- Bajracharya, S. R. and Mool, P. K.: Glaciers, glacial lakes and glacial lake outburst floods in the Mount Everest region, Nepal, *Ann. Glaciol.*, 50, 81–86, 2009.
- Bolch, T., Kulkarni, A., Kääb, A., Huggel, C., Paul, F., Cogley, J. G., Frey, H., Kargel, J. S., Fujita, K., Scheel, M., Bajracharya, S., and Stoffel, M.: The State and Fate of Himalayan Glaciers, *Science*, 336, 310–314, 2012.
- Bolch, T., Pieczonka, T., and Benn, D. I.: Multi-decadal mass loss of glaciers in the Everest area (Nepal Himalaya) derived from stereo imagery, *The Cryosphere*, 5, 349–358, doi:10.5194/tc-5-349-2011, 2011.
- Bollasina, M., Ming, Y., and Ramaswamy, V.: Anthropogenic aerosols and the weakening of the South Asian summer monsoon, *Science*, 334, 502–505, doi:10.1126/science.1204994, 2011.
- Bookhagen, B. and Burbank, D. W.: Topography, relief, and TRMM-derived rainfall variations along the Himalaya, *Geophys. Res. Lett.*, 33, L08405, doi:10.1029/2006GL026037, 2006.



---

**Mass balances of  
Mera and Pokalde  
glaciers (Nepal) since  
2007**

---

P. Wagnon et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Bookhagen, B. and Burbank, D. W.: Toward a complete Himalayan hydrological budget: spatiotemporal distribution of snowmelt and rainfall and their impact on river discharge, *J. Geophys. Res.*, 115, F03019, doi:10.1029/2009JF001426, 2010.
- 5 Burbank, D. W., Blythe, A. E., Putkonen, J., Bratt-Sitaula, B., Gabet, E., Oskin, M., Barros, A., and Ojha, T. P.: Decoupling of erosion and precipitation in the Himalayas, *Nature*, 426, 652–655, doi:10.1038/nature02187, 2003.
- Cogley, J. G., Kargel, J. S., Kaser, G., and Van der Veen, C. J.: Tracking the source of glacier misinformation, *Science*, 327, p. 522, 2010.
- 10 Cuffey, K. M. and Paterson, W. S. B.: *The Physics of Glaciers*, 4th Edn., Academic Press Inc, Amsterdam, 2010.
- Dobhal, D. P., Gergan, J. T., and Thayyen, R. J.: Mass balance studies of the Dokriani Glacier from 1992 to 2000, Garhwal Himalaya, India, *Bull. Glaciol. Res.*, 25, 9–17, 2008.
- Fujita, K. and Nuimura, T.: Spatially heterogeneous wastage of Himalayan glaciers, *P. Natl. Acad. Sci. USA*, 108, 14011–14014, 2011.
- 15 Fujita, K., Takeuchi, Y., and Seko, K.: Glaciological observations of Yala Glacier in Langtang Valley, Nepal Himalaya, 1994 and 1996, *Bull. Glacier Res.*, 16, 75–81, 1998.
- Fujita, K., Kadota, T., Rana, B., Kayastha, R. B., and Ageta, Y.: Shrinkage of Glacier AX010 in Shorong region, Nepal Himalayas in the 1990s, *Bull. Glacier Res.*, 18, 51–54, 2001a.
- Fujita, K., Nakazawa, F., and Rana, B.: Glaciological observations on Rikha Samba Glacier in Hidden Valley, Nepal Himalayas, 1998 and 1999, *Bull. Glacier Res.*, 18, 31–35, 2001b.
- 20 Gardelle, J., Berthier, E., Arnaud, Y., and Käab, A.: Region-wide glacier mass balances over the Pamir-Karakoram-Himalaya during 1999–2011, *The Cryosphere Discuss.*, 7, 975–1028, doi:10.5194/tcd-7-975-2013, 2013.
- Gardner, A. S., Moholdt, G., Cogley, J. G., Wouters, B., Arendt, A. A., Wahr, J., Berthier, E., Pfeffer, T., Kaser, G., Hock, R., Ligtenberg, S. R. M., Bolch, T., Sharp, M., Hagen, J. O., van den Broeke, M. R., and Paul, F.: A consensus estimate of glacier contribution to sea level rise: 2003 to 2009, *Science*, 340, 852, doi:10.1126/science.1234532, 2013.
- 25 Harrison, W. D., Elsberg, D. H. Cox, L. H., and March, R. S.: Different mass balances for climatic and hydrologic applications, correspondence, *J. Glaciol.*, 51, p. 176, 2005.
- 30 Haeberli, W., Maisch, M., and Paul, F.: Mountain glaciers in global climate-related observational network, *WMO Bulletin*, 51, 18–25, 2002.
- Hubbard, B. and Glasser, N.: *Field Techniques in Glaciology and Glacial Geomorphology*, Wiley and Sons Ltd, Chichester, England, 400 pp., 2005.

## Mass balances of Mera and Pokalde glaciers (Nepal) since 2007

P. Wagnon et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Huss, M.: Extrapolating glacier mass balance to the mountain-range scale: the European Alps 1900–2100, *The Cryosphere*, 6, 713–727, doi:10.5194/tc-6-713-2012, 2012.
- Huss, M., Hock, R., Bauder, A., and Funk, M.: 100yr mass changes in the Swiss Alps linked to the Atlantic multidecadal oscillation, *Geophys. Res. Lett.*, 37, L10501, doi:10.1029/2010GL042616, 2010.
- Immerzeel, W. W., van Beek, L. P. H., and Bierkens, M. F. P.: Climate change will affect the Asian water towers, *Science*, 328, 1382–1385, 2010.
- Immerzeel, W. W., Pellicciotti, F., and Shrestha, A. B.: Glaciers as a proxy to quantify the spatial distribution of precipitation in the Hunza Basin, *Mt. Res. Dev.*, 32, 30–38, doi:10.1659/MRD-JOURNAL-D-11-00097.1, 2012.
- Kääb, A., Berthier, E., Nuth, C., Gardelle, J., and Arnaud, Y.: Contrasting patterns of early 21st century glacier mass change in the Himalaya, *Nature*, 488, 495–498, doi:10.1038/nature11324, 2012.
- Kadota, T., Seko, K., Aoki, T., Iwata, S., and Yamaguchi, S.: Shrinkage of Khumbu Glacier, east Nepal from 1978 to 1995, *IAHS Publ.*, 264, 235–243, 2000.
- Kaser, G., Großhauser, M., and Marzeion, B.: Contribution potential of glaciers to water availability in different climate regimes, *P. Natl. Acad. Sci. USA*, 107, 20223–20227, doi:10.1073/pnas.1008162107, 2010.
- Kehrwald, N. M., Thompson, L. G., Tandong, Y., Mosley-Thompson, E., Schotterer, U., Alfirov, V., Beer, J., Eikenberg, J., and Davis, M. E.: Mass loss on Himalayan glacier endangers water resources, *Geophys. Res.*, 35, L22503, doi:10.1029/2008GL035556, 2008.
- Korona, J., Berthier, E., Bernard, M., Remy, F., and Thouvenot, E.: SPIRI T. SPOT 5 stereoscopic survey of polar ice: reference images and topographies during the fourth International Polar Year (2007–2009), *ISPRS J. Photogramm.*, 64, 204–212, doi:10.1016/j.isprs.2008.10.005, 2009.
- Lejeune, Y., Bertrand, J. M., Wagnon, P., and Morin, S.: A physically based model for the year-round energy and mass balance of debris-covered glaciers, *J. Glaciol.*, 50, 327–344, 2013.
- Leprince, S., Barbot, S., Ayoub, F., and Avouac, J. P.: Automatic and precise orthorectification, coregistration, and subpixel correlation of satellite images, application to ground deformation measurements, *IEEE T. Geosci. Remote*, 45, 1529–1557, 2007.
- Narod, B. B. and Clarke, G. K. C.: Miniature high-power impulse transmitter for radio-echo sounding, *J. Glaciol.*, 40, 190–194, 1994.

## Mass balances of Mera and Pokalde glaciers (Nepal) since 2007

P. Wagnon et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Nepal, S., Krause, P., Flügel, W.-A., Fink, M., and Fisher, C.: Understanding the hydrological system dynamics of a glaciated alpine catchment in the Himalayan region using the J2000 hydrological model, *Hydrol. Process.*, doi:10.1002/hyp.9627, 2013.
- 5 Nuimura, T., Fujita, K., Fukui, K., Asahi, K., Aryal, R., and Ageta, Y.: Temporal changes in elevation of the debris-covered ablation area of Khumbu Glacier in the Nepal Himalaya since 1978, *Arct. Antarct. Alp. Res.*, 43, 246–255, 2011.
- Nuimura, T., Fujita, K., Yamaguchi, S., and Sharma, R. R.: Elevation changes of glaciers revealed by multitemporal digital elevation models calibrated by GPS survey in the Khumbu region, Nepal Himalaya, 1992–2008, *J. Glaciol.*, 58, 648–656, doi:10.3189/2012JoG11J061, 10 2012.
- Nye, J. F.: The flow of a glacier in a channel of rectangular, elliptic or parabolic cross section, *J. Glaciol.*, 5, 661–690, 1965.
- Olyphant, G. A.: Longwave radiation in mountainous areas and its influence on the energy balance of alpine snowfields, *Water Resour. Res.*, 22, 62–66, 1986.
- 15 Paul, F., Kääb, A., and Haeberli, W.: Recent glacier changes in the Alps observed by satellite: consequences for future monitoring strategies, *Global Planet. Change*, 56, 111–122, 2007.
- Pomeroy, J. W. and Essery, R. L. H.: Turbulent fluxes during blowing snow: field tests and model sublimation predictions, *Hydrol. Process.*, 13, 2963–2975, 1999.
- Rabatel, A., Dedieu, J. P., and Vincent, C.: Using remote-sensing data to determine equilibrium-line altitude and mass-balance time series: validation on three French glaciers, 1994– 20 2002, *J. Glaciol.*, 51, 539–546, 2005.
- Racoviteanu, A. E., Armstrong, R., and Williams, M. W.: Evaluation of an ice ablation model to estimate the contribution of melting glacier ice to annual discharge in the Nepalese Himalaya, *Water Resour. Res.*, in press, 2013.
- 25 Schaner, N., Voisin, N., Nijssen, B., and Lettenmaier, D. P.: The contribution of glacier melt to streamflow, *Environ. Res. Lett.*, 7, 034029, doi:10.1088/1748-9326/7/3/034029, 2012.
- Scherler, D., Bookhagen, B., and Strecker, M. R.: Spatially variable response of Himalayan glaciers to climate change affected by debris cover, *Nat. Geosci.*, 4, 156–159, 2011.
- 30 Shea, J. M., Menounos, B., Moore, R. D., and Tennant, C.: An approach to derive regional snow lines and glacier mass change from MODIS imagery, western North America, *The Cryosphere*, 7, 667–680, doi:10.5194/tc-7-667-2013, 2013.

**Mass balances of  
Mera and Pokalde  
glaciers (Nepal) since  
2007**

P. Wagnon et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Thibert, E., Blanc, R., Vincent, C., and Eckert, N.: Glaciological and volumetric mass-balance measurements: error analysis over 51 years for Glacier de Sarennes, French Alps, *J. Glaciol.*, 54, 522–532, 2008.

Vincent, C., Lemeur, E., Six, D., and Funk, M.: Solving the paradox of the end of the Little Ice Age in the Alps, *Geophys. Res. Lett.*, 32, L09706, doi:10.1029/2005GL022552, 2005.

Vionnet, V., Martin, E., Masson, V., Guyomarc'h, G., Naaim-Bouvet, F., Prokop, A., Durand, Y., and Lac, C.: Simulation of wind-induced snow transport in alpine terrain using a fully coupled snowpack/atmosphere model, *The Cryosphere Discuss.*, 7, 2191–2245, doi:10.5194/tcd-7-2191-2013, 2013.

Wagnon, P., Linda, A., Arnaud, Y., Kumar, R., Sharma, P., Vincent, C., Pottakkal, J. G., Berthier, E., Ramanathan, A., Hasnain, S. I., and Chevallier, P.: Four years of mass balance on Chhota Shigri Glacier, Himachal Pradesh, India, a new benchmark glacier in the western Himalaya, *J. Glaciol.*, 53, 603–611, 2007.

Wagnon, P., Lafaysse, M., Lejeune, Y., Maisincho, L., Rojas, M., and Chazarin, J. P.: Understanding and modeling the physical processes that govern the melting of snow cover in a tropical mountain environment in Ecuador, *J. Geophys. Res.*, 114, D19113, doi:10.1029/2009JD012292, 2009.

Wang, B.: *The Asian Monsoon*, Springer, Berlin, 787 pp., 2006.

Webster, P. J., Magana, V. O., Palmer, T. N., Shukla, J., Tomas, R. A., Yanai, M., and Yasunari, T.: Monsoons: processes, predictability, and the prospects for prediction, *J. Geophys. Res.*, 103, 14451–14510, doi:10.1029/97JC02719, 1998.

Yao, T., Thompson, L. G., Yang, W. Yu, W., Gao, Y., Guo, X., Yang, X., Duan, K., Zhao, H., Xu, B., Pu, Y., Lu, A., Xiang, Y., Kattel, D. B., and Joswiak, D.: Different glacier status with atmospheric circulations in Tibetan Plateau and surroundings, *Nature Climate Change*, 2, 663–667, doi:10.1038/NCLIMATE1580, 2012.

Zhang, T., Xiao, C., Colgan, W., Qin, X., Du, W., Sun, W., Liu, Y., and Ding, M.: Observed and modelled ice temperature and velocity along the main flowline of East Rongbuk Glacier, Qomolangma (Mount Everest), Himalaya, *J. Glaciol.*, 59, 438–448, doi:10.3189/2013JoG12J202, 2013.

Zemp, M., Hoerzle, M., and Haeberli, W.: Six decades of glacier mass-balance observations: a review of worldwide monitoring network, *Ann. Glaciol.*, 50, 101–111, 2009.

## Mass balances of Mera and Pokalde glaciers (Nepal) since 2007

P. Wagnon et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Table 1.** List of geographical and topographical characteristics of Mera Glacier.  $ELA_0$  and  $AAR_0$  are the calculated equilibrium-line altitude and accumulation-area ratio values respectively for a zero glacier-wide balance given the 2012 hypsography. Mean annual temperature at  $ELA_0$  is derived from the AWS located at 5360 m a.s.l. applying a standard lapse rate of  $-0.0065\text{ }^\circ\text{C m}^{-1}$ .

General features	
Country – Region	Nepal – Everest region
Mountain range	Central Himalaya
Drainage system	Dudh Koshi basin – Ganges river
Climate	Indian monsoon zone
Glacier characteristics	
Latitude/Longitude	27.7° N/86.9° E
Max./Min. elevation	6420/~ 4940 m a.s.l.
Total glacierized area	5.1 km <sup>2</sup>
Mean orientation	North
Measurement information	
Total number of ablation stakes	28 to 44 (between 5000 and 5550 m a.s.l.)
Total number of pits/drilling sites	5 (between 5700 and 6400 m a.s.l.)
Measurements frequency	2 yr <sup>-1</sup> : Apr and Nov since 2013
Mass balance information	
$ELA_0$	~ 5550 m a.s.l.
$AAR_0$	~ 0.59
Mean annual temperature at $ELA_0$	~ $-2.6\text{ }^\circ\text{C}$

## Mass balances of Mera and Pokalde glaciers (Nepal) since 2007

P. Wagnon et al.

**Table 2.** Annual and 5 yr mean glacier-wide mass balances  $B_a$ , ELA, AAR and mass balance gradients  $db/dz$  for Mera and Pokalde glaciers, as well as some meteorological variables recorded at Pyramid Observatory (5035 m.a.s.l.). On Mera Glacier, mass balance gradients are distinguished between Mera and Naulek branches (referred as Mera and Naulek subscripts). The standard deviation (STD) is also displayed for every variable. The error range for  $B_a$  is  $\pm 0.40$  m w.e. Also shown is the cumulative centred mass balance (cum. centred  $B_a$  = annual  $B_a$  – mean value of  $B_a$  over 2009–2012, the mean 2009–2012 value for Mera Glacier is  $-0.26$  m.w.e.yr $^{-1}$ ). When there are data gaps, meteorological variables at Pyramid have been reconstructed at monthly time-step over the 2003–2012 period, applying a regression equation with the corresponding variable recorded at Pheriche (4260 m.a.s.l.) when available or otherwise at Namche Bazar (3570 m.a.s.l.). Precipitation data are missing in 2008–2009.

	2007–08	2008–09	2009–10	2010–11	2011–12	Mean	STD
Mera Glacier							
$B_a$ (m w.e.)	0.39	-0.10	-0.48	0.46	-0.77	-0.10	0.53
ELA (m)	5425	5585	5680	5335	6055	5615	280
AAR	0.74	0.55	0.42	0.89	0.10	0.54	0.31
$db/dz_{\text{Mera}}$ (m w.e. (100 m) $^{-1}$ )	0.48	0.41	0.46	0.58	0.49	0.48	0.06
$db/dz_{\text{Naulek}}$ (m w.e. (100 m) $^{-1}$ )	0.97	0.74	0.97	–	0.72	0.85	0.14
Cum. centred $B_a$ (m w.e.)			-0.22	0.72	-0.51	–	–
Pokalde Glacier							
$B_a$ (m w.e.)			-0.92	-0.11	-1.12	-0.72	0.54
ELA (m)			5625	–	> glacier	–	–
AAR			0.13	–	0	–	–
$db/dz$ (m w.e. (100 m) $^{-1}$ )			1.65	–	1.37	1.51	0.20
Cum. centred $B_a$ (m w.e.)			-0.20	0.61	-0.40	–	–
Meteorological conditions at Pyramid (5035 m.a.s.l.)							
Annual temperature (°C)	-2.6	-1.7	-2.3	-2.9	-2.8	-2.5	0.5
Summer <sup>a</sup> temperature (°C)	2.4	2.7	0.7	2.4	3.1	2.3	0.9
Winter <sup>b</sup> temperature (°C)	-5.2	-4.0	-3.9	-5.5	-5.8	-4.9	0.9
Annual precipitation (mm)	391	–	343	447	374	389	44
Summer <sup>a</sup> precipitation (mm)	372	–	295	409	330	352	50
Winter <sup>b</sup> precipitation (mm)	19	–	48	38	44	37	13

<sup>a</sup> Summer = monsoon months (June to September)

<sup>b</sup> Winter = non-monsoon months (October to May)

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

## Mass balances of Mera and Pokalde glaciers (Nepal) since 2007

P. Wagnon et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version


Interactive Discussion



**Table 3.** Ice fluxes  $Q(10^6 \text{ m}^3 \text{ ice yr}^{-1})$ , inferred from the kinematic method and from the annual mass balance data at each cross section surveyed in the field i.e. at mean elevations of 5350 m a.s.l. and 5520 m a.s.l. respectively. For sake of comparison the glacier-wide annual mass balances for the 5 surveyed years are also given (in parenthesis).

Ice fluxes $Q(10^6 \text{ m}^3 \text{ ice yr}^{-1})$ Years (annual mass balance, m w.e.)	kinematic	inferred at each cross section from annual mass-balance data					Mean (-0.10)
		2007–08 (0.39)	2008–09 (-0.10)	2009–10 (-0.48)	2010–11 (0.46)	2011–12 (-0.77)	
Cross section at 5350 m a.s.l.	$0.18 \pm 0.06$	1.73	0.62	-0.26	1.68	-1.23	0.51
Cross section at 5520 m a.s.l.	$0.38 \pm 0.11$	1.75	0.88	0.20	1.53	-0.87	0.70





**Fig. 1.** Map of the high Dudh Koshi basin (in grey in the inset map of Nepal and with limits represented by the dashed black line in the main map) where Mera and Pokalde glaciers are located (inside the red squares). Meteorological stations and summits are indicated by dots and triangles respectively. Glacierized areas from the Randolph Glacier Inventory v2.0 (Arendt et al., 2012) are represented in blue. Maps with red contours are enlargements for Pokalde and Mera glaciers, showing the ablation stake network (grey circles), the accumulation measurement sites (blue squares) and the Mera automatic weather station (AWS – black star). The backgrounds are a SPOT5 image of 4 January 2011 and a Pleiades-1A image of 25 November 2012 for Pokalde and Mera glaciers respectively (CNES 2011–2012/Distribution Astrium). From the Mera central summit (red triangle), Mera glacier flows toward the north and splits into two main branches, referred as Mera and Naulek branches respectively.

## Mass balances of Mera and Pokalde glaciers (Nepal) since 2007

P. Wagnon et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Mass balances of Mera and Pokalde glaciers (Nepal) since 2007

P. Wagnon et al.




Fig. 2. (Caption on next page.)

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

⏴

⏵

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Mass balances of Mera and Pokalde glaciers (Nepal) since 2007

P. Wagnon et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Fig. 2.** Mean annual cycle of monthly precipitation ( $P$ , histograms) and monthly air temperature ( $T$ , red line) during 2003–2012 at four meteorological stations operated by Ev-K2-CNR: Lukla (2660 m.a.s.l.), Namche Bazar (3570 m.a.s.l.), Pheriche (4260 m.a.s.l.) and Pyramid (5035 m.a.s.l.).  $T$  and  $P$  are the mean annual temperature and the cumulative annual precipitation respectively. The data gaps for Lukla station are April to September 2004, July, September and October 2006, June to September 2008, March to September 2010, August, September and December 2011, January to March 2012 (only for precipitation) April, May and October to December 2012. The data gaps for Namche Bazar station are April to June 2004, September and October 2006, September and October 2007, June to October 2009, March to September 2010, April, May and October to December 2012. The data gaps for Pheriche station are April to June 2004, June to October 2008 (only for precipitation), January to April and June 2009, April and May 2011, November 2011 to June 2012. The data gaps for Pyramid station are April to June 2004, May to September 2006 (only for precipitation), November 2008, May to October 2009 (only for precipitation), June and July 2010 (only for precipitation) October to December 2010, January and June 2011, November and December 2011 (only for temperature), January and October to December 2012.

## Mass balances of Mera and Pokalde glaciers (Nepal) since 2007

P. Wagnon et al.




Fig. 3. (Caption on next page.)

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

⏴ ⏵

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Mass balances of Mera and Pokalde glaciers (Nepal) since 2007

P. Wagnon et al.

**Fig. 3.** 5 first panels: Five years of annual point mass balance (dots) as a function of altitude derived from field measurements (stakes, drillings or pits) on Mera Glacier. Measurements were performed on 17 November 2007, 6 November 2008, 5 November 2009, 11 November 2010, 10 November 2011 and 23 November 2012  $\pm 4$  days. Dark blue is used for the Mera branch, light blue for Naulek. The measurement on Mera in 2008–2009 and 2009–2010 highlighted with a red circle refer to the same stake facing North–East. Also shown are the linear regression lines used to derive the annual glacier-wide mass balance  $B_a$ . In 2010–2011, there were not enough measurements on Naulek branch to derive a significant regression line, and therefore the mean vertical mass balance gradient obtained over the four other years on Naulek was used instead, i.e.  $0.85 \text{ m.w.e. } (100 \text{ m})^{-1}$ .  $B_a$  and ELA for every year are displayed on the corresponding panel. Bottom right panel: hypsometry of Mera Glacier showing the respective areas of Naulek (light blue) and Mera (dark blue) branches.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Mass balances of Mera and Pokalde glaciers (Nepal) since 2007


P. Wagnon et al.



**Fig. 4.** Annual (dark blue and green large histograms) and cumulative (line with big blue dots and line with big green triangles) mass balances of Mera Glacier and Pokalde Glacier respectively. Also shown are the winter and summer mass balances for 2008–2009 and 2012–2013 (narrow cyan histograms – only Mera Glacier is represented) and the corresponding cumulative mass balances (dashed line with small blue dots for Mera Glacier and dashed line with small green triangles for Pokalde Glacier).

## Mass balances of Mera and Pokalde glaciers (Nepal) since 2007

P. Wagnon et al.




**Fig. 5.** (a) Point mass balance as a function of elevation, for the entire 2008–2009 yr (red dots), for the winter season (blue dots) and for the summer season (green dots). Plain and open dots refer to Mera and Naulek branches, respectively. (b) Same as (a) but only for the winter season 2012–2013.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)



## Mass balances of Mera and Pokalde glaciers (Nepal) since 2007

P. Wagnon et al.



**Fig. 6.** Cumulative point mass balance recorded at the three lowest stakes (5505 – triangles, 5531 – squares and 5557 m.a.s.l. – dots) between November 2009 and February 2013 on Pokalde Glacier (lines). Grey areas highlight the main winter season (from November to April).

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[⏴](#)
[⏵](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)





Fig. 7. (Caption on next page.)

Mass balances of Mera and Pokalde glaciers (Nepal) since 2007

P. Wagnon et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

⏴ ⏵

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

# TCD

7, 3337–3378, 2013

## Mass balances of Mera and Pokalde glaciers (Nepal) since 2007

P. Wagnon et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Fig. 7.** Upper panel: Map of Mera Glacier showing mean annual surface velocities measured during the period 2007–2012 (green vectors). All DGPS measurements have been averaged, and the number of measuring years used to obtain annual velocities varies from 1 (1 single year of measurements) to 5 (the full period), depending on the survival time of the stakes. The blue line is the glacier outline, and the dark blue short segments are GPR survey cross sections that have been performed in November 2009 (at 5520 and 5350 m a.s.l.) and in November 2012 (at 6350 m a.s.l.). Background: 25 November 2012 Pleiades-1A image (CNES 2012/Distribution Astrium). Lower panels: Ice depths and surface topographies of the two main cross sections CS\_5350 and CS\_5520 at 5350 and 5520 m a.s.l., respectively.