

**Seasonal thaw
settlement at drained
thermokarst lake
basins, Arctic Alaska**

L. Liu et al.

Seasonal thaw settlement at drained thermokarst lake basins, Arctic Alaska

L. Liu¹, K. Schaefer², A. Gusmeroli³, G. Grosse⁴, B. M. Jones⁵, T. Zhang^{2,6},
A. D. Parsekian^{1,*}, and H. A. Zebker¹

¹Department of Geophysics, Stanford University, California, USA

²National Snow and Ice Data Center, Cooperative Institute for Research in Environmental Sciences, University of Colorado at Boulder, Colorado, USA

³International Arctic Research Center, University of Alaska Fairbanks, Alaska, USA

⁴Geophysical Institute, University of Alaska Fairbanks, Alaska, USA

⁵Alaska Science Center, US Geological Survey, Anchorage, Alaska, USA

⁶Ministry of Education Key Laboratory of West China's Environmental System, Lanzhou University, China

*currently at: Department of Geology and Geophysics, University of Wyoming, Laramie Wyoming, USA

Received: 19 November 2013 – Accepted: 25 November 2013 – Published: 6 December 2013

Correspondence to: L. Liu (liulin@stanford.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

Drained thermokarst lake basins (DTLBs) are ubiquitous landforms on arctic tundra lowlands, but their present-day dynamic states are seldom investigated. Here we report results based on high-resolution Interferometric Synthetic Aperture Radar (InSAR) measurements using space-borne data for a study area located near Prudhoe Bay, Alaska where we focus on the seasonal thaw settlement within DTLBs, averaged between 2006 and 2010. The majority (14) of the 18 DTLBs in the study area analyzed exhibited seasonal thaw settlement of 3–4 cm. However, four of the DTLBs analyzed exceeded 4 cm of thaw settlement, with one basin experiencing up to 12 cm. Combining the InSAR observations with the in situ active layer thickness measured using ground penetrating radar and mechanical probing, we calculated thaw strain, an index of thaw settlement strength along a transect across the basin that underwent large thaw settlement. We found thaw strains of 10–35 % at the basin center, suggesting the seasonal melting of ground ice as a possible mechanism for the large settlement. These findings emphasize the dynamic nature of permafrost landforms, demonstrate the capability of the InSAR technique to remotely monitor surface deformation of individual DTLBs, and illustrate the combination of ground-based and remote sensing observations to estimate thaw strain. Our study highlights the need for better description of the spatial heterogeneity of landscape-scale processes for regional assessment of surface dynamics on arctic coastal lowlands.

1 Introduction

Thermokarst lakes (or thaw lakes) are ubiquitous and dynamic landscape features on arctic tundra lowlands (Sellman et al., 1975; Mackay, 1988; Frohn et al., 2005; Hinkel et al., 2005; Grosse et al., 2013). These lakes form as ice-rich permafrost thaws. They grow laterally and deepen, often coalescing with other lakes (Hopkins, 1949; Jorgenson and Shur, 2007). Many thermokarst lakes also drain, leaving depressed wetland basins

TCD

7, 5793–5822, 2013

Seasonal thaw settlement at drained thermokarst lake basins, Arctic Alaska

L. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



known as drained thermokarst lake basins (DTLBs) (Mackay, 1988; Hinkel et al., 2007; Marsh et al., 2009). Taken together, thermokarst lakes and DTLBs cover 50–75 % of the land surface in extensive arctic lowland regions with ice-rich permafrost (Grosse et al., 2013).

Most previous DTLB studies focus on either long-term evolution (e.g., Hinkel et al., 2003; Jorgenson and Shur, 2007; Regmi et al., 2012) or abrupt drainage events (e.g., Mackay, 1988). One exception is a series of field-based studies on Lake Illisarvik in Canada, which was artificially drained in 1978 (Mackay, 1997; Mackay and Burn et al., 2002). Mackay and Burn (2002) reported their 20 yr-long leveling survey results: during the first decade following the lake drainage, the basin surface underwent ~ 10 cm of seasonal uplift (frost heave) and subsidence (thaw settlement); the magnitude of seasonal deformation dropped to ~ 3 cm during the second decade. Here we investigated the dynamic states of DTLBs with a focus on seasonal thaw settlement, which is important for the long-term landscape stability, hydrology, nutrient fluxes, and carbon cycling (Billings and Peterson, 1980; Hinkel et al., 2003; Yoshikawa and Hinzman, 2003; Schuur et al., 2007; Jones et al., 2012).

In areas underlain by permafrost, seasonal frost heave and thaw settlement occurs because of the seasonal phase change between ice and water as the active layer freezes and thaws. Specifically, active layer ice forms in three major mechanisms: (1) freezing and 9 % volume expansion of soil water in pore space, (2) segregational heave due to upward water migration through the frozen fringe to form discrete layers or lenses, defined as segregated ice, and (3) water intrusion under pressure. Seasonal melting of pore ice results in a typical surface deformation of 1–4 cm on arctic lowlands (Little et al., 2003; Liu et al., 2010; Burn, 1990; Nixon et al., 2003). Seasonal formation and melting of segregated and intrusive ice could induce larger surface deformation. For instance, maximum seasonal deformation of 10–30 cm has been observed in field studies (Burn, 1990; Nixon et al., 2003).

Recent advances in remote sensing techniques offer a viable alternative to field-based measurements to monitor thaw settlement in remote locations and across

Seasonal thaw
settlement at drained
thermokarst lake
basins, Arctic Alaska

L. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



larger regions. Interferometric Synthetic Aperture Radar (InSAR) is widely used to map surface deformation in various settings and has been recently applied to detect freeze/thaw-related vertical motion in permafrost lowlands (e.g., Liu et al., 2010; Short et al., 2011). These studies took advantage of the spatial coverage of space-borne data for regional surveys over areas of about 100 km by 100 km.

Our primary objective in this study is to use high resolution InSAR data (~ 10 m) to map and quantify seasonal thaw settlement at individual DTLBs in Arctic Alaska. In previous regional mapping efforts (e.g., Liu et al., 2010), these small landscape components have been ignored. A detailed description and a better representation of the thermokarst-related land surface dynamics are important since they are undergoing large seasonal changes, which will be presented in this study, and may have an impact on geomorphology and hydrology at regional scales.

Because the absolute magnitude of thaw settlement depends on the total water volume in the active layer and thus increases with active layer thickness (ALT) in a first-order approximation, we use thaw strain, defined as the ratio between the thaw settlement and the difference between ALT and thaw settlement, as an index to quantify the relative strength of settlement. We conducted ground-based ALT measurements using probing and ground penetrating radar (GPR) at one DTLB that experienced strong seasonal settlement. Our secondary objective is to demonstrate the combination of remote sensing and ground-based measurements to better quantify surface dynamics and to infer near-surface ground ice content.

2 Study area and methods

2.1 Site description

We studied a 20 km by 20 km area near Prudhoe Bay, on the Arctic Coastal Plain of northern Alaska. This region is underlain by 300–600 m thick permafrost with the upper 2–10 m containing large amounts of excess ground ice (Brown and Sellmann,

Seasonal thaw settlement at drained thermokarst lake basins, Arctic Alaska

L. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Seasonal thaw settlement at drained thermokarst lake basins, Arctic Alaska

L. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1973). ALT in the area varies from 30 to 80 cm (Nelson et al., 1997; Hinkel and Nelson, 2003; Walker et al., 2004; Streletskiy et al., 2008; Liu et al., 2012). The seasonal thaw settlement in this region varies between 1 and 4 cm based on survey-grade differential GPS (Little et al., 2003) and 100-m-resolution InSAR measurements (Liu et al., 2010).

DTLBs are a dominant component of the surface landforms in this region and the formation and drainage of lakes is an important landscape change mechanism (Updike and Howland, 1979; Walker et al., 1985).

We chose this study area following the discovery of very strong seasonal settlements (12 cm, four times what was observed in the surrounding tundra) in one basin centered at 70°8′20″ N, 148°38′58″ W. We informally named this DTLB as SAC basin after the institutions that the authors are affiliated with: Stanford, Alaska, and Colorado. We visited SAC basin in late summer of 2012 to examine the site and measure ALT.

SAC basin has a diameter of ~ 800 m with a small residual pond on the east side connected to a thermokarst lake to the north through a narrow drainage channel (Fig. 1). The basin is flat with the southwest portion rising only 2 m above the pond (Fig. 1b). By contrast, outside the DTLB is a typical tundra landscape with extensive ice-wedge polygon networks. An analysis of historic aerial photos indicates the lake drained before 1949 (not shown). Our field investigation found organic-rich, coarse gravelly and sandy soil along the basin margins with peat and silt in the central area. This spatial pattern is typical of DTLBs and is a result of sediment redistribution during the expansion stage of thermokarst lake development (Murton, 1996; Jorgenson and Shur, 2007).

2.2 InSAR analysis

We applied InSAR processing using a motion-compensation strategy (Zebker et al., 2010) to the Phased Array type L-band Synthetic Aperture Radar (PALSAR) data acquired by the Advanced Land Observing Satellite (ALOS). The basic InSAR product is an interferogram that maps ground surface deformation in the line-of-sight (LOS) direction between the acquisitions of two radar images. We used two ascending PALSAR paths (250 and 251) and produced 33 interferograms between 2006 and 2010,

which are shown in Fig. 2. These two interferogram networks of redundant measurements were used to reduce errors and to estimate the seasonal vertical deformation (Sect. 2.3). The y axes of the Fig. 2 plots represent the perpendicular baseline of each interferogram, which is defined as the geometric separation of satellite positions at two acquisitions perpendicular to the LOS direction. Short perpendicular baselines reduce the InSAR geometric decorrelation (Zebker and Villasenor, 1992).

We also used a LiDAR DEM to remove the topographic contribution to the interferograms. This DEM has a fine spatial posting of 1 m and a vertical accuracy of a few centimeters (Paine et al., 2013). We down-sampled the DEM to 10 m to match the interferogram posting. Following Liu et al. (2010), we used a river floodplain point ($70^{\circ}4'40''$ N, $148^{\circ}32'15''$ W) as a reference because the presence of coarse gravel results in a stable surface with little thaw settlement (Pullman et al., 2007). We also assumed surface deformation is purely vertical and converted the InSAR-measured LOS deformation to the vertical direction. This is a valid assumption within the low-lying, flat and homogeneous basin. In theory, lateral motions may occur on polygonal terrains due to thermal contraction and expansion. However, the reported lateral motions are small (less than 1 cm yr^{-1}) and confined in local scales across individual ice wedges (Mackay, 2000; Mackay and Burn, 2002).

In this study we only used radar images acquired during thaw seasons to avoid severe phase decorrelation and artifacts due to snow cover. We also ruled out the possibility of detecting water level change instead of the basin elevation change. The basin is possibly flooded during snowmelt in early thaw season. However, InSAR detection of a flooded water body requires double-bounce backscattering that occurs at the interface between water surface and woody trunks/stems, which seldom occurs over tundra wetlands where surface vegetation are randomly distributed grasses/sedges. Double-bounce backscattering also results in strong radar amplitude and InSAR coherence, neither of which was observed in the interferograms over SAC basin.

Seasonal thaw
settlement at drained
thermokarst lake
basins, Arctic Alaska

L. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.3 Estimation of thaw settlement

Because individual interferograms are contaminated by errors due to atmospheric artifacts and decorrelation, we used all 33 interferograms and a time series model to invert for the multi-year average seasonal thaw settlement over 2006–2010. Here we briefly summarize the inversion method that has been described in detail by Liu et al. (2012). We modeled the InSAR-measured ground vertical deformation as a summation of a linear trend during 2006–2010 and the seasonal subsidence during thaw months as a linear function of the square root of the accumulated degree days of thaw (ADDT). We included the linear trend component for completeness, although we expected the trend was small. One difference from Liu et al. (2012) is that we excluded the topographic error term in our model because the LiDAR DEM has a high accuracy. Mathematically, InSAR-observed deformation (D) between two thaw-season dates (t_1 and t_2), can be expressed as

$$D = R(t_2 - t_1) + E \left(\sqrt{A_2} - \sqrt{A_1} \right) + \varepsilon_{\text{InSAR}}, \quad (1)$$

where R is the long-term subsidence rate, E is the seasonal coefficient, A is the ADDT with the subscripts corresponding to the two dates, and $\varepsilon_{\text{InSAR}}$ is InSAR measurement errors that we assume are random. We used the air temperature records at the Deadhorse Airport weather station (<http://www.ncdc.noaa.gov>) to calculate ADDTs. We then fitted for both R and E from all interferograms using the least-squares method and estimated their uncertainties. This inversion approach minimizes the contributions in the estimated seasonal subsidence due to temporal variations of soil moisture and tropospheric water vapor, because these two factors are not correlated with the ADDTs.

Despite excluding winter interferograms, we assumed that the ground vertical deformation is a full seasonal cycle in each year: i.e., thaw-season subsidence and freeze-season uplift with the same magnitude. This is supported by the fact that our fitted linear trends during 2006–2010 have a magnitude $< 0.5 \text{ cm yr}^{-1}$ that is smaller than the accuracy of PALSAR interferometry (Sandwell et al., 2008). In the remainder of this

TCD

7, 5793–5822, 2013

Seasonal thaw settlement at drained thermokarst lake basins, Arctic Alaska

L. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



paper, we refer to the fitted maximum seasonal subsidence averaged between 2006 and 2010 as the InSAR-estimated thaw settlement and denote it as ΔZ .

2.4 GPR data analysis and ALT estimation

GPR is a non-invasive geophysical method that uses electromagnetic signals to im-
age the subsurface. GPR measures the travel time of radar waves transmitted into the
ground and reflected at boundaries with different dielectric permittivity back to a re-
ceiver on the surface. Multiplying the one-way travel time to a reflecting interface by
wave speed gives the estimated depth of the reflections. Frozen and unfrozen sedi-
ments have a prominent dielectric contrast and result in clear, measurable radar re-
flections. For this reason, GPR is commonly used to map the spatial variability of
thaw depth (e.g., Doolittle et al., 1990; Bradford et al., 2005; Brosten et al., 2006;
Wollschläger et al., 2010; Hubbard et al., 2013).

We conducted a GPR survey to measure the thaw depth in SAC basin on 18 Au-
gust 2012 along the red transect in Fig. 1a using a 500 MHz PulseEKKOPro 1000 sys-
tem. We also measured the thaw depth at ten locations using a metal probe to validate
the GPR estimates and calibrate the radar wave speed. The GPR transect was ac-
quired in common-offset mode, i.e., the distance between the two radar transducers
was constant during the entire survey. The GPR unit was set to collect a radar trace
every 0.2 s.

We processed the GPR data using standard routines (Neal, 2004) including remov-
ing the instrumental, low-frequency noise (“dewow”), 250–750 MHz bandpass filtering,
and an approximate spherical spreading correction by scaling the amplitude by the
travel time. Additional data analysis details can be found in Gusmeroli and Grosse
(2012). We estimated radar speed within the subsurface by comparing radar travel
times and probed thaw depth. Radar wave propagation speed within the active layer
soils is 0.042 m ns^{-1} , 0.052 m ns^{-1} , and 0.042 m ns^{-1} outside the basin, in the sandy
margin areas, and inside the basin, respectively. Because we conducted GPR mea-
surements near the end of the 2012 thaw season, the derived thaw depth is equivalent

Seasonal thaw settlement at drained thermokarst lake basins, Arctic Alaska

L. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



to the active layer thickness which we refer to as GPR ALT. Uncertainties in the GPR ALT are ± 6 cm based on calibration with the probing measurements.

2.5 Thaw strain

Because the magnitude of thaw settlement typically increases with the thaw depth, we adopted thaw strain (ε) as an index to quantify the relative strength of thaw settlement subsidence, which is defined as:

$$\varepsilon = \frac{\Delta Z}{H - \Delta Z}, \quad (2)$$

where ΔZ is the thaw settlement and H is the depth of thaw penetration (Pullman et al., 2007). In the context of examining the maximum seasonal deformation at SAC basin, ΔZ is the InSAR-measured thaw settlement and H is the GPR-estimated ALT. A similar and more commonly used index is heave strain, which is defined as the ratio of the frost heave over the depth of frost penetration minus the frost heave (Burn, 1990; French, 2007). The thaw and frost strains are equivalent if the thaw settlement and frost heave processes are reversible.

When active layer thawing occurs in a closed system, thaw settlement is only caused by the volume decrease associated with the melting of pore ice. Ignoring secondary mechanisms affecting thaw settlement (such as contraction of clay soils, consolidation of soil on thawing, surface erosion, changes in soil density, and unfrozen water in the frozen soils), the thaw settlement is simply determined by the total volume of water in the active layer (Liu et al., 2012):

$$\Delta Z = H\theta \frac{\rho_w - \rho_i}{\rho_i}, \quad (3)$$

Seasonal thaw settlement at drained thermokarst lake basins, Arctic Alaska

L. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

where θ is the volumetric water content (VWC) of the active layer, ρ_w is the density of water, and ρ_i is the density of ice. Substituting Eq. (3) into Eq. (2) gives:

$$\varepsilon = \frac{\theta(\rho_w - \rho_i)}{\rho_i - \theta(\rho_w - \rho_i)}, \quad (4)$$

- 5 which is determined by θ and independent of H . According to Eq. (4), thaw strain in a closed system increases monotonically with θ and has an upper bound of 10 % when $\theta = 100\%$ (i.e., pure water body).

2.6 Estimation of volumetric water content from radar speed

10 Water content also largely determines the dielectric constant and thus the radar speed estimated from GPR measurements. In this subsection, we describe four methods of estimating ranges of θ (i.e. VWC) from the radar speeds, which we had determined using GPR and probing measurements (Sect. 2.4). We then used the θ values in Eq. (4) to predict thaw strains and compare with the observations.

15 Engstrom et al. (2005) developed the following empirical model for the active layer over Barrow on the Arctic coast of Alaska:

$$\theta = -2.5 + 2.508\kappa - 3.634 \times 10^{-2}\kappa^2 + 2.394 \times 10^{-4}\kappa^3, \quad (5)$$

where κ is the dielectric constant of the soil, which can be directly calculated from the radar speed v as:

$$\kappa = \left(\frac{c_0}{v}\right)^2, \quad (6)$$

25 where c_0 is the speed of light in free space. We note that the study area of Engstrom et al. (2005) includes both DTLBs and upland terrains. Therefore, this empirical model is representative of the overall landscape in Barrow and may introduce some bias when applied to SAC basin.

Seasonal thaw settlement at drained thermokarst lake basins, Arctic Alaska

L. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



We also considered a few other empirical relations for specific soil types. For instance, Parsekian et al. (2012) reported a linear regression relation for peat soils under near saturation conditions ($\theta > 85\%$):

$$\theta_{\text{peat}} = 4.4 \times 10^{-1} + 7.3 \times 10^{-3} \kappa. \quad (7)$$

For mineral soils, the following empirical equation developed by Topp et al. (1980) is widely used:

$$\theta_{\text{mineral}} = -5.3 \times 10^{-2} + 2.92 \times 10^{-2} \kappa - 5.5 \times 10^{-4} \kappa^2 + 4.3 \times 10^{-6} \kappa^3. \quad (8)$$

For saturated soils, the semi-empirical two-phase mixing equation based on the Complex Refractive Index Model (CRIM) is widely used to relate the dielectric constant to θ (e.g., Greaves et al., 1996):

$$\sqrt{\kappa} = \theta \sqrt{\kappa_w} + (1 - \theta) \sqrt{\kappa_g}, \quad (9)$$

where κ_w and κ_g are the dielectric constants for pure water and dry soil grains, respectively. Solving for θ gives

$$\theta = \frac{\sqrt{\kappa} - \sqrt{\kappa_g}}{\sqrt{\kappa_w} - \sqrt{\kappa_g}}. \quad (10)$$

Eq. (10) can be equivalently expressed as a function of radar speeds in the mixed soil (ν), pure water (ν_w), and dry soil (ν_g) as

$$\theta = \frac{\nu_w (\nu_g - \nu)}{\nu (\nu_g - \nu_w)}. \quad (11)$$

We used $\nu_g = 0.09 \text{ mns}^{-1}$ for dry organic matters, and 0.11 mns^{-1} for dry silt (Davis and Annan, 1989).

Seasonal thaw settlement at drained thermokarst lake basins, Arctic Alaska

L. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3 Results

3.1 Seasonal thaw settlement

All interferograms we produced show the same general pattern in SAC basin and its surroundings. In this subsection, we first show one interferogram as an example to illustrate strong settlement within SAC basin in a regional context including other DTLBs. We then present a detailed map of the 2006–2010 averaged thaw settlement at SAC basin.

The interferogram in Fig. 3 was formed using SAR images taken at the start and end of the 2007 thaw season, therefore representing the seasonal subsidence for a single year. Of the 18 DTLBs that show sufficient high InSAR coherence on this interferogram and have a size larger than 4 ha, 14 underwent similar subsidence as the surrounding tundra while four showed larger subsidence than the surrounding tundra. In particular, SAC basin experienced 12 cm of subsidence, or as much as 8–10 cm more than the surrounding area. SAC basin is clearly exceptional in terms of the magnitude of seasonal subsidence, but it is not unique.

The Trans-Alaska Pipeline is visible next to SAC basin as a linear and segmented feature. Such multiple-scattering signature is common on SAR images for man-made structures that are parallel to the ground surface (Zebker and Goldstein, 1986). A detailed investigation on the pipeline is outside the scope of this study.

Figure 4a shows the 2006–2010 average seasonal subsidence over SAC basin and its surrounding area, estimated using all 33 interferograms. Areas of low coherence, mostly over water bodies, are masked in gray. The tundra area outside SAC basin underwent an average seasonal thaw settlement of 3–4 cm. Consistent with the 2007 snapshot (Fig. 3), SAC Basin shows a large thaw settlement of up to 12 cm occurred over the western half of the basin bounded by the dry margins. Subsidence uncertainties range from 1 to 3 cm with a small spatial variability. Figure 4b shows the relative uncertainties, defined as the ratios between the uncertainties and the absolute settle-

Seasonal thaw settlement at drained thermokarst lake basins, Arctic Alaska

L. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ment. Over the area with large seasonal subsidence, the relative uncertainties are less than 20 %.

3.2 Active layer water content and theoretical thaw strain

The active layer at the basin center is fully saturated and consists of peat and silt, although the exact proportions are unknown. We used radar speeds of 0.042 m ns^{-1} , representative for the entire active layer, to estimate VWC using the methods described in Sect. 2.6. We conducted similar calculations for the sandy basin margins, where radar speed is 0.052 m ns^{-1} . Table 1 lists the results. Given the uncertainty of the active layer soil composition and the spread results using different methods, we rounded off these estimates to represent the range of VWC as 60–80 % and 45–55 %, for the basin center/outer basin and the sandy margins, respectively. The VWC for saturated peat is about 90 % (Price et al., 2005), so the VWC we estimate for the basin center is typical for soils with a large amount of organic material. The VWC of pure mineral soil is about 45 %, depending on clay content (Clapp and Hornberger, 1978; Cosby et al., 1984), so the VWC we estimated for the sandy margins is typical of sandy soil with a small amount of organic material.

Based on these VWC estimates, we predicted thaw strains for saturated soils in a closed system using Eq. (4). The solid line in Fig. 5 denotes the full range of thaw strain with an upper limit of 10 %. The GPR-estimated VWCs of 60–80 % at the basin center and the outer basin correspond to “theoretical” thaw strains ranging from 6 % to 8 %; whereas VWCs of 45–55 % along the basin margins correspond to thaw strains of 4–5 %, both for saturated soils in a closed system.

3.3 Active layer thickness and estimated thaw strain

In this subsection, we present the ALT, the seasonal subsidence, and the thaw strains along the GPR transect (Fig. 6). We divided the transect into areas outside the basin (“AB” and “EF”), the sandy margins (“BC” and “DE”), and the basin center (“CD”). We

Seasonal thaw settlement at drained thermokarst lake basins, Arctic Alaska

L. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



calibrated the GPR ALT with the probing data, thus constraining a good match between these two measurements at all probing locations (Fig. 6a). ALT was about 0.4 m outside the basin, increasing to 0.7–1.3 m at the basin margins. At the basin center, ALT was relatively uniform with an average of 0.54 m.

5 GPR combined with in situ probing provides a continuous profile of the ALT across multiple geological units at SAC basin. The ALT probe we used in the field has a maximum length of 1 m, which did not reach the bottom of the active layer at the sandy margins. We overcame this thickness constraint and detected the > 1 m ALT by using GPR. The increased ALTs at the sandy margins from the surrounding typical tundra terrain are due to the high thermal conductivity and low latent heat (Osterkamp and Burn,
10 2003). Such characteristic pattern has been observed by in situ ALT measurements on the Arctic Coastal Plain of Alaska (e.g., Hinkel and Nelson, 2003; Shiklomanov et al., 2010).

15 The InSAR-measured seasonal thaw settlement along the transect was 2–4 cm outside the basin (“AB”), abruptly increasing to up to 12 cm at the basin center (subsidence bowl, “BD”), then decreasing to about 4 cm on the other side of the basin (“DF”). ALTs were similarly large at both margin sections (“BC” and “DE”), whereas subsidence was much larger along “BC” than “DE”.

20 The thaw strains along the same profile show a similar but inversed spatial pattern as the thaw settlement (Fig. 6c). Outside the basin, thaw strains were 5–10 % with a mean of 7 %. Along the sandy margin “DE”, thaw strains were roughly constant at 3 %. At the entire basin center and part of the sandy margin “BC”, thaw strains were systematically larger than 10 %.

4 Discussion

25 Comparisons between the calculated thaw strains (Fig. 6c) and theoretical predictions based on VWC suggest the seasonal formation of excess ground ice. Outside the basin, the averaged thaw strain was 7 %, consistent with the theoretical range (6–8 %,

TCD

7, 5793–5822, 2013

Seasonal thaw settlement at drained thermokarst lake basins, Arctic Alaska

L. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



given in Sect. 3.2). The averaged thaw strain along the margin “DE” was 3 %, slightly lower than the predicted 4–5 % for saturated sandy soils in a closed system. Thaw strains at the basin center exceeded the 10 % upper bound, suggesting the presence of ground ice whose equivalent VWC exceeds the soil pore volume in the active layer.

In wet conditions, development of segregated ice and intrusive ice can produce heave that exceeds what would be predicted from the volume expansion during the freezing of pore water, equivalently resulting in thaw/frost strains larger than 10 %. For instance, based on soil core analysis, Pullman et al. (2007) reported thaw strains that range of 20–35 % for the top 1 m of ice-rich DTLB soils on the Beaufort Coastal Plain of Alaska, where SAC basin is located. Similarly, Burn (1990) reported frost strains of 20–60 % for saturated, fine-grained lake-bottom sediments in the Mackenzie River Delta.

Ground ice is abundant on the Arctic coastal plain (e.g., Kokelj and Burn, 2005; Kanevskiy et al., 2013). Specifically for SAC basin, ground ice in the active layer possibly bears the form of segregated ice and intrusive ice, resulting from two distinct processes. Following the lake drainage and refreezing of surface soils, ice enrichment occurs in the water-saturated, fine-grained active layer in the form of segregated ice as permafrost aggrades from below (Bockheim and Hinkel, 2012). On the other hand, SAC basin has a residual pond on its east side. During the winter when the pond freezes, pressure is built within the remaining liquid water, which may inject the floodwater laterally along the base of the active layer and form ground ice, which could be a secondary ice-forming mechanism. We note that ground ice is also present in the surrounding tundra including the polygonal outer basin. Our thaw strain calculations suggest that its water equivalent is confined within the soil pore space, in contrast with the basin center where extensive ground ice exceeds the pore space.

Mapping and quantifying vertical motion at individual landscape components such as DTLBs described in this study is important for better describing regional patterns in surface deformation on arctic coastal lowlands. These landscapes are characterized by a mosaic of extent lakes, DTLBs, and remnant topography (Frohn et al., 2005; Hinkel

Seasonal thaw settlement at drained thermokarst lake basins, Arctic Alaska

L. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

et al., 2005; Grosse et al., 2013) that vary by ground-ice content and soil sediment grain size (Kanevskiy et al., 2013). These differences result in highly variable seasonal deformation that becomes averaged across the landscape when using coarser resolution imagery employed in previous studies (e.g., Liu et al., 2010). The maximum observed motion of 8–12 cm that was observed in SAC basin and the other DTLBs that experienced greater than 4 cm of seasonal deformation highlight the importance of conducting high spatial resolution InSAR studies in these settings. It is likely that landscape components of various ages and hence ground ice content will respond differently to climate change. Therefore, the ability to detect this with InSAR at the level of individual landscape components is important for documenting dynamics due to thermokarst and landscape evolution on arctic coastal lowlands. Given the large amount and extensive coverage of DTLBs on arctic lowlands and the availability of SAR data acquired by multiple satellites, it would be valuable to conduct a regional-scale InSAR analysis to identify basins that are experiencing similar seasonal deformation as SAC basin.

5 Conclusions

We have applied the InSAR technique to ALOS PALSAR data to examine the surface dynamics of DTLBs in Northern Alaska. Most of the DTLBs in our study area underwent seasonal thaw settlement of 3–4 cm. We have also found a prominent seasonal thaw settlement at one particular DTLB, SAC basin, every year from 2006 to 2010 with a maximum of 12 cm with less than 20 % of relative uncertainty. This is significantly larger than the thaw settlement observed at the surrounding tundra area and other nearby DTLBs.

Combining the InSAR and the GPR ALT measurements at SAC basin, we calculated thaw strain and used it as an index of settlement strength. The averaged thaw strain was 7 % and 3 % outside the basin and along the sandy basin margins, respectively, quantitatively consistent with the theoretical values for freeze and thaw of saturated or near-saturated active layer soils. Thaw strains at the basin center were systematically

Seasonal thaw settlement at drained thermokarst lake basins, Arctic Alaska

L. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



larger than 10 % with a maximum value of 35 %, indicating the presence of excess ground ice the active layer.

Previous InSAR studies overlooked deformation at individual DTLBs. If the observations reported in this study are related to a common process over a broad region, such large seasonal thaw settlement has significant implications for permafrost hydrology and landscape dynamics. Capturing spatial heterogeneity of landscape-scale processes as suggested in this study is important for better describing regional patterns in surface deformation in arctic coastal lowlands.

Acknowledgements. The ALOS PALSAR data are copyrighted by the Japan Aerospace Exploration Agency and provided by the Alaska Satellite Facility, University of Alaska Fairbanks. L. Liu was supported by the George Thompson Postdoctoral Fellowship from the Department of Geophysics, Stanford University and the US National Science Foundation (NSF) grant ARC-1204013. K. Schaefer and T. Zhang were in part supported by the US National Aeronautics and Space Administration (NASA) NNX10AR63G, the US National Oceanic and Atmospheric Administration NA09OAR4310063, and the NSF ARC-0901962. A. Gusmeroli was supported by the Alaska Climate Science Center, funded by Cooperative Agreement Number G10AC00588 from the US Geological Survey. G. Grosse was supported by NASA NNX11AH20G and NSF ARC-1107481. B. M. Jones was supported by the US Geological Survey, Alaska Science Center and the Land Change Science program. We thank J. Paine, A. Hosford Scheirer, E. Duncan and K. Duncan (Great Bear Petroleum) for providing the LiDAR DEM. We thank E. Pettit, J. Munk, M. Sturm, and C. Jones for providing field instruments. Any use of trade, product, or firm names is for descriptive purposes only and does not imply endorsement by the US Government.

References

- Billings, W. and Peterson, K.: Vegetational change and ice-wedge polygons through the thaw-lake cycle in Arctic Alaska, *Arctic Alpine Res.*, 12, 413–432, 1980.
- Bockheim, J. G. and Hinkel, K. M.: Accumulation of excess ground ice in an age sequence of drained thermokarst lake basins, Arctic Alaska, *Permafrost Periglac. Process.*, 23, 231–236, doi:10.1002/ppp.1745, 2012.

TCD

7, 5793–5822, 2013

Seasonal thaw settlement at drained thermokarst lake basins, Arctic Alaska

L. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Seasonal thaw settlement at drained thermokarst lake basins, Arctic Alaska

L. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Bradford, J. H., McNamara, J. P., Bowden, W., and Gooseff, M. N.: Measuring thaw depth beneath peat-lined arctic streams using ground-penetrating radar, *Hydrol. Process.*, 19, 2689–2699, 2005.
- 5 Brosten, T. R., Bradford, J. H., McNamara, J. P., Zarnetske, J. P., Gooseff, M. N., and Bowden, W. B.: Profiles of temporal thaw depths beneath two arctic stream types using ground-penetrating radar, *Permafrost Periglac. Process.*, 17, 341–355, 2006.
- Brown, J. and Sellmann, P. V.: Permafrost and coastal plain history of arctic Alaska, in: *Alaskan Arctic Tundra*, edited by Britton, M. E., no. 25 in Arctic Institute of North America, Technical Paper, 31–47, 1973.
- 10 Burn, C.: Frost heave in lake-bottom sediments, Mackenzie Delta, Northwest Territories, in: 5th Canadian Permafrost Conference, Université Laval, Québec, 6–8 June 1990, 54, 103–109, 1990.
- Clapp, R. B. and Hornberger, G. M.: Empirical equations for some soil hydraulic properties, *Water Resour. Res.*, 14, 601–604, 1978.
- 15 Cosby, B., Hornberger, G., Clapp, R., and Ginn, T.: A statistical exploration of the relationships of soil moisture characteristics to the physical properties of soils, *Water Resour. Res.*, 20, 682–690, 1984.
- Davis, J. and Annan, A.: Ground-penetrating radar for high-resolution mapping of soil and rock stratigraphy, *Geophys. Prospect.*, 37, 531–551, 1989.
- 20 Doolittle, J., Hardisky, M., and Gross, M.: A ground-penetrating radar study of active layer thicknesses in areas of moist sedge and wet sedge tundra near Bethel, Alaska, USA, *Arctic Alpine Res.*, 22, 175–182, 1990.
- Engstrom, R., Hope, A., Kwon, H., Stow, D., and Zamolodchikov, D.: Spatial distribution of near surface soil moisture and its relationship to microtopography in the Alaskan Arctic Coastal Plain, *Nord. Hydrol.*, 36, 219–234, 2005.
- 25 French, H. M.: *The Periglacial Environment*, John Wiley & Sons, Ltd., New York, 2007.
- Frohn, R., Hinkel, K., and Eisner, W.: Satellite remote sensing classification of thaw lakes and drained thaw lake basins on the North Slope of Alaska, *Remote Sens. Environ.*, 97, 116–126, 2005.
- 30 Greaves, R. J., Lesmes, D. P., Lee, J. M., and Toksöz, M. N.: Velocity variations and water content estimated from multi-offset, ground-penetrating radar, *Geophysics*, 61, 683–695, 1996.

Seasonal thaw settlement at drained thermokarst lake basins, Arctic Alaska

L. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Grosse, G., Jones, B., and Arp, C.: Thermokarst lakes, drainage, and drained basins, in: Treatise on Geomorphology, edited by Shroder, J. G. R. and Harbor, J., Elsevier Academic Press, San Diego, California, 81–29, 2013.

Gusmeroli, A. and Grosse, G.: Ground penetrating radar detection of subsnow slush on ice-covered lakes in interior Alaska, *The Cryosphere*, 6, 1435–1443, doi:10.5194/tc-6-1435-2012, 2012.

Hinkel, K., Eisner, W., Bockheim, J., Nelson, F., Peterson, K., and Dai, X.: Spatial extent, age, and carbon stocks in drained thaw lake basins on the Barrow Peninsula, Alaska, *Arctic Antarctic Alpine Res.*, 35, 291–300, 2003.

Hinkel, K., Frohn, R., Nelson, F., Eisner, W., and Beck, R.: Morphometric and spatial analysis of thaw lakes and drained thaw lake basins in the western Arctic Coastal Plain, Alaska, *Permafrost Periglac.*, 16, 327–341, 2005.

Hinkel, K. M., Jones, B. M., Eisner, W. R., Cuomo, C. J., Beck, R. A., and Frohn, R.: Methods to assess natural and anthropogenic thaw lake drainage on the western Arctic coastal plain of northern Alaska, *J. Geophys. Res.*, 112, F02S16, doi:10.1029/2006JF000584, 2007.

Hopkins, D. M.: Thaw lakes and thaw sinks in the Imuruk Lake Area, Seward Peninsula, Alaska, *J. Geol.*, 57, 119–131, 1949.

Hubbard, S. S., Gangodagamage, C., Dafflon, B., Wainwright, H., Peterson, J., Gusmeroli, A., Ulrich, C., Wu, Y., Wilson, C., Rowland, J., Tweedie, C., and Wulfschleger, S.: Quantifying and relating land-surface and subsurface variability in permafrost environments using LiDAR and surface geophysical datasets, *Hydrogeol. J.*, 21, 149–169, 2013.

Jones, M. C., Grosse, G., Jones, B. M., and Walter Anthony, K.: Peat accumulation in drained thermokarst lake basins in continuous, ice-rich permafrost, northern Seward Peninsula, Alaska, *J. Geophys. Res.*, 117, G00M07, doi:10.1029/2011JG001666, 2012.

Jorgenson, M. T. and Shur, Y.: Evolution of lakes and basins in northern Alaska and discussion of the thaw lake cycle, *J. Geophys. Res.*, 112, F02S17, doi:10.1029/2006JF000531, 2007.

Kanevskiy, M., Shur, Y., Jorgenson, M., Ping, C.-L., Michaelson, G., Fortier, D., Stephani, E., Dillon, M., and Tumskey, V.: Ground ice in the upper permafrost of the Beaufort Sea coast of Alaska, *Cold Reg. Sci. Technol.*, 85, 56–70, 2013.

Kokelj, S. and Burn, C.: Near-surface ground ice in sediments of the Mackenzie Delta, Northwest Territories, Canada, *Permafrost Periglac. Process.*, 16, 291–303, 2005.

Seasonal thaw settlement at drained thermokarst lake basins, Arctic Alaska

L. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Little, J., Sandall, H., Walegur, M., and Nelson, F.: Application of Differential Global Positioning Systems to monitor frost heave and thaw settlement in tundra environments, *Permafrost Periglac. Process.*, 14, 349–357, doi:10.1002/ppp.466, 2003.
- Liu, L., Zhang, T., and Wahr, J.: InSAR measurements of surface deformation over permafrost on the North Slope of Alaska, *J. Geophys. Res.*, 115, F03023, doi:10.1029/2009JF001547, 2010.
- Liu, L., Schaefer, K., Zhang, T., and Wahr, J.: Estimating 1992–2000 average active layer thickness on the Alaskan North Slope from remotely sensed surface subsidence, *J. Geophys. Res.*, 117, F01005, doi:10.1029/2011JF002041, 2012.
- Mackay, J.: Catastrophic lake drainage, Tuktoyaktuk Peninsula area, District of Mackenzie, Current Research, Part D, Geological Survey of Canada, Paper, 88, 83–90, 1988.
- Mackay, J. R.: A full-scale field experiment (1978–1995) on the growth of permafrost by means of lake drainage, western Arctic coast: A discussion of the method and some results, *Can. J. Earth Sci.*, 34, 17–33, 1997.
- Mackay, J. R.: Thermally induced movements in ice-wedge polygons, western Arctic coast: a long-term study, *Geogr. Phys. Quaternaire*, 54, 41–68, 2000.
- Mackay, J. R. and Burn, C. R.: The first 20 years (1978–1979 to 1998–1999) of active-layer development, Illisarvik experimental drained lake site, western Arctic coast, Canada, *Canadian J. Earth Sci.*, 39, 1657–1674, 2002.
- Marsh, P., Russell, M., Pohl, S., Haywood, H., and Onclin, C.: Changes in thaw lake drainage in the Western Canadian Arctic from 1950 to 2000, *Hydrol. Process.*, 23, 145–158, 2009.
- Murton, J. B.: Thermokarst-lake-basin sediments, Tuktoyaktuk Coastlands, western arctic Canada, *Sedimentology*, 43, 737–760, 1996.
- Neal, A.: Ground-penetrating radar and its use in sedimentology: principles, problems and progress, *Earth-Sci. Rev.*, 66, 261–330, 2004.
- Nelson, F. E., Shiklomanov, N. I., Mueller, G. R., Hinkel, K. M., Walker, D. A., and Bockheim, J. G.: Estimating active-layer thickness over a large region: Kuparuk river basin, Alaska, USA, *Arctic Alpine Res.*, 29, 367–378, 1997.
- Nixon, M., Tarnocai, C., and Kutny, L.: Long-term active layer monitoring: Mackenzie Valley, northwest Canada, in: *Proceedings of the 8th International Conference on Permafrost*, 2, 821–826, 2003.
- Osterkamp, T. E. and Burn, C. R.: Permafrost, in: *Encyclopedia of Atmospheric Sciences*, edited by Holton, J. R., Pyle, J., and Curry, J. A., 1717–1729, Academic Press, 2002.

**Seasonal thaw
settlement at drained
thermokarst lake
basins, Arctic Alaska**

L. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Paine, J. G., Andrews, J. R., Saylam, K., Tremblay, T. A., Averett, A. R., Caudle, T. L., Meyer, T., and Young, M. H.: Airborne lidar on the Alaskan North Slope: Wetlands mapping, lake volumes, and permafrost features, *The Leading Edge*, 32, 798–805, 2013.
- Parsekian, A. D., Slater, L., and Giménez, D.: Application of ground-penetrating radar to measure near-saturation soil water content in peat soils, *Water Resour. Res.*, 48, W02533, doi:10.1029/2011WR011303, 2012.
- Price, J. S., Cagampan, J., and Kellner, E.: Assessment of peat compressibility: is there an easy way?, *Hydrol. Process.*, 19, 3469–3475, 2005.
- Pullman, E. R., Jorgenson, M. T., and Shur, Y.: Thaw settlement in soils of the Arctic Coastal Plain, Alaska, *Arct. Antarct. Alp. Res.*, 39, 468–476, 2007.
- Regmi, P., Grosse, G., Jones, M. C., Jones, B. M., and Anthony, K. W.: Characterizing post-drainage succession in thermokarst lake basins on the Seward Peninsula, Alaska with TerraSAR-X backscatter and Landsat-based NDVI data, *Remote Sens.*, 4, 3741–3765, 2012.
- Sandwell, D. T., Myer, D., Mellors, R., Shimada, M., Brooks, B., and Foster, J.: Accuracy and resolution of ALOS interferometry: Vector deformation maps of the Father's Day intrusion at Kilauea, *IEEE T. Geosci. Remote*, 46, 3524–3534, 2008.
- Schuur, E. A. G., Crummer, K., Vogel, J., and Mack, M.: Plant species composition and productivity following permafrost thaw and thermokarst in Alaskan tundra, *Ecosystems*, 10, 280–292, 2007.
- Sellmann, P., Brown, J., Lewellen, R., McKim, H., and Merry, C.: The Classification and Geomorphic Implications of Thaw Lakes on the Arctic Coastal Plain, Alaska, *Tech. Rep. 344*, Cold Regions Res. and Eng. Lab., Hanover, NH, 1975.
- Shiklomanov, N. I., Streletskiy, D. A., Nelson, F. E., Hollister, R. D., Romanovsky, V. E., Tweedie, C. E., Bockheim, J. G., and Brown, J.: Decadal variations of active-layer thickness in moisture-controlled landscapes, Barrow, Alaska, *J. Geophys. Res.*, 115, G00104, doi:10.1029/2009JG001248, 2010.
- Short, N., Brisco, B., Couture, N., Pollard, W., Murnaghan, K., and Budkewitsch, P.: A comparison of TerraSAR-X, RADARSAT-2 and ALOS-PALSAR interferometry for monitoring permafrost environments, case study from Herschel Island, Canada, *Remote Sens. Environ.*, 115, 3491–3506, 2011.

Seasonal thaw settlement at drained thermokarst lake basins, Arctic Alaska

L. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Streletskiy, D. A., Shiklomanov, N. I., Nelson, F. E., and Klene, A. E.: Long-term active and ground surface temperature trends: 13 years of observations at Alaskan CALM sites, in: Ninth International Conference on Permafrost, 29, 1727–1732, 2008.
- 5 Topp, G., Davis, J., and Annan, A. P.: Electromagnetic determination of soil water content: Measurements in coaxial transmission lines, *Water Resour. Res.*, 16, 574–582, 1980.
- Updike, R. and Howland, M.: Surficial geology and processes, Prudhoe Bay oil field, with Hydrologic implications, Tech. rep., Alaska Division of Geological and Geophysical Surveys Special Report, 1979.
- 10 Walker, D. A., Walker, M. D., Everett, K. R., and Webber, P.: Pingos of the Prudhoe Bay Region, *Alaska Arctic Alpine Res.*, 17, 321–336, 1985.
- Walker, D. A., Epstein, H. E., Gould, W. A., Kelley, A. M., Kade, A. N., Knudson, J. A., Krantz, W. B., Michaelson, G., Peterson, R. A., Ping, C.-L., Raynolds, M. K., Romanovsky, V. E., and Shur, Y.: Frost-boil ecosystems: complex interactions between landforms, soils, vegetation and climate, *Permafrost Periglac.*, 15, 171–188, doi:10.1002/ppp.487, 2004.
- 15 Wollschläger, U., Gerhards, H., Yu, Q., and Roth, K.: Multi-channel ground-penetrating radar to explore spatial variations in thaw depth and moisture content in the active layer of a permafrost site, *The Cryosphere*, 4, 269–283, doi:10.5194/tc-4-269-2010, 2010.
- Yoshikawa, K. and Hinzman, L. D.: Shrinking thermokarst ponds and groundwater dynamics in discontinuous permafrost near council, Alaska, *Permafrost Periglac. Process.*, 14, 151–160, 2003.
- 20 Zebker, H. A. and Goldstein, R. M.: Topographic mapping from interferometric synthetic aperture radar observations, *J. Geophys. Res.*, 91, 4993–4999, 1986.
- Zebker, H. A. and Villasenor, J.: Decorrelation in interferometric radar echoes, *IEEE T. Geosci. Remote*, 30, 950–959, 1992.
- 25 Zebker, H., Hensley, S., Shanker, P., and Wortham, C.: Geodetically Accurate InSAR Data Processor, *IEEE T. Geosci. Remote*, 48, 4309–4321, doi:10.1109/TGRS.2010.2051333, 2010.

Seasonal thaw settlement at drained thermokarst lake basins, Arctic Alaska

L. Liu et al.

Table 1. Volumetric water content (VWC, θ) estimated based on the radar speed (0.042 m ns^{-1}) at the basin center and outer basin, and the radar speed (0.052 m ns^{-1}) at the sandy margins.

Location	VWC (%)	Method
Basin center and outer basin	63	Empirical active layer, Eq. (5)
	81	Empirical peat model, Eq. (7)
	68	Two-phase mixing model for peat, Eq. (11), $v_g = 0.09 \text{ m ns}^{-1}$
	71	Two-phase mixing model for silt, Eq. (11), $v_g = 0.11 \text{ m ns}^{-1}$
Sandy margins	50	Empirical active layer, Eq. (5)
	47	Empirical mineral model, Eq. (8)
	54	Two-phase mixing model for peat, Eq. (11), $v_g = 0.15 \text{ m ns}^{-1}$

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Seasonal thaw settlement at drained thermokarst lake basins, Arctic Alaska

L. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table A1. Notation and constants.

D	Ground vertical deformation measured in individual InSAR interferogram (m)
R	Long-term subsidence rate (m yr^{-1})
t	Time (yr)
E	Coefficient of seasonal subsidence ($\text{m}^\circ\text{C}^{-1/2} \text{ days}^{-1/2}$)
A	Accumulated degree days of thaw ($^\circ\text{C days}$)
$\varepsilon_{\text{InSAR}}$	Random errors in InSAR deformation measurements (m)
θ	Volumetric water content (–)
κ	Dielectric constant of soil (–)
κ_w	Dielectric constant of pure water (80)
κ_g	Dielectric constant of dry soil grain (–)
ν	Radar speed (m s^{-1} ; m ns^{-1} is used in this study)
c_0	The speed of light in free space ($2.9979 \times 10^8 \text{ m s}^{-1}$)
ε	Thaw strain (–)
ΔZ	Thaw settlement (m; cm is used in this study)
H	Depth of thaw penetration (m)
ρ_w	Density of water (1000 kg m^{-3})
ρ_i	Density of pure ice (917 kg m^{-3})

Seasonal thaw settlement at drained thermokarst lake basins, Arctic Alaska

L. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 1. (a) Google Earth image of SAC basin with the inset showing its location on the North Slope near Prudhoe Bay. The red line (“A” to “F”) shows the ground penetrating radar (GPR) transect across the basin and “RP” refers to the residual pond. (b) Elevation map with standing water body masked out in gray (source: Paine et al., 2013). (c) A photo of the basin center taken from point “A” looking east. (d) A photo of the residual pond.




Fig. 2. Interferograms networks produced in this study using ALOS PALSAR data from two paths. Circles represent PALSAR acquisitions and vertical grey bars represent the thaw season (June to September). Lines represent interferograms with sufficiently high coherence over SAC basin. Perpendicular baseline values are referenced to 13 June 2007 and 12 August 2006 for Paths 250 and 251, respectively.




Fig. 3. An example interferogram showing seasonal thaw settlement of DTLBs and surroundings. Color indicates the surface thaw settlement between 13 June 2007 and 13 September 2007, projected in the LOS direction, with the reference point located on the river floodplain (white cross). The white box marks SAC basin that underwent a strong settlement compared with the surrounding area. Solid circles outline DTLBs with stronger seasonal settlement than the surrounding tundra; and dashed circles outline DTLB with similar seasonal settlement as the surroundings. DTLBs smaller than 4 ha or having low interferometric coherence are not marked. The satellite flight and LOS directions are shown in the lower left.

Seasonal thaw settlement at drained thermokarst lake basins, Arctic Alaska

L. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Seasonal thaw settlement at drained thermokarst lake basins, Arctic Alaska

L. Liu et al.




Fig. 4. (a) Average seasonal thaw settlement for 2006–2010 and (b) relative uncertainties in SAC basin and the surrounding tundra area. The black line shows the basin boundary, defined by the shoreline prior to drainage. Low coherence areas are masked out in gray.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 5. Theoretical relation between the thaw strain and the volumetric water content when freezing and thawing of the soil occurs in a closed system. Thaw strain has an upper bound of 10%. The two boxes outline the ranges of thaw strains and volumetric water contents for the sandy margins and the center of the basin, respectively.

Seasonal thaw settlement at drained thermokarst lake basins, Arctic Alaska

L. Liu et al.




Fig. 6. (a) Active layer thickness along the Fig. 1a transect, estimated from in-situ (crosses) and GPR (dots). The uncertainties of the in-situ and GPR ALT are 5 cm and 6 cm, respectively, but are not shown. (b) Profile of the seasonal subsidence based on InSAR measurements, with the uncertainties shown in gray. (c) Profile of the thaw strain, with the uncertainties shown in gray. The horizontal dotted line marks the upper bound for a closed system.