

Interactive comment on “Tidally-induced velocity variations of the Beardmore Glacier, Antarctica, and their representation in satellite measurements of ice velocity” by O. J. Marsh et al.

O. J. Marsh et al.

oliver.marsh@pg.canterbury.ac.nz

Received and published: 15 July 2013

We thank the reviewer for their comments and have investigated the `t_tide` package for tidal analysis. Using this software we are able to identify tidal components with high signal to noise ratio at GPS-2 and GPS-3 but the signal is too small at GPS-1 and the data is too sparse at GPS-4. We have included a figure showing the phase of the tides at these two sites related to the CATS_2008opt tide model, similar to Figure 3 from Gudmundsson (2007) and showing the same general patterns as at the Rutford Ice Stream. As expected we could not solve for both MS_f and M_f with a high degree of confidence, although could solve each component independently. We have

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



restructured some sections in line with reviewer #1s comments and elaborated on our methods, particularly relating to the tidal analysis.

Interactive comment on The Cryosphere Discuss., 7, 1761, 2013.

TCD

7, C1082–C1084, 2013

Interactive
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



C1083




Fig. 1. Phase difference between measured horizontal displacements at GPS-2 and -3 and modelled vertical ocean tides from the CATS_2008opt tidal simulation (Padman et al., 2008).

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)