

Interactive comment on “Solving Richards Equation for snow improves snowpack meltwater runoff estimations” by N. Wever et al.

N. Wever et al.

nander.wever@slf.ch

Received and published: 13 December 2013

We thank Florent Dominé for his encouraging comment. We would like to shortly address both issues below.

Regarding specific surface area (SSA): currently, the SNOWPACK model is centered around the grain size as defined as the average of the maximum extend of snow particles in a layer (Fierz et al., 2009). There are activities underway to change the microstructural functions of SNOWPACK to reflect recent advances using SSA. However, SSA, or more precisely the optical diameter r_{opt} (Morin et al., 2013), is currently only calculated as a postprocessing step. That means that we cannot investigate properly and straightforwardly the impact of a change to r_{opt} on our results. However, once the snowpack is isothermal and wet, classical grain size and r_{opt} are equal. So the

C2727

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



most important effect may be in the early stages where melt water percolates through an initially dry snowpack. It should be noted however, that many experimental studies are forming the foundation of the parameterizations in SNOWPACK, so although SSA may allow a better simulation of microstructural processes, it is in our opinion an unanswered question right now whether this will allow a significantly better simulation of water flow in snow. We are therefore not planning to address this issue in the revised version as we cannot provide the reader with useful information regarding this topic.

Regarding the permeability calculations by Calonne et al. (2012): this question is related with the previous one. The used equivalent sphere radius in the Calonne et al. (2012) formulation of permeability roughly corresponds to r_{opt} . Thus we tested the effect of Calonne vs Shimizu, using r_{opt} in Calonne's parameterization. We found a marginal (not significant) improvement in terms of NSE coefficient and r^2 value, mainly for simulations for Col de Porte. We think that the reason for this marginal improvement is due to Shimizu's and Calonne's formula differing especially for the low density, new snow crystals. But when liquid water is involved, the snow will compact rather quickly and grains will only stay in this regime for a short period of time. Nevertheless, we will provide the results using Calonne (2012) in the revised manuscript.

Calonne, N., Geindreau, C., Flin, F., Morin, S., Lesaffre, B., Rolland du Roscoat, S., and Charrier, P.: 3-D image-based numerical computations of snow permeability: links to specific surface area, density, and microstructural anisotropy, *The Cryosphere*, 6, 939–951, doi:10.5194/tc-6-939-2012, 2012.

Fierz, C., Armstrong, R., Durand, Y., Etchevers, P., Greene, E., McClung, D., Nishimura, K., Satyawali, P., and Sokratov, S.: The International Classification for Seasonal Snow on the Ground (ICSSG), Tech. rep., IHP-VII Technical Documents in Hydrology No. 83, IACS Contribution No. 1, UNESCO-IHP, 2009.

Morin, S., Domine, F., Dufour, A., Lejeune, Y., Lesaffre, B., Willemet, J.-M., Carmagnola, C. M., and Jacobi, H.-W.: Measurements and modeling of the vertical profile

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

of specific surface area of an alpine snowpack, *Adv. Water Resour.*, 55, 111-120, <http://dx.doi.org/10.1016/j.advwatres.2012.01.010>, 2013.

Interactive comment on *The Cryosphere Discuss.*, 7, 2373, 2013.

TCD

7, C2727–C2729, 2013

Interactive
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

C2729

