

Summary of Comments on tcd-7-4813-2013-my_review.pdf

Page: 1

Number: 1 Author: Subject: Sticky Note Date: 02/01/2014 7:34:54 PM

General comment:

I was initially looking forward to read this manuscript since the work on method/model intercomparison can be insightful (as is proven to be in climatological community, e.g CMIP3, CMIP5). This work aimed to provide a comparison of different methods for assessing glacier volume over large region (Himalaya-Karakoram). The motivation and methodology is clear and well described. Unfortunately, in my opinion, this is the only positive aspect of the manuscript. The quality of presentation of results and especially their discussion progressively goes down as the manuscript reaches conclusion. At the end we are left with the study that (1) displays a bias (the authors openly oppose the volume-area scaling while even their very own results do not provide evidence for it), (2) fails to evaluate and present uncertainties of each method they use (and therefore strengthen their intercomparison method), and (3) does not provide any new insight (the pros and cons of each method have been already discussed in more detail in the previous publications to which the authors refer). Although I can accept that it is hard to provide novelty in each new scientific publication, the authors should at least seriously consider addressing the first two points (lack of scientific objectivity and adequate error analysis) before resubmitting the manuscript.

ing debates in media (Cogley et al., 2010; IPCC, 2010; Schiermeier, 2010) emphasized both the concern as well as the poor understanding of glaciers and glacier changes in this region. Recent studies focusing on mass changes of this region revealed the complex and regionally heterogeneous behavior of HK glaciers (Fujita and Nuimura, 2011; Bolch et al., 2012; Gardelle et al., 2012). Furthermore, different findings based on various methodologies underline the intricacy of understanding complex processes in such a large region (Jacob et al., 2012; Kääb et al., 2012; Gardner et al., 2013). In this study we focus on determining the ice volume for the HK region, which is a basic parameter required for estimations of future glacier evolution (e.g., Le Meur et al., 2007), runoff projections (Huss et al., 2008), and the basis for sea-level rise projections.

Various approaches are in use to estimate glacier volumes, such as volume–area (V–A) relations (e.g., Chen and Ohmura, 1990; Bahr et al., 1997); slope-dependent ice-thickness estimations (Haeberli and Hoelzle, 1995); and more recently, a variety of spatially distributed ice-thickness models have been presented (e.g., Clarke et al., 2009; Farinotti et al., 2009; Huss and Farinotti, 2012; Li et al., 2012; Paul and Linsbauer, 2012). Such knowledge of the ice-thickness distribution is a requirement for state-of-the-art physically-based models to quantify the contribution of glaciers to runoff and sea level rise.

Until recently, estimates of global glacier volumes had to rely on extrapolations of glacier size distributions from regions with good data coverage to regions where glacier data is sparse (Meier and Bahr, 1996; Raper and Braithwaite, 2005; Radić and Hock, 2010). Only with the publication of the Randolph Glacier Inventory (RGI) (Arendt et al., 2012) a globally complete dataset of glacier coverage has become available and can now be used for assessing glacier volumes without relying on data extrapolation. Volume estimations of all glaciers and ice caps of the world based on the RGI using V–A relations are given by Marzeion et al. (2012), Grinsted (2013), and Radić et al. (2013); Huss and Farinotti (2012) used the RGI to apply a model to calculate spatially distributed ice thicknesses.

4815

Available volume estimates for the HK region exhibit a large variety; but deviating definitions and delineations of regions and different regional grouping hamper comparability. Bolch et al. (2012), using the same glacier inventory dataset as the present study, highlight that estimations of ice volumes in the Himalayas are highly uncertain and range from $\sim 2300 \text{ km}^3$ to $\sim 6500 \text{ km}^3$. The IPCC AR4 even reports the total ice volume of the HK region to be $12\,000 \text{ km}^3$ (Cruz et al., 2007). Ohmura (2009) gives a value of $3800\text{--}4850 \text{ km}^3$ for Pakistan, India, Nepal and Bhutan; Cogley (2011) estimates the mass of all glaciers in the HK to be $3600\text{--}7200 \text{ km}^3$, depending on the chosen glacier inventory. Huss and Farinotti (2012) calculate a volume of glaciers in South Asia East and South Asia West of $4552 \pm 239 \text{ km}^3$, whereas for the same region Marzeion et al. (2012) obtained a volume of $5350 \pm 282 \text{ km}^3$, Grinsted (2013) of 6017 km^3 , and Radić et al. (2013) of $5595\text{--}6327 \text{ km}^3$. Currently we lack a more systematic analysis and a comparison of ice-volume estimation methods based on a consistent data basis.

Here, we present ice volume estimations for the HK region using three different V–A relations, a slope-dependent ice thickness estimation method, and two ice-thickness distribution models. As input data, the digital elevation model (DEM) from the Shuttle Radar Topography Mission (SRTM) is used, in combination with the most recent and to our knowledge most accurate glacier outline dataset available (Bolch et al., 2012). Instead of developing a new volume estimation methodology, we compare results of the different existing approaches, investigate the influence of different uncertainty factors, and discuss the potentials and limitations of the different methods.

2 Study region and data

In the past, comparisons of ice volumes calculated for the HK region were hindered by the use of different definitions of the Karakoram and the Himalayas (Gurung, 1999; Shroder, 2011). Here we divide the HK region into four sub-regions, i.e. Karakoram, Western, Central, and Eastern Himalayas. Definitions of sub-regions extents and subdivisions are given in Bolch et al. (2012) (see also Fig. 1); Cogley (2011) used the

4816

Number: 1 Author: Subject: Sticky Note Date: 02/01/2014 7:39:42 PM

Also Clarke et al 2012, using inverse modeling.

Clarke G. K. C., F. S. Anslow, A. H. Jarosch, V. Radić, B. Menounos, T. Bolch and E. Berthier (2012) Ice volume and subglacial topography for western Canadian glaciers from mass balance fields, thinning rates, and a bed stress model. *J. Climate*, e-View, doi: 10.1175/JCLI-D-12-00513.1

same study region and the identical separation between Karakoram and Himalayas, but didn't further divide the Himalayas.

The glacier outlines used for the calculations have been compiled by remote sensing, based on satellite scenes acquired between 2000 and 2010, and are as well the same than used by Bolch et al. (2012). Glacier outlines were mapped by the International Centre for Integrated Mountain Development (ICIMOD) for the southern part of the Karakoram range, parts of the Indian Himalayas (eastern Uttarakhand and Sikkim), Nepal, and Bhutan (Bajracharya and Shresta, 2011). For the western Himalayas, outlines from Frey et al. (2012), and for the Shyok basin (eastern Karakoram) from Bhambri et al. (2013), are used (Fig. 1). If no alternative datasets were available, outlines for Chinese glaciers were taken from the Chinese glacier inventory (Shi et al., 2009), as available from the GLIMS database. For many regions, these outlines are also contained by the RGI in most parts of the Karakoram, as well as in the central and eastern Himalayas (outside Nepal and China) they are more recent and – based on visual inspections – more accurate than the outlines contained by the RGI, as in these regions data from ICIMOD is used, which is not publicly available and therefore not represented in the RGI.

Required topographic parameters such as minimum and maximum elevation as well as mean slope were obtained by fusing the outlines with the void-filled SRTM version from CGIAR (Farr et al., 2007; Jarvis et al., 2008; Frey and Paul, 2012). Table 1 gives an overview of total glacier areas and mean glacier elevations per region.

3 Methods

In the following, the different ice volume estimation approaches applied in this study are described. For the existing methodologies the descriptions are restricted to short summaries, whereas the presentation of the two ice-thickness distribution models are more extensive, including newly developed modifications.

4817

3.1 Volume-area scaling

V–A scaling has been the most frequently used approach for ice volume estimations so far. Ice volume is calculated as a function of area, as large glaciers tend to have large volumes. The reason of the popularity of V–A scaling is its simple and rapid application; once the scaling parameters are determined, glacier volumes can be easily calculated for all glaciers with a known area. Equation (1) shows the general form of V–A scaling:

$$V = cA^\gamma \quad (1)$$

with V representing the glacier volume, A the glacier area and c and γ the two scaling parameters.

Here we apply three sets of scaling parameters which have been used by Cogley (2011) for the same study region: (i) Chen and Ohmura (1990), who used a set of 63 glaciers where volumes were interpolated based on radio-echo soundings to determine the related scaling parameters; (ii) Bahr et al. (1997), who confirmed these parameter values in a theoretical study; and (iii) Arendt et al. (2006) who derived their parameter set based on non-tidewater glaciers in the Western Chugach Mountains, Alaska. The applied scaling parameters are given in Table 2. Comparing our results with results from Cogley (2011) allows examining the influence of the different glacier inventory data used.

3.2 Slope-dependent thickness estimations

Haeberli and Hoelzle (1995) presented an approach for estimating glacier volume based on the average surface slope. As this parameterization scheme has been designed to estimate glaciological parameters with detailed tabular glacier inventory data, it can be directly applied using modern glacier inventories, which are available for many regions in the world (cf. the Global Land Ice Measurements from Space initiative GLIMS; e.g., Raup et al., 2007) or the Randolph Glacier Inventory (RGI; Arendt

4818

Number: 1 Author: Subject: Sticky Note Date: 02/01/2014 4:11:44 PM

Please include the version of RGI (e.g. 2.0 or 3.0). The idea behind RGI is that it will be constantly updated as new and updated data become available. RGI 3.0 already differs from RGI 2.0.

Also note that there is a paper by Pfeffer et al 'The Randolph Glacier Inventory: a globally complete inventory of glaciers' submitted to J Glaciol and close to be in press.

et al., 2012). An application of this approach to modern glacier data can be found, for instance, in Hoelzle et al. (2007) or Salzmann et al. (2013).

Haeberli and Hoelzle (1995) calculate the volume V based on the following equation:

$$V = Ah_f \quad (2)$$

5 where A is the glacier area and h_f the mean ice thickness. h_f is defined as

$$h_f = \frac{\tau}{f\rho g \sin(\alpha)} \quad (3)$$

where τ is the average basal shear stress along the central flowline; f a shape factor (chosen constant as 0.8); g the gravitational acceleration; and α the surface slope. To account for the extrapolation from the mean ice-thickness along the central flowline (10 h_f) to the mean ice-thickness of the entire glacier (h_F), in accordance with the assumption of a semi-elliptic cross sectional geometry, a multiplication with $(\pi/4)$ is added to Eq. (3):

$$h_F = h_f \left(\frac{\pi}{4} \right) \quad (4)$$

The parameterization of τ with the elevation range ΔH is based on reconstructed late-15 Pleistocene Alpine glaciers (cf. Fig. 1 in Haeberli and Hoelzle, 1995):

$$\tau[\text{bar}] = \begin{cases} 0.005 + 1.598\Delta H - 0.435\Delta H^2 & \text{for } \Delta H \leq 1600 \text{ m} \\ 1.5 & \text{for } \Delta H > 1600 \text{ m} \end{cases} \quad (5)$$

When applied to modern remotely sensed glacier inventories, a challenge arises from 20 the definition of the surface slope α : glacier length normally is included in older tabular glacier inventories, such as the World Glacier Inventory (WGI), which was used by Haeberli and Hoelzle (1995). The surface slope α can then be calculated with the glacier length l and the elevation range ΔH with:

$$\alpha = \arctan \left(\frac{\Delta H}{l} \right) \quad (6)$$

4819

By using zonal statistics in combination with a DEM, mean slope can also be directly derived for each glacier without knowing its length. However, mean surface slope derived from ΔH and l (α_l) is different – generally smaller – than mean surface slope averaged over all DEM cells of a glacier (α_{DEM}). In order to determine a correction factor for α_{DEM} , glacier length has been manually determined for 50 glaciers of the study 5 region, including the 10 largest glaciers. Resulting α_l values have then been compared to the α_{DEM} values (Fig. 2). Differences between α_{DEM} and α_l appear to be nearly constant for different glacier size classes. Based on this comparison, the correction factors to be applied to α_{DEM} have been determined as -10° , -5° , and -2.5° for glaciers with 10 an area $> 20 \text{ km}^2$, $5\text{--}20 \text{ km}^2$, $< 5 \text{ km}^2$, respectively. The standard deviations for these slope differences are 2.51° , 4.16° , and 2.02° for glaciers with areas $> 20 \text{ km}^2$, $5\text{--}20 \text{ km}^2$, and $< 5 \text{ km}^2$, respectively; and mean differences of the three size classes are all significantly different from each other and from 0 (at a 95 % confidence level).

3.3 Modeling of ice-thickness distributions

15 Based on the approach of Haeberli and Hoelzle (1995), Paul and Linsbauer (2012) developed a model for assessing the spatial distribution of ice thickness by estimating the glacier depths at several points along so-called glacier branch lines and interpolating between these points and the glacier margins. Farinotti et al. (2009) proposed an alternative approach to calculate glacier ice thickness distribution that is based on the 20 principles of ice flow dynamics. Local ice thickness is inverted from surface topography by calculating ice balance fluxes through cross profiles along the glacier and applying Glen's (1955) flow law.

A major disadvantage of these spatially distributed approaches is the time-consuming preparation of the input-data, as central flow lines (Linsbauer et al., 2012) 25 or catchment areas for each glacier branch (Farinotti et al., 2009) need to be digitized manually. Hence, we present a modified version of the approach by Paul and Linsbauer (2012), where the ice-thickness is estimated at randomly selected points on the glaciers (cf. Sect. 3.3.1). For similar reasons Huss and Farinotti (2012) further devel-

4820

oped the method by Farinotti et al. (2009) and applied it to all glaciers worldwide. The only input requirements for both approaches are glacier outline data and a DEM.

3.3.1 GlabTop2

The first approach to calculate distributed glacier thickness applied here is almost identical to the GlabTop model (Linsbauer et al., 2012; Paul and Linsbauer, 2012) but circumvents the laborious process of manually drawing branch lines. Instead, ice thickness is calculated for an automated selection of randomly picked DEM cells within the glacierized areas. Ice thickness distribution for all glacier cells is then interpolated from the ice thickness at the random cells and the glacier margins where ice thickness is known to be zero. The calculation of ice thickness is grid-based and requires a DEM and the glacier mask as input. In a first step, all groups of glaciers sharing common borders, i.e. glacier complexes, are assigned a unified ID. All following steps are performed for one ID (i.e. all cells of a glacier complex) at a time, disregarding all cells of differing IDs. Figure 3 schematically illustrates the model. A second mask is generated where a code is assigned to all non-glacier cells directly adjacent to the glacier cells (called glacier-adjacent cells). A different code is assigned to all glacier cells being located directly at the glacier margin (called marginal glacier cells). From the remaining glacier cells (inner glacier cells) a number of random cells are drawn whereas their number corresponds to a predefined percentage (r) of the inner glacier cells. An initial buffer of 3×3 cells is then laid around each random cell and each individual buffer is enlarged until the difference in elevation between the lowest and the highest DEM cell falling within the buffer is equal or greater h_{\min} . Thereby all glacier cells in the buffer (marginal and non-marginal) are considered. This procedure avoids very small slope values with corresponding extremely high ice thicknesses and thus makes a slope cut-off (i.e. a minimum local slope considered) obsolete. The mean surface slope of all glacier cells in the buffer is used to calculate local ice thickness according to Eq. (3) and the result is assigned to the corresponding random cell. From ice thickness at all random cells and an ice thickness value h_{ga} assigned to all glacier adjacent cells ice


4821

thickness is interpolated to all glacier cells using Inverse Distance Weighting (IDW). The ice thickness calculation for each ID is repeated n -times with different sets of random points and the n ice thickness distributions are averaged into a final estimate of ice thickness distribution.

The ice-thickness calculation with GlabTop2 requires estimating the parameters τ and f as well as the model parameters h_{\min} , r , h_{ga} and n . The chosen values and their influence on the output is explained in the following. Parameters from Eq. (3) are set identical to Linsbauer et al. (2012) and Haeberli and Hoelzle (1995): for all glaciers f is set to 0.8 and τ is calculated as a function of vertical glacier extent (ΔH) according to Eq. (5). With the approach from Huss and Farinotti (2012) (see below), these parameters can be calculated for each glacier individually, which in turn can be used for a comparison.

Linsbauer et al. (2012) calculate the mean slope α over 50 m elevation intervals along the branch lines to ensure averaging of α over a reference distance of approximately one order of magnitude larger than local ice thickness (cf. Kamb and Echelmeyer, 1986). Consequently $h_{\min} = 50$ m was applied here as well. The parameters r and h_{ga} govern the shape of the glacier cross section. The latter has been measured only for a few glaciers but the cross sections of formerly glaciated valleys are generally described by a parabolic function (e.g., Graf, 1970). In the original model by Linsbauer et al. (2012) the modeled cross section is controlled by deciding manually how many branch lines run parallel and where to draw them (cf. Paul and Linsbauer, 2012). Here, random points are chosen automatically and their number and spatial distribution across the glacier-width determine the modeled cross-sectional profile of ice thickness. A high density of random points r reduces the influence of the glacier adjacent cells (set to ice thickness h_{ga}) in the interpolation and vice versa. Thereby the cross-section also changes from more v-shaped at low r to a parabolic profile with very steep sidewalls at high r . Both h_{ga} and r were calibrated by adjusting calculated cross sections to theoretical parabolic glacier cross sections and set to $h_{ga} = 15$ m and $r = 0.3$ (i.e., in each model run, ice thickness is calculated for 30 % of all inner glacier

4822

 Number: 1 Author: Subject: Sticky Note Date: 02/01/2014 4:12:35 PM
How do you make sure that the sensitivity tests among different methods are comparable?

 Number: 2 Author: Subject: Sticky Note Date: 02/01/2014 2:41:41 PM
for each individual glacier?

used for V–A calculations were increased and decreased by +5% and –5% for the sensitivity analysis.

For sensitivity tests of both the slope-dependent thickness estimations and the GlabTop2 model, the two scaling parameters f and τ were modified. f was altered by ± 0.1 . Two alternative parameterizations of τ with DH were performed; a high shear-stress version with an upper limit for τ_{\max} (i.e., the basal shear-stress for all glaciers with $\Delta H > 1600$ m) of 1.8 bar, and a low shear-stress version with a lower limit with $\tau_{\max} = 1.2$ bar. The parameterization of τ for glaciers with $\Delta H \leq 1600$ m have been adapted accordingly (Fig. 4). For uncertainty and sensitivity analyses of the HF model, see Huss and Farinotti (2012).

3.5 Evaluation and validation

Since direct measurement of glacier volume is not possible, glacier volumes or mean glacier thicknesses obtained from V–A approaches and slope-dependent thickness estimations can only be evaluated on an imperfect basis. Reference values build on the interpretation of ground penetrating radar (GPR) measurements and interpolations between measured profiles. Considering the uncertainties related to interpretation and analysis of GPR data, the comparison of estimated vs. “measured” total glacier volumes is rather an evaluation than a validation. However, to our knowledge only two published measurement-based estimates of total glacier volumes exist in the entire HK region (Gergan et al. (1999) for Dokriani and Ma et al. (2010) for Kangwure), hence it was not possible to perform such an evaluation. Furthermore, V–A relations are designed to estimate the volume of a larger glacier ensemble (Farinotti and Huss, 2013), but are not suitable to assess the volume of individual glaciers, which further hampers their comparison with measurements.

Results from spatially distributed ice-thickness models, in contrast, can be directly validated with local ice-thickness measurements from GPR profiles. This is a major advantage and reduces the uncertainties of the reference values. In the entire HK region, however, only a very limited number of ice-thickness measurements are available

4825

(Bolch et al., 2012). To evaluate our estimates, we compared the results of the ice-thickness distribution model to published point measurements of ice thicknesses at one glacier in the Karakoram (Baltoro, measured by Marussi, 1964), one in the western Himalayas (Chhota Shigri, Azam et al., 2012), and four glaciers in the central Himalayas (Dokriani, Gergan et al., 1999; Kangwure, Ma et al., 2010; and Khumbu and Lirung, both from Gades et al., 2000). For the eastern Himalayas no ice-thickness measurements suitable for such a validation were found in the literature.

4 Results

Volumes were calculated for all 28 100 glaciers of the glacier dataset using six approaches: three V–A relations, the adapted approach from Haeberli and Hoelzle (1995), and two ice-thickness distribution models. Figure 5 shows the total volume as well as resulting ice volumes for all sub-regions. The calculated volumes for the entire HK region vary between 2955 and 6455 km³, depending on the method applied. By far the largest glacier volumes exist in the Karakoram region, amounting to about 50–60% of the total ice volume in the HK region. The smallest volumes are found in the Eastern Himalaya (194–408 km³), while the Western and Central Himalayas show comparable glacier volumes (504–1128 km³ each). All V–A approaches lead to larger volumes in all sub-regions than the other three approaches. The largest differences are found between the V–A relation using the scaling parameters of Arendt et al. (2006), and the slope-dependent thickness estimation and the two ice-thickness distribution models, with differences of up to 118% for the entire HK region and up to 139% for the Karakoram.

The slope-dependent thickness estimation and the two ice-thickness distribution models, however, lead to comparable results, with total absolute differences of < 13%. For the sub-regions, largest deviations are observed for the Karakoram, where the largest ice volumes are stored and which contains the largest individual glaciers of the region. Figure 6 shows the modeled ice thickness distribution of Chhota Shigri in

4826

Number: 1 Author: Subject: Sticky Note Date: 02/01/2014 2:36:13 PM
what dictated this choice?

Number: 2 Author: Subject: Sticky Note Date: 02/01/2014 4:13:42 PM
Is this because Arendt et al used parameters best suited for Alaskan glaciers? After all, $c=0.28$, which is much larger (by 40%) than c in the other two V-A scaling methods.

Number: 3 Author: Subject: Sticky Note Date: 02/01/2014 4:15:11 PM
Am not sure if the % is best way to show the differences here. % for a small region means much less than % for a large (large ice volume). If large region is chopped into several small regions your % would have completely different meaning. Can you report the errors in SLE as well?

Number: 1 Author: Subject: Sticky Note Date: 02/01/2014 4:16:49 PM

Radic & Hock 2010, or better to say Radic et al 2013 used parameter c from Chen and Ohmura 1990. Note that your estimate with these parameters are the closest to the estimates from other methods you used. Also SLE estimates are not using those studies (Raper, Radi 2010) any more since those studies suffered from incomplete glacier inventories. See IPCC AR5 for the current studies used for SLE. Please do not mix tools (V-A scaling) with assessments of SLE, since it is much more in the assessment (e.g. data) than just the tool that can cause the uncertainties.

Number: 2 Author: Subject: Sticky Note Date: 02/01/2014 4:17:35 PM

I do not understand why you use Arendt et al as a representative for V-A scaling large bias. This is your upper band estimate since c is 40% larger than c in other methods. Please mention this somewhere in the text because it seems to me that the authors do not point out the obvious.

(2006), and to 7.4 mm for GlabTop2 (assuming a density of 900 kg m^{-3} for the entire glacier volume and a total ocean area of $362 \text{ km} \times 106 \text{ km}$). It is thus likely that the global total potential contribution of glaciers and ice caps to sea-level rise might be smaller than previously assumed, as was also found by Huss and Farinotti (2012) at the global scale. Potential sources for this difference are (i) the input data, i.e. the glacier inventory used for the calculation; (ii) conceptual aspects of the different methodologies; and (iii) the values used for the required model parameters. These points are addressed in the following.

5.1 Input data

The glacier inventory used for this study is the most detailed dataset currently available for this region. Previous studies had to rely on extrapolation of glacier size classes (Raper and Braithwaite, 2005; Radić and Hock, 2010) while others used glacier data of older date (e.g., Cogley, 2011). In many of these cases glacier accumulation areas were too large, especially if the inventories are based on the digitalization of topographic maps. Smaller glacier areas in the new glacier inventory (and resulting lower ice volume estimates) are thus rather due to improved mapping accuracy than real glacier shrinkage (Bolch et al., 2012). The separation of individual glacier entities has as well a large influence on volume estimates, in particular on results from V–A relations (see conceptual aspects below).

The quality of the glacier inventory input data is also of major importance for potential ice volume estimations in other regions of the world. The RGI, the most recent global-level inventory for instance, is of heterogeneous quality (Arendt et al., 2012) and especially the separation of glacier complexes into individual glacier entities is still a challenge and source of errors in some regions (e.g., Huss and Farinotti, 2012; Grinsted, 2013). Nevertheless, extrapolations of glacier size classes from one region to another are not required anymore.

4829

5.2 Conceptual aspects

Estimating a glacier's volume based on its area has three major shortcomings: (i) mean ice thickness and area have a weak correlation. Plotting the two variables against each other, a scatter of about one order of magnitude is evident (Fig. 8). The high correlation of volume and area comes from the self-correlation in the relationship of area and volume, as volume is the product of area and mean thickness. The correlation of glacier area and thickness is rather weak (cf. Fig. 8), and modified by the surface slope of the glacier. (ii) Large uncertainties are introduced as (a) the “measured” volumes of the glaciers are based on the interpolation of GPR measurements, which in general are biased towards flat and crevasse-free glacier parts (Binder et al., 2009; Fischer, 2009); and (b) the scaling parameters are determined on the basis of only a few hundred glaciers with measurements at most. There is a variety of possible combinations for the scaling parameters c and d , regardless whether they are calculated with a set of measured or idealized glaciers (e.g., Adhikari and Marshall, 2012; Grinsted, 2013; Farinotti and Huss, 2013). And (iv) due to non-linearity of the scaling relationships, the separation of glacier complexes into individual glaciers has major impact on resulting ice volumes: in the extreme case of a glacier complex not separated at all, an overestimation of the volume by 70–80 % can result (Huss and Farinotti, 2012; Grinsted, 2013).

5.3 Model parameters

Besides modeled regional glacier volumes, Huss and Farinotti (2012) also provide region-specific thickness-area scaling parameters according to the best fit to their calculated glacier volumes (see their Fig. 7). Applied to our inventory dataset, their thickness-area relations lead to considerably smaller ice volumes, which are comparable to our results from the slope-dependent thickness estimations and GlabTop2. The total volume for the HK region based on these scaling parameters from Huss and Farinotti (2012) is 3113 km^3 (1935 km^3 and 479 km^3 for the Karakoram and Western

4830

-
- Number: 1 Author: Subject: Sticky Note Date: 02/01/2014 2:37:04 PM
superscript
-
- Number: 2 Author: Subject: Sticky Note Date: 02/01/2014 7:45:08 PM
so basically this result was already found and there is no novelty in this study at this point.
-
- Number: 3 Author: Subject: Sticky Note Date: 02/01/2014 7:46:27 PM
But isn't the same data used for each method in this study? Why don't you discuss in detail the results from your study rather than generality stating what we know from before (i.e. that the input data differs).
-
- Number: 4 Author: Subject: Sticky Note Date: 02/01/2014 4:19:35 PM
This section is irrelevant for this study since the whole point in the study is to use the same data (inventory) and apply different tools.
-
- Number: 5 Author: Subject: Sticky Note Date: 02/01/2014 2:37:46 PM
except for evaluation of statistical methods which proved to be useful.
-
- Number: 6 Author: Subject: Sticky Note Date: 02/01/2014 2:37:56 PM
Why is only conceptual aspect of V-A scaling discussed here? All other methods are also based on concepts.
-
- Number: 7 Author: Subject: Sticky Note Date: 02/01/2014 4:20:37 PM
Isn't the volume-area relation what you are explaining here, not ice thickness and area relation?
-
- Number: 8 Author: Subject: Sticky Note Date: 02/01/2014 4:21:48 PM
This argument is flawed (see e.g. Lüthi et al 2008). Also, rather than trying to reinvent the wheel here (i.e. finding theoretical basis for V-A scaling in order to identify misconceptions) why not properly read Bahr et al (1998) who correctly points out the pros and cons of the scaling in the conceptual light (e.g. each closure in the scaling analysis is discussed).
-
- Number: 9 Author: Subject: Sticky Note Date: 02/01/2014 4:22:48 PM
Scaling parameters are also determined theoretically (using scaling analysis of mass continuity equation and ice rheology). Please read publications of Bahr from late 1990ies.
-
- Number: 10 Author: Subject: Sticky Note Date: 02/01/2014 4:23:18 PM
Wrong for gamma. Gamma is well constrained parameter (only differs between glacier and ice cap). Please read Bahr papers.
-
- Number: 11 Author: Subject: Sticky Note Date: 02/01/2014 4:23:57 PM
It would make sense that the authors performed this sensitivity test (e.g. delineating the glacier complexes and not delineating, and delineating in different ways) to quantify the uncertainty of this factor. Especially since the effect can be large. This sensitivity test would make one valuable contribution to this study that so far has not much new to offer.

Himalayas, respectively, calculated with the parameters for the “South Asia West” region; and 511 km^3 and 188 km^3 for the Central and Eastern Himalayas, respectively, calculated with the parameters for “South Asia East”). This corroborates the hypothesis that the overestimations of the V–A approaches are mainly caused by the lack of measurement-based volume estimates of glaciers in the HK region, which is obvious as well from the large volumes resulting from the scaling parameters of Arendt et al. (2006), which were calibrated with (maritime) Alaskan glaciers (Table 5, volume estimates are given following the V–A approaches from Marzeion et al. (2012), Grinsted (2013), and Radić et al. (2013), applied to the glacier inventory used in this study. Results following Marzeion et al. (2012) are virtually identical to the results according to Bahr et al. (1997), as their approach was directly used. Results obtained by the V–A approach of Grinsted (2013) correspond for most sub-regions to the lower-bound range of the three V–A approaches applied here (between the volumes following Chen and Ohmura (1990) and Bahr et al. (1997), except for the Karakoram, where the volume estimate is in the range of the results from the slope-dependent ice-thickness estimations and the ice-thickness distribution models. Results obtained by the approach from Radić et al. (2013) exceed the other estimates and are only surpassed by the estimates following Arendt et al. (2006).

For the slope-dependent thickness estimations following Haeberli and Hoelzle (1995), here, mean slope had to be obtained in a different manner than originally proposed, as glacier length is not available from the glacier data set used here. Glacier length information is, however, missing in many modern glacier inventories (Paul et al., 2009), due to the lack of a suitable methodology that automatically determines glacier length without manual corrections (e.g., Le Bris and Paul, 2013). If α_{DEM} (the slope from the DEM averaged per glacier) is directly used for Eq. (4) without the suggested correction (see Sect. 3.2 and Fig. 2), the resulting total ice volume is only 49 % of the volume calculated with corrected slope values (55 % of the GlabTop2 estimate), since alpha DEM is larger than the average slope along the central flowline which has to be used in the parameterization of Haeberli and Hoelzle (1995). The slope-dependent

4831

ice thickness estimations by Bolch et al. (2012) are considerably smaller (total volume of the HK region 2330 km^3), as different correction factors had been applied, which were based on fewer measurements. Obviously, the corrections applied here are also a generalization and thus introduce considerable uncertainty. Hence, we assume that results from this approach could be improved, if (automatically derived) glacier length values were available for all glaciers in the inventory.

Results of both the slope-dependent thickness estimations and GlabTop2 strongly depend on the parameterization of the average basal shear stress τ , in particular the upper limit of τ , which is used for glaciers with elevation extents of more than 1.6 km. As the HF-model calculates the average basal shear stress for each glacier, these τ -values can be used for an independent comparison to the τ -parameterization used for GlabTop2 and the slope-dependent thickness estimation: averaged by the number of glaciers, τ -values used for GlabTop2 are 7.7 % lower than as derived using the HF method, but if weighted by the glacier area 4.1 % higher than in the HF-model. In Fig. 9 τ -values from the HF-model are plotted for each region.

According to this comparison, the parameterization of Haeberli and Hoelzle (1995) overestimates τ for HK glaciers with ΔH between about 0.5 km and 2 km, but slightly underestimates it for glaciers with $\Delta H > 2.5 \text{ km}$. The values of τ_{max} used for the uncertainty assessment cover almost all τ -values as calculated with the HF-method, in particular those of glaciers larger than 100 km^2 . However, an independent assessment of the plausibility of average basal shear-stress values is not possible without a representative set of measured glaciers. Taking these considerations into account together with the validation, we conclude that the results from the GlabTop2 model runs with the modified parameters ($\tau_{\text{max}} = 1.8 \text{ bar}$ and 1.2 bar , and $f = 0.7$ and 0.9 , respectively) constitute upper and lower bound estimates (Table 4). On the other hand it also indicates that calculated volumes can exhibit relatively large differences on the level of individual glaciers. Since the same equation (Eq. 3) is used for the slope-dependent ice thickness estimations, these findings apply as well to this approach.

4832

Number: 1 Author: Subject: Sticky Note Date: 02/01/2014 4:25:33 PM

It would be worth providing the full range of estimates from V-A scaling (as you start doing it here), instead on focusing rigidly on the selected three methods one of which is an outlier (Arendt et a) for this region. There is no reason to stick to the outlier just because initially you thought it would be good to use it. But not only do you stick with the outlier (which I can understand) but you use the outlier to bring your main conclusions about the V-A scaling overestimated results. So the study is biased and lacks scientific objectivity.

Number: 2 Author: Subject: Sticky Note Date: 02/01/2014 4:29:46 PM

Do these scaling approaches, in addition to different data sets and glacier delineation method, differ only in the parameters c and γ ? If yes, it would be useful to specify the scaling parameters for each estimate (add in Table 2). It seems that all what you are restating here is that the results will depend on the choice of c and γ . Again, this is not new. It would be much better to provide systematic sensitivity tests for each method you use and provide a full range of results (and present them in Figure 5).

Number: 3 Author: Subject: Sticky Note Date: 02/01/2014 2:39:14 PM

Which should have been quantified. Again provide the uncertainty range for each method.

6 Conclusions and perspectives

Glacier volumes were calculated for all glaciers in the HK region, based on recent and up-to-date dataset of glacier outlines, containing more than 28 100 glaciers covering an area of 40 775 km², using three V–A scaling relationships, a slope-dependent mean ice thickness estimation approach, and two models for assessing the ice-thickness distribution. Results from the latter three approaches are in good agreement and indicate a total ice volume of 2955–3360 km³ (1683–2122 km³ in the Karakoram, 504–543 km³ in the W-Himalayas, 512–560 km³ in the C-Himalayas, and 194–215 km³ in the E-Himalayas), corresponding to a mean ice thickness of 72.5–82.5 m. These volumes are lower than estimates based on V–A relations, which are by average 50 % larger (ranging from 15 % to over 100 %, depending on the scaling parameters). These differences are caused by the lack of measurement-based volume estimates of HK glacier as well as by conceptual aspects of the different methods. The results are important in the context of improved estimates of water stored in the Himalaya region, both in view of fresh water resources and sea level rise.

Comparisons of ice-thickness measurements with modeled ice thicknesses in general showed good agreement, and thus increase the confidence in the results from the distributed ice-thickness models. Although they require more computational effort, approaches that take the three-dimensional shape of the glacier surface into account are considered as superior, since they are based on ice-mechanical and ice-dynamical considerations on the one hand, but also because they offer the possibility for validation with ice-thickness measurements at points or in GPR profiles. Furthermore, knowledge about ice-thickness distribution is important for several other fields of glaciology, hydrology or natural hazards. Due to the good agreement with local ice-thickness measurements, we have higher confidence in the results of ice-thickness distributions models such as GlabTop2 and the HF-method. Looking at these advantages and the numerous potential for further applications, we recommend using distributed ice-thickness modeling approaches for ice volume estimations ranging from a single-glacier up to

4833

the global scale, as they solely require digital glacier outlines and terrain information, which today are both globally available. Previous estimates of regional or global ice volumes, and sea level equivalent may result in systematic deviations of up to 100 % or even more, and thus may need thorough revision.

To further validate and improve ice-thickness distribution models, more ice-thickness measurement, for instance from GPR is needed, in particular in regions with sparse data coverage, such as the HK region (Bolch et al., 2012). New technologies, such as airborne GPR (Blindow et al., 2011) offer new possibilities for such measurements, which are less biased towards flat and easy accessible glacier parts and, hence, can serve as a basis for improved estimations of total glacier volumes as well.

Acknowledgements. This study was funded by the EU FP7 project HighNoon and by the Swiss Agency for Development and Cooperation (SDC) Indian Himalaya Climate Adaptation Programme (IH-CAP).

References

- Adhikari, S. and Marshall, S. J.: Glacier volume–area relation for high-order mechanics and transient glacier states, *Geophys. Res. Lett.*, 39, L16505, doi:10.1029/2012GL052712, 2012. 4830
- Arendt, A. A., Echelmeyer, K., Harrison, W., Lingle, C., Zirnheld, S., Valentine, V., Ritchie, B., and Druckenmiller, M.: Updated estimates of glacier volume changes in the western Chugach Mountains, Alaska, and a comparison of regional extrapolation methods, *J. Geophys. Res.*, 111, F03019, doi:10.1029/2005JF000436, 2006. 4818, 4824, 4826, 4828, 4831, 4842, 4843
- Arendt, A., Bolch, T., Cogley, J. G., et al.: Randolph Glacier inventory: a dataset of global glacier outlines, Version: 2.0, Tech. rep., GLIMS, available at: http://www.glims.org/RGI/RGI_Tech_Report_V2.0.pdf, 2012. 4815, 4818, 4829
- Azam, M. F., Wagnon, P., Ramanathan, A., Vincent, C., Sharma, P., Arnaud, Y., Linda, A., Pottakkal, J. G., Chevallier, P., Singh, V. B., and Berthier, E.: From balance to imbalance: a shift in the dynamic behaviour of Chhota Shigri glacier, western Himalaya, India, *J. Glaciol.*, 58, 315–324, doi:10.3189/2012JoG11J123, 2012. 4826

4834

Number: 1 Author: Subject: Sticky Note Date: 02/01/2014 4:31:55 PM

This point is not logical: You can not do 'by average' where 1 out of the sample of 3 is an outlier. The conclusion displays bias. Also, considering the emphasis on the importance of better estimates of SLE one would expect more effort from the authors in performing the **uncertainty analysis**, displaying its results and discussing their results in this light. It became obvious that the authors had some other agenda (e.g. blacking the V-A scaling by using flawed arguments) rather than submitting results of a scientific study.

Number: 2 Author: Subject: Sticky Note Date: 02/01/2014 4:32:53 PM

The authors did not show any effort to propagate the errors from these individual assessment into regional estimates. This should have been done to arrive at a bulk uncertainty in these methods, in addition to sensitivity tests (whose results are also ignored in the conclusion). Again, this shows a bias in the presentation of the results.

Number: 3 Author: Subject: Sticky Note Date: 02/01/2014 2:39:43 PM

How is this 100% deviation assessed?

Number: 4 Author: Subject: Sticky Note Date: 02/01/2014 4:36:14 PM

It would be nice that this paper reveals some insight from an error analysis of the discussed methods, and propose a way for future progress (e.g. what kind of measurements are needed and on what kind of glaciers, on how many glaciers etc) and what kind of measurements would help improve the parameterizations in the 'advanced' methods. We all know that more volume measurements are needed so this is not news. So, not only that the study displays bias (lacks scientific objectivity), under-performs in terms of uncertainty analysis (having all the data it needs!) but also lacks novelty.

- Van den Broeke, M. R., and Paul, F.: A reconciled estimate of glacier contributions to sea level rise: 2003 to 2009, *Science*, 340, 852–857, doi:10.1126/science.1234532, 2013. 4815
- Gergan, J. T., Dobhal, D. P., and Kaushik, R.: Ground penetrating radar ice thickness measurements of Dokriani bamak (glacier), Garhwal Himalaya, *Curr. Sci. India*, 77, 169–173, 1999. 4825, 4826
- Glen, J. W.: The creep of polycrystalline ice, *P. Roy. Soc. A-Math. Phys.*, 228, 519–538, doi:10.1098/rspa.1955.0066, 1955.
- Graf, W. L.: The geomorphology of the glacial valley cross section, *Arctic Alpine Res.*, 2, 303–312, 1970. 4822
- Grinsted, A.: An estimate of global glacier volume, *The Cryosphere*, 7, 141–151, doi:10.5194/tc-7-141-2013, 2013. 4815, 4816, 4828, 4829, 4830, 4831, 4845, 4853
- Gurung, H.: Mountains of Asia, *Tech. rep.*, ICIMOD, Kathmandu, doi:10.5194/tc-7-141-2013, 1999. 4816
- Haeberli, W. and Hoelzle, M.: Application of inventory data for estimating characteristics of and regional climate-change effects on mountain glaciers: a pilot study with the European Alps, *Ann. Glaciol.*, 21, 206–212, 1995. 4815, 4818, 4819, 4820, 4822, 4824, 4826, 4831, 4832, 4849
- Hoelzle, M., Chinn, T., Stumm, D., Paul, F., Zemp, M., and Haeberli, W.: The application of glacier inventory data for estimating past climate change effects on mountain glaciers: a comparison between the European Alps and the Southern Alps of New Zealand, *Global Planet. Change*, 56, 69–82, doi:10.1016/j.gloplacha.2006.07.001, 2007. 4819
- Huss, M. and Farinotti, D.: Distributed ice thickness and volume of all glaciers around the globe, *J. Geophys. Res.*, 117, F04010, doi:10.1029/2012JF002523, 2012. 4815, 4816, 4820, 4822, 4823, 4824, 4825, 4827, 4829, 4830
- Huss, M., Farinotti, D., and Bauder, A.: Modelling runoff from highly glacierized alpine drainage basins in a changing climate, *Hydrol. Process.*, 22, 3888–3902, doi:10.1002/hyp.7055, 2008. 4815
- Immerzeel, W. W., van Beek, L. P. H., and Bierkens, M. F. P.: Climate change will affect the Asian water towers, *Science*, 328, 1382–1385, doi:10.1126/science.1183188, 2010. 4814
- IPCC: IPCC statement on the melting of Himalayan glaciers, *Press release*, 2010. 4815
- Jacob, T., Wahr, J., Pfeffer, W. T., and Swenson, S.: Recent contributions of glaciers and ice caps to sea level rise, *Nature*, 482, 514–518, doi:10.1038/nature10847, 2012. 4815

4837

- Jarvis, A., Reuter, H. I., Nelson, A., and Guevara, E.: Hole-filled seamless SRTM data V4, International Centre for Tropical Agriculture (CIAT), available at: <http://srtm.csi.cgiar.org>, 2008. 4817
- Kääb, A., Berthier, E., Nuth, C., Gardelle, J., and Arnaud, Y.: Contrasting patterns of early twenty-first-century glacier mass change in the Himalayas, *Nature*, 488, 495–498, doi:10.1038/nature11324, 2012. 4815
- Kamb, B. and Echelmeyer, K.: Stress-gradient coupling in glacier flow. I: Longitudinal averaging of the influence of ice thickness and surface slope, *J. Glaciol.*, 32, 267–284, 1986. 4822
- Kaser, G., Grosshauser, M., and Marzeion, B.: Contribution potential of glaciers to water availability in different climate regimes, *P. Natl. Acad. Sci. USA*, 107, 20223–20227, doi:10.1073/pnas.1008162107, 2010. 4814
- Le Bris, R. and Paul, F.: An automatic method to create flow lines for determination of glacier length A pilot study with Alaskan glaciers, *Comput. Geosci.*, 52, 234–245, doi:10.1016/j.cageo.2012.10.014, 2013. 4831
- Le Meur, E., Gerbaux, M., Schäfer, M., and Vincent, C.: Disappearance of an Alpine glacier over the 21st Century simulated from modeling its . . . , *Earth Planet. Sc. Lett.*, 261, 367–374, doi:10.1016/j.epsl.2007.07.022, 2007. 4815
- Li, H., Ng, F., Li, Z., Qin, D., and Cheng, G.: An extended “perfect-plasticity” method for estimating ice thickness along the flow line of mountain glaciers, *J. Geophys. Res.*, 117, F01020, doi:10.1029/2011JF002104, 2012. 4815
- Linsbauer, A., Paul, F., and Haeberli, W.: Modeling glacier thickness distribution and bed topography over entire mountain ranges with GlabTop: application of a fast and robust approach, *J. Geophys. Res.*, 117, F03007, doi:10.1029/2011JF002313, 2012. 4820, 4821, 4822
- Ma, L. L., Tian, L. D., Pu, J. C., and Wang, P. L.: Recent area and ice volume change of Kangwure Glacier in the middle of Himalayas, *Chinese Sci. Bull.*, 55, 2088–2096, doi:10.1007/s11434-010-3211-7, 2010. 4825, 4826
- Marussi, A.: Geophysics of the Karakorum, Italian Expeditions to the Karakorum (K2) and Hindu Kush, E. J. Brill, Leiden, 1964. 4826
- Marzeion, B., Jarosch, A. H., and Hofer, M.: Past and future sea-level change from the surface mass balance of glaciers, *The Cryosphere*, 6, 1295–1322, doi:10.5194/tc-6-1295-2012, 2012. 4815, 4816, 4828, 4831, 4845

4838

- Meier, M. F. and Bahr, D. B.: Counting glaciers: use of scaling methods to estimate the number and size distribution of the glaciers of the world, in: *Glaciers, Ice Sheets and Volcanoes: A Tribute to Mark F. Meier*, edited by: Colbeck, S. C., DTIC Document, 1–120, 1996. 4815
- Ohmura, A.: Completing the World Glacier Inventory, *Ann. Glaciol.*, 50, 144–148, 2009. 4816
- 5 Paul, F. and Linsbauer, A.: Modeling of glacier bed topography from glacier outlines, central branch lines, and a DEM, *Int. J. Geogr. Inf. Sci.*, 26, 1173–1190, doi:10.1080/13658816.2011.627859, 2012. 4815, 4820, 4821, 4822, 4824
- Paul, F., Barry, R., Cogley, J., Frey, H., Haeberli, W., Ohmura, A., Ommanney, C., Raup, B., Rivera, A., and Zemp, M.: Recommendations for the compilation of glacier inventory data from digital sources, *Ann. Glaciol.*, 50, 119–126, 2009. 4831
- 10 Radić, V. and Hock, R.: Regional and global volumes of glaciers derived from statistical upscaling of glacier inventory data, *J. Geophys. Res.*, 115, F01010, doi:10.1029/2009JF001373, 2010. 4815, 4828, 4829
- Radić, V., Bliss, A., Beedlow, A. C., Hock, R., Miles, E., and Cogley, J. G.: Regional and global projections of twenty-first century glacier mass changes in response to climate scenarios from global climate models, *Clim. Dynam.*, 1–22, doi:10.1007/s00382-013-1719-7, 2013. 4815, 4816, 4828, 4831, 4845
- 15 Raper, S. C. B. and Braithwaite, R. J.: The potential for sea level rise: new estimates from glacier and ice cap area and volume distributions, *Geophys. Res. Lett.*, 32, L05502, doi:10.1029/2004GL021981, 2005. 4815, 4828, 4829
- 20 Raup, B., Racoviteanu, A., Khalsa, S., Helm, C., Armstrong, R., and Arnaud, Y.: The GLIMS geospatial glacier database: a new tool for studying glacier change, *Global Planet. Change*, 56, 101–110, 2007. 4818
- Salzmann, N., Huggel, C., Rohrer, M., Silverio, W., Mark, B. G., Burns, P., and Portocarrero, C.: Glacier changes and climate trends derived from multiple sources in the data scarce Cordillera Vilcanota region, southern Peruvian Andes, *The Cryosphere*, 7, 103–118, doi:10.5194/tc-7-103-2013, 2013. 4819
- 25 Schiermeier, Q.: Glacier estimate is on thin ice, *Nature*, 463, 276–277, 2010. 4815
- Shi, Y., Liu, C., and Kang, E.: The glacier inventory of China, *Ann. Glaciol.*, 50, 1–4, doi:10.3189/172756410790595831, 2009. 4817
- 30 Shroder, J. F.: Himalaya, in: *Encyclopedia of Snow, Ice, and Glaciers*, edited by: Singh, V. P., Singh, P., and Haritashya, U. K., Springer Science+Business Media, Dordrecht, 510–520, 2011. 4816

4839

- Slangen, A. B. A. and van de Wal, R. S. W.: An assessment of uncertainties in using volume-area modelling for computing the twenty-first century glacier contribution to sea-level change, *The Cryosphere*, 5, 673–686, doi:10.5194/tc-5-673-2011, 2011. 4827


4840

Table 1. Number, area, and mean elevation of glaciers; see also supplementary materials in Bolch et al. (2012).

	Number of glaciers	Total area (km ²)	mean elevation (m a.s.l.)
Karakoram	7820	17 946	5326
W-Himalayas	9760	8943	5155
C-Himalayas	7547	9940	5600
E-Himalayas	3054	3946	5395
Entire HK-region	28 181	40 775	5362

4841

Table 2. Parameters of the applied V–A relations. Cogley (2011) used the same relationships for the glacier volume estimations of the same region. Volumes are calculated in m³ for input glacier areas measured in m.

Source	 1	γ
Chen and Ohmura (1990)	0.2055	1.36
Bahr et al. (1997)	0.191	1.375
Arendt et al. (2006)	0.28	1.375

4842

Table 3. Total ice volumes (km³) calculated with V–A relations, using glacier areas modified by $\pm 5\%$; relative difference to calculations with original areas in brackets. Abbreviations in the second line refer to: C&O90 to Chen and Ohmura (1990), B97 to Bahr et al. (1997), and A06 to Arendt et al. (2006)

	glacier areas +5%			glacier areas –5%		
	C&O90	B97	A06	C&O90	B97	A06
Karakoram	2387.9 (+6.86%)	2935.2 (+6.94%)	4303.0 (+6.94%)	2084.0 (–6.74%)	2557.9 (–6.81%)	3749.8 (–6.81%)
W-Himalayas	550.4 (+6.86%)	652.6 (+6.94%)	956.7 (+6.94%)	480.4 (–6.74%)	568.7 (–6.81%)	833.7 (–6.81%)
C-Himalayas	691.6 (+6.86%)	823.1 (+6.94%)	1206.7 (+6.94%)	603.6 (–6.74%)	717.3 (–6.81%)	1051.5 (–6.81%)
E-Himalayas	251.2 (+6.86%)	297.9 (+6.94%)	436.7 (+6.94%)	219.3 (–6.74%)	259.6 (–6.81%)	380.6 (–6.81%)

4843

Table 4. Total ice volumes (km³) calculated with the slope-dependent thickness estimation approach and GlabTop2, using modified parameters: $f = 0.7$ and $\tau_{\max} = 1.8$ bar for the large ice-volume combination; and $f = 0.9$ and $\tau_{\max} = 1.2$ bar for the small ice-volume combination. Parameterizations of τ as shown in Fig. 4, relative change to ice volumes calculated with original parameterization given in brackets.

	$f = 0.7$ and $\tau_{\max} = 1.8$ bar		$f = 0.9$ and $\tau_{\max} = 1.2$ bar	
	slope-dependent thickness est.	GlabTop2	slope-dependent thickness est.	GlabTop2
Karakoram	2965.1 (+39.71%)	2299.9 (+36.67%)	1537.5 (–27.56%)	1232.9 (–26.37%)
W-Himalayas	759.2 (+44.08%)	687.5 (+36.51%)	393.7 (–25.28%)	383.8 (–23.79%)
C-Himalayas	723.1 (+41.09%)	749.8 (+35.62%)	375.0 (–26.84%)	414.4 (–25.05%)
E-Himalayas	283.3 (+43.01%)	293.5 (+36.33%)	146.9 (–25.84%)	159.9 (–25.71%)
Entire HK-region	4730.7 (+40.80%)	4030.7 (+36.67%)	2453.1 (–26.99%)	2191.0 (–25.84%)

4844

Table 5. Ice-volume estimates (km^3) of the HK region according to the V–A approaches from Marzeion et al. (2012), Grinsted (2013), and Radić et al. (2013) applied to the glacier inventory used in this study.

	Marzeion et al. (2012)	Grinsted (2013)	Radić et al. (2013)
Karakoram	2748	1896	2953
W-Himalayas	611	584	657
C-Himalayas	771	714	828
E-Himalayas	279	265	300
Entire HK-region	4409	3459	4738

4845




Fig. 1. Study region and sources of the glacier inventory used for this study as well as by Bolch et al. (2012).

4846




Fig. 2. Differences between α_l (mean slope from $\arctan(\Delta H/l)$) and α_{DEM} (mean slope from DEM) for selected test glaciers, including the linear regression lines that are used as correction factors for glaciers $> 20 \text{ km}^2$ (blue), glaciers between 5 km^2 and 20 km^2 (purple), and glaciers $< 5 \text{ km}^2$ (orange). Glaciers were randomly selected, but the 20 largest glaciers of the inventory are included.

4847




Fig. 3. Schematic sketch of the functionality of GlabTop2: glacier polygons (blue curved line) are converted to a raster matching the DEM cells (red outline). Cells are discriminated as inner glacier cells (light blue), glacier marginal cells (powder-blue), glacier adjacent cells (yellow), and non-glacier cells (white). Pink cells represent randomly selected cells (r) for which local ice thickness is calculated; the blue square symbolizes the buffer of variable size, which is enlarged (dashed blue square), until an elevation extent of h_{min} is reached within the buffer.

4848

Number: 1 Author: Subject: Sticky Note Date: 02/01/2014 2:30:17 PM
What does different color shading refer to?




Fig. 4. Parameterizations of the basal shear-stress τ with the elevation extent ΔH . Original parameterization from Haeberli and Hoelzle (1995) (blue curve), and the parameterizations used for the uncertainty assessment (with $\tau_{max} = 1.8$ bar (red) and 1.2 bar (orange)).

4849




Fig. 5. Resulting volumes (km^3) for the entire HK region (left) and the four sub-regions.



4850

Number: 1 Author: Subject: Sticky Note Date: 02/01/2014 2:40:12 PM

Would be good to add the uncertainty range for each bar, and to assess statistical significance of the differences in volume estimates among the methods.




Fig. 6. Ice-thickness distribution of Chhota Shigri (W-Himalayas) modeled with GlabTop2 (left) and the HF-method (right).

4851




Fig. 7. Measured vs. modeled local ice thicknesses for results from GlabTop2 (blue pluses) and the HF-model (red circles). Numbers on top-left indicate the mean difference, the RMSE, and, in brackets, the RMSE divided by the mean of the measurements.

4852




Fig. 8. Mean thickness vs. glacier area for “measured” glaciers, including the scaling relations used in this study. Note the variability of mean thickness of up to an order of magnitude for a given glacier size. (Data collected by Graham Cogley and Regine Hock, published in Cogley, 2012, downloaded from Grinsted, 2013).

4853




Fig. 9. Basal shear stresses calculated by the HF-model compared to the applied τ -parameterizations for GlabTop2 (solid blue line) and the slope-dependent thickness estimations (solid black line). Dashed blue lines represent τ -parameterizations used for the sensitivity analysis.

4854