

TCD

8, 1227–1256, 2014

Union Glacier

A. Rivera et al.

This discussion paper is/has been under review for the journal The Cryosphere (TC).  
Please refer to the corresponding final paper in TC if available.

# Union Glacier: a new exploration gateway for the West Antarctic Ice Sheet

A. Rivera<sup>1,2</sup>, R. Zamora<sup>1</sup>, J. A. Uribe<sup>1</sup>, R. Jaña<sup>3</sup>, and J. Oberreuter<sup>1</sup>

<sup>1</sup>Centro de Estudios Científicos, P.O. Box 5110466, Valdivia, Chile

<sup>2</sup>Departamento de Geografía, Universidad de Chile, P.O. Box 3387, Santiago, Chile

<sup>3</sup>Instituto Antártico Chileno, Punta Arenas, Chile

Received: 31 December 2013 – Accepted: 24 January 2014 – Published: 20 February 2014

Correspondence to: A. Rivera (arivera@cecs.cl)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Abstract

Union Glacier (79°46' S/83°24' W) in the West Antarctic Ice Sheet (WAIS), has been used by the private company Antarctic Logistic and Expeditions (ALE) since 2007 for their landing and commercial operations, providing a unique logistic opportunity to perform glaciological research in a vast region, including the Ice divide between Institute and Pine Island glaciers and the Subglacial Lake Ellsworth. Union glacier is flowing into the Ronne Ice Shelf, where future migrations of the grounding line zone (GLZ) in response to continuing climate and oceanographic changes have been modelled. In order to analyse the potential impacts on Union glacier of this scenario, we installed an array of stakes, where ice elevation, mass balance and ice velocities have been measured since 2007, resulting in near equilibrium conditions with horizontal displacements between 10 and 33 myr<sup>-1</sup>. GPS receivers and three radar systems have been also used to map the subglacial topography, the internal structure of the ice and the presence of crevasses along surveyed tracks. The resulting radar data showed a subglacial topography with a minimum of 858 m below sea level, much deeper than estimated before. The below sea level subglacial topography confirms the potential instability of the glacier in foreseen scenarios of GLZ upstream migration during the second half of the XXI century.

## 1 Introduction

WAIS, the West Antarctic Ice sheet (Fig. 1) has been considered potentially unstable because its bedrock is located well below sea level (Humbert, 2012), and its total disintegration could contribute up to 4.3 m (Fretwell et al., 2013) to global sea-level rise. In many cases, the topography underneath WAIS is inversed (ice is deeper upstream) and steeper than at the present grounding line zones (GLZ) (Ross et al., 2012). This topographic condition, in a context of ongoing global changes, especially oceanic warming in areas of the Southern Ocean, is leading to GLZ upstream migration, as ob-

TCD

8, 1227–1256, 2014

Union Glacier

A. Rivera et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Union Glacier

A. Rivera et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

served for example at the Amundsen Sea Embayment Area (ASEA) glaciers, like Pine Island (PIG) and Thwaites (Rignot et al., 2002). This migration process in a context of inversed subglacial topographies, has an impact on glacier dynamics, since bottom melting at the GLZ is higher in deeper waters, provoking higher ice fluxes, ice thinning and, in general, a more negative mass balance (Rignot and Jacobs, 2002). However, there are WAIS areas where the changes are not as dramatic as observed at ASEA, with some regions thickening rather than thinning (Joughin and Bamber, 2005).

One of the WAIS areas where glaciological changes are not strong at present is the Weddell Sea sector (Rignot and Thomas, 2002), where the Ronne Ice Shelf (RIS) is located (Fig. 1). This floating platform has been relatively stable in recent decades, with different behaviours of the GLZ and very small net mass balance changes (Rignot et al., 2011b). The relative stability of the RIS is partly explained by the Weddell Seas oceanographic and atmospheric conditions, and the associated sea ice extension (Mayewski et al., 2009). The sea ice volume in this area has been stable (Cavalieri and Parkinson, 2008), but recent studies (Hellmer et al., 2012) forecast a sea ice volume reduction for the XXI century resulting in incremental circulation of warm water beneath the Filchner ice shelf toward the GLZ of the Slessor, Support Force, Moeller and Foundation ice streams. The intrusion of warmer waters will certainly increase the basal melt rate of the ice shelf, and possibly lead to retreat of the GLZ. This, in turn, would favour higher ice fluxes along tributary glaciers, and thinning that will spread upstream along deep channels (Rignot et al., 2011a).

Between two of the main ice streams in the region (Rutford and Institute), there are two smaller glaciers also draining into the RIS; Union glacier (79°46' S/83°24' W) flowing into the Constellation inlet and the Horseshoe valley (80°18' S/81°22' W) flowing into the Hercules inlet. The study of these glaciers could provide an important clue about ongoing changes taking place in the region, especially considering that local glacier conditions can be significantly affected by RIS changes.

In this context, the main aim of this paper is to present recent glaciological results obtained at the Union glacier and nearby areas that provide a base line for possible ice dynamic responses to ongoing and modelled future changes of RIS.

## 2 Study area

5 The Horseshoe valley ( $80^{\circ}18' S/81^{\circ}22' W$ ) has been studied since the 1990s (Wendt et al., 2009), when Chilean Air Force C-130 airplanes landed on wheels at the Patriot Hills Blue Ice Area (BIA). BIAs have long been recognized as suitable places for landing airplanes (Mellor and Swithinbank, 1989). Since the 1980s, the private company Antarctic Logistics and Expeditions (ALE) also operated at this part of the Ellsworth  
10 Mountains using the Patriot Hills BIA for their commercial operations. However, the landing of heavy airplanes has been frequently disrupted due to strong prevailing cross winds.

In order to increase the number of airplane operational days and improve access for heavy cargo airplanes, in 2007/8 ALE moved to Union Glacier ( $79^{\circ}46' S/83^{\circ}24' W$ ) (Fig. 2), where the prevailing wind direction is in line with the landing strip (tail-head katabatic winds), helping airplanes to operate even with strong gusts. Thanks to ALE logistic support, four scientific campaigns have been conducted on Union glacier since  
15 2008, where a glaciological program was established, including ice dynamic, mass balance and geophysical surveys (Rivera et al., 2010).

20 Union glacier has an estimated total area of  $2561 \text{ km}^2$ , with a total length of 86 km from the ice divide with Institute ice stream down to the grounding line of the Constellation Inlet on the RIS. The glacier has several glacier tributaries, the main trunks being located in the Union and Schanz valleys (Figs. 1 and 2), which are feed through narrow glacial valleys (9 km wide) flowing from the interior plateau until they join together at the  
25 Union “gate”. This gate or narrowest section of the glacier (7 km wide), has a central moraine line comprised of clasts and small-sized debris. The central moraine is visible

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



along the local BIA that is used for landing Ilyushin IL76 airplanes on wheels under a contract with ALE llc.

An array of 21 stakes was installed at the gate in 2007 for ice dynamic and surface mass balance studies. The Union gate has been re-surveyed by using GPS and radar systems installed onboard convoys pulled by Camoplast tractors travelling along pre-designated routes. Results of the first campaigns of 2007 and 2008 were published by Rivera et al. (2010).

### 3 Methods

#### 3.1 Meteorological data

The meteorological data analysed here was collected at the Union Glacier base camp (Fig. 2) Automatic Weather Station (AWS), which has been operating since 2008. The AWS includes air temperature, humidity, air pressure, and wind speed and direction sensors. The AWS data series show several data gaps due to malfunction or low power supply problems during the winter. In spite of the above, the available data are good enough for weather forecast during landing operations in the summer, and for describing general temperature conditions in the area, as well as predominant winds and directions.

#### 3.2 Remote sensing

Mid-summer ASTER satellite images acquired between 2002 and 2013 were analyzed and classified via manual digitalization in order to delineate the local Union Glacier main BIA. This task allowed areal changes in snow cover to be quantified and analysed for possible relationships with local meteorological data.

The ASTER images were geometrically corrected using the internal parameters of each scene. A true colour composition (Band 1, 2 and 3N) for each year was pro-

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



duced and the contrast and brightness modified by means of histogram equalization to facilitate the delimitation of the blue ice area.

The BIA outline mapping was based upon a similar procedure established by Rivera et al. (2013), where the boundary between snow and ice is manually identified assuming the maximum extent criterion of identification. This is easily done thanks to the spectral differences between snow and ice.

### 3.3 Glaciological mass balance

The protruding heights of every stake were measured yearly between 2007 and 2011 (Figs. 1 and 2). The annual heights were compared and local mass balance was subsequently calculated. In order to convert these values into water equivalent, a mean density of  $910 \text{ kg m}^{-3}$  (Paterson, 1994) was used for stakes drilled on ice. In those cases where the stakes were located at snow surfaces, the densities were measured by using a Mount Rose snow probe with penetration capacity of near 50 cm of snow-firn.

### 3.4 GPS

In December 2007, the initial observation network of 21 bamboo-stakes was installed on a Blue Ice Area on the Union Glacier (Fig. 2). For each of these stakes GPS measurements were made using Topcon GR3 dual frequency GPS receivers with measurement times of less than 30 s per stake, yielding metre scale errors. In December 2008, 2009 and 2010, the same network, as well as 35 new stakes were surveyed with dual frequency Javad GPS model Lexon GGD receivers. In order to apply a differential correction procedure, a similar GPS receiver was continuously collecting data at a rock location (RX1 in Fig. 2). The precision obtained by these surveys increased to less than 10 cm, by measuring between 15–30 min per stake. Additionally, a dual frequency Javad GPS receiver was installed on snow (Bstat in Fig. 2) collecting data between 11 December 2009 and 31 January 2010 for high resolution studies (each 30 s), with

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the aim of detecting possible tidal modulated ice velocities. In January 2011, a Real Time Kinematic procedure was applied to data collected at the Union glacier gate by a Leica SR 9500 receiver, with estimated decimetre accuracy. All GPS data collected were processed using the commercial software GrafNav 8.20.

### 5 3.5 Surface elevation changes and geodetic mass balance estimation

The data obtained by the GPS measurements were compared in order to calculate the local mass balance and the absolute surface elevation changes at the BIA.

Using the measured height differences obtained by GPS and considering the surface topographic slope effect, the submergence/emergence velocity ( $w_e$ ) was calculated:

$$10 \quad w_e = w_s - u_s \tan \alpha \quad (1)$$

where  $w_s$  is the vertical ice velocity,  $u_s$  is the ice velocity along the surface flow direction and  $\alpha$  is the slope. The emergence velocity represents the vertical flow of ice relative to the glaciers surface and allows the estimation of the net balance if it is assumed that density does not change with depth during the period (Hooke, 2005).

15 By subtracting the emergence velocity of the measured specific balance (stake height difference) the surface elevation change with time at a fixed position can be obtained.

$$\frac{\partial h}{\partial t} = b - w_e \quad (2)$$

20 If the glacier is in a steady state, there would be no surface changes since accumulation and ablation equals out the submergence and emergence velocities so that the surface profile remains unchanged (Hooke, 2005). In this case  $b = w_e$ , however most glaciers are not perfectly in steady state. Only the stake array measured at the Union gate was considered for this analysis.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

### 3.6 Radar

A pulse compression radar depth sounder designed at CECS, was used to measure ice thickness. The radar operates at a central frequency of 155 MHz, a bandwidth of 20 MHz, and 200 W of peak power. Yagi antennae were used for both the transmitter and receiver. The radar includes a digital waveform generator, up converter transmitter, a timing and synchronization controller and an 80 dB variable gain digital receiver.

A Frequency Modulated – Continuous Wave (FM-CW) radar, was also used to measure surface snow accumulation (up to 450 m thick), with high vertical resolution (1 m). This radar works at a frequency from 550 MHz to 900 MHz using two separated log periodic antennae for the transmitter and receiver. The transmit power was 21 dBm, the intermediate frequency (IF) amplifier gain was 70 dB and the whole system operated at a PRF (Pulse Repetition Frequency) 10 kHz. A Direct Digital Synthesis (DDS) system was used to generate an extremely linear frequency sweep transmitted signal.

The third radar system used along these tracks was a commercial Ground Penetrating Radar (GPR), a Geophysical Survey Systems Inc. (GSSI) model SIR-3000, working at 400 MHz. The digital control unit of the radar triggers pulses at 100 kHz repetition rate. The range was set between 190 and 300 ns to record subsurface reflection with the aim to find crevasses in real time. The GPR antennae were mounted on a 7 m long rod which was attached to a tractor and had rubber car tire tube installed at the opposite end. The radar data were analysed in real time while the tractor was moving, in order to alert the driver to the presence of hyperbolae, potentially related to crevasses. This system was used particularly in areas previously detected on ASTER satellite imagery as being highly crevassed.

Post processing and data analyses were carried out using Reflex-Win V5.6 (Sandmeier Scientific Software) for the three radar systems. A background removal, dewow filter and adjustment of the gain function, among other procedures, were applied to the raw data. For crevasses analysis, migration correction procedures were also applied to collapse the hyperbolic diffractions to their proper point origins (Plewes and Hub-

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





bard, 2001). In order to convert travel time to ice thickness in metres, it was assumed that the electromagnetic wave travelled through the ice at  $0.168 \text{ mns}^{-1}$ , a mean value representative of cold ice (Glen and Paren, 1975) and at  $0.194 \text{ mns}^{-1}$  assumed to be representative for snow/firm (Woodward and King, 2009).

## 4 Results

### 4.1 Meteorological data and BIA area changes

The meteorological data collected at Union Glacier between 2008 and 2013 contains several gaps and non valid records, but in general provide a good idea of the local conditions, which are especially useful for landing operations. In terms of climatological analysis, the series is too short and noisy, but is the only one available in the region.

The mean daily air temperature at this location (Fig. 3) from 2008 to 2012 was  $-20.6^\circ\text{C}$ , with an absolute minimum of  $-42.7^\circ\text{C}$  recorded on 12 August 2008 at 2 a.m. and an absolute maximum of  $0.5^\circ\text{C}$  registered on 15 January 2010 at 8 p.m. Daily air temperatures have maximums in December–January following direct solar radiation, however between April and August (included) when the Sun is below the horizon, the lowest temperatures have a mean of  $-25.5 \pm 3.5^\circ\text{C}$ . Temperatures close to the melting point have only been observed in three summer events: 7–8 January 2008, 14–15 January 2010 and 25–26 December 2010, but no surface melting has been observed on this site. The mean air temperature in January is  $-10.3^\circ\text{C}$ .

Regarding wind speed and direction, the data are very consistent between 2008–2012 with a mean value 16.3 knots and predominant direction from  $224^\circ$ . The maximum wind speed recorded on site was near 60 knots and the predominant wind speeds are higher than 25 knots (Fig. 4). This predominant wind direction explains the extension of the main BIA which has experienced less than 10% area changes between 2002 and 2013. The series is however too short to detect a trend (Table 1).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 4.2 Mass balance

Snow densities measured at stakes drilled on snow surfaces have a mean value of  $400 \pm 3 \text{ kg m}^{-3}$ , with small variability between all surveyed stakes. No fresh and soft snow was detected due to the lack of precipitation in the region during our surveys.

5 Table 2 indicates the details of each stake, the type of surface and period of measurements.

The mass balance estimations between 2007 and 2011 at the stakes installed on the BIA gate (V00 to V21, Table 2), are typically negative (Fig. 5), with a mean inter annual mass balance of  $-0.097 \text{ m w.eq. a}^{-1}$ .

10 The stakes installed along the traverse between Patriot Hills and Union Glacier (Fig. 1) were only measured between 2008 and 2009 and showed differing results depending on their location. Stakes installed on snow surfaces (B1–B12) have positive mass balances with a mean snow accumulation of  $0.3 \text{ m a}^{-1}$  ( $0.12 \text{ m w.eq. a}^{-1}$ ). Just outside the BIA, stake B12 showed the maximum net balance ( $0.2 \text{ m w.eq. a}^{-1}$ ),  
15 indicating that snow drift coming from the BIA, due to the predominant katabatic wind direction (Fig. 4) is having a positive downstream accumulation effect. The only negative mass balance along the track Patriot Hills–Union Glacier (excluding the Union glacier BIA) was detected at stake B5, which is located at the edge of the local BIA of the Plumber glacier (Fig. 1). Here a modest  $-0.07 \text{ m w.eq. a}^{-1}$  (Table 2) was obtained  
20 between 2008 and 2009. Stakes B13–B19 (Table 2, Fig. 2), also had negative values with a mean surface balance of  $0.1 \text{ m w.eq. a}^{-1}$ , coincident with the other stakes located at a local BIA.

## 4.3 Surface ice velocity

25 Surface ice velocities were obtained for 19 stakes (B1–B19) located along the track between Patriot Hills and Union Glacier base camp (Figs. 3 and 6), with measurements periods of near 1 yr, and resulting ice velocities between  $0.1$  and  $34.6 \text{ m a}^{-1}$ . Minimum values were observed at local ice divides (B2–B8) while the maximum velocity was

TC D

8, 1227–1256, 2014

Union Glacier

A. Rivera et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



observed at stake B10 located at the steepest area of Union Glacier, between two crevasse fields (Figs. 2 and 6).

Between 11 December 2009 and 31 January 2010 a dual frequency GPS receiver was attached to stake Bstat, taking continuous measurements every 15 s. The receiver was powered by batteries and solar panels, providing a high detailed record of daily ice dynamic during the spanned period of time. The main aim of this survey was to test the hypothesis that the ice dynamics of Union glacier are affected by Ronne Ice Shelf tides, as has been observed at the Rutford ice stream where a modulated ice flow was detected (Gudmundsson, 2006). This stake (Bstat, Table 2) is located  $\sim 38$  km upstream the local grounding line zone of the Constellation inlet at the Ronne Ice Shelf (Figs. 1 and 2). The resulting ice velocity was  $33 \text{ m a}^{-1}$ , and thanks to the analysis carried out by H. Gudmunsson (personal communication, 2010), no tidal effect was detected.

The neighbouring stake to Bstat (B9, located within a distance of 10 m) was only measured for 30 min in December 2009 and in December 2010, resulting in a annual velocity of  $33.26 \pm 0.7 \text{ m a}^{-1}$ . The good correspondence with the Bstat results ( $33 \text{ m a}^{-1}$ ) indicates that Union Glacier does not show important seasonal ice velocity changes.

At the Union Glacier gate stake array (Fig. 2), 22 stakes were drilled into its Blue Ice Area in 2007 and have been annually resurveyed up until 2011. Unfortunately, not all the original stakes have survived, or were re-measured every year. However, from the remaining stake network a mean velocity of  $20 \text{ m a}^{-1}$  was estimated for the measurement period (Fig. 6). The orientation of the overall ice movement shows an angle of  $65.42^\circ$  at the gate, similar to the main wind direction (Fig. 4) observed at the BIA (from  $224^\circ$  to  $44^\circ$ ). No temporal velocity changes were observed during the studied period.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## 4.4 Surface elevation change

Considering the surface velocities obtained for the stakes (Table 2) and by applying Eq. (1), a mean vertical velocity for the blue ice of  $-0.07 \pm 0.007 \text{ m a}^{-1}$  was found at the Union Gate. For a steady state glacier this would be equal to the local mass balance, however this is not the case for the studied area (there is a negative mass balance at the BIA) and a mean local elevation change of  $-0.012 \text{ m a}^{-1}$  was found. In any case, this result is close to the estimated error of the measurements and indicates near equilibrium conditions.

## 4.5 Glacier thickness and internal structure

More than 450 km long radar tracks were measured between 2008 and 2010 at the Union Glacier area, including oversnow traverses between Patriot Hills, Union Glacier and the high Antarctic plateau.

The  $\sim 80$  km long profile between the Antarctic Plateau and Union Glacier (Fig. 7 upper), including the transit along Balish, Schneider, Schanz and Driscoll glaciers, was surveyed with a FM-CW radar for snow accumulation data, a compression pulse radar for ice thickness data and a GPR system for crevasse detection purposes.

The deep ice of the plateau is seen at the beginning of the survey profile (A in Fig. 7), then is followed by a passage characterized by shallow ice (B) where the steepest section of the whole traverse was crossed. After this section, the thickness increases sharply until a local maximum thickness of 1120 m was observed at Balish Glacier. The local divide between the Balish and Schneider glaciers (C in Fig. 7) has a prominent subglacial peak (C in Fig. 7) where the mountain range dividing the two glaciers is also visible underneath the ice. The maximum thickness measured at Schneider Glacier was 900 m, 1050 m at Schanz Glacier, 1510 m in Driscoll Glacier and 1540 m in Union Glacier, close to the base camp.

At the beginning of the radar profile (A and B in Fig. 7), it is possible to see the shallowest ice at the Gifford Peaks pass (passage), with a thickness between 45 and

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



140 m (Fig. 8) obtained with the low gain channel. This section (B in Fig. 7 bottom) corresponds to a ~ 5 km long FM-CW radargram where 400 m of dynamic range were processed detecting the snow/ice stratigraphy with a resolution of 1 m.

All along the whole A–F profile (Fig. 7 bottom), a detailed internal stratigraphy was detected in the upper hundred metres, comprised of isochronous layers that are particularly prominent at local ice divides (e.g. at C in Fig. 7 bottom). In those areas, the internal structures are characterized by typical minimum and divergent flow and the presence of Raymond Bumps (Paterson, 1994) and other internal inflections that can lead to the understanding of long-term changes in the ice flow.

The collected ice thickness data at Union glacier has a mean ice thickness of 1450 m. The subglacial topography in the valley is smooth, with “U” shape flanks. The FM-CW radar detected multiple internal snow-ice layers and the snow-ice boundary layer with a high resolution (Fig. 8). Also, the FM-CW radar was capable of detecting the bedrock up to 350 m, improving the bedrock detection from the pulse-compression radar when surveying shallow ice (B at Fig. 7 bottom).

The GPR data collected along the track from Union Glacier to the Antarctic Plateau allowed the detection of many crevasses both near the runway area, and at the Gifford Peaks passage, just before reaching the plateau. Along the survey route from the Union Glacier to the Antarctic Plateau the GPR system was able to detect the upper 20 m of the internal structure of the ice. Within this upper layer annual snow/ice layers suffered (in places) discontinuities due to crevasses that appeared as hyperbolae on the radar traces.

In general, the route from Union glacier to the Antarctic plateau is almost totally free of crevasses. An exception to this was the Gifford passage (B in Fig. 7) where a minor system of crevasses was detected corresponding to steepening slopes. These crevasses have widths between 1 and 5 m, with snow bridges between 1.5 and 3 m thick. The Gifford passage is a narrow and steep valley, where the longitudinal stresses and velocities are probably much higher than the rest of the area. In spite of these crevasses, the Gifford passage is the shortest (235 km) gateway for oversnow tra-

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

verses from Union glacier to the Subglacial Lake Ellsworth (Vaughan et al., 2007) at the Antarctic plateau, and is therefore the preferred route. The alternative route to this subglacial lake is near 520 km long, starting at Union glacier, passing along Patriot Hills, then travelling around the Tree Sails and finally turning west toward the Subglacial lake Ellsworth (Fig. 1 bottom).

## 5 Discussion

At Union glacier, the local BIA is shaped by strong snowdrift caused by katabatic winds accelerated by the slope at the main junction between the two glacial valley arms feeding Union glacier from the upper Antarctic plateau. As a result, the net mass balance at this BIA is negative, the driving factor explaining the mass losses being sublimation of ice during summer months, as no melting event has been observed since 2007. However, melting events have been observed in the region, especially at Patriot Hills, where a small water pond was formed at the boundary between the BIA and the nunatak during a very warm 1997 summer (Carrasco et al., 2000).

In spite of this negative surface mass balance, the resulting ice elevation changes between 2008–2011 at the Union gate are very close to the error of the measurements (combined error of 0.01 m), with a mean of  $-0.012 \text{ ma}^{-1}$  and high spatial variability ( $\pm 0.044 \text{ ma}^{-1}$ ) among the gate stakes. Accordingly, the glacier must be considered in equilibrium without significant changes compared with previous data (Rivera et al., 2010).

The ice velocities at the BIA fluctuate between 11 and  $24 \text{ ma}^{-1}$  without significant changes between 2007 and 2011. However, downstream of the Union gate, the velocities increase up to  $33 \text{ ma}^{-1}$  at the continuous GPS stake (Bstat), where no seasonal variations or tidal modulated variability were detected.

The GPR survey allowed the detection of many more crevasses than were previously mapped with the ASTER imagery. The 400 MHz GPR is capable of identifying in real time surface and buried crevasses. The 400 MHz GPR crevasses data were compared

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



to the 500–900 MHz FMCW records and in most of the cases the wider crevasses (2 to 4 m width) could be detected in both radars. However, the FMCW radar did not provide the best information in regards to snow bridge thicknesses (less than 1 m), its resolution being insufficient to detect the first meters of the snow pack.

Comparisons of surface and bedrock topography along transect A–F (Fig. 9), resulted in a maximum ice thickness difference of 1447 m with a mean difference of 477 m (standard deviation: 348 m) between the data presented here and BEDMAP2 (Fretwell et al., 2013). The surface topography of the surveyed traverse was found to differ from BEMAP2 and IceSat (2003–2005) by a mean of 91 m and 169 m, respectively. These differences are understandable due to the coarse resolution of BEDMAP2 and the footprint of IceSat.

The subglacial topography (Fig. 9) at the main trunk of the Union glacier toward the local GLZ (Segment E–G in Fig. 7), showed a subglacial topography well below sea level (–858 m), much deeper than previously thought. However, between Union glacier and the local GLZ, the subglacial topography shows a maximum altitude of –190 m at F in Fig. 9. The subglacial topography then deepens toward the GLZ, where the bedrock is estimated to be 1050 m below sea level (Fretwell et al., 2013). This subglacial condition implies that an upstream migration of the GLZ until point F in Fig. 9 will not have a strong effect on Union glacier. However, the glacier can respond in a more dynamic way if this possible migration is affecting Union glacier upstream this point.

## 6 Conclusions

Several glaciological oversnow campaigns have been undertaken to Union Glacier and nearby areas since 2007, where the surface and subglacial topographies were mapped in detail. These results were compared to the Bedmap dataset, showing much deeper bedrock and a much more complex subglacial topography than previously estimated. The obtained results determined a maximum ice thickness of 1540 m in Union Glacier,

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Union Glacier

A. Rivera et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



with a maximum snow ice boundary layer at 120 m. The internal structure of the ice was also mapped, including the detection of isochronous layers and crevasses, allowing logistic operators and scientists to work along safer routes.

Ice dynamics were also recorded thanks to the measurement of ice velocities and ice thickness. Maximum ice velocity values of  $34.6 \text{ m a}^{-1}$  were obtained. Near equilibrium conditions were calculated at the BIA, where mean velocities of  $20 \text{ m a}^{-1}$  were measured. The snow accumulation among the studied stakes outside BIAs showed values of up to  $0.2 \text{ m w.eq. a}^{-1}$  (near  $0.5 \text{ m a}^{-1}$  of snow). At the BIA, a local negative mass balance was detected as expected, with mean ablation rates of  $0.1 \text{ m w.eq. a}^{-1}$ .

Due to the below sea level subglacial topography upstream the local GLZ, it is expected that Union Glacier can experience some thinning and acceleration in future scenarios of GLZ migration. However these impacts will not affect the whole glacier, because at near 55 km upstream the present GLZ, the bedrock topography is close to sea level.

Overall the collected data allowed a ground route from Union glacier to the upper Antarctic plateau to be satisfactory mapped. The route is important as it traverses the Subglacial lake Ellsworth and the main ice streams flowing into the Amundsen Sea Embayment area (Pine Island Glacier) and toward the Weddell Sea (Institute and Rutford Ice stream). This new 235 km long route toward Subglacial Lake Ellsworth is shorter than half of the previously traversed track of 517 km, providing a much direct and short gateway into inner Antarctica.

*Acknowledgements.* This research was supported by Antarctic Logistic Expeditions (ALE) Ltd. Special thanks to M. Sharp, M. and P. McDowell, C. Jacobs, Eddy, Boris and all ALE personnel at Union. CECs is funded by the Basal fund of CONICYT, among other grants. J. A. Neto collaborated with GPS data collection in 2011. A. Wendt, C. Rada, T. Kohoutek and M. Barandun helped with data processing. H. Gudmundsson analysed the GPS data at Bstat in order to detect possible tidal modulated effects on the ice flow. J. Carrasco and F. Burger helped with the met data. D. Carrión helped with figures and satellite image analysis. R. Wilson edited the text. A. Rivera is a Guggenheim fellow.



## References

- Carrasco, J., Casassa, G., and Rivera, A.: A warm event at Patriot Hills, Antarctica: an ENSO related phenomena?, in: Sixth International Conference on Southern Hemisphere Meteorology and Oceanography, 240–241, Santiago, Chile, 3–7 April, 2000. 1240
- 5 Cavalieri, D. and Parkinson, C.: Antarctic sea ice variability and trends, 1979–2006, *J. Geophys. Res.*, 113, C07004, doi10.1029/2007JC004564, 2008. 1229
- Fretwell, P., Pritchard, H. D., Vaughan, D. G., Bamber, J. L., Barrand, N. E., Bell, R., Bianchi, C., Bingham, R. G., Blankenship, D. D., Casassa, G., Catania, G., Callens, D., Conway, H., Cook, A. J., Corr, H. F. J., Damaske, D., Damm, V., Ferraccioli, F., Forsberg, R., Fujita, S., Gim, Y., Gogineni, P., Griggs, J. A., Hindmarsh, R. C. A., Holmlund, P., Holt, J. W., Jacobel, R. W., Jenkins, A., Jokat, W., Jordan, T., King, E. C., Kohler, J., Krabill, W., Riger-Kusk, M., Langley, K. A., Leitchenkov, G., Leuschen, C., Luyendyk, B. P., Matsuoka, K., Mouginit, J., Nitsche, F. O., Nogi, Y., Nost, O. A., Popov, S. V., Rignot, E., Rippin, D. M., Rivera, A., Roberts, J., Ross, N., Siegert, M. J., Smith, A. M., Steinhage, D., Studinger, M., Sun, B., Tinto, B. K., Welch, B. C., Wilson, D., Young, D. A., Xiangbin, C., and Zirizzotti, A.: Bedmap2: improved ice bed, surface and thickness datasets for Antarctica, *The Cryosphere*, 7, 375–393, doi:10.5194/tc-7-375-2013, 2013. 1228, 1241, 1256
- 15 Glen, J. and Paren, J.: The electrical properties of snow and ice, *J. Glaciol.*, 15, 15–38, 1975. 1235
- 20 Gudmundsson, G.: Fortnightly variations in the flow velocity of Rutford Ice Stream, West Antarctica, *Nature*, 444, 1063–1064, 2006. 1237
- Hellmer, H., Kauker, F., Timmermann, R., Determann, J., and Rae, J.: Twenty-first-century warming of a large Antarctic ice-shelf cavity by a redirected coastal current, *Nature*, 485, 225–228, 2012. 1229
- 25 Hooke, R.: *Principles of Glacier Mechanics*, 2nd Edn., Cambridge University Press, Cambridge, UK, 2005. 1233
- Humbert, A.: Vulnerable ice in the Weddell Sea, *Nat. Geosci.*, 5, 370–371, 2012. 1228
- Jezek, K., and RAMP Product Team: RAMP AMM-1 SAR Image Mosaic of Antarctica. Version 2. Fairbanks, AK: Alaska Satellite Facility, in association with the National Snow and Ice Data Center, Boulder, CO, Digital media, 2002. 1254
- 30

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## Union Glacier

A. Rivera et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Joughin, I. and Bamber, J.: Thickening of the ice stream catchments feeding the Filchner-Ronne Ice Shelf, Antarctica, *Geophys. Res. Lett.*, 32, L17503, doi:10.1029/2005GL023844, 2005. 1229
- Mayewski, P., Meredith, M., Summerhayes, C., Turner, J., Worby, A., Barrett, P., Casassa, G., Bertler, N., Bracegirdle, T., Naveira, A., Bromwich, D., Campbell, H., Hamilton, G., Lyons, W., Maasch, K., Aoki, S., Xiao, C., and van Ommen, T.: State of the Antarctic and Southern Ocean climate system, *Rev. Geophys.*, 47, RG1003, doi:10.1029/2007RG000231, 2009. 1229
- Mellor, M. and Swithinbank, C.: Airfields on Antarctic glacier ice, CRREL Rep. 89–21, US Army Cold Regions Research and Engineering Laboratory, USA, 97 pp., 1989. 1230
- Paterson, W.: *The Physics of Glaciers*, 2nd Edn., Pergamon Press, Exeter, Great Britain, 1994. 1232, 1239
- Plewes, A. and Hubbard, B.: A review of the use of radio-echo sounding in glaciology, *Prog. Phys. Geog.*, 25, 203–236, 2001. 1234
- Rignot, E. and Jacobs, S.: Rapid bottom melting widespread near Antarctic Ice Sheet grounding lines, *Science*, 296, 2020–2023, 2002. 1229
- Rignot, E. and Thomas, R.: Mass balance of polar ice sheets, *Science*, 297, 1502–1506, 2002. 1229
- Rignot, E., Vaughan, D., Schmelz, M., Dupont, T., and MacAyeal, D.: Acceleration of Pine Island and Thwaites Glaciers, West Antarctica, *Ann. Glaciol.*, 34, 189–194, 2002. 1229
- Rignot, E., Mouginot, J., and Scheuchl, B.: Ice flow of the Antarctic Ice Sheet, *Science*, 333, 1427–1430, 2011a. 1229
- Rignot, E., Mouginot, J., and Scheuchl, B.: Antarctic grounding line mapping from differential satellite radar interferometry, *Geophys. Res. Lett.*, 38, L10504, doi:10.1029/2011GL047109, 2011b. 1229
- Rivera, A., Zamora, R., Rada, C., Walton, J., and Proctor, S.: Glaciological investigations on Union Glacier, Ellsworth Mountains, West Antarctica, *Ann. Glaciol.*, 51, 91–96, 2010. 1230, 1231, 1240
- Rivera, A., Cawkwell, F., Wendt, A., and Zamora, R.: Mapping blue ice areas and crevasses in West Antarctica using ASTER images, GPS and radar measurements, in: *Global Land Ice Measurements from Space*, edited by: Kargel, J., Leonard, G., Bishop, M., Käab, A., and Raup, B., Springer-Praxis, Heidelberg, 743–757, 2013. 1232

Ross, N., Bingham, Corr, H., Ferraccioli, F., Jordan, T., Le Brocq, A., Rippin, D., Young, D., Blankenship, D., and Siegert, J.: Steep reverse bed slope at the grounding line of the Weddell Sea sector in West Antarctica, *Nat. Geosci.*, 5, 393–396, 2012. 1228

5 Scambos, T., Haran, T., Fahnestock, M., Painter, T., and Bohlander, J.: MODIS-based Mosaic of Antarctica (MOA) data sets: continent-wide surface morphology and snow grain size, *Remote Sens. Environ.*, 111, 242–257, 2007. 1248

Vaughan, D., Rivera, A., Woodward, J., Corr, H., Wendt, J., and Zamora, R.: Topographic and hydrological controls on Subglacial Lake Ellsworth, West Antarctica, *Geophys. Res. Lett.*, 34, L18501, 2007. 1240

10 Wendt, A., Casassa, G., Rivera, A., and Wendt, J.: Reassessment of ice mass balance at Horseshoe Valley, Antarctica, *Antarct. Sci.*, 21, 505–513, 2009. 1230

Woodward, J. and King, E.: Radar surveys of the Rutford Ice Stream onset zone, West Antarctica: indications of flow (in)stability?, *Ann. Glaciol.*, 50, 57–62, 2009. 1235

TCD

8, 1227–1256, 2014

## Union Glacier

A. Rivera et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)**Table 1.** BIA extension based upon ASTER images collected since 2002.

Date	Area (km <sup>2</sup> )
24 Nov 2002	95.6
9 Dec 2004	104.7
27 Jan 2007	112.0
17 Jan 2009	111.8
11 Feb 2012	88.2
1 Feb 2013	96.7


**Table 2.** Stake co-ordinates, velocities and mass balance per period. See Figs. 1 and 2 for locations.

Stakes	Lat°	Lon°	H. (m.a.e)	Velocity (m a <sup>-1</sup> )	Mass Balance (m w.eq. a <sup>-1</sup> )	Period	Type of surface
B1	-80.211	-81.230	803.12	12.7	0.18		
B2	-80.124	-81.113	813.51	1.8	0.19		
B3	-80.045	-80.937	759.07	0.6	0.04		snow
B4	-79.985	-81.080	573.57	1.8	0.06		
B5	-79.922	-81.211	538.16	2.0	-0.07	2008–2009	snow/BIA
B6	-79.840	-81.298	669.59	0.1	0.03		
B7	-79.801	-81.348	553.87	1.2	0.04		
B8	-79.757	-81.965	459.99	2.9	0.14		
B9	-79.709	-82.455	531.04	33.3	0.14		
Bstat	-79.708	-82.454	531.34	33.0	-	2009/11/12 2010/31/01	snow
B10	-79.717	-82.643	639.79	34.6	0.17		
B11	-79.757	-82.803	672.46	20.5	0.13		
B12	-79.760	-82.931	690.56	20.9	0.20		
B13	-79.764	-83.091	703.53	22.9	-0.09	2008–2009	ice
B14	-79.768	-83.266	738.39	22.4	-0.06		
B15	-79.798	-83.281	735.19	21.6	-0.16		
B16	-79.796	-83.316	742.20	23.2	-0.13		
B17	-79.765	-83.343	739.36	17.0	-0.08		
B18	-79.760	-83.370	747.68	14.0	-0.11		ice/crevasses
B19	-79.764	-83.372	743.72	11.1	-0.10		
V00	-79.769	-83.370	739.47	18.1	-0.08	2008–2010	
V01	-79.770	-83.370	738.44	19.0	-0.08	2007–2011	
V02	-79.772	-83.368	738.13	20.2	-0.04	2007–2010	
V03	-79.774	-83.365	739.69	21.7	-0.11		
V04	-79.775	-83.363	736.40	22.3	-0.07	2007–2011	
V05	-79.777	-83.361	733.57	23.0	-0.22		
V06	-79.779	-83.358	735.12	23.5	-0.11		
V07	-79.781	-83.355	736.61	23.9	-0.08	2007–2010	
V08	-79.784	-83.352	739.05	24.3	-0.13	2007–2009	ice
V09	-79.787	-83.343	738.60	24.3	-0.11		
V10	-79.790	-83.334	737.63	24.3	-0.13	2007–2011	
V11	-79.792	-83.325	738.82	24.0	-0.07	2007–2010	
V12	-79.771	-83.337	738.51	21.9	-0.09		
V13	-79.773	-83.335	735.87	22.3	-0.08	2007–2011	
V14	-79.774	-83.332	733.20	22.4	-0.08		
V15	-79.777	-83.327	731.27	22.7	-0.09		
V16	-79.779	-83.323	731.72	23.1	-0.10	2007–2009	
V17	-79.782	-83.316	731.16	23.4	-0.11		
V18	-79.784	-83.313	730.33	23.5	-0.13		
V19	-79.786	-83.309	729.48	23.6	-0.07	2007–2011	
V20	-79.788	-83.303	728.57	23.3	-0.11		
V21	-79.791	-83.298	733.66	23.0	-0.07		




**Fig. 1.** Left: WAIS Location map. Right: union glacier basin (light blue area). In yellow, the main ice divides in the region. In dashed white the 2005 track to Subglacial lake Ellsworth (517 km). In green, the new track surveyed in 2010 (235 km). Stakes B1 to B9 are described in the text. The black box is shown at Fig. 2. Background image: MODIS mosaic of Antarctic (Scambos et al., 2007).

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

**Fig. 2.** Yellow dots show the stakes described in the text (Fig. 1 and Table 2). RX1 is the location of the static GPS on rock. Crevasses are shown in red. The Union Glacier “gate” is a transversal profile between B18 and B15 (inset) where the ice runway is shown in purple. Mean ice velocities area shown as arrows. Summer base camp is located near B11. The background image is an ASTER false composite 321 acquired on 1 February 2013.




**Fig. 3.** Mean daily temperature ( $^{\circ}\text{C}$ ) during 2008 (black triangles) and modeled solar direct radiation (red line) at the AWS site ( $\text{kW m}^{-2}$ ).

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	







**Fig. 4.** Predominant wind speed and direction 2008–2012 at the Union Glacier Automatic Weather Station.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶


Back Close

Full Screen / Esc

Printer-friendly Version


Interactive Discussion






**Fig. 5.** Mass balance (m w.eq. a<sup>-1</sup>) 2007–2011 at stake located at the local BIA. Stakes locations are shown in Fig. 2.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)





**Fig. 6.** Ice velocities ( $\text{m a}^{-1}$ ) at the Union Glacier gate by indicated years. Stakes location is shown in Fig. 2.


[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

**Fig. 7.** Upper: location map of the radar and GPS survey measured between the Antarctic Plateau (A) and the Union Glacier (F) in 2010 (green line). In blue, the available data between F and the local Grounding Line Zone at the Constellation Inlet (G). The background image is the RAMP AMM-1 SAR Image Mosaic of Antarctica (Jezek and RAMP Product Team, 2002). Bottom: surface (red line) and subglacial topography (blue line) interpreted from the radar data collected in 2010 along the A–F track.



**Fig. 8.** FM-CW data obtained at Union glacier. Upper: location of the surveyed track near the base camp of Union glacier. The background image is an ASTER false composite 321 acquired on 14 March 2013. Bottom: radargram showing the surface topography (red line), internal annual snow/firn layers and the firn/ice boundary (blue line). Letters A and B indicate the base camp where the snow/firn layers are near 30–40 m thick in total. Letter C, corresponds to the central moraine line of the glacier. Letter D, indicates the appearance of the BIA at the surface.



**Fig. 9.** Comparisons of surface and bedrock topography of different data sources along transect A–G (see Fig. 8 upper for location of transect). CECs surface and bedrock from year 2010. IceSat surface from 2003–2005, Bedmap 2 surface and Bedrock from Fretwell et al. (2013). Bedmap2 bottom dashed line is interpolated.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)