

How much snow falls on the Antarctic ice sheet?

C. Palerme^{1,2}, J. E. Kay³, C. Genthon^{1,2}, T. L'Ecuyer⁴, N. B. Wood⁵, and C. Claud⁶

¹CNRS, LGGE (UMR5183), F-38041 Grenoble, France

²Univ. Grenoble Alpes, LGGE (UMR5183), F-38041 Grenoble, France

³NCAR CGD - National Center for Atmospheric Research Climate and Global Dynamics, Boulder, Colorado, USA

⁴Department of Atmospheric and Oceanic Sciences, University of Wisconsin-Madison, USA

⁵Cooperative Institute for Meteorological Satellite Studies, University of Wisconsin-Madison, USA

⁶Laboratoire de Météorologie Dynamique/IPSL, CNRS, Ecole Polytechnique, Palaiseau, France

Correspondence to: Cyril Palerme
(cyril.palerm@lgge.obs.ujf-grenoble.fr)

Abstract. Climate models predict Antarctic precipitation to increase during the 21st century, but their present day Antarctic precipitation differs. A model-independent climatology of the Antarctic precipitation characteristics, such as snowfall rates and frequency, is needed to assess the models, but is not yet available. Satellite observations of precipitation by active sensors has been possible in the polar regions since the launch of CloudSat in 2006. Here we use two CloudSat products to produce the first multi-year model-independent climatology of Antarctic precipitation. The first product is used to determine the frequency and the phase of precipitation, while the second product is used to assess the snowfall rate. The mean snowfall rate from August 2006 to April 2011 is 171 mm/year over the Antarctic ice sheet north of 82°S. While uncertainties on individual precipitation retrievals from CloudSat data are potentially large, the mean uncertainty should be much smaller but cannot be easily estimated. There are no in situ measurements of Antarctic precipitation to directly assess the new climatology. However distributions of both precipitation occurrences and rates generally agree with the ECMWF ERA-Interim dataset, the production of which is constrained by various in situ and satellite observations but does not use any data from CloudSat. The new dataset thus offers unprecedented capability to quantitatively assess Antarctic precipitation statistics and rates in climate models.

1 Introduction

Evaluating Antarctic snow accumulation, the sum of precipitation, sublimation / evaporation, melt-water run-off, and blowing snow (Eisen et al., 2008), is a major challenge with relevance to sea level rise. While no significant change in the total Antarctic snow accumulation has been found in ice

cores and reanalysis products over the last 50 years (Monaghan et al., 2006a; Frezzotti et al., 2013), future changes are likely to occur, with global consequences : a projected 25% increase in accumulation over the 21st century would result in a drop of approximately 160 mm/century in global sea level (Gregory and Huybrechts, 2006).

25 Climate models consistently predict Antarctic precipitation to increase in a warming climate (Church et al., 2013), however their present day mean Antarctic precipitation differs widely, ranging from 150 to 550 mm/year in the CMIP3 archive (Genthon et al., 2009a). The Antarctic precipitation rates have also been evaluated from regional atmospheric models (Bromwich et al., 2004; Van de Berg et al., 2005; Monaghan et al., 2006b; Lenaerts et al., 2012). The mean solid precipitation rate
30 over the grounded ice sheet reported by Van de Berg et al. (2005) is 164 mm/year using the model RACMO2/ANT. Moreover, Monaghan et al. (2006b) found a precipitation rate of 178 mm/year and 200 mm/year with Polar MM5 using the reanalysis NCEP-II and ERA40 respectively for the initial and boundary conditions. Reanalyses have also been used for assessing Antarctic precipitation (Monaghan et al., 2006a; Bromwich et al., 2011). Bromwich et al. (2011) have compared 6 re-
35 analysis datasets and found that the mean precipitation rate on the grounded ice sheet varies from 145 to 203 mm/year depending on the reanalysis. There is therefore a need to document Antarctic precipitation from observations to benchmark climate models.

While accumulation rates have been assessed using in situ observations (Arthern et al., 2006; Eisen et al., 2008), characteristics of Antarctic precipitation (the dominant positive term of the accumulation) such as the frequency and the rate remain poorly known. Ground-based measurements
40 are sparse and difficult to make in Antarctica. In coastal areas, katabatic winds are strong, which makes the distinction between blowing snow and precipitation difficult. On the Antarctic plateau, the annual accumulation is small (few centimeters per year, Bromwich et al., 2004), and the instrumentation must be able to detect very light precipitation. In addition, low temperatures and hoarfrost
45 negatively impact instruments that are not designed for harsh environments.

Precipitation characteristics depend greatly on the region in Antarctica. In coastal areas, precipitation is influenced by synoptic scale features such as cyclones and fronts (Bromwich, 1988). In the interior (> 2500 m), a potentially significant part of the precipitation falls in the form of "diamond dust" (ice crystals) under clear sky conditions (Bromwich, 1988; Fujita and Abe, 2006).

50 In the past, passive microwave remote sensing has been used to detect new snow accumulation, using changes in surface emissivity (Bindschadler et al., 2005). However the method was not quantitative, and was found to be affected by other processes such as temperature and surface roughness.

Observations from the Cloud Profiling Radar (CPR) on CloudSat provide the first opportunity to estimate precipitation in polar regions from a spaceborne radar (Stephens et al., 2008; Liu, 2008).
55 With data available from August 2006 to April 2011, CloudSat directly observes snow precipitating through the atmosphere, rather than after it has been accumulated on the surface. Several algorithms have been tested for precipitation over polar regions using CloudSat (Kulie and Bennartz, 2009;

Hiley et al., 2010). Moreover, Boening et al. (2012) have already shown that there is good agreement between CloudSat, GRACE, and ERA Interim for Antarctic precipitation variability over Dronning Maud Land (30°W-60°E, and 65°S-80°S). However, no precipitation climatology has been done over Antarctica at the continent scale. In this study, we used two CloudSat products to make the first multi-year climatology of Antarctic precipitation north of 82°S from spaceborne observations.

2 Data and Methods

The CPR, onboard CloudSat, is a nadir-looking radar at 94 GHz which measures the power backscattered by cloud particles and hydrometeors according to the distance from the sensor. It provides radar reflectivity profiles divided into 150 bins at a vertical resolution of 240 m, with a 1.7 km x 1.3 km footprint, and up to 82° of latitude. Its minimum detectable radar reflectivity is around -28 dBZ.

In this study, two CloudSat products are used to determine the characteristics of Antarctic precipitation. The first product, 2C-PRECIP-COLUMN (Haynes et al., 2009), is used to assess the phase and occurrence frequency of Antarctic precipitation. 2C-PRECIP-COLUMN provides a precipitation flag based on the near-surface reflectivity (dBZ) at the fourth bin over the ocean (between 600 and 840 m above the surface), and at the sixth bin over land (about 1300 m) to remove surface contamination (ground clutter). The radar bin containing the surface is determined with a digital elevation model. The phase is obtained using the temperature at 2 m predicted by the European Centre for Medium Range Weather Forecasts operational weather analysis, and a model of melting layer with a constant lapse rate of 6 °C/km. According to the phase, different thresholds are applied to the near surface reflectivity to determine a likelihood of precipitation (possible or certain). Thus, the precipitation flags inform about the likelihood and the phase of precipitation.

The second product, 2C-SNOW-PROFILE (Wood, 2011; Wood et al., 2013) is used to assess the snowfall rates. 2C-SNOW-PROFILE retrieves estimates of liquid-equivalent snowfall rate for profiles where 2C-PRECIP-COLUMN indicates 'snow possible' or 'snow certain', or where 2C-PRECIP-COLUMN indicates 'mixed possible' or 'mixed certain' and the estimated melted mass fraction at the surface is less than or equal to 0.1. Using a priori estimates of snow particle size distribution, microphysical and scattering properties, an optimal estimation retrieval (Rodgers, 2000) is performed for the contiguous layer of snow-containing radar bins nearest the surface, with exclusions for likely ground clutter contamination. With this approach, the so-called Z-S relationship between radar reflectivity and snowfall rate is not fixed, but can vary subject to the constraints of the reflectivity profile and the a priori expectations.

The retrieval also provides estimates of uncertainties for the retrieved snowfall rates. The uncertainties depend on the uncertainties in the observed reflectivities as well as those in the simulated reflectivities provided by the retrieval's radar forward model. These uncertainties arise due to measurement error and due to the approximate nature of the forward model and its a priori assumptions.

To the extent they can be characterized, systematic errors are removed. Within the context of the retrieval algorithm the remaining uncertainties are considered to be unbiased and random; however, these likely consist of some combination of systematic and random uncertainties since, for example, the algorithm's a priori assumptions are not tuned to the particular characteristics of Antarctic snowfall. Thus, while multi-year averaging of retrieved snowfall rates reduces the truly random component of the uncertainties, some indeterminate bias is likely also present, and its evaluation is an ongoing area of research.

In this study, both CloudSat datasets are processed over a grid of 1° of latitude by 2° of longitude between 63°S and 82°S . The number of orbits per grid cell for the period August 2006 - April 2011 is shown in Fig. 1. Over the Antarctic continent, the number of orbits per grid cell is at least 350 for the entire period, which represents one orbit every 5 days. This does not seem to be an issue to produce a climatology of precipitation (figure 5, discussed below). Moreover, the ratio of the surface directly observed by CloudSat over the surface of the ice sheet is shown in figure S1. Even for latitudes less than 82°S , the surface directly covered by CloudSat is only a fraction of the total surface of the ice sheet. However, the spatial scale of precipitation events and the overpass frequency ensure adequate statistical sampling over the duration of the study (supplementary material S1).

CloudSat products provide the data along their orbit. In order to map the 2C-PRECIP-COLUMN data over a grid of 1° by 2° , one sample per grid cell overflown is retained for each orbit. Thus, we redefined new samples from the original 2C-PRECIP-COLUMN flags. First, for the precipitation frequency, flags are sorted into three classes : no precipitation, precipitation possible, and precipitation certain. Then, if all the flags in the same grid cell indicate no precipitation, no precipitation is retained. If at least one flag indicates precipitation certain in the grid cell, precipitation certain is retained. And if there is no flag indicating precipitation certain, and at least one flag indicating precipitation possible, precipitation possible is retained. It is relevant to note that this method tends to inflate the precipitation occurrence.

To map the precipitation phase, flags are sorted into four classes : no precipitation, liquid, mixed, and solid precipitation. If the flags in the grid cell indicate no precipitation and precipitation, but only one precipitation phase, this phase is retained. If the flags in the same grid cell indicate rain and mixed precipitation, mixed precipitation and snow, or rain and snow, mixed precipitation is retained. For the precipitation phase, this method tends to inflate the mixed precipitation class. For the snowfall rate and its uncertainty from the 2C-SNOW-PROFILE product, the mean value in the grid cell has been retained for each orbit.

CloudSat observations have been compared to ERA Interim reanalysis in this study. ERA Interim is the latest global atmospheric reanalysis produced by the European Centre for Medium Range Weather Forecast (Simmons et al., 2006; Dee et al., 2011). ERA Interim provides data from 1979 to present at a 6-hourly resolution. Its coverage is global at a spatial resolution of about $0.75^\circ \times 0.75^\circ$. The 6-hour and 12-hour forecasts of precipitation are used here. Data from surface observations and

130 radiosondes, commercial aircraft observations, and satellites measurements are assimilated in the numerical model to improve and constrain the forecasts (Dee et al., 2011). Direct precipitation observations are not assimilated into the model, but precipitation is influenced by the four-dimensional variational assimilation of other variables such as temperature and humidity (www.ecmwf.int).

ERA Interim has been chosen in this study because it likely offers the most realistic depiction of 135 Antarctic precipitation variability among reanalyses (Bromwich et al., 2011). However, it has been shown that ERA Interim could have a dry bias over the East Antarctic plateau (Bromwich et al., 2011; Favier et al., 2013).

3 Results

3.1 Precipitation characteristics from CloudSat

140 Fig. 2 shows two maps of the precipitation frequency assessed from the 2C-PRECIP-COLUMN flags, and a map of the ratio of the number of samples indicating precipitation possible over the number of samples indicating precipitation possible and certain. The map a) represents the proportion of samples indicating precipitation certain, and the map b), the proportion of samples indicating precipitation certain and possible.

145 The mean precipitation frequency (% of time) observed by CloudSat over the Antarctic continent (latitude $< 82^{\circ}\text{S}$) is 14 % when the samples precipitation possible are not taken into account, and 26 % with the samples precipitation possible included. The spatial pattern of the precipitation frequency shows two distinct regions. The first area includes the West Antarctic ice sheet and the peripheral part of the East Antarctic ice sheet, which corresponds approximately to the part of the 150 continent with surface elevation below 2250m. In this region, relatively high precipitation frequency is observed by CloudSat (between 22 % and 34 % depending on whether the samples precipitation possible are taken into account). The second region is the Antarctic plateau in East Antarctica (with surface elevation $> 2250\text{m}$), where the precipitation frequency observed by CloudSat is much lower (between 5 % and 19 %). Fig. 1 shows the parts of the ice sheet with surface elevation over and 155 below 2250m derived from combined satellite radar and laser data (Bamber et al., 2009). Each part represents 50 % of the surface of the Antarctic ice sheet (including the part of the continent between 82°S and 90°S).

In figure 2, the map c) shows the number of samples indicating precipitation possible over the number of samples indicating precipitation possible and certain. While over the periphery of the ice 160 sheet, most of the precipitation events detected are sorted as certain, most of the samples indicating precipitation are sorted as possible in the interior. Near surface reflectivity is sensitive to the size of hydrometeors, and on the plateau, particles are probably too small to increase the near surface reflectivity above the threshold "precipitation certain". The reflectivity thresholds applied in this algorithm could be too high for this kind of precipitation.

165 Precipitation phase has also been studied from the 2C-PRECIP-COLUMN flags. Only samples
indicating precipitation certain were taken into account. Over the Antarctic ice sheet (latitude <
82°S), solid precipitation represents 99.60 %, mixed precipitation 0.32 %, and rain 0.08 % of the
precipitation occurrence (similar results have been found with the samples indicating precipitation
possible included). In peripheral areas (surface elevation < 2250m), mixed precipitation represents
170 0.63 % and rain 0.15 % of the precipitation occurrence. Relatively more liquid and mixed precipita-
tion occurs over the Peninsula compared to the rest of the ice sheet (mixed precipitation represents
4.10 %, and rain 1.32 % of the precipitation occurrence over this region). Furthermore, on the
Antarctic plateau (surface elevation > 2250m), all the precipitation is solid.

Because snowfall rate in the 2C-SNOW-PROFILE product is only estimated when the melted
175 fraction is assessed to be less than or equal to 0.1, this product is well-suited to examining precip-
itation over Antarctica. Fig. 3 shows the mean annual snowfall rate, the single retrieval snowfall
rate uncertainty, and the ratio of the single retrieval uncertainty over the snowfall rate from the 2C-
SNOW-PROFILE data. The mean snowfall rate observed by CloudSat on the Antarctic continent
(latitude < 82°S) is 171 mm water equivalent (w.e.) per year. However, the spatial pattern of the
180 snowfall rate shows considerable differences between West Antarctica and East Antarctica. In West
Antarctica, the mean annual snowfall rate is 303 mm w.e. per year, compared to 118 mm w.e. per
year in East Antarctica. Furthermore, the mean snowfall rate over the peripheral part of the ice sheet
(with surface elevation < 2250m) is 303 mm w.e. per year, compared to 36 mm w.e. per year for the
interior of the ice sheet (with surface elevation > 2250m).

185 The map of the snowfall rate uncertainty in Fig. 3 represents the mean value of the single retrieval
uncertainty for all the snowfall rate retrievals from August 2006 to April 2011. These are the ex-
pected uncertainties for individual snowfall rate retrievals and as noted earlier likely consist of both
random and systematic components. Considering the maps in Fig. 3, 2C-SNOW-PROFILE product
provides a snowfall rate uncertainty between 1.5 and 2.5 times the snowfall rate for 4.7 years of cur-
190 tain data accumulated on the 1° x 2° grid boxes. This relative uncertainty is particularly high on the
Antarctic plateau and the Peninsula, and it is lower on the peripheral part of the ice sheet and in West
Antarctica. When calculating mean values with large number of observations, the standard error of
the mean decreases as the number of samples increases. Therefore, in this study, the uncertainty on
a 4.7 years mean snowfall rate should be fairly small. However, the real snowfall rate uncertainty on
195 the entire CloudSat period is difficult to assess because the relative contribution of systematic and
random errors remains unknown.

3.2 Comparison of the CloudSat products to ERA Interim reanalysis

Table 1 shows a comparison between ERA Interim reanalysis and the precipitation samples from
CloudSat at the French station Dumont d'Urville (Fig. 1). The nearest grid cell of the ERA Interim
200 reanalysis has been taken into account for comparing the datasets. The ability of ERA Interim to

represent precipitation in Antarctica is poorly known, but it is expected that observations assimilated in the model help to constrain the forecasts. It is important to note that CloudSat observations are not used to produce ERA Interim reanalysis (Dee et al., 2011). Surface observations and radiosoundings are performed at Dumont d'Urville and assimilated in ERA Interim. Humidity profiles obtained
205 by radiosounding influence the forecasts used to predict precipitation in ERA Interim. Therefore, precipitation predicted by ERA Interim should be relatively more reliable at Dumont d'Urville than other sites where no observation is available.

Comparisons of the ERA Interim reanalysis data at Dumont D'Urville station against the precipitation samples from CloudSat were used to establish a precipitation rate threshold for comparing
210 the datasets. This threshold is necessary because the ERA Interim precipitation rates were strictly positive 60 % of the time between 2006 and 2011 at Dumont D'Urville station (for comparison the precipitation frequency from CloudSat at Dumont d'Urville is between 22 % and 32 %). A threshold of 0.07 mm/6h for the ERA Interim precipitation rates was found empirically to give good agreement with the CloudSat precipitation samples (0.07 mm/6h corresponds to the threshold for which
215 the highest Heidke Skill Score (Barnston, 1992) has been obtained).

From 2006 to 2011, 67 % of the time, the ERA Interim precipitation rates at Dumont D'Urville were below this rate. During this period, for the 265 samples that indicate no precipitation, 92 % match with no precipitation in ERA Interim. Furthermore, for the 38 samples indicating precipitation possible, 55 % match with a precipitation event in ERA Interim. And for the 85 samples indicating
220 precipitation certain, the success rate was 91 %.

A similar comparison at the continent scale is shown in Fig. 4. For comparing the datasets, ERA Interim reanalysis have been interpolated on the same $1^\circ \times 2^\circ$ grid as CloudSat. The threshold used in ERA Interim was the same as for Dumont d'Urville (0.07 mm/6h). The value of the appropriate threshold seems to depend on the location in Antarctica, and it is probably lower where the precipi-
225 tation rate is small.

The method used for determining the threshold at Dumont d'Urville (highest Heidke skill score obtained for this threshold) has been tested in the interior of the ice sheet. This method tends to maximize the success rate of the different classes according to the number of samples in each class. Because there are less samples indicating precipitation certain in the interior of the ice sheet compared to the total number of samples (Fig. 2 c)), it tends to maximize the success rate for the class
230 "periods without precipitation". Thus, the threshold in the interior of the ice sheet tends to be higher than 0.07 mm/6h, while the precipitation rate is smaller in this region. That is why, we chose to keep the same threshold for the whole continent as the threshold determined at Dumont d'Urville.

Overall, the success rate for the samples "precipitation certain" and "precipitation possible" is
235 better near the coast than in the interior of the ice sheet. It could be due to the threshold applied to ERA Interim precipitation which could be too high for the Antarctic interior. Moreover shallow precipitation missed by CloudSat, and the weak reflectivity of small hydrometeors in the interior

could contribute to this difference.

Fig. 4 shows also a map of the Heidke skill score. The Heidke skill score measures the accuracy
240 of forecasts relative to random forecasts (Barnston, 1992). It can vary between -1 and 1. A Heidke
skill score equal to 0 means that forecasts are as good as random draw, and it is equal to 1 for perfect
forecasts. If the Heidke skill score is positive, the forecasts are better than random forecasts. Here,
samples indicating precipitation certain and period without precipitation are used. Fig. 4 shows
better agreement between CloudSat and ERA Interim (higher Heidke skill score) over peripheral
245 areas than over the interior.

Even if precipitation is not assimilated in ERA Interim, observations assimilated such as humidity
profiles are more numerous in peripheral areas than in the Antarctic interior. Observations of wind,
pressure, and temperature further help constrain the strength and timing of the perturbations and thus
that of the precipitation events. Perturbations are fainter when they reach the interior and, as there
250 are less observations available they are less efficient at controlling the occurrence and timing of pre-
cipitation through assimilation. Therefore, ERA Interim should be more reliable in peripheral areas
than on the Antarctic plateau. This could help explain why there is better agreement between Cloud-
Sat and ERA Interim in peripheral areas than in the interior of the ice sheet. However, agreement
between CloudSat and ERA Interim is good over West Antarctica, where there are less observations
255 assimilated than over peripheral areas of East Antarctica. Overall, there is better agreement where
the snowfall rate is high.

A comparison between the snowfall rate observed by CloudSat and simulated by ERA Interim is
shown in Fig. 5 and in Table 2. For this comparison, a map with the same temporal sampling as
CloudSat has been created. Every time a grid cell has been overflowed by CloudSat, the corresponding
260 time step in ERA Interim has been retained. This dataset has been created in order to test if the
temporal sampling of CloudSat may result in a bias for the period August 2006-April 2011.

The snowfall rates from ERA Interim with the same temporal sampling as CloudSat and from
the full ERA Interim dataset are similar (163 mm w.e. per year for both datasets over the Antarctic
continent north of 82°S). The snowfall rate from ERA Interim with the same temporal sampling as
265 CloudSat is slightly stronger over the interior of the ice sheet (53 compared to 49 mm w.e. per year),
and slightly lower over the periphery (271 compared to 273 mm w.e. per year). These are considered
as marginal differences, and temporal sampling of CloudSat does not seem to be an issue.

Over the Antarctic continent (latitude < 82 °S), the mean snowfall rate is 171 mm w.e. per year for
CloudSat and 163 mm w.e. per year for ERA Interim. The snowfall rate observed by CloudSat and
270 predicted by ERA Interim are relatively similar over the ice sheet, except over parts of the Peninsula,
the Vinson massif (78°S, 85°W), and the Prince Charles Mountains (around 72°S, 65°E), where the
snowfall rate from CloudSat is significantly stronger. Orographic precipitation could be seen by
CloudSat, but not predicted by ERA Interim due to the difference in spatial resolution between both
datasets. However, ground clutter should be stronger over mountainous areas than over flat terrain,

275 and may induce a spuriously high snowfall rate.

3.3 Comparison of the snowfall rate from CloudSat to surface mass balance observations

Table 2 and Fig 6 show a comparison between the snowfall rate obtained by CloudSat for the period August 2006–April 2011, and the accumulation rate assessed by Arthern et al. (2006) for the period 1950–2000. Arthern et al. (2006) used in-situ glaciological measurements to assess the accumula-
280 tion, and passive radiometer data (AMSR-E) sensitive to snowpack characteristics for interpolating their results.

Assuming that accumulation has not significantly changed during the last 50 years (Monaghan et al., 2006a; Frezzotti et al., 2013), the accumulation from Arthern et al. (2006) represents 95 % of the snowfall over the Antarctic ice sheet north of 82 °S. The snowfall rate observed by CloudSat
285 is higher than the accumulation over the periphery of the ice sheet, which is expected due to the negative contribution to accumulation of sublimation, melt/run-off, and blowing snow.

However, the snowfall rate observed by CloudSat is lower than the accumulation in the interior. Snowfall rate assessed by CloudSat over the interior of the ice sheet may be underestimated due to shallow precipitation missed by CloudSat and the weak reflectivity of small hydrometeors. Addi-
290 tionally, modelling studies have suggested that deposition (inverse sublimation) could be stronger than evaporation at some locations in the interior of the ice sheet (Genthon and Krinner, 2001). Thus, hoarfrost formation could contribute significantly to the accumulation, and precipitation could be lower than accumulation in these regions.

Ground-based measurements used to produce the accumulation map from Arthern et al. (2006)
295 were not filtered according to their accuracy, and some measurements have been found to be unreliable (Magand et al., 2007). Genthon et al. (2009b) have shown that the unreliability of some in situ observations used by Arthern et al. (2006) would lead to an overestimated accumulation in the interior of the ice sheet. Thus the accumulation from Arthern et al. (2006) could be overestimated in this region. Moreover, Magand et al. (2008) have shown that the interpolation based on microwave
300 surface emission used by Arthern et al. (2006) can be inaccurate in coastal areas affected by melt during the summer.

Favier et al. (2013) assembled a surface mass balance database in which ground-based measure-
ments have been sorted into three classes according to their accuracy. Observations sorted in the most reliable class for the 20th century by Favier et al. (2013) have been used in this study. The
305 ratio of the snowfall rate observed by CloudSat over the accumulation from Favier et al. (2013) is reported on the map of Fig. 6. When several values of accumulation are given in the database of Favier et al. (2013) for the same grid cell of 1° x 2°, the mean value for the grid cell is shown in Fig. 6. Overall the comparison between the snowfall rate from CloudSat and the accumulation from Favier et al. (2013) confirms the results from the comparison with the accumulation map of Arthern
310 et al. (2006). However, in some grid cells, accumulation is not spatially homogeneous, and a few in

situ measurements can be sometimes not representative of the mean accumulation in the grid cell. For instance, there is only one value of accumulation for the three red dots of Fig. 6 showing the largest ratio between the snowfall rate from CloudSat and the accumulation from Favier et al. (2013).

4 Discussion and conclusion

315 A climatology of the Antarctic precipitation, the single most important positive term of the ice sheet mass balance, has yet to be established from observations. Filling this gap, Antarctic precipitation features such as the frequency, the phase, and the snowfall rate have been determined here using CloudSat products. CloudSat is the first spaceborne radar able to observe precipitation in Antarctica (Boening et al., 2012), and its potential is confirmed here. The mean snowfall rate from August
320 2006 to April 2011 is 171 mm/year over the Antarctic ice sheet north of 82°S. Expectedly, the observed surface accumulation of snow is on average less than snowfall. However, it appears to exceed snowfall in areas of lesser precipitation where uncertainties on both precipitation and accumulation reports are largest. A significant contribution of hoarfrost to the surface mass balance of these areas may not be excluded.

325 However, due to assumptions about particle size distribution, particle masses, shapes and fall-speeds, snowfall rates assessed in the 2C-SNOW-PROFILE product have large uncertainties. In the 2C-PRECIP-COLUMN dataset, a large number of samples are sorted as "precipitation possible". This leads to a considerable range of precipitation frequency, even if the frequency estimated is probably more reliable than the snowfall rate. Moreover, on the Antarctic plateau, the 2C-PRECIP-
330 COLUMN algorithm may have difficulties in distinguishing precipitating from non precipitating hydrometeors due to their small particle size.

CloudSat is likely more accurate in peripheral areas than in the interior. Shallow precipitation (< 1300m), missed by CloudSat, could be an important contribution to precipitation on the Antarctic plateau. Therefore, the precipitation frequency and the snowfall rate could be underestimated over
335 this region. On the other hand, because near surface reflectivity is measured about 1300m over the surface, blowing snow is not confounded with precipitation in peripheral areas, which is usually the main problem for precipitation measurements over this region.

Due to the difficulties for CloudSat to detect precipitation in the interior of the ice sheet, CloudSat precipitation products are more useful in the periphery of the ice sheet than in the interior. However,
340 precipitation in the periphery of the ice sheet is quite important. Three quarters of the total Antarctic precipitation falls in this region, and it is where the models predict the largest precipitation increase over the 21st century (Genthon et al., 2009a).

The lack of ground-based measurements prevents direct validation of CloudSat data. Nevertheless, agreement between CloudSat data and ERA Interim reanalysis is encouraging for reliability of
345 both datasets. This is consistent with the study of Boening et al. (2012) who have already found

good agreement between CloudSat and ERA Interim for the snowfall rate in Antarctica in the region 30°W-60°E, 65°S-80°S. Even if the spatiotemporal sampling of CloudSat is relatively low (between 350 and 500 orbits per grid cell over the Antarctic periphery for the period August 2006 - April 2011), the snowfall rate obtained with CloudSat is similar to the snowfall rate predicted by ERA
350 Interim during the same period (Fig. 5). Therefore, the spatiotemporal sampling of CloudSat seems to be sufficient to reproduce characteristics of Antarctic precipitation for the period August 2006 - April 2011. Extrapolating CloudSat snowfall observations to the region south of 82°S using ERA Interim data, the mean snowfall rate over the full Antarctic ice sheet amounts to 148 mm/year.

CloudSat does not provide any data during the night since April 2011. The EarthCARE satellite
355 scheduled for launch in 2015 into a polar orbit will carry a Cloud Profiling Radar (Kumagai et al., 2003). In addition to the reflectivity profiles, EarthCARE will measure the vertical Doppler velocity which will allow to get new informations about the cloud particles. Moreover, it will have a better sensitivity than CloudSat (-35dBZ compared to -28dBZ for CloudSat), and a better sampling interval (100m compared to 250m for CloudSat) (Nakatsuka et al., 2008). In situ observations are highly
360 desirable to evaluate and improve remote sensing techniques for Antarctic precipitation studies, and could be very useful during the EarthCARE mission. Future spaceborne radar missions should allow us to determine if Antarctic precipitation is increasing due to global warming as predicted by models.

Acknowledgements. This study was supported by the LEFE/CLAPA programme (Institut National des Sciences de l'Univers). This work was partially supported by funding by the ICE2SEA programme from the
365 European Union 7 framework programme, grant number 226375. Ice2Sea contribution number 176. Funding for this project was provided by a grant from la Région Rhône-Alpes. We are grateful to the Observatoire des Sciences de l'Univers de Grenoble (OSUG) which provided fundings for visits in France and in the USA. Parts of this research by N.B.W. and T.L. were performed at the University of Wisconsin - Madison for the Jet Propulsion Laboratory, California Institute of Technology, sponsored by the National Aeronautics and Space Administration. CloudSat data were obtained from the CloudSat Data Processing Center
370 (<http://www.cloudsat.cira.colostate.edu>). Finally, we thank the two anonymous reviewers for their comments which significantly improved the manuscript.

References

- Arthern, R. J., Winebrenner, D. P., and Vaughan, D. G.: Antarctic snow accumulation mapped using polarization of 4.3-cm wavelength microwave emission, *Journal of Geophysical Research: Atmospheres*, 111, doi:<http://dx.doi.org/10.1029/2004JD005667>, 2006.
- 375 Bamber, J. L., Gomez-Dans, J. L., and Griggs, J. A.: A new 1 km digital elevation model of the Antarctic derived from combined satellite radar and laser data – Part 1: Data and methods, *The Cryosphere*, 3, 101–111, doi:<http://dx.doi.org/10.5194/tc-3-101-2009>, 2009.
- 380 Barnston, A. G.: Correspondence among the Correlation, RMSE, and Heidke Forecast Verification Measures; Refinement of the Heidke Score, *Wea. Forecasting*, 7, 699–709, 1992.
- Bindschadler, R., Choi, H., Shuman, C., and Markus, T.: Detecting and measuring new snow accumulation on ice sheets by satellite remote sensing, *Remote Sensing of Environment*, 98, 388 – 402, 2005.
- Boening, C., Lebsack, M., Landerer, F., and Stephens, G.: Snowfall-driven mass change on the East Antarctic ice sheet, *Geophysical Research Letters*, 39, 2012.
- 385 Bromwich, D. H.: Snowfall in high southern latitudes, *Reviews of Geophysics*, 26, 149–168, 1988.
- Bromwich, D. H., Guo, Z., Bai, L., and Chen, Q.-S.: Modeled Antarctic Precipitation. Part I: Spatial and Temporal Variability, *J. Climate*, 17, 427–447, 2004.
- Bromwich, D. H., Nicolas, J. P., and Monaghan, A. J.: An Assessment of Precipitation Changes over Antarctica and the Southern Ocean since 1989 in Contemporary Global Reanalyses*, *J. Climate*, 24, 4189–4209, doi:<http://dx.doi.org/doi:10.1175/2011JCLI4074>.doi:10.1175/2011JCLI4074.1, 2011.
- 390 Church, J. A., Clark, P. U., Cazenave, A., Gregory, J. M., Jevrejeva, S., Levermann, A., Merrifield, M. A., Milne, G. A., Nerem, R. S., Nunn, P. D., Payne, A. J., Pfeffer, W. T., Stammer, D., and Unnikrishnan, A. S.: Sea Level Change., in: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by Stocker, T. F., Qin, D., Plattner, G. K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA., 2013.
- 395 Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, I., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J.-N., and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Quarterly Journal of the Royal Meteorological Society*, 137, 553–597, 2011.
- 400 Eisen, O., Frezzotti, M., Genthon, C., Isaksson, E., Magand, O., van den Broeke, M. R., Dixon, D. A., Ekaykin, A., Holmlund, P., Kameda, T., Karlöf, L., Kaspari, S., Lipenkov, V. Y., Oerter, H., Takahashi, S., and Vaughan, D. G.: Ground-based measurements of spatial and temporal variability of snow accumulation in East Antarctica, *Reviews of Geophysics*, 46, 2008.
- 405 Favier, V., Agosta, C., Parouty, S., Durand, G., Delaygue, G., Gallée, H., Drouet, A.-S., Trouvilliez, A., and Krinner, G.: An updated and quality controlled surface mass balance dataset for Antarctica, *The Cryosphere*, 7, 583–597, doi:<http://dx.doi.org/10.5194/tc-7-583-2013>, 2013.
- 410 Frezzotti, M., Scarchilli, C., Becagli, S., Proposito, M., and Urbini, S.: A synthesis of the Antarctic surface

- mass balance during the last 800 yr, *The Cryosphere*, 7, 303–319, doi:http://dx.doi.org/10.5194/tc-7-303-201310.5194/tc-7-303-2013, 2013.
- 415 Fujita, K. and Abe, O.: Stable isotopes in daily precipitation at Dome Fuji, East Antarctica, *Geophysical Research Letters*, 33, doi:http://dx.doi.org/10.1029/2006GL026936.1029/2006GL026936, 2006.
- Genthon, C. and Krinner, G.: Antarctic surface mass balance and systematic biases in general circulation models, *Journal of Geophysical Research: Atmospheres*, 106, 20 653–20 664, 2001.
- Genthon, C., Krinner, G., and Castebrunet, H.: Antarctic precipitation and climate change predictions: Horizontal resolution and margin vs plateau issues, *Annals of Glaciology*, 50, 55–60, 2009a.
- 420 Genthon, C., Magand, O., Krinner, G., and Fily, M.: Do climate models underestimate snow accumulation on the Antarctic plateau? A re-evaluation of/from in situ observations in East Wilkes and Victoria Lands, *Annals of Glaciology*, 50, 61–65, 2009b.
- Gregory, J. and Huybrechts, P.: Ice-sheet contributions to future sea-level change, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 364, 1709–1732, 2006.
- 425 Haynes, J. M., L'Ecuyer, T. S., Stephens, G. L., Miller, S. D., Mitrescu, C., Wood, N. B., and Tanelli, S.: Rainfall retrieval over the ocean with spaceborne W-band radar, *Journal of Geophysical Research: Atmospheres*, 114, doi:http://dx.doi.org/10.1029/2008JD009973.1029/2008JD009973, 2009.
- Hiley, M. J., Kulie, M. S., and Bennartz, R.: Uncertainty Analysis for CloudSat Snowfall Retrievals, *J. Appl. Meteor. Climatol.*, 50, 399–418, 2010.
- 430 Kulie, M. S. and Bennartz, R.: Utilizing Spaceborne Radars to Retrieve Dry Snowfall, *J. Appl. Meteor. Climatol.*, 48, 2564–2580, 2009.
- Kumagai, H., Kuroiwa, H., Kobayashi, S., and Orikasa, T.: Cloud profiling radar for EarthCARE mission, *Proc. SPIE*, 4894, 118–125, 2003.
- 435 Lenaerts, J. T. M., van den Broeke, M. R., van de Berg, W. J., van Meijgaard, E., and Kuipers Munneke, P.: A new, high-resolution surface mass balance map of Antarctica (1979–2010) based on regional atmospheric climate modeling, *Geophysical Research Letters*, 39, doi:http://dx.doi.org/10.1029/2011GL050713.1029/2011GL050713, 2012.
- Liu, G.: Deriving snow cloud characteristics from CloudSat observations, *Journal of Geophysical Research: Atmospheres*, 113, doi:http://dx.doi.org/10.1029/2007JD009766.1029/2007JD009766, 2008.
- 440 Magand, O., Genthon, C., Fily, M., Krinner, G., Picard, G., Frezzotti, M., and Ekaykin, A. A.: An up-to-date quality-controlled surface mass balance data set for the 90°–180°E Antarctica sector and 1950–2005 period, *Journal of Geophysical Research: Atmospheres*, 112, doi:http://dx.doi.org/10.1029/2006JD007691.1029/2006JD007691, 2007.
- 445 Magand, O., Picard, G., Brucker, L., Fily, M., and Genthon, C.: Snow melting bias in microwave mapping of Antarctic snow accumulation, *The Cryosphere*, 2, 109–115, doi:http://dx.doi.org/10.5194/tc-2-109-200810.5194/tc-2-109-2008, 2008.
- Monaghan, A. J., Bromwich, D. H., Fogt, R. L., Wang, S.-H., Mayewski, P. A., Dixon, D. A., Ekaykin, A., Frezzotti, M., Goodwin, I., Isaksson, E., Kaspari, S. D., Morgan, V. I., Oerter, H., Van Ommen, T. D., Van der Veen, C. J., and Wen, J.: Insignificant Change in Antarctic Snowfall Since the International Geophysical Year, *Science*, 313, 827–831, 2006a.
- 450 Monaghan, A. J., Bromwich, D. H., and Wang, S.-H.: Recent trends in Antarctic snow accumulation from Polar

MM5 simulations, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 364, 1683–1708, doi:<http://dx.doi.org/10.1098/rsta.2006.1795>, 2006b.

455 Nakatsuka, H., Okada, K., Horie, H., Kimura, T., Iida, Y., Kojima, M., Sato, K., Ohno, Y., Takahashi, N., and Kumagai, H.: System Design of Cloud Profiling Radar for Earthcare, in: *Geoscience and Remote Sensing Symposium*, 2008. *IGARSS 2008. IEEE International*, vol. 5, pp. V – 93–V – 96, doi:<http://dx.doi.org/10.1109/IGARSS.2008.4780035>, 2008.

Rodgers, C.: *Inverse methods for atmospheric sounding*, vol. 2 of *Series on Atmospheric, Oceanic and Planetary*
 460 *Physics*, World Scientific, 2000.

Simmons, A., Uppala, S., Dee, D., and Kobayashi, S.: ERA-interim: new ECMWF re-analysis products from 1989 onwards, *ECMWF Newslett*, 110, 25–35, 2006.

Stephens, G. L., Vane, D. G., Tanelli, S., Im, E., Durden, S., Rokey, M., Reinke, D., Partain, P., Mace, G. G., Austin, R., L'Ecuyer, T., Haynes, J., Lebsock, M., Suzuki, K., Waliser, D., Wu,
 465 D., Kay, J., Gettelman, A., Wang, Z., and Marchand, R.: CloudSat mission: Performance and early science after the first year of operation, *Journal of Geophysical Research: Atmospheres*, 113, doi:<http://dx.doi.org/10.1029/2008JD009982>, 2008.

Van de Berg, W., Van den Broeke, M., Reijmer, C., and Van Meijgaard, E.: Characteristics of the Antarctic surface mass balance, 1958-2002, using a regional atmospheric climate model, *Annals of Glaciology*, 41,
 470 97–104, doi:<http://dx.doi.org/doi:10.3189/172756405781813302>, 2005.

Wood, N. B.: Estimation of snow microphysical properties with application to millimeter-wavelength radar retrievals for snowfall rate., Ph.d. dissertation, Colorado State University, Fort Collins, Colorado, 231pp, available from <http://hdl.handle.net/10217/48170>., 2011.

Wood, N. B., L'Ecuyer, T., Vane, D. G., Stephens, G. L., and Partain, P.: Level 2C snow profile process description and interface control document, version 0., Tech. rep., http://www.cloudsat.cira.colostate.edu/ICD/2C-SNOW-PROFILE/2C-SNOW-PROFILE_PDICD_P_R04.pdf., 2013.

475




Fig. 1. a) Total number of orbits per grid cell from August 2006 to April 2011. b) The 2250m elevation contour derived from the digital elevation model of Bamber et al., 2009. The part of the ice sheet with surface elevation over 2250m (red), and below 2250m (purple). The black dot indicates the location of Dumont d'Urville station.




Fig. 2. a) Precipitation frequency (%) with the samples indicating precipitation certain for the period August 2006 - April 2011. b) Precipitation frequency (%) with the samples indicating precipitation certain and possible for the period August 2006 - April 2011. c) Ratio of the number of samples indicating precipitation possible over the number of samples indicating precipitation certain and possible.




Fig. 3. a) Mean annual snowfall rate (mm water equivalent / year) from the 2C-SNOW-PROFILE product for the period August 2006 - April 2011. b) Mean single retrieval uncertainty from the 2C-SNOW-PROFILE product for the period August 2006 - April 2011. c) The ratio of the single retrieval uncertainty over the snowfall rate.




Fig. 4. Comparison between the precipitation samples from CloudSat and ERA Interim reanalysis for the period August 2006 - April 2011. In ERA Interim reanalysis, precipitation events were defined for a precipitation rate over 0.07 mm / 6 h. a) Heidke Skill Score assessed using samples indicating precipitation certain and period without precipitation. b) Proportion of samples indicating periods without precipitation that match with periods without precipitation in ERA Interim. c) Proportion of samples indicating precipitation certain that match with precipitation events in ERA Interim. d) Proportion of samples indicating precipitation possible that match with precipitation events in ERA Interim.




Fig. 5. a) Mean annual snowfall rate (mm water equivalent / year) from CloudSat from August 2006 to April 2011. b) Mean annual snowfall rate (mm water equivalent / year) from ERA Interim reanalysis from August 2006 to April 2011. c) Mean annual snowfall rate (mm water equivalent / year) from ERA Interim with the same temporal sampling as CloudSat.




Fig. 6. Map : ratio of the snowfall rate observed by CloudSat for the period August 2006 - April 2011 over the accumulation rate determined by Arthern et al. (2006) for the period 1950 - 2000 (the isoline 1 is shown on the map). Dots : ratio of the snowfall rate observed by CloudSat for the period August 2006 - April 2011 over the accumulation rate from Favier et al. (2013).

Table 1. Comparison between the precipitation samples from CloudSat and ERA Interim reanalysis at Dumont d’Urville for the period August 2006 - April 2011. In ERA Interim reanalysis, precipitation events were defined for a precipitation rate over 0.07 mm / 6 h. The success rate is the proportion of samples indicating a situation (precipitation / no precipitation) that match with the same situation in ERA Interim. For the precipitation possible, the success rate is the proportion of samples indicating precipitation possible that match with precipitation events in ERA Interim.

Detection	Number of samples	Success rate
Period without precipitation	265	92%
Precipitation certain	85	91%
Precipitation possible	38	55%

Table 2. Comparison between the snowfall rate (mm/year) from CloudSat and ERA Interim reanalysis for the period August 2006 - April 2011, and the accumulation rate (mm/year) from Arthern et al. (2006) for the period 1950 - 2000. All the rates given in this table are averaged over the surface observed by CloudSat (latitude < 82°S).

	Continent	Altitude > 2250m	Altitude < 2250m
Snowfall rate from CloudSat	171	36	303
Snowfall rate from ERA Interim	163	49	273
Snowfall rate from ERA Interim (sampling of CloudSat)	163	53	271
Accumulation rate from Arthern et al.	163	81	243