



## Abstract

This study investigates the soil organic carbon (SOC) storage in Tarfala Valley, Northern Sweden. Field inventories upscaled based on land cover show that this alpine permafrost environment does not store large amounts of SOC, with an estimate mean of  $0.9 \pm 0.2 \text{ kg C m}^{-2}$  for the upper meter of soil. This is one to two orders of magnitude lower than what has been reported for lowland permafrost terrain. The SOC storage varies for different land cover classes and ranges from  $0.05 \text{ kg C m}^{-2}$  for stone-dominated to  $8.4 \text{ kg C m}^{-2}$  for grass-dominated areas. No signs of organic matter burial through cryoturbation or slope processes were found and radiocarbon dated SOC is generally of recent origin ( $< 2000 \text{ cal yr BP}$ ). An inventory of permafrost distribution in Tarfala Valley, based on bottom temperature of snow measurements and a logistic regression model, showed that at an altitude where permafrost is probable, the SOC storage is very low. In the high altitude permafrost zones (above 1500 m), soils store only ca  $0.1 \text{ kg C m}^{-2}$ . Under future climate warming an upward shift of vegetation zones may lead to a net ecosystem C uptake from increased biomass and soil development. As a consequence, alpine permafrost environments could act as a net carbon sink in the future, as there is no loss of older or deeper SOC from thawing permafrost.

## 1 Introduction

The permafrost-affected area in the northern circumpolar region is widespread, occupying about 18.78 million  $\text{km}^2$  (Tarnocai et al., 2009). The soils in the northern permafrost region store large amounts of soil organic carbon (SOC), which are vulnerable to climate change. With a warming climate, which is expected to be most pronounced in northern high latitudes, thawing permafrost soils may cause remobilization of soil organic matter (SOM) previously protected in permafrost (Gruber et al., 2004; Schuur et al., 2008). This can lead to an increased microbial decomposition of SOM and a release of carbon dioxide ( $\text{CO}_2$ ) and methane ( $\text{CH}_4$ ) to the atmosphere. As a

### Low soil organic carbon storage in a subarctic alpine permafrost environment

M. Fuchs et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Low soil organic carbon storage in a subarctic alpine permafrost environment

M. Fuchs et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

consequence, permafrost soils may act as a future carbon (C) source and lead to a positive climate feedback. However, the total storage of SOC within the northern permafrost region and the amount of greenhouse gases that can be released to the atmosphere and trigger accelerated climate warming are still uncertain (Schuur et al., 2009; Kuhry et al., 2010; McGuire et al., 2010; Schuur et al., 2013).

Several local to regional scale studies have been carried out to investigate stocks of SOC in northern permafrost environments, e.g. Michaelson et al. (1996), Kuhry et al. (2002), Zimov et al. (2006a), Ping et al. (2008), Horwath Burnham and Sletten (2010) and Hugelius et al. (2010, 2011). Tarnocai et al. (2009) estimated the 0–300 cm SOC stock in the northern permafrost region to 1024 Pg C, based on the Northern Circumpolar Soil Carbon Database (NCSCD). However, many regions in the NCSCD are under-represented and contain few sampled pedons, leading to a much generalized estimation of the C stocks for some remote areas (Mishra et al., 2013). Especially in the Eurasian and the High Arctic sectors, confidence for C estimates is low (Tarnocai et al., 2009).

Limited data are available for the more mountainous areas in the northern permafrost region. This study presents a detailed SOC inventory for a subarctic alpine permafrost environment by investigating the C stocks in soils of the Tarfala Valley, Northern Sweden. It is essential to establish to what extent these type of environments contribute to the large SOC storage in the northern permafrost region. Mountain areas and alpine permafrost are sensitive to climate change due to steep ecoclimatic gradients. The aim of this study is to assess the permafrost extent and SOC pools in a subarctic alpine environment, and evaluate their potential fate under conditions of future global warming.

## 2 Study area

Tarfala Valley is located in the Scandes mountains of northern Sweden, at c. 67°55' N and 18°37' E. The study area (31.2 km<sup>2</sup>) is delineated based on the catchment of Tarfala River (Tarfalajåkk), which drains into the broader Ladstjovagge. It includes the

alluvial fan of the Tarfala River to encompass the entire altitudinal gradient from the source to the outlet of Tarfala River. The area ranges between 550 and 2100 m above present sea level (a.p.s.l.) and is characterized in the upper part by six glaciers that drain into Tarfala River (Fig. 1).

The mean annual air temperature (MAAT) at Tarfala Research Station is  $-3.4^{\circ}\text{C}$  (1965–2009) and the mean annual precipitation for the Tarfala River catchment is 1997 mm (Dahlke et al., 2012). The MAAT in Tarfala has increased by  $0.54^{\circ}\text{C}$  per decade for the period 1969–2009, whereas the mean annual precipitation did not change significantly (Dahlke et al., 2012). The mean altitudinal lapse rate between Tarfala Research Station (1135 m a.p.s.l.) and the mountain saddle (Tarfalaryggen) along the eastern border of the study area (1540 m a.p.s.l.) is c.  $4.5^{\circ}\text{C km}^{-1}$ , however, the lapse rate in the summer months (JJA) of around  $5.8^{\circ}\text{C km}^{-1}$  is significantly higher than the winter lapse rate (DJF) of around  $2.7^{\circ}\text{C km}^{-1}$  (Jonsell et al., 2013).

The vegetation cover in the study area is generally sparse. In high elevation areas there is mostly barren ground. The middle part of the valley, around the Tarfala Research Station (1135 m a.p.s.l.), is characterized by patchy boulder fields and shallow soils with a mix of bare rocks, grasses, mosses and lichen. Further down the valley, dwarf shrubs (mainly *Salix* species and *Empetrum hermaphroditum*) appear up to 1000 m a.p.s.l. and the mountain birch forest (*Betula pubescens* ssp. *czerepanovii*) reaches up to c. 750 m a.p.s.l. On the alluvial fan, in the lowest part of the study area, the vegetation consists of a mix of deciduous and evergreen shrubs, graminoids and herbs.

The Tarfala Valley is characterized by little and very shallow soil development. The predominant soil types in the study area are classified as Leptosols and Regosols (IUSS Working Group WRB, 2006). On Tarfalaryggen, frost-thaw cycles have led to patterned ground formation and in these areas soils are Turbic Cryosols. In riverbeds of glacial streams (e.g. in the glacier forefield of Isfallsglaciären) soils are classified as Fluvisols.

## Low soil organic carbon storage in a subarctic alpine permafrost environment

M. Fuchs et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Low soil organic carbon storage in a subarctic alpine permafrost environment

M. Fuchs et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Extensive research has been carried out in Tarfala Valley, focusing mainly on glaciology and permafrost. Glaciers are the main subject of studies, with Storglaciären having the longest ongoing glacier mass balance measurements in the world (Holmlund et al., 2005). According to Brown et al. (1997) Tarfala Valley is located in the discontinuous permafrost zone. A permafrost borehole installed by the PACE (Permafrost and Climate in Europe) project is situated at 1540 m a.p.s.l. on Tarfalaryggen (Harris et al., 2001). The borehole measures the soil temperature down to 100 m every 6 h (Sollid et al., 2000). Mean annual ground temperature at the depth of zero annual amplitude is  $-2.8^{\circ}\text{C}$ , with a mean active layer depth of 1.5–1.6 m. Permafrost is currently not present in a 15 m deep borehole located at an elevation of 1135 m a.p.s.l. near Tarfala Research Station (Bolin Centre for Climate Research, 2013). King (1984) reports an active layer depth of 2.5–4 m in the valley floor around 1200 m a.p.s.l. Even though many scientific studies have been carried out in Tarfala Valley (e.g., Stork, 1963 on vegetation cover; King, 1984; Isaksen et al., 2007 on permafrost; Holmlund et al., 2005; Jansson and Pettersson, 2007 on glaciology; Dahlke et al., 2012 on hydrology), there are no previous studies on SOC storage from this area.

### 3 Methods

#### 3.1 Soil sampling

In August 2012, a stratified-random sampling program was executed in the Tarfala Valley during which soil profiles were collected along five transects. Transects were chosen to represent the altitudinal zones and vegetation types in the valley. Individual profiles were placed at strict equidistant distances along the transects to introduce a degree of randomness in the sampling. Whenever possible, near surface organic layers and fine textured soils were collected from pits dug into the soils by cutting out samples of known volume. Deeper soil layers were sampled by hammering a steel pipe of c. 4 cm diameter into the soil at 5–10 cm increments. Most of the collected soil

profiles were shallow as the stony soils did in most cases not enable a sampling to the full reference depth of 100 cm. Furthermore, permafrost was never encountered during coring, even at high elevations, indicating generally deep active layers in Tarfala Valley. In total, 56 profile sites were sampled and described and 295 individual soil samples collected.

### 3.2 Land cover classification

A description of the vegetation cover in a ground truth plot (diameter 10 m) was made around each profile site, with special attention paid to the occurrence of stones and boulders (see description of SOC mass calculation below). For upscaling purposes, a land cover classification (LCC) was compiled from remotely sensed data. For this LCC, an orthophoto (compiled with ERDAS Imagine LPS from CIR aerial photographs with 0.5 m spatial resolution) (Lantmäteriet, 2008), a WorldView2 satellite image (European Space Imaging GMBH, 2012), and a Landsat 5TM (USGS, 2011) satellite image were used. The remote mountainous area as well as cloud- and snow cover in the images made a usage of different datasets unavoidable to cover the whole valley. The LCC includes nine different classes which have been separated by a combination of a 3-D stereo analysis and supervised classification (maximum likelihood). The requirements for a supervised classification in general and the training areas in particular followed Campbell (2011). To verify the classification, the kappa index of agreement was calculated based on the 56 ground truth plots. The nine land cover classes recognized for Tarfala Valley are presented in the Supplement (Table S1).

### 3.3 Geochemical analyses

Soil samples of known volume were weighed in the laboratory after oven drying at 60 °C (for 48 h) to calculate dry bulk density (DBD,  $\text{g cm}^{-3}$ ). For loss on ignition (LOI), samples were burned at 550 °C for 6 h to determine the organic carbon content and at 950 °C for 2 h to determine the carbonate content (Dean, 1974; Heiri et al., 2001). In addition,

TCD

8, 3493–3524, 2014

## Low soil organic carbon storage in a subarctic alpine permafrost environment

M. Fuchs et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





## Low soil organic carbon storage in a subarctic alpine permafrost environment

M. Fuchs et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

made based on field observations. The mean SOC storage for each of the recognized land cover types is calculated as the arithmetic mean of all soil profiles representing those land cover types. To avoid overestimation of the C content, each LCC mean SOC kg C m<sup>-2</sup> value was weighted by the mean percentage of large stones (> 4 cm diameter) visible at the surface. These areas were considered to have no soil development and to contain no SOC. The coverage of large stones was derived by field observations at every sample spot within a radius of five meters. Thereafter, the mean SOC storage in Tarfala Valley was calculated, based on the proportions of the land cover classes in the LCC. These calculations were performed for all land cover classes together (including glaciers, barren grounds and lakes) and for the vegetated classes only.

### 3.5 Statistical methods

The results from the geochemical analyses and the upscaling were further analyzed with statistical methods. All statistical analyses were carried out with the open source statistical analysis package PAST 2.17 (Hammer et al., 2001). Three main statistical analyses were carried out: (1) confidence intervals (CI) for the mean C estimates of the total study area were calculated according to Hugelius (2012); (2) linear correlations (Pearson's correlation) between soil depth and the different geochemical parameters (DBD, %C, LOI, C/N-ratio,  $\delta^{13}\text{C}$ ,  $\delta^{15}\text{N}$ ) were calculated to examine whether the different parameters decrease or increase significantly with increasing depth; (3) the student's *t* test was applied to examine if there is a statistically significant difference between the organic rich top soil and the underlying mineral samples for all the different geochemical parameters. In all cases, the probability limit of  $p \leq 0.01$  was chosen for statistical significance.

### 3.6 Permafrost mapping

In addition to the SOC inventory, the permafrost distribution in Tarfala Valley was mapped. Bottom temperature of snow (BTS) measurements were carried out in







be considered that only a very minor to negligible amount of organic C is stored within the permafrost layer. It should be noted that permafrost was never reached during field coring due to the occurrence of bare rock and stones.

The soils of Tarfala Valley display no signs of cryoturbation of the organic rich top soil layer into the deeper mineral soil horizons. Likewise, no burial of the organic rich layer due to solifluction processes on slopes was observed.

### 4.3 SOM quality and age

The soils in Tarfala Valley are characterized by a steady, statistically significant ( $p < 0.01$ ) increase in bulk density with depth (Fig. 3a; Table 2). However, LOI (550 °C) and percentage C show strong, statistically significant ( $p < 0.01$ ) negative correlations with depth (Fig. 3b; Table 2). As a result, there is less SOM with greater depth in the soil. There is also a statistically significant ( $t$  test,  $p < 0.01$ ) difference in the C content of the organic-rich top soil layer and the underlying mineral layer (Table 2).

Besides C content, other geochemical analyses of the soil samples also show a coherent picture. The C/N ratio and stable isotopic composition of SOM reflect its relative state of decomposition (e.g. Mariotti and Balesdent, 1990; Kuhry and Vitt, 1996; Ping et al., 1998; Hugelius et al., 2012). There is a statistically significant ( $t$  test,  $p < 0.01$ ) difference between the mean C/N ratio of the organic rich top soil layer ( $23.3 \pm 11.4$ ) and that of the mineral layer ( $14.6 \pm 4.05$ ). The C/N ratio decreases with increasing depth ( $p < 0.01$ ), indicating progressively more decomposed SOM (Fig. 3c). Ping et al. (1998) pointed out that the C/N ratio is dependent on vegetation cover and that trends need to be interpreted carefully. In the Tarfala Valley, the decrease of C/N ratio with depth is consistent across all land cover classes. However these trends are not statistically significant for the separate land cover classes, probably due to the limited number of replicates within each class (data not shown). The stable isotope composition of  $\delta^{13}\text{C}$  vs. PDB and  $\delta^{15}\text{N}$  vs. air show statistical significant ( $p < 0.01$ ) enrichment of stable isotopes with increasing soil depth (Fig. 3d; Table 2). The enrichment of  $\delta^{13}\text{C}$

## Low soil organic carbon storage in a subarctic alpine permafrost environment

M. Fuchs et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and  $\delta^{15}\text{N}$  with depth can be considered an indication for SOM degradation through microbial respiration (Mariotti and Balesdent, 1990; Ping et al., 1998).

Four bulk soil samples (living roots removed) from two profiles belonging to the class “Patchy Boulder Grass/Moss”, located close to the floor of the central Tarfala Valley, have been radiocarbon-dated (Table 3). These profiles were selected because they had the thickest organic-rich top soil layer among the collected profiles in the study area and displayed a slight, but highly unusual for this area, C-enrichment in the underlying mineral soil (weak B-horizon development). Results indicate that the SOM close to the surface is recent in age (< 100 yr old), whereas the mineral soil at greater depths contain slightly older SOM, with ages of 1269 and 1919 cal yr BP (Table 3). Considering the fact that the two dated profiles are among the most well-developed soils in the study area, it is most likely that most of the SOM in Tarfala Valley is of very young age. The geochemistry of these two dated profiles, which reflect the general trends described for the whole dataset, is presented in the Supplement (Fig. S2).

#### 4.4 Permafrost mapping

Permafrost zones are commonly separated into the classes continuous, discontinuous, sporadic and isolated patches (e.g. Brown et al., 1997). However with the logistic regression approach, not the areal extent of permafrost but the probability for the occurrence of permafrost was used to map the permafrost distribution into the conventional classes (Fig. 4). This was already applied by Lewkowicz and Ednie (2004) in their study in the Yukon Territory, Canada. However, with this approach, the permafrost distribution has to be interpreted carefully, especially in a highly heterogeneous alpine environment like Tarfala Valley. Areas with a > 90 % probability for the occurrence of permafrost are considered as continuous, which in Tarfala Valley includes all areas above 1561 m a.p.s.l. The discontinuous permafrost zone (probability between 50–90 %) occurs at an altitude between 1218 and 1561 m a.p.s.l., while the sporadic permafrost zone commences at an altitude above 875 m a.p.s.l. (probability > 10 %).

### Low soil organic carbon storage in a subarctic alpine permafrost environment

M. Fuchs et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion







## Low soil organic carbon storage in a subarctic alpine permafrost environment

M. Fuchs et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Geochemical indicators, such as C/N ratios and stable isotopes ( $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$ ), indicate that the SOM in Tarfala soils becomes gradually more decomposed with depth and age. Cryoturbation of C-enriched material is one of the mechanisms that significantly increase SOC storage in permafrost soils (e.g., Ping et al., 2008). In Tarfala, we did not find evidence for burial of relatively undecomposed SOM from the organic rich top soil layer deeper into the profiles. The two dated soil profiles are exceptional for Tarfala Valley as they have the thickest organic-rich top soil layer and relatively high carbon values in greater depths. But basal dates for even these thickest organic rich top soil layers are recent and the SOC at greater depth is also quite young ( $< 2000$  cal yr BP). Therefore, much of the SOM in Tarfala Valley seems to be cycled within 100 yr or less and does not accumulate at greater depths. This is in stark contrast with permafrost soils from lowland regions, which are reported to have extensive cryoturbation of relatively undecomposed SOM that has been preserved at greater soil depths for thousands of years (e.g., Bockheim, 2007; Hugelius et al., 2010).

### 5.2 Future developments

Our results indicate that there are no large amounts of SOC stored in the soils of Tarfala Valley. The relatively highest mean SOC storage is found in vegetated ground at lower elevations. A further analysis that takes into account the permafrost zonation shows that the potential for SOC storage in permafrost affected soils is very small (Fig. 5). The mean SOC value at an elevation of 1250 m a.p.s.l., where the probability for permafrost is just above 50 %, is  $0.7 \text{ kg C m}^{-2}$  (for 0–100 cm) and at an altitude of 1500 m a.p.s.l. (permafrost probability 85 %) it is only  $0.1 \text{ kg C m}^{-2}$ . Therefore, most of the SOC in Tarfala Valley is stored at lower elevations where the probability for permafrost affected soils is low. Taking into account that the active layer is 2.5–4 m thick in the valley floor around 1200 m a.p.s.l. (King, 1984) and the fine soil is only rarely deeper than 1 m, the amount of SOC stored in the permafrost layer is assumed to be negligible.



## Low soil organic carbon storage in a subarctic alpine permafrost environment

M. Fuchs et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

The vegetation and SOC distribution in Tarfala Valley allow some considerations about future total ecosystem C storage in the area under conditions of global change. Climate warming will result in an upwards shift of vegetation zones with the corresponding initiation of soil development in currently high-alpine barren areas. Upwards altitudinal shifts of plants due to increased temperatures have been observed in alpine regions (e.g., Walther et al., 2005), including the Scandinavian mountain range (e.g., Klanderud and Birks, 2003; Kullman, 2002, 2010). Kullman and Öberg (2009) report an altitudinal upward shift of trees of about 200 m in the past 100 yr in the Swedish Scandes, in accordance with observed temperature increases. For a first rough estimation of potential upwards shifts of vegetation zones, the mean summer temperature change was taken as a first indicator, even though many other factors will affect the vegetation (e.g. winter temperatures, precipitation, windexposition, etc., Kullman, 2010). The projected mean summer (JJA) temperature increase for the Tarfala mountain region until 2100 is 2.8 °C (SRES A1B scenario, SMHI, 2013). Considering a summer lapse rate of 5.8 °C km<sup>-1</sup> (Jonsell et al., 2013), the potential altitudinal upward shift for the vegetation cover is c. 500 m. Grace et al. (2002) and Kullman (2010) calculated a similar potential treeline shift in the region by the end of this century. However, not the entire Tarfala Valley will be suitable for plant colonization, because of steep slopes, a lack of fine soil matrix and wind-exposed ridges.

Schuur et al. (2009) showed that in the Alaskan tundra, increased plant productivity is eventually outweighed by increased decomposition of deeper and older SOM following permafrost thaw. For projections of permafrost degradation in Tarfala Valley, the mean annual air temperature has to be considered. A climate scenario for the Tarfala mountain region estimates a mean annual temperature increase of c. 4.6 °C until 2100 (SRES A1B scenario, SMHI, 2013). Taking into account a mean annual lapse rate of 4.5 °C km<sup>-1</sup> (Jonsell et al., 2013) the 0 °C air temperature isotherm could rise with c. 1000 m, which would greatly affect permafrost occurrence in the area. Data from the PACE borehole at Tarfalaryggen shows that the permafrost temperature at the zero annual amplitude depth of 20 m has already experienced a warming of 0.047 °C yr<sup>-1</sup>



## Low soil organic carbon storage in a subarctic alpine permafrost environment

M. Fuchs et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

(Jonsell et al., 2013). Even though future permafrost degradation is highly plausible for most of the upper Tarfala Valley, only a negligible amount of SOC is currently stored in the area and could be affected by thaw. Under future climate warming and permafrost thawing little or no SOC will be remobilized from permafrost soils in Tarfala Valley. On the contrary, increased temperatures will lead to an upward vegetation shift, phytomass production and soil development, with the result of an increased C uptake in Tarfala Valley in the future. The only way that projected permafrost thaw might negatively affect C uptake is through an initial increased slope instability in steep terrain (Gregory and Goudie, 2011; French, 2007).

Compared to lowland permafrost regions in the northern circumpolar region (see e.g. Gruber et al., 2004; Zimov et al., 2006b; Schuur et al., 2009), a subarctic high-alpine permafrost environment like the upper Tarfala Valley cannot be considered a future source of C to the atmosphere. In general, alpine permafrost environments above the contiguous vegetation limit have the potential of becoming a C sink in the future and therefore stand out as an exception in the general assessment of thawing permafrost soils representing an important positive feedback to future climate warming (e.g., Schuur et al., 2013).

## 6 Conclusion

The SOC inventory in Tarfala Valley, with a mean storage of  $0.9 \text{ kg C m}^{-2}$  for the upper meter of soil, shows that this area cannot be considered a C-rich permafrost environment. This low value is a result of the high amount of barren ground and stony surfaces in the study area, low plant productivity, shallow soils, and lack of SOM burial through cryoturbation or slope processes. The low SOC storage leads to the conclusion that environments like Tarfala Valley cannot become significant sources of C with future permafrost thawing. Instead, they could act as net C sinks following an upward shift of vegetation zones causing increased phytomass production, soil development and SOM accumulation. The potential magnitude of an increased C uptake in this type of

mountainous permafrost region remains to be addressed by further studies. Nevertheless, this study shows that there is a need to include alpine environments to estimate the total SOC stock in permafrost soils of the northern circumpolar region and to fully assess the permafrost thaw-C feedback.

5 **The Supplement related to this article is available online at doi:10.5194/tcd-8-3493-2014-supplement.**

*Acknowledgements.* We wish to thank E. L. Palmtag, J. Palmtag, N. Weiss and M. Brandel for help with the fieldwork and A. Bergström and his team for the support at Tarfala Research Station. The study was supported by the Nordic Center of Excellence DEFROST project.

## 10 **References**

- Bockheim, J. G.: Importance of cryoturbation in redistributing organic carbon in permafrost-affected soils, *Soil Sci. Soc. Am. J.*, 71, 1335–1342, doi:10.2136/sssaj2006.0414N, 2007. Bolin Centre for Climate Research: Bolin Centre Database, Tarfala data, permafrost monitoring data, <http://bolin.su.se/data/tarfala/permafrost.php>, last access: 30 October 2013.
- 15 Bronk Ramsey, C.: OxCal v4.1.7, Radiocarbon Calibration Software, Research Lab for Archaeology, Oxford, UK, available at: <http://c14.arch.ox.ac.uk> (last access: 13 February 2013), 2010.
- Brown, J., Ferrians Jr., O. J., Heginbottom, J. A., and Melnikov, E. S.: Circum-Arctic map of permafrost and ground-ice conditions, 1 : 10 000 000, Map CP-45, United States Geological Survey, International Permafrost Association, 1997.
- 20 Campbell, J. B.: Introduction to remote sensing, 5. Edn., Guilford Press, New York, 667 pp., 2011.
- Dahlke, H. E., Lyon, S. W., Stedinger, J. R., Rosqvist, G., and Jansson, P.: Contrasting trends in floods for two sub-arctic catchments in northern Sweden – does glacier presence matter?, *Hydrol. Earth Syst. Sci.*, 16, 2123–2141, doi:10.5194/hess-16-2123-2012, 2012.
- 25

### Low soil organic carbon storage in a subarctic alpine permafrost environment

M. Fuchs et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Low soil organic carbon storage in a subarctic alpine permafrost environment**

M. Fuchs et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Dean, W. E.: Determination of carbonate and organic matter in calcareous sediments and sedimentary rocks by loss on ignition: comparison with other methods, *J. Sediment. Petrol.*, 44, 242–248, 1974.

Doerfer, C., Kuehn, P., Baumann, F., He, J.-S., and Sholten, T.: Soil organic carbon pools and stocks in permafrost-affected soils on the Tibetan Plateau, *PLOS ONE*, 8, e57024, doi:10.1371/journal.pone.0057024, 2013.

Dostmann Electronic: Temperature Probe P400, <http://www.dostmann-electronic.de/englisch/Produkte/produkt.php?pid=205&lang=eng>, last access: 6 May 2013.

European Space Imaging GMBH: WorldView2 satellite image, date of acquisition: 4 July 2012, <http://www.euspaceimaging.com/> (last access: 6 May 2013), 2012.

French, H. M.: *The Periglacial Environment*, 3rd Edn., John Wiley & Sons Ltd, West Sussex, England, 480 pp., doi:10.1002/9781118684931, 2007.

Grace, J., Berninger, F., and Nagy, L.: Impacts of climate change on tree line, *Ann. Bot.-London*, 90, 537–544, 2002.

Gregory, K. J. and Goudie, S. A.: *The SAGE Handbook of Geomorphology*, SAGE Publications Ltd, London, England, 648 pp., 2011.

Gruber, N., Friedlingstein, P., Field, C. B., Valentini, R., Heimann, M., Richey, J. E., Romero-Lankao, P., Schulze, D., and Chen, C. T. A.: The vulnerability of the carbon cycle in the 21st century: an assessment of carbon-climate-human interactions, in: *The Global Carbon Cycle: Integrating Humans, Climate, and the Natural World*, edited by: Field, C. and Raupach, M., Island Press, Washington, DC, 45–76, 2004.

Haeberli, W.: Die Basis-Temperatur der winterlichen Schneedecke als möglicher Indikator für die Verbreitung von Permafrost in den Alpen, *Zeitschrift für Gletscherkunde und Glaziologie*, 9, 221–227, 1973.

Hammer, Ø., Harper, D. A. T., and Ryan, P. D.: PAST: paleontological statistics software package for education and data analysis, *Palaeontol. Electron.*, 4, 9 pp., [http://palaeo-electronica.org/2001\\_1/past/issue1\\_01.htm](http://palaeo-electronica.org/2001_1/past/issue1_01.htm), 2001.

Harris, C., Haeberli, W., Vonder Muehll, D., and King, L.: Permafrost monitoring in the high mountains of Europe: the PACE project in its global context, *Permafrost Periglac.*, 12, 3–11, doi:10.1002/ppp.377, 2001.

Heiri, O., Lotter, A. F., and Lemcke, G.: Loss on ignition as a method for estimating organic and carbonate content in sediments: reproducibility and comparability of results, *J. Paleolimnol.*, 25, 101–110, doi:10.1023/A:1008119611481, 2001.

## Low soil organic carbon storage in a subarctic alpine permafrost environment

M. Fuchs et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Holmlund, P., Jansson, P., and Pettersson, R.: A re-analysis of the 58 year mass-balance record of Storglaciären, Sweden, *Ann. Glaciol.*, 42, 389–394, 2005.
- Horwath Burnham, J. and Sletten, R. S.: Spatial distribution of soil organic carbon in northwest Greenland and underestimates of high arctic carbon stores, *Global Biogeochem. Cy.*, 24, GB3012, doi:10.1029/2009GB003660, 2010.
- Hugelius, G.: Spatial upsampling using thematic maps: an analysis of uncertainties in permafrost soil carbon estimates, *Global Biogeochem. Cy.*, 26, GB2026, doi:10.1029/2011GB004154, 2012.
- Hugelius, G., Kuhry, P., Tarnocai, C., and Virtanen, T.: Soil organic carbon pools in a periglacial landscape: a case study from the central Canadian Arctic, *Permafrost Periglac.*, 21, 16–29, doi:10.1002/ppp.677, 2010.
- Hugelius, G., Virtanen, T., Kaverin, D., Pastukhov, A., Rivkin, F., Marchenko, S., Romanovsky, V., and Kuhry, P.: High-resolution mapping of ecosystem carbon storage and potential effects of permafrost thaw in periglacial terrain, European Russian Arctic, *J. Geophys. Res.*, 116, G03024, doi:10.1029/2010JG001606, 2011.
- Hugelius, G., Routh, J., Kuhry, P., and Crill, P.: Mapping the degree of decomposition and thaw remobilization potential of soil organic matter in discontinuous permafrost terrain, *J. Geophys. Res.*, 117, G02030, doi:10.1029/2011JG001873, 2012.
- Hugelius, G., Tarnocai, C., Broll, G., Canadell, J. G., Kuhry, P., and Swanson, D. K.: The Northern Circumpolar Soil Carbon Database: spatially distributed datasets of soil coverage and soil carbon storage in the northern permafrost regions, *Earth Syst. Sci. Data*, 5, 3–13, doi:10.5194/essd-5-3-2013, 2013.
- Isaksen, K., Sollid, J. L., Holmlund, P., and Harris, C.: Recent warming of mountain permafrost in Svalbard and Scandinavia, *J. Geophys. Res.*, 112, F02S04, doi:10.1029/2006JF000522, 2007.
- IUSS Working Group WRB: World reference base for soil resources, 2006, 2nd Edn., World Soil Resources Reports, 103, FAO, Rome, 143 pp., 2006.
- Jansson, P. and Pettersson, R.: Spatial and temporal characteristics of a long mass balance record, Storglaciären, Sweden, *Arct. Antarct. Alp. Res.*, 39, 432–437, 2007.
- Jonsell, U., Hock, R., and Duguay, M.: Recent air and ground temperature increases at Tarfala Research Station, Sweden, *Polar Res.*, 32, 19807, doi:10.3402/polar.v32i0.19807, 2013.
- King, L.: High mountain permafrost in Scandinavia, in: *Permafrost: Fourth International Conference, Proceedings*, Nat. Acad. Press, Washington, DC, 612–617, 1983.

## Low soil organic carbon storage in a subarctic alpine permafrost environment

M. Fuchs et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



King, L.: Permafrost in Skandinavien, Untersuchungsergebnisse aus Lappland, Jotunheimen und Dovre/Rondane, Heidelberger Geographische Arbeiten, Heidelberg, Germany, 76, 174 pp., 1984.

Klanderud, K. and Birks, H. J. B.: Recent increases in species richness and shifts in altitudinal distribution of Norwegian mountain plants, *The Holocene*, 13, 1–6, 2003.

Kuhry, P. and Vitt, D. H.: Fossil carbon/nitrogen ratios as a measure of peat decomposition, *Ecology*, 77, 271–275, doi:10.2307/2265676, 1996.

Kuhry, P., Mazhitova, G. G., Forest, P.-A., Deneva, S. V., Virtanen, T., and Kultti, S.: Upscaling soil organic carbon estimates for the Usa Basin (Northeast European Russia) using GIS-based landcover and soil classification schemes, *Dan. J. Geogr.*, 102, 11–25, 2002.

Kuhry, P., Dorrepaal, E., Hugelius, G., Schuur, E. A. G., and Tarnocai, C.: Potential remobilization of belowground permafrost carbon under future global warming, *Permafrost Periglac.*, 21, 208–214, doi:10.1002/ppp.684, 2010.

Kullman, L.: Rapid recent range-margin rise of tree and shrub species in the Swedish Scandes, *J. Ecol.*, 90, 68–77, 2002.

Kullman, L.: A richer, greener and smaller alpine world: review and projection of warming-induced plant cover change in the Swedish Scandes, *Ambio*, 39, 159–169, doi:10.1007/s13280-010-0021-8, 2010.

Kullman, L. and Öberg, L.: Post-little ice age tree line rise and climate warming in the Swedish Scandes: a landscape ecological perspective, *J. Ecol.*, 97, 415–429, doi:10.1111/j.1365-2745.2009.01488.x, 2009.

Lantmateriet: CIR aerial images 08o48zu50\_31~ 2008-09-10\_105654\_98–08o48zu50\_31~ 2008-09-10\_105654\_103, <http://www.lantmateriet.se/> (last access: 6 May 2013), 2008.

Lewkowicz, A. G. and Ednie, M.: Probability mapping of mountain permafrost using the BTS Method, Wolf Creek, Yukon Territory, Canada, *Permafrost Periglac.*, 15, 67–80, doi:10.1002/ppp.480, 2004.

Mariotti, A. and Balesdent, J.:  $^{13}\text{C}$  natural abundance as a tracer of soil organic matter turnover and paleoenvironment dynamics, *Chem. Geol.*, 84, 217–219, 1990.

Marklund, P.: Alpin permafrost i Kebnekaisefjällen: Modellering med logistic regression och BTS-data, Självständigt arbete i geovetenskap, Nr. 22, Uppsala University, Disciplinary Domain of Science and Technology, Earth Sciences, Department of Earth Sciences, LUVAL, 2011.

## Low soil organic carbon storage in a subarctic alpine permafrost environment

M. Fuchs et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

McGuire, A. D., Macdonald, R. W., Schuur, E. A. G., Harden, J. W., Kuhry, P., Hayes, D. J., Christensen, T. R., and Heimann, M.: The carbon budget of the northern cryosphere region, *Current Opinion in Environmental Sustainability*, 2, 231–236, 2010.

Michaelson, G. J., Ping, C. L., and Kimble, J. M.: Carbon storage and distribution in tundra soils of Arctic Alaska, USA, *Arctic Alpine Res.*, 28, 414–424, 1996.

Mishra, U., Jastrow, J. D., Matamala, R., Hugelius, G., Koven, C. D., Harden, J. W., Ping, C. L., Michaelson, G. J., Fan, Z., Miller, R. M., McGuire, A. D., Tarnocai, C., Kuhry, P., Riley, W. J., Schaefer, K., Schuur, E. A. G., Jorgenson, M. T., and Hinzman, L. D.: Empirical estimates to reduce modelling uncertainties of soil organic carbon in permafrost regions: a review of recent progress and remaining challenges, *Environ. Res. Lett.*, 8, 035020, doi:10.1088/1748-9326/8/3/035020, 2013.

Ohtsuka, T., Hirota, M., Zhang, X., Shimono, A., Senga, Y., Du, M., Yonemura, S., Kawashima, S., and Tang, Y.: Soil organic carbon pools in alpine to nival zones along an altitudinal gradient (4400–5300 m) on the Tibetan Plateau, *Polar Science*, 2, 277–285, 2008.

Ping, C. L., Bockheim, J. G., Kimble, J. M., Michaelson, G. J., and Walker, D. A.: Characteristics of cryogenic soils along a latitudinal transect in Arctic Alaska, *J. Geophys. Res.*, 103, D22, 28917–28928, doi:10.1029/98JD02024, 1998.

Ping, C. L., Michaelson, G. J., Jorgenson, M. T., Kimble, J. M., Epstein, H., Romanovsky, V. E., and Walker, D. A.: High stocks of soil organic carbon in the North American arctic region, *Nat. Geosci.*, 1, 615–619, doi:10.1038/ngeo284, 2008.

Schuur, E. A. G., Bockheim, J., Canadell, J. G., Euskirchen, E., Field, C. B., Goryachkin, S. V., Hagemann, S., Kuhry, P., Fafleur, P. M., Lee, H., Mazhitova, G., Nelson, F. E., Rinke, A., Romanovsky, V. E., Shiklomanov, N., Tarnocai, C., Venevsky, S., Vogel, J. G., and Zimov, S. A.: Vulnerability of permafrost carbon to climate change: implications for the global carbon cycle, *Bioscience*, 58, 701–714, doi:10.1641/B580807, 2008.

Schuur, E. A. G., Vogel, J. G., Crummer, K. G., Lee, H., Sickman, J. O., and Osterkamp, T. E.: The effect of permafrost thaw on old carbon release and net carbon exchange from tundra, *Nature*, 459, 556–559, doi:10.1038/nature08031, 2009.

Schuur, E. A. G., Abbott, B. W., Bowden, W. B., Brovkin, V., Camill, P., Canadell, J. G., Chanton, J. P., Chapin III, F. S., Christensen, T. R., Ciais, P., Crosby, B. T., Czimczik, C. I., Grosse, G., Harden, J., Hayes, D. J., Hugelius, G., Jastrow, J. D., Jones, J. B., Kleinen, T., Koven, C. D., Krinner, G., Kuhry, P., Lawrence, D. M., McGuire, A. D., Natali, S. M., O'Donnell, J. A., Ping, C. L., Riley, W. J., Rinke, A., Romanovsky, V. E., Sannel, A. B. K.,

## Low soil organic carbon storage in a subarctic alpine permafrost environment

M. Fuchs et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Schädel, C., Schaefer, K., Sky, J., Subin, Z. M., Tarnocai, C., Turetsky, M. R., Waldrop, M. P., Walter Anthony, K. M., Wickland, K. P., Wilson, C. J., and Zimov, S. A.: Expert assessment of vulnerability of permafrost carbon to climate change, *Climatic Change*, 119, 359–374, doi:10.1007/s10584-013-0730-7, 2013.

5 SMHI, Sveriges meteorologiska och hydrologiska institut: Klimatscenarier, <http://www.smhi.se/klimatdata/Framtidens-klimat/Klimatscenarier/2.2252/2.2264>, last access: 8 November 2013.

Sollid, J. L., Holmlund, P., Isaksen, K., and Harris, C.: Deep permafrost boreholes in western Svalbard, northern Sweden, and southern Norway, *Norsk. Geogr. Tidsskr.*, 54, 186–191, 2000.

10 Stork, A.: Plant immigration in front of retreating glaciers, with examples from the Kebnekajse area, Northern Sweden, *Geogr. Ann.*, 45, 1–22, 1963.

Tarnocai, C., Canadell, J. G., Mazhitova, G., Schuur, E. A. G., Kuhry, P., and Zimov, S.: (2009), Soil organic carbon pools in the northern circumpolar permafrost region, *Global Biogeochem. Cy.*, 23, GB2023, doi:10.1029/2008GB003327, 2009.

15 US Geological Survey: Earth Resources Observation and Science Center (EROS), USGS Global Visualization Viewer, Landsat 5 TM satellite image, acquisition date: 18 September 2011, <http://glovis.usgs.gov/> (last access: 25 October 2012), 2011.

Walker, M.: *Quaternary Dating Methods*, John Wiley & Sons Ltd, England, 306 pp., 2005.

20 Walther, G.-R., Beissner, S., and Burga, C. A.: Trends in the upward shift of alpine plants, *J. Veg. Sci.*, 16, 541–548, 2005.

Wang, G., Li, Y., Wang, Y., and Wu, Q.: Effects of permafrost warming on vegetation and soil carbon pool losses on the Qinghai-Tibet Plateau, China, *Geoderma*, 143, 143–152, 2008.

25 Yang, Y., Fang, J., Tang, Y., Ji, C., Zheng, C., He, J., and Zhu, B.: Storage, patterns and controls of soil organic carbon in the Tibetan grasslands, *Glob. Change Biol.*, 14, 1592–1599, doi:10.1111/j.1365-2486.2008.01591.x, 2008.

Zimov, S. A., Schuur, E. A. G., and Chapin, F. S.: Permafrost and the global carbon budget, *Science*, 312, 1612–1613, doi:10.1126/science.1128908, 2006a.

30 Zimov, S. A., Davydov, S. P., Zimova, G. M., Davydova, A. I., Schuur, E. A. G., Dutta, K., and Chapin, F. S. II I.: Permafrost carbon: stock and decomposability of a globally significant carbon pool, *Geophys. Res. Lett.*, 33, L20502, doi:10.1029/2006GL027484, 2006b.

Zollinger, B., Alewell, C., Kneisel, C., Meusburger, K., Gärtner, H., Brandová, D., Ivy-Ochs, S., Schmidt, M. W. I., and Egli, M.: Effect of permafrost on the formation of soil organic carbon

pools and their physical-chemical properties in the Eastern Swiss Alps, Catena, 11, 70–85, doi:10.1016/j.catena.2013.06.010, 2013.

Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper

**TCD**

8, 3493–3524, 2014

**Low soil organic carbon storage in a subarctic alpine permafrost environment**

M. Fuchs et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





## Low soil organic carbon storage in a subarctic alpine permafrost environment

M. Fuchs et al.

**Table 1.** Mean soil organic carbon (SOC) storage and sample site characteristics for the different land cover classes in Tarfala Valley.

Land Cover Classes	Mean SOC Storage				Mean Profile Depth				Profile Distribution					
	0–30 cm (kg C m <sup>-2</sup> ± std)	0–100 cm (kg C m <sup>-2</sup> ± std)	Organic Rich Top Soil Layer (kg C m <sup>-2</sup> ± std)	Mineral Layer (kg C m <sup>-2</sup> ± std)	Profile Site (cm ± std)	Organic Rich Top Soil Layer (cm ± std)	Mineral Layer (cm ± std)	Area (km <sup>2</sup> )	Area (%)	Mean Altitude of Profiles incl. Range (m a.p.s.l.)	Number of Sites	Sites in Permafrost <sup>d</sup> (cont./discont./ sporad./isol. or none)		
Birch Forest	5.7 ± 3.5	6.6 ± 5.0	2.0 ± 0.6	4.6 ± 4.6	25 ± 22	3.4 ± 0.9	22 ± 22	0.3	1.0	656 (636–675)	3	–/–/3		
Tundra Meadow	6.0 ± 3.0	7.2 ± 5.5	2.7 ± 1.4	4.5 ± 4.7	24 ± 24	4.3 ± 2.1	20 ± 23	1.3	4.3	652 (562–864)	8	–/–/8		
Shrub	4.6 ± 4.3	4.6 ± 4.3	1.5 ± 0.8	3.1 ± 3.6	16 ± 11	4.0 ± 1.8	12 ± 10	0.3	0.9	688 (581–824)	5	–/–/5		
Pat. Bould. Grass/Moss	6.2 ± 4.0	8.4 ± 5.4	1.4 ± 0.8	7.0 ± 5.2	38 ± 25	4.8 ± 1.8	33 ± 25	0.8	2.6	1129 (993–1202)	11	–/–/11/–		
Patchy Boulder Moss	1.8 ± 1.7	2.3 ± 1.9	0.6 ± 0.6	1.8 ± 1.5	40 ± 23	2.9 ± 1.5	37 ± 24	3.1	9.9	1181 (1076–1485)	12	–/1/11/–		
Sand/Gravel	0.7 ± 0.8	1.0 ± 0.9	0.1 ± 0.1	0.9 ± 0.9	53 ± 27	1.3 ± 1.0	52 ± 27	0.3	1.0	1293 (1122–1559)	9	–6/3/–		
Stones	0.05 ± 0.1	0.05 ± 0.1	0.05 ± 0.1	0.0	3 ± 2	3.0 ± 1.6	0	18.6	59.8	1304 (1154–1542)	8	–5/3/–		
Water	–	–	–	–	–	–	–	–	0.7	1194 <sup>d</sup>	(–)	0	–	
Permanent Snow/Ice	–	–	–	–	–	–	–	–	5.8	18.5	1530 <sup>d</sup>	(–)	0	–
Study Area <sup>a</sup>	0.7 ± 0.2	0.9 ± 0.2	0.3 ± 0.1	0.6 ± 0.2	8.6	2.4	6.2	31.2	100	1059 (562–1559)	56	–9/31/16		
Vegetated Area <sup>a, b</sup>	3.7 ± 0.8	4.6 ± 1.2	1.3 ± 0.3	3.3 ± 1.0	33.7	3.5	30.2	5.8	18.7	954 (562–1485)	39	–/1/22/16		

<sup>a</sup> Mean SOC storage is based on the land cover classification upscaling. The second number in each column is not the standard deviation like in the land cover classes, but the 95 % confidence interval (calculated according to Hugelius, 2012) which is based on the SOC variance and areal extent of each LCC.

<sup>b</sup> Only the vegetated area is considered. The following classes have been excluded from the calculations: “Sand/Gravel”, “Stones”, “Water”, “Permanent Snow/Ice”.

<sup>c</sup> The permafrost table was not reached during sampling at any of the sample sites.

<sup>d</sup> The mean altitude of the classes “Water” and “Permanent Snow/Ice” is based on the land cover classification and not on profile sites.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Low soil organic carbon storage in a subarctic alpine permafrost environment

M. Fuchs et al.

**Table 2.** Statistics of the geochemical analyses of soil samples.

Geochemical analysis	All samples, mean $\pm$ std	Organic rich top soil layer samples, mean $\pm$ std	Mineral layer samples, mean $\pm$ std	Significant difference between organic and mineral samples, student's <i>t</i> test	Correlation with increasing depth, Pearson's correlation
DBD ( $\text{g cm}^{-3}$ ) <sup>*</sup>	0.9 $\pm$ 0.8	0.4 $\pm$ 0.3	1.6 $\pm$ 0.7	yes ( $p < 0.01$ )	0.71 ( $p < 0.01$ )
LOI <sub>550</sub> (%) <sup>*</sup>	21.6 $\pm$ 27.0	40.3 $\pm$ 28.5	4.8 $\pm$ 8.1	yes ( $p < 0.01$ )	-0.47 ( $p < 0.01$ )
LOI <sub>950</sub> (%) <sup>*</sup>	0.4 $\pm$ 0.4	0.4 $\pm$ 0.4	0.3 $\pm$ 0.4	no ( $p = 0.06$ )	-0.11 ( $p = 0.05$ )
% C	11.4 $\pm$ 13.8	25.8 $\pm$ 13.7	3.8 $\pm$ 5.2	yes ( $p < 0.01$ )	-0.54 ( $p < 0.01$ )
C/N ratio (-)	17.6 $\pm$ 8.5	23.3 $\pm$ 11.4	14.6 $\pm$ 4.1	yes ( $p < 0.01$ )	-0.38 ( $p < 0.01$ )
$\delta^{13}\text{C}_{\text{tot}}$ vs. PDB (‰)	-26.1 $\pm$ 1.2	-26.8 $\pm$ 1.0	-25.6 $\pm$ 1.0	yes ( $p < 0.01$ )	0.42 ( $p < 0.01$ )
$\delta^{15}\text{N}$ vs. air (‰)	1.8 $\pm$ 2.6	-0.54 $\pm$ 2.0	3.2 $\pm$ 1.8	yes ( $p < 0.01$ )	0.53 ( $p < 0.01$ )

<sup>\*</sup> calculations carried out with all 295 samples; other calculations based on 96 samples from elemental analysis.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Low soil organic carbon storage in a subarctic alpine permafrost environment

M. Fuchs et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




**Table 3.** Results from the radiocarbon analysis.

Site	Depth (cm)	Lab. No.	Site and sample description	Age <sup>14</sup> C	Age cal yr BP
TA T1–9B	19–20	Poz-51853	Grass/moss patch, base of top organics	123.48 ± 0.4 pMC	modern
TA T1–9B	50–60	Poz-51854	Grass/moss patch, silty sand and stones	2035 ± 35 BP	1919
TA T2–11	10–15	Poz-51856	Grass/moss patch, base of top organics	95 ± 30 BP	20
TA T2–11	33–37	Poz-51857	Grass/moss patch, silty sand and small stones	1380 ± 30 BP	1269

Low soil organic carbon storage in a subarctic alpine permafrost environment

M. Fuchs et al.




**Figure 1.** The Tarfala Valley study area, including an overview location map, a map of the whole study area with land cover classification and detailed maps showing transect and sample point locations in the central (A) and lower (B) parts of the valley.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



**Low soil organic carbon storage in a subarctic alpine permafrost environment**

M. Fuchs et al.




**Figure 2.** Partitioning of total SOC storage and proportional area coverage of land cover classes in Tarfala Valley (31.2 km<sup>2</sup>).

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



## Low soil organic carbon storage in a subarctic alpine permafrost environment

M. Fuchs et al.



**Figure 3.** Results of the geochemical analyses of the soils samples of Tarfala Valley. DBD: dry bulk density (a); %C: percentage C (b); C/N weight ratio (c);  $\delta^{13}\text{C}_{\text{tot}}$  vs. PDB: stable isotope  $\delta^{13}\text{C}$  analyzed to the international standard PeeDeeBelemnite (d). Lines are best-fit power-, polynomial- or exponential regressions, shown for graphic representation of mean trends only.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Low soil organic carbon storage in a subarctic alpine permafrost environment

M. Fuchs et al.



**Figure 4.** Permafrost probability in relation to altitude – the probability is based on a logistic regression model with the altitude as single independent variable. The grey corridor shows the range of the permafrost probability if outliers (red dots) are removed from the model.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Low soil organic carbon storage in a subarctic alpine permafrost environment

M. Fuchs et al.



**Figure 5.** Fraction of vegetation cover and probability for permafrost presence in relation to altitude in Tarfala Valley, including the mean SOC storage per altitude (calculated in 50 m altitudinal intervals). The permafrost probability is based on the BTS-measurements and a logistic regression with the altitude as single independent variable. The vegetation fraction is based on the altitudinal distribution of vegetated classes in the land cover classification. Relatively low SOC values at elevations below 700 m are related to exposed streambeds in the Tarfala River alluvial fan.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion