

1 **Organic carbon pools in permafrost regions on the**
2 **Qinghai-Xizang (Tibetan) Plateau**

3 Cuicui Mu¹, Tingjun Zhang^{1*}, Qingbai Wu², Xiaoqing Peng¹, Bin Cao¹, Xiankai
4 Zhang¹, Bo Cao¹, Guodong Cheng²

5 ¹ College of Earth and Environmental Sciences, Lanzhou University, Lanzhou Gansu
6 730000, China

7 ² State Key Laboratory of Frozen Soil Engineering, Cold and Arid Regions
8 Environmental and Engineering Research Institute, CAS, Lanzhou Gansu 730000,
9 China

10

11 Correspondence to: Tingjun Zhang (tjzhang@lzu.edu.cn).

12 **Abstract**

13 The current Northern Circumpolar Soil Carbon Database did not include organic carbon storage in
14 permafrost regions on the Qinghai-Xizang (Tibetan) Plateau (QXP). In this study, we reported a
15 new estimation of soil organic carbon (SOC) pools in the permafrost regions on the QXP up to 25
16 m depth using a total of 190 soil profiles. The SOC pools were estimated to be 17.3 ± 5.3 Pg for the
17 0~1 m depth, 10.6 ± 2.7 Pg for the 1~2 m depth, 5.1 ± 1.4 Pg for the 2~3 m depth and 127.2 ± 37.3 Pg
18 for the layer of 3~25 m depth. The percentage of SOC storage in deep layers (3~25 m) on the QXP
19 (80%) was higher than that (39%) in the yedoma and thermokarst deposits in arctic regions. In
20 total, permafrost regions on the QXP contain approximately 160 ± 87 Pg SOC, of which
21 approximately 132 ± 77 Pg (83%) stores in perennially frozen soils and deposits. Total organic
22 carbon pool in permafrost regions on the QXP was approximately 8.7% of that in northern
23 circumpolar permafrost region. The present study demonstrates that the total organic carbon
24 storage is about 1832 Pg in permafrost regions on northern hemisphere.

25

26 **1. Introduction**

27 Soil organic carbon (SOC) storage in permafrost regions has received worldwide attention due to
28 its direct contribution to the atmospheric greenhouse gas contents (Ping et al., 2008a; Tarnocai et
29 al., 2009; Zimov et al., 2009). Climate warming will thaw permafrost, which can cause previously
30 frozen SOC become available for mineralization (Zimov et al., 2006). Permafrost has potentially
31 the most significant carbon-climate feedbacks not only due to the intensity of climate forcing, but
32 also the size of carbon pools in permafrost regions (Schuur et al., 2008; Mackelprang et al., 2012;
33 Schneider von Deimling et al., 2012).

34 Recently, carbon stored in permafrost regions has created many concerns because of the
35 implication on global carbon cycling (Ping et al., 2008; Burke et al., 2012; Zimov et al., 2006;
36 Michaelson et al., 2013; Hugelius et al., 2013). It has been estimated that permafrost regions of
37 circum-Arctic areas contain approximately 1672 Pg of organic carbon, which includes 495.8 Pg
38 for the 0~1 m depth, 1024 Pg for the 0~3 m depth and 648 Pg for 3~25 m depth (Tarnocai et al.,
39 2009). Based on newly available regional soil maps, the estimated storage of SOC in 0~3 m depth
40 is estimated to 1035 ± 150 Pg (Hugelius et al., 2014), about 1% higher than the previous estimate
41 by Tarnocai et al. (2009). The thawing of permafrost would expose the frozen organic carbon to
42 microbial decomposition, and thus may initiate a positive permafrost carbon feedback on climate
43 (Schuur et al., 2008). The strength and timing of permafrost carbon feedback greatly depend on
44 the distribution of SOC in permafrost regions. Therefore, understanding soil carbon storage in
45 permafrost regions is critical for better predicting future climate change. However, the present
46 knowledge of SOC pool in permafrost regions only limited to the circum-Arctic areas. Little is
47 known about the SOC pools in the low-altitude permafrost regions.

48 The Qinghai-Xizang (Tibetan) Plateau (QXP) in China has the largest extent of permafrost in
49 the low-middle latitudes of the world, with permafrost regions of about 1.35×10^6 km² and
50 underlying ~67% of the QXP area (Ran et al., 2012). It has been suggested that SOC in permafrost
51 regions on the QXP was very sensitive to global warming, due to the permafrost characteristics of
52 high temperature ($< -2.0^\circ\text{C}$), thin thickness (< 100 m) and unstable thermal states (Cheng and Wu,
53 2007; Li et al., 2008; Wu and Zhang, 2010). Mean annual permafrost temperatures at 6.0 m depth
54 increased by a range of 0.12°C to 0.67°C from 1996 to 2006 (Wu and Zhang, 2008), and increased
55 $\sim 0.13^\circ\text{C}$ from 2002 to 2012 (Wu et al., 2015). Active layer thickness increased, on average,

56 approximately $\sim 4.26 \text{ cm y}^{-1}$ along the Qinghai-Tibetan Highway from 2002 to 2012 (Wu et al.,
57 2015). In addition, the carbon stored in permafrost area was labile and a great part of the carbon
58 was mineralizable (Mu et al., 2014; Wu, et al., 2014).

59 Some studies have been conducted on SOC pools in 0~1 m depth on the QXP (Wang et al.,
60 2002; 2008; Yang et al., 2008; 2010; Liu et al., 2012; Wu et al., 2012). It was estimated that total
61 SOC for the top 0.7 m was about 30~40 Pg in the grassland of the plateau. The disagreement
62 among the studies on the SOC pools was attributed to the limited sampling points and the quality
63 of the SOC data gathered to date. Despite the importance of SOC in permafrost areas, there are
64 still few reports to the SOC storage in permafrost regions of the QXP. So far, the current Northern
65 Circumpolar Soil Carbon Database (Tarnocai et al., 2009) does not include the SOC in permafrost
66 regions on the QXP (Tarnocai et al., 2009).

67 Perennially frozen soils are important earth system carbon pools because of their vulnerability
68 to climate change (Koven et al., 2011). Some of the movement of SOC from surface to few meter
69 depth is accomplished through cryoturbation (Bockheim et al., 1998), which is caused by cracking
70 due to soil freeze-thaw cycles and by soil hydrothermal gradients (Ping et al., 2008b). It was
71 reported that the total yedoma region contains $211+160/-153 \text{ Pg C}$ in deep soil deposits (Strauss et
72 al., 2013). Current studies have shown the importance of deep organic carbon in permafrost
73 regions and its feedback with climate change (Hobbie et al., 2000; Davidson and Janssens, 2006;
74 Schuur et al., 2009). Deep organic carbon can be more sensitive to temperature increasing
75 compared with that in the active layer (Waldrop et al., 2010). Therefore, it is essential to study the
76 distribution of organic carbon content in deep layers of permafrost regions.

77 For the top layer, important factors controlling SOC pools are vegetation type and climate

78 (Jobbagy and Jackson, 2000). The vegetation type and climate conditions related closely to each
79 other on the QXP (Wang et al., 2002). Thus it is possible to calculate the SOC pools at 0~2 m
80 depth according to the area of vegetation type (Chinese Academy of Sciences, 2001) in the
81 permafrost regions (LIGG/CAS, 1988). For deep layers, the geomorphology and lithological
82 conditions play an important role in the distribution of SOC pools (Hugelius et al., 2013). Thus it
83 is reasonable to estimate the SOC pools at 2~25 m depth according to the area of Quaternary
84 geological stratigraphy in permafrost regions on the QXP.

85 The objective of this study is to assess the SOC pools in permafrost regions on the QXP, based
86 on the published data and new field sampling through deep drilling from this study. The new
87 estimation focuses on the permafrost regions and includes deeper layers, down to 25 m. SOC
88 storages of the plateau were estimated using the published data of 190 soil profiles and 11 deep
89 sampling sites from this study in combination with the vegetation map, permafrost map and
90 geological stratigraphy map of the QXP (Figs. 1, 2, 3). The result would update current estimation
91 of surface organic carbon pool and deep organic carbon storage in permafrost regions of the QXP,
92 which can provide new insights in permafrost carbon on the global scale.

93

94 **2. Materials and methods**

95 **2.1 Soil carbon database in previous reports**

96 The soil carbon databases in 0~1 m depth were retrieved from the previous reports (Yang et al.,
97 2010; Liu et al., 2012; Wu et al., 2012; Dorfer et al.,2013; Mu et al., 2013) (Table 1). We
98 integrated the databases from Yang et al. (2010), Dorfer et al. (2013) and Ohtsuka et al. (2008)
99 because these studies were all performed in the middle and eastern part of the QXP. The data of

100 Wu et al. (2012), Liu et al. (2012) and Mu et al. (2013) in the soil carbon database in 0~1 m depth
101 were calculated separately, since their study regions of western QXP, Shulehe river basin (SLRB)
102 and Heihe river basin (HHRB) belonged to the isolated permafrost zone and the climate conditions
103 differed greatly with the continuous permafrost zones of the QXP. The total organic carbon pools
104 in 0~1 m depth in permafrost regions on the QXP were calculated using 190 profile sites from
105 published sources.

106 **2.2 Field sampling**

107 To calculate the deep carbon pools (2~25 m) in permafrost regions, 11 boreholes on the QXP were
108 drilled from 2009 to 2013 (Fig. 1). Geographic location for the 11 boreholes, together with the
109 active layer depth, sampling depth, vegetation type, geological stratigraphies, SOC contents, bulk
110 density, water contents and soil texture are provided in the supplement materials.

111 The deep sampling sites were mainly located in three vegetation types of alpine meadow, alpine
112 stepper and alpine desert (Fig. 2). Three sampling sites (KXL, HLH-1, HLH-2) were located in the
113 vegetation type of alpine steppe. Another site (ZEH) was near to the *Zhouerhu Lake* in *Kekexili*,
114 with soil formed from lacstrine deposits. It was typical alpine desert and perennially frozen,
115 containing less amounts of organic carbon. Five sampling sites (KL150, KL300, KL450, WDL,
116 XSH) were located in the vegetation type of alpine meadow. In addition, two sites in permafrost
117 regions of the Heihe river basin (HHRB: Heihe-1, Heihe-2) with vegetation type of alpine
118 meadow were rich in organic carbon with high soil water contents (Mu et al., 2013).

119 The deep sampling sites were mainly distributed in three geological stratigraphies: ZEH, WDL,
120 XSH, Heihe-1 and Heihe-2 were in the Quaternary stratigraphy, KL150, KL300, KL450, HLH-1
121 and HLH-2 were in the Triassic stratigraphy, and KXL was in the Permian stratigraphy (Fig. 3).

122 **2.3 Analytical methods**

123 For SOC analyses, the homogenized samples were quantified by dry combustion on a Vario EL
124 elemental analyzer (Elemental, Hanau, Germany). During measurement, 0.5 g dry soil samples
125 were pretreated by HCl (10 mL 1 mol L⁻¹) for 24 h to remove carbonate (Sheldrick, 1984). Bulk
126 density was determined by measuring the volume (length, width, height) of a section of frozen
127 core, and then drying the segment at 105°C (for 48 h) and determining its mass.

128 **2.4 Calculation of soil carbon pools**

129 For the Stock of Soil Organic Carbon (SSOC, kg m⁻²), it was calculated using the formula (1)
130 (Dorfer et al., 2013):

$$131 \quad SSOC = C \times BD \times T \times (1 - CF) \quad (1)$$

132 where C was the organic carbon content (wt%), BD was the bulk density (g cm⁻³), T was the soil
133 layer thickness and CF was the coarse fragments (wt%). Using this information, the SSOC was
134 calculated for the 0~1m, 1~2 m, 2~3 m and 3~25 m depths, respectively. Then, SOC storage (Pg)
135 was estimated by multiplying the SSOC at different depth by the distribution area.

136 For the organic carbon storage in 0~1 m depth, the reported SOC densities data of 190 sampling
137 sites were collected through their distribution in permafrost regions (Fig. 1). The area of alpine
138 meadow, alpine steppe and alpine desert in permafrost regions was calculated through overlaying
139 the vegetation map over the QXP permafrost regions (Fig. 2). For the organic carbon storage in
140 1~2 m depth, the organic carbon densities of 11 boreholes were extrapolated to the located
141 vegetation type area.

142 For the organic carbon storage in 2~3 m and 3~25 m depths, the area of permafrost regions in
143 the Quaternary, Triassic and Permian stratigraphies on the QXP was calculated through overlaying

144 the distribution of geological stratigraphies over the permafrost map (Fig. 3). The organic carbon
145 pools of 2~3 m and 3~25 m depth were estimated through deep organic carbon densities
146 multiplied by the area of geological stratigraphies. The three geological stratigraphies had thick
147 sediments of about 25 m (Fang et al., 2002; 2003; Qiang et al., 2001). As for other geological
148 stratigraphies, the poor soil development was reported and soil thickness was usually less than 3 m
149 (Wu et al., 2012; Yang et al., 2008; Hu et al., 2014). Thus other stratigraphies were not considered
150 in the estimation of deep organic carbon pool in the permafrost regions.

151

152 **3. Results**

153 **3.1 Organic carbon pools in the 0~1 m depth**

154 Based on the vegetation data on the QXP (Figs. 1, 2), the area of permafrost regions in the alpine
155 meadow, alpine steppe and alpine desert are $0.302 \times 10^6 \text{ km}^2$, $0.772 \times 10^6 \text{ km}^2$ and $0.175 \times 10^6 \text{ km}^2$
156 respectively, with a total area of approximately $1.249 \times 10^6 \text{ km}^2$.

157 Organic carbon storage of the permafrost regions in the 0~1 m depth on the QXP was
158 approximately $17.3 \pm 5.3 \text{ Pg}$, of which approximately $11.3 \pm 4.0 \text{ Pg}$ (65%) in the alpine meadow,
159 $5.3 \pm 2.8 \text{ Pg}$ (31%) in the alpine steppe, and $0.7 \pm 0.3 \text{ Pg}$ (4%) in the alpine desert, respectively
160 (Table 1). There were great variations in SOC contents among the sites under alpine meadow area.
161 SOC store in the HHRB ($39.0 \pm 17.5 \text{ kg m}^{-2}$) was much higher than that of most sites in the
162 predominately continuous permafrost zone on the QXP. In contrast, the SOC stores showed little
163 variation over the sites in the alpine steppe and alpine desert areas, with the ranges of $6.9 \pm 3.6 \text{ kg}$
164 m^{-2} and $3.9 \pm 1.5 \text{ kg m}^{-2}$, respectively.

165 **3.2 Distribution of deep organic carbon**

166 According to the distribution of sampling sites at the geological stratigraphies, for the Quaternary
167 stratigraphy, average SOC contents at 2~3 m and 3~25 m depths were $0.8\pm0.6\%$ and $0.8\pm0.7\%$.
168 For the Triassic stratigraphy, average SOC contents at 2~3 m and 3~25 m depths were $1.1\pm0.3\%$
169 and $1.2\pm0.6\%$. For the Permian stratigraphy, average SOC contents at 2~3 m and 3~25 m depths
170 were $1.5\pm0.4\%$ and $1.1\pm0.3\%$. As for the permafrost regions in HHRB, the SOC contents (Heihe-1,
171 Heihe-2) were higher than those of predominately continuous permafrost zone on the QXP, with a
172 range of $5.1\pm3.7\%$ and $2.7\pm2.4\%$ to depth of 19 m. SOC contents decreased with depth in most
173 deep boreholes, while SOC contents in deeper layers were higher than those in the top layer at the
174 XSH, KL150 and KL300 (Fig. 4).

175 With the deep soil data, a relationship between SOC contents (SOC%) and soil depth (h) in
176 deep soils of permafrost regions can be characterized by a power function (2) (Fig. 4):

$$177 \quad \text{SOC}\% = 14.11h^{-1.20} \quad (R^2 = 0.68, p < 0.01, n = 362) \quad (2)$$

178 **3.3 Deep organic carbon pools**

179 Based on the Quaternary stratigraphies data in permafrost regions of the QXP (Fig. 3), the area of
180 permafrost regions in the Quaternary, Triassic and Permian stratigraphies are $0.194\times10^6 \text{ km}^2$,
181 $0.238\times10^6 \text{ km}^2$ and $0.135\times10^6 \text{ km}^2$ respectively, with a total area of approximately $0.567\times10^6 \text{ km}^2$,
182 about 45% of permafrost regions on the QXP.

183 Organic carbon storages in permafrost regions on the QXP were approximately $10.6\pm2.7 \text{ Pg}$ in
184 the 1~2 m depth, $5.1\pm1.4 \text{ Pg}$ in the 2~3 m depth and $127.2\pm37.3 \text{ Pg}$ in deep depth of 3~25 m
185 (Table 2). In total, it contains approximately $160\pm87 \text{ Pg}$ of organic carbon at depth of 25 m in
186 permafrost regions on the QXP.

187 Active layer thickness on the QXP varies from 0.8 to 4.6 m, and in most regions, active layer

188 thickness was about 2 m (Cheng and Wu, 2007; Wu and Zhang, 2008; Zhao et al., 2010; Wu et al.,
189 2012). Thus we consider the upper 2 m as the active layer. According to this depth, the organic
190 carbon storage in permafrost layers of 132 ± 77 Pg was approximately five times of that (28 ± 6 Pg)
191 in the active layer.

192 SOC storages in Quaternary, Triassic and Permian stratigraphies were 31 ± 17 Pg, 69 ± 53 Pg and
193 32 ± 20 Pg at depth of 2~25 m, respectively. More than a half of organic carbon is stored in
194 permafrost layers which belonged to the Triassic stratigraphy.

195

196 **4. Discussions**

197 Our estimates indicate that organic carbon storage in permafrost regions in the 0~1 m depth on the
198 QXP was approximately 17.3 ± 5.3 Pg. However, previous soil carbon pools on the alpine
199 grasslands of the whole QXP was estimated to be 33.5 Pg of 0~0.75 m (Wang et al., 2002), and
200 10.5 Pg of 0~0.30 m (Yang et al., 2010). The difference, in large part, between our new estimate
201 and previous reports can be explained as follows: i) Area of vegetation types in permafrost regions
202 was recalculated. The area of permafrost regions of about 1.249×10^6 km² was smaller than that of
203 Wang et al. (2002) (1.63×10^6 km²) and Yang et al. (2010) (1.26×10^6 km²). ii) Carbon density data
204 of sampling sites located in permafrost regions was collected. The integration of carbon data from
205 the results of recent publications (Ohtsuka et al., 2008; Dorfer et al., 2013; Wu et al., 2012) and
206 our field data resulted in a higher carbon density than those of previous reports (Wang et al., 2002;
207 Yang et al., 2010). iii) The regions of SLRB and HHRB were not considered in previous SOC
208 pool estimate. The organic carbon storages of 0.43 ± 0.11 Pg in SLRB and 0.25 ± 0.11 Pg in HHRB
209 were added in the present study.

210 It is worth to mention that there were wide variations in organic carbon contents in permafrost
211 regions on the QXP in previous reports (Wang et al., 2002; Yang et al., 2010; Liu et al., 2012; Wu
212 et al., 2012; Dorfer et al., 2013; Ohtsuka et al., 2008; Mu et al., 2013). A possible explanation is
213 the spatial heterogeneity of SOC contents in permafrost regions of the QXP. In addition, the
214 different analytical methods may also contribute to the differences of carbon contents (Table 1). It
215 has been demonstrated that if taking the dry combustion method as standard, the recovery of
216 organic carbon was 99% for wet combustion and 77% for Walkley-Black (Kalembasa and
217 Jenkinson, 1973; Nelson and Sommers, 1996).

218 The SOC stocks at 0~1 m depth (17.3 kg m^{-2}) in the alpine meadow on the QXP is higher than
219 that in subarctic alpine permafrost (0.9 kg m^{-2}) (Fuchs et al., 2014), and similar with that of the
220 lowland and hilly upland soils in North American Arctic region (55.1 kg m^{-2} , 40.6 kg m^{-2}) (Ping et
221 al., 2008a). It implies that SOC of the alpine meadow in permafrost regions has a large proportion
222 in permafrost carbon pools. The SOC contents at 0~1 m depth ($3.9 \pm 1.5 \text{ kg m}^{-2}$) in the alpine desert
223 on the QXP was similar with that (3.4 kg m^{-2} , 3.8 kg m^{-2}) in rubbleland and mountain soils in
224 North American Arctic region (Ping et al., 2008a). These results suggest that the SOC stocks are
225 closely related to the vegetation type in the permafrost regions.

226 SOC decreases with the depth on the QXP (Fig. 4), which is in good agreement with those
227 reported in circum-Arctic regions (Strauss et al., 2013; Zimov et al., 2006). This could be
228 explained by the dynamics of Quaternary deposit and SOC formation in permafrost regions
229 (Strauss et al., 2013). However, the organic carbon contents of deep layers in some sites (XSH,
230 KL150 and KL300) were higher than those in the top layers (Fig. 4), which may be caused by the
231 cryoturbation and sediment burying process (Ping et al., 2010), and Quaternary deposits following

232 the uplift of Tibetan Plateau (Li et al., 1994; 2014). Overall, SOC decreases exponentially with
233 depth (equation 1) in permafrost regions on the QXP, which is in agreement with results from
234 other regions (Don et al., 2007). Certainly, more efforts are still needed in studying the distribution
235 of deep organic carbon density in permafrost regions.

236 In the present study, it is the first time to study the deep organic carbon in permafrost regions,
237 and quantify the carbon storage below 1.0 m depth on the QXP. The mean SOC content of 11
238 boreholes in permafrost regions on the QXP (2.5 wt%) was similar with that in the yedoma
239 deposits (3.0 wt%) (Strauss et al., 2013), and that of lowland steppe-tundra soils in Siberia and
240 Alaska (2.6 wt%) (Zimov et al., 2006). Since it has been pointed out that yedoma deposits contain
241 a large amount of organic carbon, it would be reasonable to infer that deep soil carbon in
242 permafrost regions on the QXP may also have a great contribution to carbon pools. Our
243 estimations indicate that the soils on the QXP contains 33.0 ± 13.2 Pg of organic carbon in the top
244 3.0 m of soils, with an additional 127.2 ± 37.3 Pg C distributed in deep layers (3~25 m) of the
245 Quaternary, Triassic and Permian stratigraphies in permafrost regions. In northern circumpolar
246 permafrost region, 1024 Pg of organic carbon was in the 0~3 m depth and 648 Pg (39%) of carbon
247 was stored in deep layers of yedoma and deltaic deposits (Tarnocai et al., 2009). The percentage of
248 SOC storage in deep layers (3~25 m) on the QXP (80%) is much higher than that (39%) in the
249 yedoma and thermokarst deposits in Siberia and Alaska. This could be explained as that the
250 paleoenvironment of the QXP was wet and warm, or lacustrine sediment in most regions (Zhang et
251 al., 2003; Lu et al., 2014), which always links to the well formation of soil organic matter (Kato et
252 al., 2004; Piao et al., 2006; Chen et al., 1990).

253 In total, there is approximately 160 ± 87 Pg of organic carbon stored at 0~25 m depth in

254 permafrost regions on the QXP, which would update the total carbon pools to 1832 Pg in
255 permafrost regions of northern hemisphere. The total carbon pools on the QXP permafrost regions
256 account for approximately 8.7% of the total carbon pools in permafrost regions in northern
257 hemisphere. Since the permafrost region on the QXP was about 6% of northern permafrost area
258 (Ran et al., 2012), it could be seen that SOC in permafrost regions on the QXP should be paid
259 more attention in the future studies.

260

261 **5. Conclusions**

262 (1) According to the organic carbon data in previous analysis and field exploration of deep
263 boreholes in permafrost regions, the organic carbon storages in permafrost regions on the QXP
264 were estimated to approximately 17.3 ± 5.3 Pg in the 0~1 m, 10.6 ± 2.7 Pg in the 1~2 m, 5.1 ± 1.4 Pg
265 in the 2~3 m and 127.2 ± 37.3 Pg in deep depth of 3~25 m.

266 (2) The percentage of SOC storage in deep layers (3~25 m) of permafrost regions on the QXP was
267 80%, which was higher than that in the yedoma and thermokarst deposits in Siberia and Alaska.

268 (3) In total, organic carbon pools in permafrost regions on the QXP are approximately 160 ± 87 Pg,
269 of which 132 ± 76 Pg occurs in permafrost layers. The total carbon pools in permafrost regions in
270 northern hemisphere are now updated to 1832 Pg.

271

272 **Acknowledgements**

273 This work was supported by the National Key Scientific Research Project (Grant 2013CBA01802),
274 National Natural Science Foundation of China (Grants 91325202, 41330634), and the Open
275 Foundations of State Key Laboratory of Cryospheric Sciences (Grant SKLCS-OP-2014-08) and

276 State Key Laboratory of Frozen Soil Engineering (Grant SKLFSE201408). The authors gratefully
277 acknowledge the reviewers, professor Gustaf Hugelius and professor Chien-Lu Ping, as well as the
278 editor, Dr. M. Noe Steffen, for their constructive comments and suggestions.

279

280 **References**

281 Bockheim, J. B., Walker, D. A., Everett, L. R., Nelson, F. E., and Shikolmanov, N. I.: Soils and
282 cryoturbation in moist nonacidic and acidic tundra in the Kuparuk river basin arctic Alaska,
283 USA, *Arct. Alp. Res.*, 30, 166–174, 1998.

284 Burke, E. J., Hartley, I. P., and Jones, C. D.: Uncertainties in the global temperature change caused
285 by carbon release from permafrost thawing, *The Cryosphere*, 6(5), 1063–1076, 2012.

286 Editorial Board of Vegetation Map of China, Chinese Academy of Sciences, *Vegetation Atlas of*
287 *China (1:1 000 000)*, Beijing: Science Press, 2001.

288 Chen, K. Z., Bowler, J. M., and Kelts, K.: Palaeoclimate evolution within the Qinghai-Xizang
289 (Tibet) plateau in the last 40000 years, *Quaternary Sciences*, 1, 22–30, 1990.

290 Cheng, G. D., and Wu, T. H.: Responses of permafrost to climate change and their environmental
291 significance, Qinghai-Tibet Plateau, *Journal of Geophysical Research*, 112, F02S03, DOI:
292 10.1029/2006JF000631, 2007.

293 Davidson, E. A., and Janssens, I. A.: Temperature sensitivity of soil carbon decomposition and
294 feedbacks to climate change, *Nature*, 440, 165–173, 2006.

295 Dorfer, C., Kuhn, P., Baumann, F., He, J. S., and Scholten, T.: Soil Organic Carbon Pools and
296 Stocks in Permafrost-Affected Soils on the Tibetan Plateau, *PLoS ONE*, 8(2), e57024,
297 doi:10.1371/journal.pone.0057024, 2013.

298 Don, A., Schumacher J., Scherer-Lorenzen, M., Scholten, T., and Schulze, E.D.: Spatial and
299 vertical variation of soil carbon at two grassland sites — Implications for measuring soil carbon
300 stocks, *Geoderma*, 141, 272–282, 2007.

301 Fang, X. M., Lu, L. Q., Mason, J. A., Yang, S. L., An, Z. S., Li, J. J., and Guo, Z. L.: Pedogenic
302 response to millennial summer monsoon enhancements on the Tibetan Plateau, *Quaternary*
303 *International*, 106–107, 79–88, 2003.

304 Fang, X. M., Lu, L. Q., Yang, S. L., Li, J. J., An, Z. S., Jiang, P.A., and Chen, X.L.: Loess in
305 Kunlun Mountains and its implications on desert development and Tibetan Plateau uplift in
306 west China, *Science in China*, 45, 291–298, 2002.

307 Fuchs, M., Kuhry, P., and Hugelius, G.: Low soil organic carbon storage in a subarctic alpine
308 permafrost environment, *The Cryosphere Discuss*, 8, 3493–3524, 2014.

309 Hobbie, S. E., Schimel, J. P., Trumbore, S. E., and Randerson, I. R.: Controls over carbon storage
310 and turnover in high-latitude soils, *Glob. Change Biol.*, 6, 196–210, 2000.

311 Hu, G. L., Fang, H. B., Liu, G. M., Zhao, L., Wu, T. H., Li, R., and Wu, X. D.: Soil carbon and
312 nitrogen in the active layers of the permafrost regions in the Three Rivers' Headstream, *Environ.*
313 *Earth. Sci*, 72, 5113–5122, 2014.

314 Hugelius, G., Tarnocai, C., Broll, G., Canadell, J. G., Kuhry, P., and Swanson, D. K.: The Northern
315 Circumpolar Soil Carbon Database: spatially distributed datasets of soil coverage and soil
316 carbon storage in the northern permafrost regions, *Earth Syst. Sci. Data*, 5, 3–13, 2013.

317 Hugelius, G., Strauss, J., Zubrzycki, S., Harden, J. W., Schuur, E. A. G., Ping, C. L., Schirmermeister,
318 L., Grosse, G., Michaelson, G. J., Koven, C. D., O'Donnell, J. A., Elberling, B., Mishra, U.,
319 Camill, P., Yu, Z., Palmtag, J., Kuhry, P.: Estimated stocks of circumpolar permafrost carbon

320 with quantified uncertainty ranges and identified data gaps, *Biogeosciences*, 11, 6573–6593,
321 2014.

322 Jobbagy, E. G., and Jackson, R. B.: The vertical distribution of soil organic carbon and its relation
323 to climate and vegetation, *Ecological Applications*, 10(2), 423–436, 2000.

324 Kalembasa, S. J., and Jenkinson, D. D.: A comparative study of titrimetric and gravimetric
325 methods for the determination of organic carbon in soil, *Journal of the Science of Food and*
326 *Agriculture*, 24, 1085–1090, 1973.

327 Kato, T., Tang, Y., and Gu, S.: Carbon dioxide exchange between the atmosphere and an alpine
328 meadow ecosystem on the Qinghai–Tibetan Plateau, China, *Agric For Meteorol*, 124, 121–34,
329 2004.

330 Koven, C. D., Ringeval, B., Friedlingstein, P., Ciais, P., Cadule, P., Khvorostyanov, D., Krinner, G.,
331 and Tarnocai, C.: Permafrost carbon-climate feedbacks accelerate global warming, *P. Natl. Acad.*
332 *Sci. USA*, 108, 14769–14774, 2011.

333 Lanzhou Institute of Glaciology and Geocryology, Chinese Academy of Sciences, Map of Snow,
334 Ice and Frozen Ground in China (1:4,000,000). Cartographic Publishing House: Beijing, China
335 (in Chinese), 1988.

336 Li, J. J., Zhang, Q. S., and Li, B. Y.: Main processes of geomorphology in China in the past fifteen
337 years, *Acta Geographical Sinica*, 49, 642–648, 1997.

338 Li, J. J., Fang, X. M., Song, C. H., Pan, B. T., Ma, Y. Z., Yan, M. D.: Late Miocene–Quaternary
339 rapid stepwise uplift of the NE Tibetan Plateau and its effects on climatic and environmental
340 changes, *Quaternary Research*, 81, 400–423, 2014.

341 Li, X., Cheng, G. D., Jin, H. J., Kang, E. S., Che, T., Jin, R., Wu, L. Z., Nan, Z. T., Wang, J., and

342 Shen, Y. P.: Cryospheric Change in China, *Global and Planetary Change*, 62(3–4), 210–218,
343 2008.

344 Liu, W. J., Chen S. Y., Qin, X., Baumann, F., Scholten, T., Zhou, Z. Y., Sun, W. J., Zhang, T. Z.,
345 Ren, J. W., and Qin, D. H.: Storage, patterns, and control of soil organic carbon and nitrogen in
346 the northeastern margin of the Qinghai–Tibetan Plateau, *Environ. Res. Lett.*, 7, 1–12, 2012.

347 Lu, H. Y. and Guo, Z. T.: Evolution of the monsoon and dry climate in East Asia during late
348 Cenozoic: A review, *Science China Earth Science*, 57, 70–79, 2014.

349 MacDougall, A. H., Avis, C. A., and Weaver, A. J.: Significant contribution to climate warming
350 from the permafrost carbon feedback, *Nat. Geosci.*, 5(10), 719–721, doi:10.1038/ngeo1573,
351 2012.

352 Michaelson, G. J., Ping, C. L., Clark, M.: Soil pedon carbon and nitrogen data for Alaska: An
353 analysis and update, *Open Journal of Soil Science*, 3, 132–142, 2013.

354 Mu, C. C., Zhang, T. J., Wu, Q. B., Zhang, X. K., Cao, B., Wang, Q. F., Peng, X. Q., and Cheng,
355 G. D.: Stable carbon isotopes as indicators for permafrost carbon vulnerability in upper reach of
356 Heihe River basin, northwestern China, *Quaternary International*, 321, 71–77: 2014.

357 Mu, C. C., Zhang, T. J., Cao, B., Wan, X. D., Peng, X. Q., and Cheng, G. D.: Study of the organic
358 carbon storage in the active layer of the permafrost over the Eboiling Mountain in the upper
359 reaches of the Heihe River in the Eastern Qilian Mountains, *Journal of Glaciology and*
360 *Geocryology*, 35(1), 1–9, 2013.

361 Nelson, D. E., and Sommers, L. E.: Total carbon, organic carbon, and organic matter, *Methods of*
362 *soil analysis, Part 3 - chemical methods*, 961–1010, 1996.

363 Ohtsuka, T., Hirota, M., Zhang, X., Shimono, A., Senga, Y., Du, M., Yonemura, S., Kawashima, S.,

364 and Tang, Y.: Soil organic carbon pools in alpine to nival zones along an altitudinal gradient
365 (4400–5300 m) on the Tibetan Plateau, *Polar Science*, 2, 277–285, 2008.

366 Ping, C. L., Michaelson, G. J., Jorgenson, T., Kimble, J. M., Epstein, H., Romanovsky, V. E., and
367 Walker, D. A.: High stocks of soil organic carbon in the North American arctic region, *Nature*
368 *Geosci.*, 1, 615–619, 2008a.

369 Ping, C. L., Michaelson, G. J., Kimble, J. M., Romanovsky, V. E., Shur, Y. L., Swanson, D. K., and
370 Walker, D. A.: Cryogenesis and soil formation along a bioclimate gradient in Arctic North
371 America, *J. Geophys. Res.*, 113, G03S12, doi:10.1029/2008JG000744, 2008b.

372 Piao, S. L., Fang, J. Y., and He, J. S.: Variations in vegetation net primary production in the
373 Qinghai–Xizang Plateau, China, from 1982 to 1999, *Clim Chang*, 74, 253–67, 2006.

374 Qiang, X. K., Li, Z. X., Powell, C. McA., and Zheng, H. B.: Magnetostratigraphic record of the
375 Late Miocene onset of the East Asian monsoon, and Pliocene uplift of northern Tibet, *Earth and*
376 *Planetary Science Letters*, 187, 83–93, 2001.

377 Ran, Y. H., Li, X., Cheng, G. D., Zhang, T. J., Wu, Q. B., Jin, H. J., and Jin, R.: Distribution of
378 Permafrost in China: An Overview of Existing Permafrost Maps, *Permafrost and Periglacial*
379 *Processes*, 23, 322–333, 2012.

380 Schneider von Deimling, T., Meinshausen, M., Levermann, A., Huber, V., Frieler, K., Lawrence, D.
381 M., and Brovkin, V.: Estimating the near-surface permafrost-carbon feedback on global
382 warming, *Biogeosciences*, 9(2), 649–665, doi:10.5194/bg-9-649-2012, 2012.

383 Schuur, E. A. G., Vogel, J. G., Crummer, K. Lee, G. H., Sickman, J. O., and Osterkamp, T. E.: The
384 effect of permafrost thaw on old carbon release and net carbon exchange from tundra, *Nature*,
385 459, 556–559, 2009.

386 Sheldrick, B. H.: Analytical Methods Manual, 212 pp., Land Resour. Res. Inst., Res. Branch,
387 Agric. Can., Ottawa, 1984.

388 Strauss, J., Schirrmeister, L., Grosse, G., Wetterich, S., Ulrich, M., Herzsuh, U., and Hubberten,
389 H. W.: The deep permafrost carbon pool of the yedoma region in Siberia and Alaska, *Geophys.*
390 *Res. Lett.*, 40, 6165–6170, doi:10.1002/2013GL058088, 2013.

391 Tarnocai, C., Canadell, J. G., Schuur, E. A. G., Kuhry, P., Mazhitova, G., and Zimov, S.: Soil
392 organic carbon pools in the northern circumpolar permafrost region, *Global Biogeochem.*
393 *Cycles*, 23, GB2023, doi:10.1029/2008GB003327, 2009.

394 Wang, G. X., Qian, J., Cheng, G. D., and Lai, Y. M.: Soil organic carbon pool of grassland soils
395 on the Qinghai-Tibetan Plateau and its global implication, *The Science of the Total*
396 *Environment*, 291, 207–217, 2002.

397 Waldrop, M. P., Wickland, K. P., White, R., Berhe, A. A., Harden, J. W., and Romanovsky, V. E.:
398 Molecular investigations into a globally important carbon pool: permafrost-protected carbon in
399 Alaskan soils, *Global Change Biology*, doi: 10.1111/j.1365-2486.2009.02141.x, 2010.

400 Wang, G., Li, Y., Wang, Y., and Wu, Q.: Effects of permafrost thawing on vegetation and soil
401 carbon losses on the Qinghai-Tibet Plateau, China, *Geoderma*, 143, 143–152, 2008.

402 Wu, X. D., Zhao, L., Fang, H. B., Yue, G. Y., Chen, J., Pang, Q. Q., Wang, Z. W., and Ding, Y. J.:
403 Soil Organic Carbon and Its Relationship to Vegetation Communities and Soil Properties in
404 Permafrost of Middle-western Qinghai-Tibet Plateau, *Permafrost and Periglacial Process*, 23,
405 162–169, 2012.

406 Wu, X. D., Fang, H. B., Zhao, L., Wu, T. H., Li, R., Ren, Z. W., Pang, Q. Q., and Ding, Y. J.:
407 Mineralization and Fractions Changes in Soil Organic Matter in Soils of Permafrost Region in

408 Qinghai-Tibet Plateau, Permafrost and Periglacial Processes, DOI:10.1002/ppp.1796, 2014.

409 Wu, Q., Hou, Y., Yun, H., Liu, Y. Changes in active-layer thickness and near-surface permafrost
410 between 2002 and 2012 in alpine ecosystems, Qinghai–Xizang (Tibet) Plateau, China. *Global
411 and Planetary Change*, 124, 149–155, 2015.

412 Wu, Q. B., and Zhang, T. J.: Changes in active layer thickness over the Qinghai-Tibetan Plateau
413 from 1995 to 2007, *Journal of Geophysical Research*, 115, D09107, doi:10.1029/2009JD012974,
414 2010.

415 Wu, Q. B., and Zhang, T. J.: Recent permafrost warming on the Qinghai-Tibetan Plateau, *Journal
416 of Geophysical Research*, 113, D13108, doi:10.1029/2007JD009539, 2008.

417 Yang, Y. H., Fang, J. Y., Tang, Y. H., Ji, C. J., Zheng, C. Y., He, J. S., and Zhu, B.: Storage,
418 patterns and controls of soil organic carbon in the Tibetan grasslands, *Global Change Biology*,
419 14, 1592–1599, 2008.

420 Yang, M., Nelson, F. E., Shiklomanov, N. I., Guo, D., and Wan, G.: Permafrost degradation and its
421 environmental effects on the Tibetan Plateau: A review of recent research, *Earth-Science
422 Reviews*, 103 (1–2), 31–44, 2010.

423 Zimov, N. S., Zimov, S. A., Zimova, A. E., Zimova, G. M., Chuprynin, V. I., and Chapin III, F. S.:
424 Carbon storage in permafrost and soils of the mammoth tundra-steppe biome: Role in the global
425 carbon budget, *Geophys. Res. Lett.*, 36, L02502, doi:10.1029/2008GL036332, 2009.

426 Zimov, S. A., Schuur, E. A. G., and Chapin, F. S.: Permafrost and the global carbon budget,
427 *Science*, 312, 1612–3, 2006.

428 Zhang, Q. B., Cheng, G. D., Yao, T. D., Kang, X. C., and Huang, J. G.: A 2,326-year tree-ring
429 record of climate variability on the northeastern Qinghai-Tibetan Plateau, *Geophysical Research*

430 Letter, 30, doi:10.1029/2003GL017425, 2003.

431 Zhao, L., Wu, Q. B., Marchenko, S. S., Sharkhuu, N.: Thermal state of permafrost and active layer

432 in Central Asia during the international polar year, *Permafrost and Periglacial Processes*, 21(2),

433 198–207, 2010.

434

435 **Supplement**

436 Dataset of the geographic location for 11 boreholes on the Qinghai-Xizang (Tibetan) Plateau,
437 together with the active layer depth, sampling depth, vegetation type, geological stratigraphies,
438 soil organic carbon (SOC) contents, bulk density, water contents and soil texture.

439

440 **Table captions**

441 Table 1 Organic carbon pools in the 0~1 m depth with different vegetation type on the QXP

442 Table 2 Permafrost organic carbon storage to the depth of 25 m on the QXP

443

444 **Figure captions**

445 Figure 1 Location of sampling sites on the QXP, shown on the background of QXP permafrost
446 distribution (blue points were sampling sites in Yang et al., (2010); orange points were in Wu et
447 al., (2012); red box was Shule river basin (SLRB) in Liu et al., (2012); black box was Heihe river
448 basin (HHRB) in Mu et al., (2013))

449 Figure 2 Location of sampling sites on the QXP, shown on the background of QXP vegetation
450 atlas at a scale of 1:400 0000 (Chinese Academy of Sciences, 2001) (Sampling sites were same
451 with those shown on the background of permafrost distribution)

452 Figure 3 Location of sampling sites on the QXP, shown on the background of QXP Quaternary
453 geological map (Sampling sites were same with those shown on the background of permafrost
454 distribution).

455 Figure 4 Distributions of soil organic carbon contents in deep soils in permafrost regions on the
456 QXP.

457 **Table 1** Organic carbon pools in the 0~1 m depth with different vegetation type on the QXP


Vegetation types	References	Analytical methods	Study area	Site data (n)	Area ($\times 10^6 \text{ km}^2$)	SOC stock (kg m^{-2})	SOC storage (Pg)
Alpine meadow	Yang et al., (2010)	Wet oxidation	QXP	22		9.3 \pm 3.9	
	Ohtsuka et al., (2008)	Heat combustion	QXP	1	0.224	13.7	10.7 \pm 3.8
	Dorfer et al., (2013)	Heat combustion	QXP	2		10.4	
	Mu et al., (2013)	Heat combustion	HHRB	11	0.0065	39.0 \pm 17.5	0.3 \pm 0.1
	Liu et al., (2012)	Wet oxidation	SLRB	~42	0.013	8.7 \pm 1.2	0.1 \pm 0.02
Alpine steppe	Yang et al., (2010)	Wet oxidation	QXP	33		3.7 \pm 2.0	
	Wu et al., (2012)	Wet oxidation	Western QXP	52	0.772	7.7 \pm 3.2	5.3 \pm 2.8
	Liu et al., (2012)	Wet oxidation	SLRB	~42		9.2 \pm 1.1	
Alpine desert	Wu et al., (2012)	Wet oxidation	Western QXP	25		3.3 \pm 1.5	
	Liu et al., (2012)	Wet oxidation	SLRB	~42	0.175	4.4 \pm 0.7	0.7 \pm 0.3

458 **Table 2** Permafrost organic carbon storage to the depth of 25 m on the QXP

Vegetation types	Alpine meadow			Alpine steppe		Alpine desert		Total (Pg)
	SOC (kg m ⁻²)	SOC storage (Pg)		SOC (kg m ⁻²)	SOC storage (Pg)	SOC (kg m ⁻²)	SOC storage (Pg)	
Soil depth (m)		QTP	HHRB					
0~1 m	~	11.0±3.9	0.3±0.1	6.9±3.6	5.3±2.8	3.8±1.5	0.7±0.3	17.3±5.3
1~2 m	16.7±4.7	4.9±1.4	0.2±0.1	6.5 ±2.2	5.0±1.7	3.0±1.3	0.5±0.2	10.6±2.7
Total (Pg)		16.4±5.2			10.3±2.7		1.2±0.3	27.9±6.2

Geological stratigraphies	Quaternary			Triassic		Permian		Total (Pg)
	SOC (kg m ⁻²)	SOC storage (Pg)		SOC (kg m ⁻²)	SOC storage (Pg)	SOC (kg m ⁻²)	SOC storage (Pg)	
Soil depth (m)		QTP	HHRB					
2~3 m	9.8±8.4	1.9±1.6	0.1±0.06	9.6±4.5	2.3±1.1	5.6±0.9	0.8±0.1	5.1±1.4
3~25 m	134.9±115.3	26.2±22.4	2.3±1.4	281.9±191.7	67.1±45.6	234.2±86.0	31.6±11.6	127.2±37.3
Total (Pg)		30.5±16.6			69.4±52.8		32.4±20.2	132.3±76.8


459 Figure 1



460

461


462 Figure 2



463

464


465 Figure 3



466

467

468 Figure 4



469