

This discussion paper is/has been under review for the journal The Cryosphere (TC).  
Please refer to the corresponding final paper in TC if available.

# Thin-ice dynamics and ice production in the Storfjorden polynya for winter-seasons 2002/2003–2013/2014 using MODIS thermal infrared imagery

A. Preußer, S. Willmes, G. Heinemann, and S. Paul

Dept. of Environmental Meteorology, Fac. of Regional and Environmental Sciences,  
University of Trier, Trier, Germany

Received: 10 October 2014 – Accepted: 26 October 2014 – Published: 13 November 2014

Correspondence to: A. Preußer (preusser@uni-trier.de)

Published by Copernicus Publications on behalf of the European Geosciences Union.

TCD

8, 5763–5791, 2014

Thin-ice dynamics  
and ice production in  
the Storfjorden  
polynya

A. Preußer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Abstract

Spatial and temporal characteristics of the Storfjorden polynya, which forms regularly in the proximity of the islands Spitsbergen, Barentsøya and Edgeøya in the Svalbard archipelago under the influence of strong north-easterly winds, have been investigated for the period 2002/2003 to 2013/2014 using thermal infrared satellite imagery. Thin-ice thicknesses were calculated from MODIS ice-surface temperatures, combined with ECMWF ERA-Interim atmospheric reanalysis data in an energy-balance model. Based on calculated thin-ice thicknesses, associated quantities like polynya area and total ice production were derived and compared to previous remote sensing and modeling studies. A basic coverage-correction scheme was applied to account for cloud-gaps in the daily composites. The sea ice in the Storfjorden area experiences a late fall freeze-up in several years over the 12 winter-period, with an increasing frequency of large polynya events until the end of December. During the regarded period, the mean polynya area is  $4555.7 \pm 1542.9 \text{ km}^2$ . The average ice production in the fjord is estimated with  $28.3 \pm 8.5 \text{ km}^3$  per winter and therefore lower than in previous studies. Despite this comparatively short record of 12 winter-seasons, a significant positive trend of  $20.2 \text{ km}^3$  per decade could be detected, which contrasts earlier reports of a slightly negative trend in accumulated ice production prior to 2002. Derived estimates underline the importance of this relatively small coastal polynya system considering its contribution to the cold halocline layer through salt release during ice formation processes.

## 1 Introduction

Recurrent polynyas are considered to play an important role for sea ice production and ocean circulation (Barber and Massom, 2007). The knowledge of ice production in the Arctic is of vital interest for the understanding of the Arctic climate system and the verification of climate and ocean models. Divergent ice motions due to wind are the main cause for the opening of near-coastal polynyas (Smith et al., 1990). During winter, these

TCD

8, 5763–5791, 2014

### Thin-ice dynamics and ice production in the Storfjorden polynya

A. Preußner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



areas of open water or thin ice, exposed to the cold atmosphere, are characterized by strong ice production and brine release and modify the atmospheric boundary layer (Ebner et al., 2011; Bauer et al., 2013). The timing, duration and size of a polynya can often have large interannual variations because of the general interannual variability of atmosphere-ice-ocean interaction processes (Morales-Maqueda et al., 2004). A precise derivation of the thin-ice thickness (TIT) distribution, the areal extent of a polynya (here defined as open water and thin-ice below 0.2 m) and the amount of local sea ice production is therefore an important step towards a comprehensive understanding of physical processes in the Arctic shelf regions and the Arctic sea ice cover in general. Ice production in thin-ice regions can either be estimated by the use of remote sensing methods (Willmes et al., 2010), or by using model approaches. The use of thermal infrared and microwave remote sensing data offers the opportunity for daily monitoring of thin-ice thicknesses and their distribution on large spatial scales (Yu and Rothrock, 1996; Willmes et al., 2010; Adams et al., 2013).

Storfjorden (see Fig. 1), situated in the southeastern part of the Svalbard archipelago between the main island of Spitsbergen in the west and Barentsøya/Edgeøya in the east, is a region of high coastal polynya activity during winter, as reported by a number of earlier studies (e.g. Haarpaintner et al., 2001; Skogseth et al., 2004). Two shallow straits, Heleysundet and Freemansundet, connect the northern part of Storfjorden with the Barents Sea and show tidally induced currents as well as exchange processes between water masses inside the fjord and the Barents Sea (Skogseth et al., 2013).

Persistent and strong northerly to north-easterly winds are the main cause for larger polynya openings inside Storfjorden. These opening events may result in large volumes of brine-enriched shelf water, which can be significant regarding the total amount of bottom water in the Arctic (Skogseth et al., 2004; Hendricks et al., 2011). Besides thermodynamically induced ice growth in Storfjorden during wintertime, the dynamical part due to rafting and deformation processes may play a substantial role in the overall ice thickness distribution, as southerly (onshore) winds can potentially advect sea ice from the Barents Sea into the fjord (Hendricks et al., 2011). An estimation of

TCD

8, 5763–5791, 2014

## Thin-ice dynamics and ice production in the Storfjorden polynya

A. Preußner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



thermodynamic ice production (IP) within the open water and thin-ice area (i.e. Storfjorden polynya) has been conducted in previous studies using mainly microwave remote sensing (both passive and active systems) in combination with different model approaches. Therein, values are ranging between 20 and 35–45 km<sup>3</sup> for single winter-seasons (November–May) between 1997/1998–2000/2001 (Haarpaintner et al., 2001; Skogseth et al., 2004, 2005) and 47 km<sup>3</sup> when averaged over the winter-seasons (October/November–June/July) from 2002/2003–2011/2012 (Jardon et al., 2014).

Based on daily empirically derived ice thickness distributions in thin-ice areas at a comparatively high spatial resolution, we perform a broad investigation on the associated quantities polynya area and ice-production for the period between 2002/2003 and 2013/2014 using high-resolution Moderate Resolution Imaging Spectroradiometer (MODIS) thermal infrared data. The presented results will aid to get a more comprehensive understanding of the polynya dynamics in Storfjorden as well as associated processes over both annual and interannual timescales (in the course of this paper, the term “annual” refers to one winter-season from November–March). Further, results will contribute to pan-arctic studies of polynya dynamics.

## 2 Data and methods

### 2.1 MODIS ice-surface temperatures

Thin-ice thicknesses are computed using the MOD/MYD29 sea-ice product (Hall et al., 2004; Riggs et al., 2006), derived from MODIS satellite data. We used data from both MODIS instruments onboard Terra and Aqua polar-orbiting satellite platforms. It is available from the National Snow and Ice Data Center (NSIDC) located in Boulder, USA (<ftp://n5eil01u.ecs.nsidc.org/SAN/>). The product contains swath data of ice-surface temperatures (IST). IST swaths have a spatial resolution of 1 km × 1 km at nadir. Under ideal conditions (clear sky and low amount of water vapor in the atmosphere), the accuracy is given with ±1.6 K (Hall et al., 2004). An important remark to

TCD

8, 5763–5791, 2014

## Thin-ice dynamics and ice production in the Storfjorden polynya

A. Preußner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the accuracy-assessment conducted by Riggs et al. (2006) is the restriction to daytime pixels only, as no nighttime validation has been done.

For our analysis, swaths covering the Storfjorden area were extracted using meta-data information for each MODIS swath, resulting in 27 individual swaths per day on average for later composite generation (nighttime scenes). The total number of incorporated MODIS swaths is illustrated in Table 1. As MODIS swath data presumably suffer from inherent panoramic distortion effects, all IST-swaths were mapped onto a  $2\text{ km} \times 2\text{ km}$  ( $0.018^\circ$ ) equirectangular grid covering the southern part of Svalbard ( $75\text{--}80^\circ\text{ N}$ ,  $10\text{--}30^\circ\text{ E}$ ).

A polynya mask (red area in Fig. 1) is applied in order to cover an area comparable to previous studies, as well as to limit the probability of remaining hidden cloud artifacts. The mask comprises a total area of  $12\,594\text{ km}^2$  and has its southern limit at  $77^\circ\text{ N}$  (following Skogseth et al., 2004), where a 120 m deep sill serves as kind of a boundary for the accumulation of brine-enriched shelf water in the Storfjorden basin (see Fig. 1).

## 2.2 ERA-interim reanalysis

For the calculation of TIT, atmospheric variables from the ERA-Interim reanalysis product (Dee et al., 2011) are used to provide 2 m temperature, 2 m dew point temperature, 10 m wind speed components ( $u$  and  $v$ ) and the mean sea-level pressure at a resolution of 6 h. The data set is provided by the European Center for Medium-Range Weather Forecasts (ECMWF) in a horizontal resolution of  $0.75^\circ$  ( $\sim 79\text{ km}$ ). As the satellite data set is by far spatially higher resolved than the atmospheric data set, a linear interpolation of the ERA-Interim data is performed to match the MODIS data. In the course of the TIT-calculations, each MODIS swath is linked to the closest time step of the atmospheric fields.

TCD

8, 5763–5791, 2014

### Thin-ice dynamics and ice production in the Storfjorden polynya

A. Preußer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 2.3 Thin-ice thickness retrieval using a 1-D thermodynamic surface energy model

In order to derive daily TIT-distributions, we use an approach that follows the work of Yu and Rothrock (1996), Yu and Lindsay (2003) and Drucker et al. (2003), with most recent improvements and modifications made by Willmes et al. (2010, 2011) and Adams et al. (2013). The applied method utilizes the distinct relationship between ice surface temperature and thickness of thin ice depending on atmospheric energy fluxes. The IST of very thin ice (with no snow cover) is close to the freezing point of the upper ocean layer and decreasing IST can be observed for thicker ice (Drucker et al., 2003; Kwok et al., 2007). However, IST is governed by atmospheric radiation fluxes and turbulent fluxes of heat, which does not allow for a simple relationship between IST and ice thickness. The thin-ice thickness  $h_{ice}$  is therefore calculated using a 1-D thermodynamic sea ice model.

Several assumptions have to be made to apply this model. First, the temperature profile through the ice is assumed to be linear. This approximation is valid for  $h_{ice} \leq 0.5$  m (Drucker et al., 2003). Second, the water temperature at the boundary between ice and ocean is constant and at its freezing point ( $T_f = 271.35$  K). Third, the ice is assumed to be free of snow. The ice thickness is calculated by using MODIS IST and ERA-Interim reanalysis data. The method is based on the condition, that the conductive heat flux through the ice ( $Q_{ice}$ , Eq. 1) equals the total heat flux to the atmosphere ( $Q_{atm}$ , Eq. 2). This implies, that all energy loss at the ice-surface is compensated by the conductive heat flux  $Q_{ice}$ .

$$Q_{ice} = \kappa_{ice} \cdot \frac{(T_{surf} - T_f)}{h_{ice}} \quad (1)$$

$$Q_{atm} = Q_0 - H_0 - E_0 \quad (2)$$

where  $Q_0$  is the net radiation balance,  $H_0$  and  $E_0$  are the turbulent fluxes of sensible heat and latent heat, respectively,  $\kappa_{ice}$  is the thermal conductivity of sea ice,  $T_{surf}$  is the

TCD

8, 5763–5791, 2014

### Thin-ice dynamics and ice production in the Storfjorden polynya

A. Preußner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



# Thin-ice dynamics and ice production in the Storfjorden polynya

A. Preußner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



surface temperature,  $T_f$  the freezing point of sea water and  $h_{ice}$  is the ice thickness. Because of difficulties with the parametrization of shortwave radiation terms (Adams et al., 2013), only nighttime scenes were incorporated into the calculations (by checking the solar incidence angle on a pixel-by pixel basis), so that the radiation balance  $Q_0$  simplifies to the long wave radiation terms  $L \downarrow$  and  $L \uparrow$ . Upwelling longwave radiation  $L \uparrow$  is calculated using the IST from MOD/MYD29 with the surface emissivity  $\epsilon_s$  set to 1.  $L \downarrow$  can be obtained using ERA-Interim 2 m temperatures and the atmospheric emissivity  $\epsilon_{atm}$ , calculated using the formulation of Jin et al. (2006). Turbulent fluxes of sensible ( $H_0$ ) and latent ( $E_0$ ) heat in Eq. (2) are calculated using an iterative bulk approach by Launiainen and Vihma (1990), where near surface stratification is considered using Monin–Obukhov similarity theory and associated universal functions for the heat transfer coefficients  $C_H$  and  $C_E$ . The 10 m wind speed is interpolated to the 2 m level. A constant roughness length for momentum  $z_0 = 1 \times 10^{-3}$  m is used. As could be shown by Adams et al. (2013), the standard approach used in previous studies using constant heat transfer coefficients tends to underestimate the ice thickness by approximately 7 cm on average, when compared to an iteratively calculated heat transfer coefficient. To obtain the ice thickness  $h_{ice}$ , the total atmospheric flux  $Q_{atm}$  (Eq. 2) is set equal to the conductive heat flux through the ice ( $Q_{ice}$ ; Eq. 1) and Eq. (1) is solved for the ice thickness  $h_{ice}$ .

TIT is calculated using a value for the thermal conductivity of  $\kappa_{ice} = 2.03 \text{ Wm}^{-2} \text{ K}^{-1}$  (Drucker et al., 2003). It is assumed to be uniform at a given pixel location and therefore hypothetical (Tamura and Ohshima, 2011). The described method is only applicable to clear sky conditions, as clouds and fog strongly influence the accuracy of the recorded IST (Riggs et al., 2006). We constrain our analysis to TIT-values  $\leq 0.2$  m, as this range is regarded sufficient to get reliable results for ice production (Yu and Rothrock, 1996; Adams et al., 2013).

In a sensitivity analysis of the above described method, Adams et al. (2013) state an uncertainty for the TIT retrieval of  $\pm 1.0$ ,  $\pm 2.1$  and  $\pm 5.3$  cm for TIT classes 0–5, 5–10 and 10–20 cm, respectively.

### 2.3.1 Calculation of daily TIT-composites

From all available MODIS swaths covering the Storfjorden polynya region, daily composites of IST and TIT were computed. The TIT is first calculated from each swath on its own with the procedure described in Sect. 2.3. Subsequently, the mean TIT is computed pixel-wise and stored with its corresponding IST value.

By using daily composites, the daily IST-coverage within the polynya mask is significantly increased compared to single swaths, reaching an average value of 82 % for the winter-seasons 2002/2003 to 2013/2014. Figure 2 shows the seasonal and inter-annual variability of IST-coverage. Noteworthy is a very low coverage in November to early January 2009/2010, as well as generally less frequent coverage at the beginning (November) and end (March) of each winter-season. The latter effect originates from the previously stated restriction to nighttime scenes and consequently a reduction in the amount of available MODIS swaths. Hence, it is also the main reason why the months of October and April were left out in our investigations.

### 2.3.2 Derivation of ice production and polynya area

Following Tamura et al. (2007, 2008) and Willmes et al. (2011), daily ice production rates are calculated from the heat loss at the ice surface by assuming that the entire heat loss to the overlying atmosphere contributes to new ice formation.

$$\frac{\partial h}{\partial t} = \frac{-\bar{Q}_{\text{ice}}}{\rho_{\text{ice}} \cdot L_f} \quad (3)$$

Therein,  $\frac{\partial h}{\partial t}$  stands for the ice production rate,  $\bar{Q}_{\text{ice}}$  is the daily mean conductive heat flux through the ice,  $\rho_{\text{ice}}$  is the density of sea ice (taken as  $\rho_{\text{ice}} = 910 \text{ kg m}^{-3}$ ) and  $L_f$  is the latent heat of fusion of ice ( $L_f = 0.334 \text{ MJ kg}^{-1}$ ; e.g. Tamura and Ohshima, 2011). Note that the negative sign in the right side of Eq. (3) handles the convention that  $Q_{\text{atm}}$  is considered positive when the surface gains energy, as well as considering

TCD

8, 5763–5791, 2014

## Thin-ice dynamics and ice production in the Storfjorden polynya

A. Preußner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





that ice production only takes place when there is a net energy loss from the surface. Multiplying  $\frac{\partial h}{\partial t}$  with the areal extent of each pixel in the regarded region yields the volume ice production rate  $\frac{\partial V}{\partial t}$  (IP). Ice production rates are calculated for each pixel with a TIT  $\leq 0.2$  m, and afterwards extrapolated to daily rates for later accumulation and/or averaging. Considering the total polynya area (POLA), a threshold of  $h_{\text{ice}} \leq 0.2$  m is applied to identify TIT-pixels from the daily composites as part of the POLA. The total area of those pixels is then accumulated to obtain the total daily POLA in km<sup>2</sup>.

### 2.3.3 IST-coverage correction

Cloud-induced gaps in our daily TIT and IST composites are a serious problem when we want to compare our initially calculated POLA and IP to estimates from other studies, in which the use of active and passive microwave remote sensing systems largely reduces the effect of atmospheric disturbances.

To overcome these difficulties, a simple scaling approach is applied that works under the assumption, that pixels in the uncovered part of the masked (Fig. 1) daily composites also contribute to the total POLA by approximately the same proportion as those areas that are covered by a IST-signal.

The initially calculated daily POLA is therefore scaled by a factor  $F$ :

$$F = 1/\text{coverage} \quad (4)$$

$$\text{POLA}_{\text{CC}} = \text{POLA} \cdot F. \quad (5)$$

The coverage in Eq. (4) could range from 0–1, but we apply the correction only for coverages exceeding 0.5. Similarly, estimated ice production (IP) values are also scaled by the same factor yielding a corrected ice production  $\text{IP}_{\text{CC}}$ .

Application of this correction scheme is limited to cases where the daily IST-coverage (compare Fig. 2) within the fjord (i.e. the predefined polynya mask) surpasses a threshold of 50 %. By using this threshold, unrealistically high  $\text{POLA}_{\text{CC}}$  and  $\text{IP}_{\text{CC}}$  values are avoided. Otherwise, both  $\text{POLA}_{\text{CC}}$  and  $\text{IP}_{\text{CC}}$  are linearly interpolated using bounding days with a coverage fulfilling this precondition.

### 3 Results

Figure 3 shows histograms of the mean relative TIT-distributions for ice thicknesses below 0.2 m, based on daily composites for the winters 2002/2003 to 2013/2014. A temporal differentiation between the beginning of the freezing season (November–December) and the end of the freezing season (January–March) as well as a combined distribution for the whole winter period is given. Error-bars refer to  $\pm 1$  SD of the interannual averaged distribution frequencies. Between November and March, ice thinner than 0.02 m contributes around 15 % to the total polynya area in an average winter-season. Due to a higher sensitivity of open water and very thin ice for sea smoke effects and consequently an exclusion of thin-ice pixels by the cloud mask, this thickness class is potentially biased (Willmes et al., 2011). The average ice thickness within the polynya mask is  $10 \pm 1.8$  cm. Thicker ice classes ( $> 0.1$  m) cover around half of the entire Storfjorden polynya in January to March and show overall lower contributions at the beginning of the freezing season. This follows from the high contribution of very thin ice in November to December, where over 60 % of the total POLA consists of ice thinner than 0.1 m. SD are highest in the lowest thickness classes ( $< 0.04$  m) and appear to be lower towards the end of the freezing season, possibly indicating a stronger seasonal variability of ice thickness distributions within the polynya in November and December over the twelve year record (also apparent in relative TIT-frequencies for November–December; Fig. 4).

Spatial distributions of the relative TIT-frequency for 2002/2003 to 2013/2014 are presented as a seasonal comparison between the beginning of each freezing season (November–December, Fig. 4) and the end of each freezing season (January–March, Fig. 5). Therein, a pixel-value of e.g. 0.25 means that on 25 % of all days, a TIT-threshold of 0.2 m was not exceeded. Following the previously stated definition of a polynya, these pixels represent the number of polynya-occurrences in the given period. Figure 5 shows, that the main regions for polynya development towards the end of the freezing season are at the lee-sides of Barentsøya and Edgeøya at the eastern

TCD

8, 5763–5791, 2014

## Thin-ice dynamics and ice production in the Storfjorden polynya

A. Preußner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



side of the fjord, extending more (2006–2008, 2010, 2012–2014) or less (2003–2005, 2009, 2011) further south-westwards.

At the beginning of the freezing season (Fig. 4), annual distributions show a more variable pattern of TIT-frequencies. In some years (e.g. 2007, 2009, 2012), frequencies exceeding 0.3 cover large areas within the masked area in Storfjorden, which might be an effect of a late fall freeze-up and subsequently longer periods with thin ice and open water in the fjord. In other years (e.g. 2003, 2004), the observed pattern of high TIT-frequencies is similar to typical polynya locations between January and March, indicating an earlier freeze-up of the ice cover in Storfjorden.

Frequent TIT-occurrences in the southern part of Storfjorden could originate from a generally northwards retreating ice edge in Storfjorden in some of the regarded winter-seasons. This becomes most obvious in 2006, 2008 and 2012 to 2014 (Fig. 5). Low values ( $< 0.05$ ) at the (north-) western side of Storfjorden towards the end of a freezing season (Fig. 5) further indicate typical fast-ice areas, which are most pronounced in the years 2004, 2005, 2007 and 2011.

The daily  $POLA_{CC}$  for all winters (November–March) between 2002/2003 and 2013/2014 is presented in Fig. 6. Highest  $POLA_{CC}$  values are generally found in November and December, which might be an effect of the already mentioned late fall freeze-up and therefore very thin newly formed ice (compare Figs. 3a and 4) in large parts of Storfjorden. From 2006 to 2008 and 2012 to 2013, this effect even extends well into January with a series of large  $POLA_{CC}$  events ( $> 5000 \text{ km}^2$ ) developing in February and March.

The mean (November to March) POLA for the entire regarded period amounts to  $3511.8 \pm 999.6 \text{ km}^2$ , with an increase of 29 % for  $POLA_{CC}$  when the presented coverage correction is applied. Around 60 % of this area consists of ice thinner than 10 cm. A complete overview on wintertime mean POLA and  $POLA_{CC}$  values is given in Table 2.

A spatial overview of the resulting accumulated daily ice production rates (in  $\text{m winter}^{-1}$ ), averaged between 2002/2003 and 2013/2014 is presented in Fig. 7. Largest IP occurs in regions which also show high TIT-frequencies (Fig. 5). Average IP

TCD

8, 5763–5791, 2014

## Thin-ice dynamics and ice production in the Storfjorden polynya

A. Preußner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



values in the western part of Storfjorden are generally lower, as thin-ice in these areas is mainly present at the beginning of the freezing season.

Maximum daily ice production rates in the fjord can reach as high as  $26 \text{ cm d}^{-1}$  (winter-season 2006/2007), with the highest rates occurring on average in the northern part of Storfjorden (north of  $77.5^\circ \text{ N}$ ).

Figure 8 shows values of accumulated ice production per winter-season from November to March, both with and without applied coverage-correction. On average, the correction scheme increases the wintertime ice production by approximately 30 %. While average IP-values range between  $21.7 \pm 5.9 \text{ km}^3$  (no CC) and  $28.3 \pm 8.5 \text{ km}^3$  (CC), a high interannual variability of ice production is found, with values as low as  $16.1 \text{ km}^3$  (CC; no CC  $-16\%$ ) in 2002/2003 and up to  $47.2 \text{ km}^3$  (CC; no CC  $-31\%$ ) in 2012/2013. A detailed overview for each winter-season is given in Table 2. Regardless of an applied coverage correction, a significant ( $p \leq 0.05$ ) positive trend of  $1.2\text{--}2.0 \text{ km}^3 \text{ yr}^{-1}$  over the examined 12-winter period can be observed.

## 4 Discussion

In Skogseth et al. (2004, 2005), a manual classification of several ERS-2 synthetic aperture radar (SAR) images from 1998–2002 was performed to derive parameters for a wind-driven polynya width model (Haarpaintner et al., 2001), which gives estimates of ice production and brine-enriched shelf water based on satellite imagery, wind data from nearby weather stations and surface hydrography. Ice production was calculated from surface heat balance and includes contributions from open water (frazil ice), thin ice as well as fast and pack ice inside the Storfjorden basin north of  $77^\circ \text{ N}$ . This setup was later extended to a 33 year record of modeled total ice production covering the winters 1970 to 2002 using model parameters derived from the examined 5-winter period (Skogseth et al., 2005). While the average total ice production for 1998 to 2002 was estimated to  $43.6 \pm 9.7 \text{ km}^3$ , it was slightly lower for the 33 year time series with only  $39.9 \pm 11.7 \text{ km}^3$ . When comparing these numbers to our average estimate of

TCD

8, 5763–5791, 2014

### Thin-ice dynamics and ice production in the Storfjorden polynya

A. Preußer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



28.3 ± 8.5 km<sup>3</sup> (CC) for 2002/2003 to 2013/2014, one has to keep in mind that Skogseth et al. (2004, 2005) also include ice production under fast- and pack-ice inside the fjord, while our study concentrates on open water and thin-ice areas only. Assuming an average contribution of 25 % from fast-ice and pack-ice areas (Table 3 in Skogseth et al., 2005) and after reducing annually accumulated total ice production accordingly, the average IP (1970 to 2002) inside open water and thin-ice areas of Skogseth et al. (2005) is lowered to approximately 30 ± 9 km<sup>3</sup> which is within the here presented SD. Skogseth et al. (2005) give some further information on the interannual variability of ice production within the Storfjorden polynya, although it is stated that the uncertainty for the 33 year time series is increased. The presented 33 year time series shows a slightly negative and non-significant trend of approximately -2 km<sup>3</sup> decade<sup>-1</sup>, while the here presented last 12 winter-seasons show a positive trend of 20.2 km<sup>3</sup> decade<sup>-1</sup>. However, the long-term model-estimates by Skogseth et al. (2005) indicate the presence of multi-decadal fluctuations.

Jardon et al. (2014) developed an ice production model which uses Advanced Microwave Scanning Radiometer – EOS (AMSR-E) sea ice concentration data to derive daily open water fractions. Heat flux calculations over water were performed using ERA interim reanalysis data in a bulk algorithm. In contrast to our results, their estimates of POLA and IP incorporate both frazil ice growth as well as ice growth under ice thicker than 20 cm. In comparison, our results should profit from the enhanced resolution of MODIS and daily derived TIT-distributions. Nevertheless, derived POLA from TIT ≤ 0.2 m in our study compares well with POLA derived from AMSR-E sea ice concentration data. Both wintertime mean POLA and POLA<sub>CC</sub> are within the 25th and 75th percentile of the estimated open water area by Jardon et al. (2014) in almost every winter-season, except for 2004/2005 and 2010/2011, where both values derived by MODIS exceed the 75th percentile. Discrepancies in derived total ice production are obvious, but can be partly explained by differences in the regarded time frame of each winter-season. While Jardon et al. (2014) analyze the full period with sea ice in the fjord (roughly October/November to June/July), we focus on the November to March

# Thin-ice dynamics and ice production in the Storfjorden polynya

A. Preußner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





spans the months from September to May, could therefore explain part of the difference.

## 5 Summary and conclusions

In this study, we presented a comprehensive overview of daily thin-ice thickness distributions and resulting thermodynamic ice-production rates for the Storfjorden polynya, located in the southern part of the Svalbard archipelago. The investigation period covers the winter-seasons 2002/2003–2013/2014 from November to March. Our results are complementary to previous studies in the Storfjorden region (e.g. Skogseth et al., 2004, 2005; Hendricks et al., 2011) and underline the importance of this polynya considering the total amount of ice production and accompanied salt release each winter-season, despite its comparatively small spatial extent. Within the polynya, ice thicknesses below 10 cm are dominating in the course of an average winter-season and cover larger areas in the eastern part of Storfjorden. This enables ice-production rates, which sum up to an average value of  $28.3 \pm 8.5 \text{ km}^3$  per winter-season. There is a positive trend in accumulated ice production over the last 12 winter-seasons which contrasts earlier indications of a slightly negative trend prior to 2002 (Skogseth et al., 2005).

Compared to other studies which rely on active and passive microwave remote sensing, the use of MODIS IST has distinct drawbacks considering the effect of clouds, which produce data gaps if correctly identified by the MODIS cloud mask and possibly erroneous IST-values if nighttime clouds remain undetected. By composing daily composites, data gaps in the Storfjorden region were minimized resulting in an average IST-coverage of 82 % within the applied polynya mask. The application of a coverage-correction scheme yielded plausible adjustments of approximately 30 % to the total polynya area and ice production, although its current basic approach offers room for future improvements like e.g. including long-term spatial statistics. Still, this comprehensive data set will be of high value for a variety of climate and ocean applications,

### Thin-ice dynamics and ice production in the Storfjorden polynya

A. Preußner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





including the provision of quasi-daily high-resolution thin-ice thickness charts for regional climate and ocean models.

*Acknowledgements.* The authors want to thank the National Snow and Ice Data Center (NSIDC) as well as the European Center for Medium-Range Weather Forecasts (ECMWF) for providing the MODIS Sea Ice product and the ERA-Interim atmospheric reanalysis data. This work was part of the German-Russian cooperation “WTZ RUS: System Laptev Sea: TRANSDRIFT” funded by the Federal Ministry of Education and Research (BMBF) under grant 03G0833D.

## References

- Adams, S., Willmes, S., Schroeder, D., Heinemann, G., Bauer, M., and Krumpen, T.: Improvement and sensitivity analysis of thermal thin-ice retrievals, *IEEE T. Geosci. Remote*, 51, 3306–3318, 2013. 5765, 5768, 5769
- Barber, D. G. and Massom, R. A.: The role of sea ice in Arctic and Antarctic polynyas, in: *Polynyas – Windows to the World*, edited by: Smith, W. O. and Barber, D. G., Elsevier Oceanography Series, 1–54, doi:10.1016/s0422-9894(06)74001-6, 2007. 5764
- Bauer, M., Schröder, D., Heinemann, G., Willmes, S., and Ebner, L.: Quantifying polynya ice production in the Laptev Sea with the COSMO model, *Polar Res.*, 32, 20922, doi:10.3402/polar.v32i0.20922, 2013. 5765
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, I., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J.-N., and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Q. J. Roy. Meteorol. Soc.*, 137, 553–597, 2011. 5767
- Drucker, R., Martin, S., and Moritz, R.: Observations of ice thickness and frazil ice in the St. Lawrence Island polynya from satellite imagery, upward looking sonar, and salinity/temperature moorings, *J. Geophys. Res.*, 108, 3149, doi:10.1029/2001jc001213, 2003. 5768, 5769

TCD

8, 5763–5791, 2014

## Thin-ice dynamics and ice production in the Storfjorden polynya

A. Preußner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





# Thin-ice dynamics and ice production in the Storfjorden polynya

A. Preußner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Ebner, L., Schröder, D., and Heinemann, G.: Impact of Laptev Sea flaw polynyas on the atmospheric boundary layer and ice production using idealized mesoscale simulations, *Polar Res.*, 30, 7210, doi:10.3402/polar.v30i0.7210, 2011. 5765
- Haarpaintner, J., Gascard, J.-C., and Haugan, P. M.: Ice production and brine formation in Storfjorden, Svalbard, *J. Geophys. Res.-Oceans*, 106, 14001–14013, doi:doi/10.1029/1999JC000133/full, 2001. 5765, 5766, 5774
- Hall, D., Key, J., Casey, K., Riggs, G., and Cavalieri, D.: Sea ice surface temperature product from MODIS, *IEEE T. Geosci. Remote*, 42, 1076–1087, 2004. 5766
- Hendricks, S., Gerland, S., Smedsrud, L., Haas, C., Pfaffhuber, A., and Nilsen, F.: Sea-ice thickness variability in Storfjorden, Svalbard, *Ann. Glaciol.*, 52, 61–68, 2011. 5765, 5777
- Iwamoto, K., Ohshima, K. I., and Tamura, T.: Improved mapping of sea ice production in the Arctic Ocean using AMSR-E thin ice thickness algorithm, *J. Geophys. Res.-Oceans*, 119, 3574–3594, doi:10.1002/2013jc009749, 2014. 5776
- Jakobsson, M., Mayer, L., Coakley, B., Dowdeswell, J. A., Forbes, S., Fridman, B., Hodnesdal, H., Noormets, R., Pedersen, R., Rebesco, M., Schenke, H. W., Zarayskaya, Y., Accettella, D., Armstrong, A., Anderson, R. M., Bienhoff, P., Camerlenghi, A., Church, I., Edwards, M., Gardner, J. V., Hall, J. K., Hell, B., Hestvik, O., Kristoffersen, Y., Marcussen, C., Mohammad, R., Mosher, D., Nghiem, S. V., Pedrosa, M. T., Travaglini, P. G., and Weatherall, P.: The international bathymetric chart of the Arctic Ocean (IBCAO) version 3.0, *Geophys. Res. Lett.*, 39, L12609, doi:10.1029/2012gl052219, 2012. 5784
- Jardon, F., Vivier, F., Bouruet-Aubertot, P., Lourenço, A., Cuypers, Y., and Willmes, S.: Ice production in Storfjorden (Svalbard) estimated from a model based on AMSR-E observations: Impact on water mass properties, *J. Geophys. Res.-Oceans*, 119, 377–393, 2014. 5766, 5775, 5776
- Jin, X., Barber, D., and Papakyriakou, T.: A new clear-sky downward longwave radiative flux parameterization for Arctic areas based on rawinsonde data, *J. Geophys. Res.*, 111, D24104, doi:10.1029/2005jd007039, 2006. 5769
- Kwok, R., Comiso, J. C., Martin, S., and Drucker, R.: Ross sea polynyas: response of ice concentration retrievals to large areas of thin ice, *J. Geophys. Res.*, 112, C12012, doi:10.1029/2006jc003967, 2007. 5768
- Launiainen, J. and Vihma, T.: Derivation of turbulent surface fluxes – an iterative flux-profile method allowing arbitrary observing heights, *Environ. Softw.*, 5, 113–124, 1990. 5769

- Morales-Maqueda, M., Willmott, A., and Biggs, N.: Polynya dynamics: a review of observations and modeling, *Rev. Geophys.*, 42, 1–37, 2004. 5765
- Riggs, G., Hall, D., and Salomonson, V.: MODIS Sea Ice Products User Guide to Collection 5, NSIDC, University of Colorado, Boulder, CO, USA, 2006. 5766, 5767, 5769
- 5 Skogseth, R., Haugan, P., and Haarpaintner, J.: Ice and brine production in Storfjorden from four winters of satellite and in situ observations and modeling, *J. Geophys. Res.-Oceans*, 109, 73–88, doi:10.1029/158gm07, 2004. 5765, 5766, 5767, 5774, 5775, 5777
- Skogseth, R., Fer, I., and Haugan, P. M.: Dense-water production and overflow from an arctic coastal polynya in Storfjorden, in: *The Nordic Seas: An Integrated Perspective*, 73–88, doi:10.1029/158GM07, 2005. 5766, 5774, 5775, 5776, 5777
- 10 Skogseth, R., McPhee, M. G., Nilsen, F., and Smedsrud, L. H.: Creation and tidal advection of a cold salinity front in Storfjorden: 1. Polynya dynamics, *J. Geophys. Res.-Oceans*, 118, 3278–3291, doi:10.1002/jgrc.20231/full, 2013. 5765
- Smith, S. D., Muench, R. D., and Pease, C. H.: Polynyas and leads: an overview of physical processes and environment, *J. Geophys. Res.*, 95, 9461–9479, 1990. 5764
- Tamura, T. and Ohshima, K. I.: Mapping of sea ice production in the Arctic coastal polynyas, *J. Geophys. Res.*, 116, C07030, doi:10.1029/2010jc006586, 2011. 5769, 5770, 5776
- Tamura, T., Ohshima, K. I., Markus, T., Cavalieri, D. J., Nishihashi, S., and Hirasawa, N.: Estimation of thin ice thickness and detection of fast ice from SSM/I data in the Antarctic Ocean, *J. Atmos. Ocean. Tech.*, 24, 1757–1772, 2007. 5770
- 20 Tamura, T., Ohshima, K. I., and Nishihashi, S.: Mapping of sea ice production for Antarctic coastal polynyas, *Geophys. Res. Lett.*, 35, L07606, doi:10.1029/2007gl032903, 2008. 5770
- Willmes, S., Krumpen, T., Adams, S., Rabenstein, L., Haas, C., Hoelemann, J., Hendricks, S., and Heinemann, G.: Cross-validation of polynya monitoring methods from multisensor satellite and airborne data: a case study for the Laptev Sea, *Can. J. Remote Sens.*, 36, S196–S210, 2010. 5765, 5768
- 25 Willmes, S., Adams, S., Schröder, D., and Heinemann, G.: Spatio-temporal variability of polynya dynamics and ice production in the Laptev Sea between the winters of 1979/80 and 2007/08, *Polar Res.*, 30, 5971, doi:10.3402/polar.v30i0.5971, 2011. 5768, 5770, 5772
- 30 Yu, Y. and Lindsay, R.: Comparison of thin ice thickness distributions derived from RADARSAT Geophysical Processor System and advanced very high resolution radiometer data sets, *J. Geophys. Res.*, 108, 3387, doi:10.1029/2002jc001319, 2003. 5768

# Thin-ice dynamics and ice production in the Storfjorden polynya

A. Preußner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Yu, Y. and Rothrock, D. A.: Thin ice thickness from satellite thermal imagery, J. Geophys. Res., 101, 25753–25766, 1996. 5765, 5768, 5769

TCD

8, 5763–5791, 2014

Thin-ice dynamics  
and ice production in  
the Storfjorden  
polynya

A. Preußner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Thin-ice dynamics  
and ice production in  
the Storfjorden  
polynya**

A. Preußner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Table 1.** Total number of individual MODIS swaths, which were incorporated into the TIT-calculations. Numbers are given for each winter-season from November to March.

Winter-season	Number of MODIS swaths
2002–2003	4054
2003–2004	4023
2004–2005	4062
2005–2006	4073
2006–2007	3990
2007–2008	4069
2008–2009	4135
2009–2010	4142
2010–2011	4205
2011–2012	4129
2012–2013	4137
2013–2014	4244
Sum	49 263

# Thin-ice dynamics and ice production in the Storfjorden polynya

A. Preußner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion




**Table 2.** Accumulated ice production in  $\text{km}^3 \text{winter}^{-1}$  and average polynya area (in  $\text{km}^2$ ) each winter-season (November to March) in the Storfjorden polynya, together with the interannual average (Mean) and its SD. CC denote values after coverage correction, with relative increase compared to uncorrected values in brackets. All values are calculated within the predefined polynya mask (Fig. 1).

	Acc. IP ( $\text{km}^3$ )	Acc. IP <sub>CC</sub> ( $\text{km}^3$ )	POLA ( $\text{km}^2$ )	POLA <sub>CC</sub> ( $\text{km}^2$ )
2002–2003	13.9	16.1 (+16 %)	2185.2	2429.6 (+11 %)
2003–2004	15.2	16.7 (+10 %)	2186.6	2382.0 (+9 %)
2004–2005	19.2	23.7 (+24 %)	2684.7	3296.6 (+23 %)
2005–2006	20.6	25.1 (+22 %)	3934.8	4750.6 (+21 %)
2006–2007	19.4	25.3 (+31 %)	3473.9	4524.2 (+30 %)
2007–2008	27.4	33.1 (+21 %)	4491.0	5472.5 (+22 %)
2008–2009	21.6	24.6 (+14 %)	2965.4	3353.3 (+13 %)
2009–2010	17.3	30.0 (+73 %)	3390.1	6357.0 (+88 %)
2010–2011	23.2	31.9 (+38 %)	2996.0	4024.7 (+34 %)
2011–2012	21.5	29.6 (+38 %)	4087.6	5001.5 (+22 %)
2012–2013	36.0	47.2 (+31 %)	5592.4	7388.0 (+32 %)
2013–2014	25.4	36.5 (+44 %)	4154.3	5688.4 (+37 %)
Mean	21.7	28.3 (+30 %)	3511.8	4555.7 (+29 %)
SD	5.9	8.5	999.6	1542.9




**Figure 1.** Map of the Svalbard archipelago in western Barents Sea. The study region is located in Storfjorden, which is surrounded by the main island Spitsbergen in the west and Barentsøya/Edgeøya in the east. The applied polynya mask is marked in red, enclosing the Storfjorden basin north of the sill at 77° N. Bathymetric data by Jakobsson et al. (2012) (IBCAO v3.0).





**Figure 3.** Relative TIT-distribution in the Storfjorden polynya, with ice thickness classes of 2 cm range ( $x$  axis). Input data is based on daily TIT-composites. The bars indicate the mean relative distribution of each thickness class from the total number of TIT  $\leq 0.2$  m appearances between the winter-seasons 2002/2003 to 2013/2014. Error bars refer to  $\pm 1$  SD. Besides an overview of the complete freezing season from November to March **(c)**, it is further separated between the months November to December **(a)** as well as January to March **(b)**.





**Figure 4.** Early freezing season (November to December) relative frequency distribution of  $TIT \leq 0.2$  m for 2002 to 2013, based on daily TIT-composites.

# Thin-ice dynamics and ice production in the Storfjorden polynya

A. Preußner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back


Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Figure 5.** Late freezing season (January to March) relative frequency distribution of TIT  $\leq 0.2$  m for 2003 to 2014, based on daily TIT-composites.

# Thin-ice dynamics and ice production in the Storfjorden polynya

A. Preußer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back


Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Figure 6.** Hovmoeller diagram of daily  $POLA_{CC}$  ( $TIT \leq 0.2$  m) in the Storfjorden polynya between 2002/2003 and 2013/2014.

# Thin-ice dynamics and ice production in the Storfjorden polynya

A. Preußer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



# Thin-ice dynamics and ice production in the Storfjorden polynya

A. Preußer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶


Back

Close

Full Screen / Esc

Printer-friendly Version


Interactive Discussion



**Figure 7.** Spatial distribution of the interannual mean accumulated IP rate in  $\text{m winter}^{-1}$ . Values are first accumulated for each winter-season from November to March and afterwards averaged for 2002/2003 to 2013/2014.

# Thin-ice dynamics and ice production in the Storfjorden polynya

A. Preußer et al.



**Figure 8.** Annual wintertime accumulated ice production in the Storfjorden polynya, given in  $\text{km}^3 \text{ winter}^{-1}$ . Estimations are based on daily heat flux calculations using the daily derived TIT-composites. Special emphasis is given to the effect of an applied coverage correction (CC). Dotted lines show linear trend estimations for both IP and  $\text{IP}_{\text{CC}}$ .

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)
