

1

2

Brief Communication:

3

**Contending estimates of 2003-2008 glacier mass balance
over the Pamir-Karakoram-Himalaya**

4

5

Andreas Kääb¹, Christopher Nuth¹, Désirée Treichler¹, Etienne Berthier²

7

[1] {Department of Geosciences, University of Oslo, P.O. Box 1047, Oslo, Norway}

8

[2] {CNRS, Université de Toulouse, LEGOS, 14 avenue Ed. Belin, Toulouse 31400, France }

9

Correspondence to: Andreas Kääb (kaeab@geo.uio.no)

10

11

Abstract

12

We present glacier thickness changes over the entire Pamir-Karakoram-Himalaya arc based on ICESat satellite altimetry data for 2003-2008. We highlight the importance of C-band penetration for studies based on the SRTM elevation model. To the very east and west of our study area, this penetration seems to be of larger magnitude and variability than previously assumed. The most negative rate of region-wide glacier elevation change ($< -1 \text{ m yr}^{-1}$) is observed for the East Nyainqêntanglha Shan. Conversely, glaciers of the West Kunlun Shan are slightly gaining volume, and Pamir and Karakoram seem to be on the western edge of this mass gain anomaly rather than its centre. For the Ganges, Indus and Brahmaputra basins, the glacier mass change reaches $-22 \pm 3 \text{ Gt yr}^{-1}$, about 10% of the current glacier contribution to sea-level rise. For selected catchments, we estimate glacier imbalance contributions to river runoff from a few percent to greater than 10%.

23

24

25

1 Introduction and Methods

26

27

28

29

30

31

32

33

34

Region-wide measurements of glacier volume or mass change are limited for the Pamir-Karakoram-Himalaya region, leaving room for speculation about the glacier response to climate change and its hydrological significance. Glacier mass change in high mountain Asia (or some part of it) have been obtained by (i) extrapolating the few existing in-situ mass balance series (Cogley, 2011; Bolch et al., 2012; Yao et al., 2012), (ii) using space gravimetry (Jacob et al., 2012; Gardner et al., 2013), (iii) laser altimetry (Kääb et al., 2012; Gardner et al., 2013; Neckel et al., 2014), and (iv) the differencing of digital elevation models (Gardelle et al., 2013). Between this handful of studies that narrow down the range of uncertainties for core parts of this remote mountain region, significant inconsistencies remain.

35

36

37

38

39

The aims of this study are to provide (i) a new consistent regional-scale data set from the ICESat autumn laser campaigns (2003-2008) by extending Kääb et al. (2012) to completely cover the study region by Gardelle et al. (2013) and several major river basins, (ii) to compare the results to other previous estimates of the Pamir-Karakoram-Himalaya glacier volume change, and (iii) to evaluate the contribution of glacier mass change to river runoff.

40

41

42

43

44

45

46

47

48

49

50

51

52

We follow the methods explained in Kääb et al. (2012) with a considerable extension towards the East Nyainqêntanglha Shan, the Pamir and part of the Tibetan Plateau (Fig. 1). In short, ICESat footprints are intersected with the February 2000 SRTM DEM and overlaid on the most snow-free multispectral Landsat images over ~2000-2013 to manually classify footprints into three classes; glaciers, non-glaciers and water. Glacier elevation difference trends are then estimated regionally and at a $1^\circ \times 1^\circ$ geographic grid by fitting a robust linear temporal trend to the time series of elevation differences between the SRTM DEM and individual ICESat footprint elevations. Trends are derived from autumn ICESat campaigns only (2009 ICESat winter campaigns excluded), because combined autumn and winter trends are sensitive to temporal variations in accumulation amount and timing, potentially introducing bias (see Supplement of Kääb et al., 2012). We confirm that our trends are not due to sampling bias of ICESat elevations by comparing ICESat elevation histograms with glacier hypsometry. The resulting elevation difference trends for all our zones are given in Tab. 1.

2 Glacier thickness changes

2.1 Thickening in the Karakoram and West Kunlun Shan

A first striking feature in the regional map of elevation difference trends (Fig. 1) is glacier thickness gain in the West Kunlun Shan ($\sim +0.1 \text{ m yr}^{-1}$), agreeing with in-situ mass balance and length change measurements (Yao et al., 2012). There is a southwest to northeast gradient from considerably negative glacier mass balances in Hindu Kush and Spiti Lahaul to positive values in the Pamir-Karakoram-West Kunlun Shan region (Fig. 1). This suggests the so-called Karakoram glacier mass-balance anomaly (Hewitt, 2011; Gardelle et al., 2012), or Pamir-Karakoram anomaly (Gardelle et al., 2013), is rather the edge or southwest limit of an anomaly centred more to the northeast over the West Kunlun Shan, or Tarim Basin. The anomaly seems thus indeed the result of a larger-scale meteorological or climatic feature, and peculiarities of the Karakoram topography or glaciers (e.g., surge type, hypsometry, avalanche contribution; Hewitt, 2011) do not necessarily play a decisive role. Combined, the results by Gardner et al. (2013), Neckel et al. (2014), and the glacier elevation change pattern of Fig. 1 suggest the centre of the anomaly could be located over the Tibetan Plateau.

Direct precipitation measurements in this region are scarce thus trends are uncertain. Satellite-retrieved precipitation and gauge data (Global Precipitation Climatology Project) suggest an increase of precipitation over the study region north of Karakoram and east of Pamir (Yao et al., 2012). Chinese measurements show increased precipitation over the Tibetan Plateau (personal communication Chong-Yu Xu), and Tao et al. (2014) suggest wetter conditions over the Tarim Basin since the mid 1980s. A number of abnormally wet years occurred during the early 21st century over the Tarim Basin and the Tibetan Plateau (Becker et al., 2013), in particular for the hydrological years 2003/4 and 2005/6. A recent climate modelling study proposes that stable or increasing snowfalls characterise the Karakoram anomaly on a background of increasing air temperatures (Kapnick et al., 2014). Despite the available studies and data, further research seems necessary to consolidate the precipitation and temperature trends and the reason behind the slight glacier volume gains.

2.2 Massive thinning in East Nyainqêntanglha Shan and Jammu-Kashmir

The other striking feature in Fig. 1 is the massive glacier thickness loss in the East Nyainqêntanglha Shan to the very east (between -1 m yr^{-1} and -1.7 m yr^{-1}), also consistent with the large negative mass balances and frontal retreats in this zone (Yao et al., 2012). The glaciers of East Nyainqêntanglha Shan have the smallest total elevational range in our study

85 region, indicating a large sensitivity to fluctuations in the equilibrium line altitude (Pelto,
86 2010; Loibl et al., 2014). The few available in-situ mass balance measurements in the area
87 suggest that the equilibrium line was over the vertical limits of the monitored glaciers in the
88 late 2000s, and precipitation in this zone shows the strongest long-term decrease over the
89 entire Pamir-Karakoram-Himalaya region (Yao et al., 2012; Becker et al. 2013). A similar
90 pattern of glacier shrinkage, though less distinct, is found in Jammu Kashmir within our Spiti
91 Lahaul zone and forms the cluster of second-largest thickness loss rates in this study (-0.5 to -
92 0.7 m yr⁻¹). Also here, Landsat data indicate that firn lines have risen towards high glacier
93 elevations resulting in very small accumulation areas or even their complete loss.

94 The 2003-2008 glacier thickness changes in the other study zones are all similar, on the order
95 of ~ -0.4 to -0.5 m yr⁻¹ (Tab.1), with more negative values in the Bhutan zone at the transition
96 between the East Nyaiqêntanglha and Everest zones. We note that glaciers dominated by the
97 summer monsoon (i.e. east of the Spiti Lahaul) all show thickness losses (summer-
98 accumulation type glaciers; Fujita, 2008; Kapnick et al., 2014; Maussion et al., 2014). East
99 Nyaiqêntanglha Shan, the zone with strongest glacier thickness loss, receives most
100 accumulation during March-May (spring-accumulation type; Maussion et al., 2014). The
101 glaciers with considerable winter accumulation under influence of the Westerlies show a more
102 mixed picture with stable or growing thicknesses in the Karakoram and West Kunlun Shan,
103 but thickness losses for instance in the Hindu Kush.

104 **2.3 Comparison to previous thickness change studies**

105 The following comparison to other studies uses average glacier thickness changes rather than
106 total mass changes in order to minimize effects from different delineations of study zones,
107 glacier cover areas, and density assumptions. From Hindu Kush and Karakoram in the west to
108 Nepal in the east, results of all studies agree within their errors (Tab.1). Results are most
109 sensitive to zone delineation in the Hindu Kush, reflecting the strong spatial variability of
110 glacier thickness change rates in this area (Fig. 1) and presumably also locally heterogeneous
111 glacier behaviour (Sarikaya et al., 2012; see also below for Pamir).

112 Significant differences between the results of all studies are found over East Nyainqêntanglha
113 Shan. Our results and those from Neckel et al. (2014) agree within the errors, but not with
114 Gardner et al. (2013) although all three studies are based on ICESat. While our study and
115 Neckel et al. use ICESat footprint classifications from contemporary satellite images, Gardner
116 et al. use Randolph Glacier Inventory outlines (RGI version 2.0; Pfeffer et al., 2014), which
117 contain considerable errors of commission and omission in this zone (see Table 1 in Gardelle

118 et al., 2013). Repeating our analysis with footprint classifications based on the Randolph
119 Glacier Inventory results in less negative elevation difference trends on glaciers (~ 20% less
120 negative) due to inclusion of non-glacier footprints. Vice versa, the elevation difference trends
121 on land, very close to 0 when using our own footprint classification, become negative if
122 ICESat footprints are classified using RGI, due to inclusion of glacier footprints. The
123 remaining discrepancy is presumably due to the fact that the ICESat-based results of Gardner
124 et al. are averaged from three different methods. Their results based on autumn footprints only
125 (method B, Gardner et. al., 2013) suggest a thickness change rate of -0.86 m yr^{-1} , which is in
126 closer agreement with our results.

127 At a first glance, East Nyainqêntanglha Shan results from Gardelle et al. (2013; zone called
128 there Hengduan Shan) and Gardner et al. (2013) seem to agree, but we believe this might be a
129 coincidence. First, above we argue why the Gardner et al. results might be less negative.
130 Second, the results in Gardelle et al. (2013) rely crucially on an estimate of SRTM C-band
131 penetration. Over any glacier globally, the SRTM radar waves will typically have penetrated
132 into the snow and ice, with potential largest penetration depths through snow and firn, and
133 smallest through ice (Kääb et al, 2012; Dall et al., 2001; Rignot et al. 2001). As a
134 consequence, SRTM glacier elevations do not, in general, reflect real mid-February 2000
135 glacier surface elevations but some lower horizon, the elevation of which depends among
136 others on the dielectric properties and structure of the penetrated glacier volume during the
137 SRTM campaign. For elevation difference studies where one of the elevation data sets is the
138 SRTM, its penetration depth needs to be estimated for correction, and biases in this estimate
139 translate directly into offsets in thickness change. Gardelle et al. (2013) used an average C-
140 band penetration of 1.7 m for East Nyainqêntanglha Shan estimated from the difference of
141 SRTM C-band and X-band DEMs (Gardelle et al., 2012). Here, we extrapolate our ICESat
142 elevation trends over 2003-2008 and their uncertainty back in time to the SRTM acquisition
143 period in February 2000. Under the coarse assumption that the 2000-2003 trends equal the
144 2003-2008 ones, the extrapolation should at February 2000 result in a zero elevation
145 difference to ICESat since the SRTM DEM was used as elevation reference. Offsets in this
146 elevation difference for February 2000 are mainly attributed to SRTM radar penetration into
147 ice and snow (for method and discussion see Kääb et al., 2012). For East Nyainqêntanglha
148 Shan this analysis indicates an average penetration of 8-10 m (7-9 m if based on the winter
149 trends that might alternatively be assumed to reflect February conditions), much more than
150 the 1.7 m assumed in Gardelle et al. (2013), while the corresponding off-glacier penetration is
151 not discernible from zero. Clearly, our penetration depth lies at the high end, but remains

152 within the range of possible C-band phase-centre penetrations (Kääb et al., 2012, Dall et al.,
153 2001, Rignot et al. 2001). Sakai et al. (2014) suggest the highest accumulation rates of the
154 entire study region occur in East Nyainqêntanglha Shan, together with Hindu Kush.
155 Correction of the Gardelle et al. (2013) results by our present C-band penetration estimate
156 completely reconciles their results with ours. Note, however, that extrapolation of our 2003-
157 2008 elevation difference trend back to 2000 is based on the risky assumption that the 2000-
158 2003 trend equals the 2003-2008 trend.

159 For the Bhutan zone, Gardelle et al. (2013) estimated a C-band penetration for February 2000
160 of 2.4 m whereas our extrapolation of ICESat trends suggests around 6 m, which again
161 reconciles the results of both studies for this zone.

162 In the Pamir, our results are more negative than Gardner et al. (2013) and in particular
163 Gardelle et al. (2013). As above, we suggest that our manual classification of ICESat
164 footprints versus the Randolph Glacier Inventory contributed to the difference with Gardner et
165 al. (Gardelle et al. used their own inventory). Also, the difference between our study and
166 Gardner et al. is reduced if only the results from their Method B (similar to ours) is
167 considered. Gardelle et al. (2013) find glacier thickness changes of $+0.16 \pm 0.15 \text{ m yr}^{-1}$ over
168 the Pamir whereas the present study suggests $-0.48 \pm 0.08 \text{ m yr}^{-1}$. Again, we find larger
169 SRTM C-band penetration of 5-6 m compared to 1.8 m (Gardelle et al., 2013). Applying the
170 average C-band penetration from the present study again reconciles the results of both studies.
171 However, comparison of both studies in Pamir is complicated by a number of glacier surges
172 (Gardelle et al., 2013) in connection with particularly sparse ICESat glacier coverage.
173 Superimposing ICESat tracks over Landsat images and the elevation change map of Gardelle
174 et al. (2013) reveals that they cross areas of either strongly positive or negative elevation
175 change zones from surge waves. The ICESat trends thus become biased depending on where
176 they sample surges, and the total ICESat sample size over Pamir is not large enough to
177 compensate for these effects. The different observation periods for both studies (2000-2011
178 versus 2003-2008) may also have considerable impact due to surge activities and climate
179 inter-annual variability (Yi and Sun, 2014).

180 **3 Glacier mass changes and water resources**

181 We assume an average density of 850 kg m^{-3} for all 2003-2008 volume changes to convert the
182 thickness changes to water equivalent quantities (Huss, 2013; see Kääb et al. for different
183 density scenarios). The total glacier area is estimated using a simple cross-product: we

184 multiply the number of ICESat glacier footprints in each zone with the ratio between the total
185 zone area and total number of ICESat footprints. Our method to estimate the total glacier
186 areas is certainly open to discussion, but we prefer the above procedure over using areas from
187 the Randolph Glacier Inventory because of the large deviations to our estimates, mainly for
188 East Nyainqêntanglha and Pamir, from the obviously outdated glacier outlines and voids in
189 the Randolph inventory (Nuimura et al., 2014). The uncertainty of water equivalent quantities
190 includes the standard error of the elevation difference trend fit, the off-glacier trends, an error
191 due to temporal offset of the ICESat autumn campaigns from maximum cumulative ablation
192 conditions, an uncertainty of $\pm 20\%$ for the glacier cover areas, and an uncertainty of $\pm 60 \text{ kg}$
193 m^{-3} for density (Kääb et al. 2012; Huss, 2013). The effects of these individual sources of
194 uncertainty, all converted to error in mass change, are combined through the root sum of
195 squares to arrive at the total uncertainty. Note that water equivalent results from this study are
196 not identical to Kääb et al. (2012), even if elevation difference trends agree, due to the
197 simplified density assumption and the different glacier area estimates used.

198 **3.1 Comparison to gravimetric mass loss**

199 For the Pamir, Kunlun Shan and Karakoram (zone 8b of Jacob et al., 2012; note that the
200 Karakoram is part of their zone 8b, not 8c as suggested by their zone names) we estimate a
201 glacier mass change of $-6 \pm 2 \text{ Gt yr}^{-1}$ for 2003-2008 that agrees well within the error with Jacob
202 et al. results from satellite gravimetry of $-5 \pm 10 \text{ Gt yr}^{-1}$ (Jan 03-Dec 07) and $-8 \pm 9 \text{ Gt yr}^{-1}$ (Jan
203 04-Dec 08). For the Himalayas and East Nyainqêntanglha Shan (zone 8c of Jacob et al. 2012)
204 we estimate a 2003-2008 glacier mass change of $-19 \pm 3 \text{ Gt yr}^{-1}$ that compares to $-3 \pm 12 \text{ Gt yr}^{-1}$
205 (Jan 03-Dec 07) and $-2 \pm 10 \text{ Gt yr}^{-1}$ (Jan 04-Dec 08) from satellite gravimetry. Given their
206 fundamentally different approaches, it is challenging to discuss potential sources of
207 disagreement between the two studies in the Himalayas and East Nyainqêntanglha.
208 Groundwater depletion (Rodell et al., 2009), glacier imbalance runoff into endorheic basins
209 (Zhang et al., 2013), and errors and biases in the ICESat-derived trends as discussed above
210 and in Kääb et al. (2012) are all likely explanations. Note that Gardner et al., 2013 offer a
211 second, more negative gravimetric estimate for the entire combined High Mountain Asia that
212 is, though, not spatially resolved enough to compare to our results. The uncertainties of our
213 results in this entire paragraph are given at 2σ confidence level to better agree with the
214 uncertainty level in Jacob et al. (2012), whereas elsewhere in this contribution uncertainty is
215 provided at 1σ confidence level.

3.2 River runoff

The glaciers of the Tarim Basin (only 40% of total glacier area is covered here, with notably Tien Shan missing) and the Amu Darya basin (all glacier areas covered) drain into endorheic basins and thus their mass changes do not contribute to sea-level (Tab. 2). The glacier mass changes in the Indus, Ganges and Brahmaputra basins from the present study contributed together $\sim 0.06 \pm 0.01 \text{ mm yr}^{-1}$ to eustatic sea-level rise, that is $\sim 10\%$ of the current sea level contribution of $0.71 \pm 0.08 \text{ mm yr}^{-1}$ from glaciers outside the ice sheets (Gardner et al., 2013).

The discharge equivalent of these mass changes, that is the annual average glacier imbalance contribution to river runoff, is given in Tab. 2 for the major river basins covered. Note that computation of our discharge equivalents is a pure unit conversion from Gt yr^{-1} to $\text{m}^3 \text{ s}^{-1}$, neglecting any hydrological processes and with the sole aim to roughly evaluate the relative importance of glacier mass changes for river flow in the catchments.

The Tarim Basin glaciers most likely stored water over 2003-2008 ($+24 \pm 33 \text{ m}^3 \text{ s}^{-1}$ discharge equivalent, DE). The glacier imbalance contribution to runoff is largest for Brahmaputra ($-400 \pm 60 \text{ m}^3 \text{ s}^{-1}$ DE), followed by the Indus ($-220 \text{ m}^3 \text{ s}^{-1}$ DE), and Ganges and Amu Darya ($-130 \text{ m}^3 \text{ s}^{-1}$ DE). Comparison of the discharge equivalent of glacier imbalance to measured river runoff is biased the further downstream the gauging stations are situated from the glaciers due to cumulative natural and man-made losses. It is important to note that the available runoff data from literature and databases refer to various time periods, in parts considerably older than the ICESat period. Figure 2 illustrates thus only roughly the hydrological significance of the 2003-2008 glacier mass change in selected gauged catchments. (For details on the gauging stations used and the uncertainty of the contributions see Supplement). As an example, the 2003-2008 glacier imbalance within the Upper Indus basin at Besham Qila contributes $\sim 6\%$ to annual average river discharge (Fig. 2; Supplement), and we roughly estimate a very similar number for the Amu Darya (Supplement). For the Upper Indus basin, the hydrological balance is under ongoing discussion (cf. Reggiani and Rientjes, 2014) and we hope that our glacier mass change estimates can contribute towards balance closure and better understanding of spatial-temporal patterns of run-off or high-elevation precipitation amounts in the region (e.g. Immerzeel et al., 2012).

The modelling results for “non-renewable glacier runoff” of Savoskul and Smakhtin (2013) agree well with ours for Amu Darya, less for Indus (they obtain $-0.55 \text{ m w.e. yr}^{-1}$ specific mass loss rate over 2001-2010, we $-0.28 \text{ m w.e. yr}^{-1}$) and Ganges (they obtain $-0.77 \text{ m w.e. yr}^{-1}$)

248 ¹, we $-0.37 \text{ m w.e. yr}^{-1}$), and not very well for Brahmaputra (they obtain $-0.36 \text{ m w.e. yr}^{-1}$, we
249 $-0.90 \text{ m w.e. yr}^{-1}$).

250 **4 Conclusions**

251 From 2003-2008 ICESat-derived elevation difference trends over Pamir-Karakoram-
252 Himalaya and from comparison to geographically overlapping studies we draw the following
253 conclusions:

- 254 • Glacier thickness loss over the study region is most pronounced for the East
255 Nyainqêntanglha Shan, followed by Jammu-Kashmir. Glaciers in and around the West
256 Kunlun Shan are in balance or even gaining volume, and Pamir and Karakoram seem to
257 be on the western limit of this mass balance anomaly rather than its centre. This suggests
258 it is a meteorological or climatic anomaly (rise in precipitation). But the cause and
259 duration of this regional glacier anomaly is not fully understood yet.
- 260 • Our glacier volume changes seem especially uncertain in Pamir and, to a lesser extent
261 Hindu Kush. The heterogeneous behaviour of individual glaciers in these two zones, for
262 instance from glacier surges, may lead to biases when extrapolating elevation difference
263 trends from particularly sparse ICESat tracks, or areas covered by differential DEMs, to
264 the entire zones.
- 265 • Extrapolation of ICESat trends back in time to the SRTM acquisition date suggests a
266 much larger potential magnitude and variability of SRTM C-band phase-centre
267 penetration than often assumed. Given the crucial importance of radar penetration for
268 glacier thickness change studies based on radar DEMs, such as the SRTM or the
269 upcoming TanDEM-X, we recommend to be critical against penetration assumptions
270 used in previous studies and to investigate the issue more extensively and systematically
271 (Langley et al., 2007; chapter 7 in Müller, 2011). The problem is complicated by the fact
272 that radar penetration has to be known specifically for certain dates from the past..
- 273 • The glacier mass changes in the Tarim and Amu Darya Basins of $+0.7 \pm 1.0 \text{ Gt yr}^{-1}$ and $-$
274 $4.0 \pm 0.8 \text{ Gt yr}^{-1}$ do not contribute to sea level rise. The combined Ganges, Indus and
275 Brahmaputra basin glacier mass change is $-23.7 \pm 2.1 \text{ Gt yr}^{-1}$, almost 10% of the glacier
276 contribution to sea-level rise during 2003-2009.
- 277 • Neglecting water losses downstream of the glaciers, the 2003-2008 glacier imbalances
278 amount to $\sim 6\%$ of the annual discharge of Amu Darya and Upper Indus where they leave

279 the mountains. This is a considerable amount given the significance of the rivers for the
280 Aral Sea (Amu Darya), and massive irrigation schemes and household use in these dry
281 climate regions. Maximum glacier imbalance contributions to annual average river runoff
282 of up to ~17% are found for the Shyok (Indus) and ~10% Vaksh (Amu Darya), minimum
283 contributions are only ~1-3% for the monsoon-type catchments in Nepal.

- 284 • Our results on glacier mass loss agree with those from satellite gravimetry (Jacob et al.
285 2012) over Pamir, West Kunlun Shan and Karakoram, but significantly diverge over the
286 Himalaya and East Nyainqêntanglha Shan.

287 It is important to note that our results only cover 5 yr, 2003-2008, and it remains open to what
288 extent those years are representative for longer periods, such as the 10 yr covered by Gardelle
289 et al. (2013). For short mass balance series, single anomalous years may have large impacts
290 on trends. Our water equivalent results are also sensitive to density and glacier area
291 assumptions. We find that glacier outlines and areas in the study region are still quite
292 uncertain and invite the reader to use improved glacier area estimates for upscaling our
293 results, and their own assumptions for the conversion of volume changes to mass changes.

294 **Acknowledgement**

295 The study was funded by the European Research Council under the European Union's
296 Seventh Framework Programme (FP/2007-2013) / ERC grant agreement no. 320816, the ESA
297 project Glaciers_cci (4000109873/14/I-NB), and the Department of Geosciences, University
298 of Oslo. E. Berthier acknowledges support from TOSCA (CNES). We are very grateful to
299 NASA and NSIDC for free provision for the ICESat data, and the USGS for the SRTM DEM
300 and Landsat imagery.


301 **Author contributions**

302 A.K. designed the study, performed the data analysis and wrote the paper. C.N., D.T. and E.B.
303 contributed to data analysis, performed supporting analyses and edited the paper.

305

306

Figures and Tables



307

308

309

310

311

312

313


314

315

316

317

Fig. 1 Study region and trends of elevation differences during 2003–08. Data are shown on a 1° grid with overlapping rectangular geographic averaging cells of 2° × 2°. Trends are based on autumn ICESat acquisitions. Only ICESat footprints over glaciers are indicated. The zones indicated by black outlines are equivalent to the ones of Gardelle et al. (2013) with the W Kunlun Shan-Tarim zone (dashed outline) being the only additional one. Trends for all cells (coloured data circles) are statistically significant except for the cells that are marked with grey centres. The uncertainty of the temporal trends per cell is indicated through circle sizes indirectly proportional to the standard error of trends at 68% level.



319

320

321

322

323

324

325

326

Fig. 2 The percentage of discharge equivalent from annual glacier imbalance to measured average river runoff for selected catchments. Note that the actual numbers will be somewhat lower due to unaccounted water losses such as from evaporation or to groundwater. For details on the gauging stations used and the uncertainty of the contributions see Supplement.

327

328 **Table 1:** Glacier elevation difference trends over the Pamir-Karakoram-Himalaya from
 329 this and other studies. Note that Gardelle et al. (2013) cover the period 2000 to ~2010, while
 330 the other studies cover 2003 to 2008/9. Note also that the zones of this study and Gardelle et
 331 al. coincide, whereas the zones of the other do so only roughly, which can potentially explain
 332 parts of the disagreements. See text in sections 3 and 4 for an explanation of how the glacier
 333 areas were estimated. * named Hengduan Shan in Gardelle et al.; ** two zones of Gardner et
 334 al. overlap with our zone and both their values are given.

335

Zone	Glacier area (km ²)	This study (m yr ⁻¹ , ± at 1σ-level)	Gardner et al. (m yr ⁻¹ , ± at 2σ-level)	Neckel et al. (m yr ⁻¹ , ± at 1σ-level)	Gardelle et al. (m yr ⁻¹ , ± at 1σ-level)
East Nyainqêntanglha *	6000	-1.34 ± 0.29	-0.30 ± 0.13 -0.40 ± 0.41 **	-0.81 ± 0.32	-0.39 ± 0.16
Bhutan	3500	-0.89 ± 0.16	-0.89 ± 0.18	-0.78 ± 0.27	-0.26 ± 0.15
Everest	8500	-0.37 ± 0.10	-0.44 ± 0.20		-0.30 ± 0.16
West Nepal	7500	-0.43 ± 0.09		-0.44 ± 0.26	-0.38 ± 0.16
Spiti Lahaul	9500	-0.49 ± 0.12	-0.53 ± 0.13		-0.53 ± 0.16
Karakoram	21000	-0.10 ± 0.06	-0.12 ± 0.15		+0.12 ± 0.19
Hindu Kush	5500	-0.49 ± 0.10			-0.14 ± 0.19
Pamir	6500	-0.48 ± 0.14	-0.13 ± 0.22		+0.16 ± 0.15
West Kunlun Shan - Tarim	12500	+0.05 ± 0.07	+0.17 ± 0.15	+0.04 ± 0.29	

336

337

338

339

340

341

342

343

Table 2: Glacier thickness and mass changes over the major river basins of the study area. The discharge equivalent is a unit conversion from mass change and neglects any losses such as by evaporation or to groundwater. (i) The Tarim Basin is endorheic. Only parts of the glacier area (~40%) within the Tarim Basin are covered in this study. (ii) Endorheic basin.

Major river basin	Glacier area (km ²)	Elevation difference trend (m yr ⁻¹)	Mass change (Gt yr ⁻¹)	Discharge equivalent DE (m ³ s ⁻¹)
Tarim ⁽ⁱ⁾	15000	+0.06 ± 0.08	+0.7 ± 1.0	+24 ± 33
Amu Darya ⁽ⁱⁱ⁾	11000	-0.43 ± 0.08	-4.0 ± 0.8	-128 ± 25
Indus	25000	-0.33 ± 0.04	-7.0 ± 0.8	-220 ± 26
Ganges	11000	-0.44 ± 0.07	-4.1 ± 0.6	-130 ± 20
Brahmaputra	14000	-1.06 ± 0.15	-12.6 ± 1.9	-400 ± 60

344

345

References

- Becker, A., Finger, P., Meyer-Christoffer, A., Rudolf, B., Schamm, K., Schneider, U., and Ziese, M.: A description of the global land-surface precipitation data products of the Global Precipitation Climatology Centre with sample applications including centennial (trend) analysis from 1901–present, *Earth Syst. Sci. Data*, 5, 71–99, 10.5194/essd-5-71-2013, 2013.
- Bolch, T., Kulkarni, A., Kääb, A., Huggel, C., Paul, F., Cogley, J.G., Frey, H., Kargel, J.S., Fujita, K., Scheel, M., Bajracharya, S. and Stoffel, M. The state and fate of Himalayan glaciers. *Science*. 336, 310–314, 2012.
- Cogley, J. G.: Present and future states of Himalaya and Karakoram glaciers, *Ann. Glaciol.*, 52, 69–73, 2011.
- Dall, J., Madsen, S. N., Keller, K. and Forsberg, R.: Topography and penetration of the Greenland Ice Sheet measured with airborne SAR interferometry, *Geophysical Research Letters*, 28(9), 1703–1706, 2001.
- Fujita, K.: Effect of precipitation seasonality on climatic sensitivity of glacier mass balance, *Earth Planet. Sc. Lett.*, 276, 14–19, 10.1016/j.epsl.2008.08.028, 2008.
- Gardelle, J., Berthier, E. and Arnaud, Y. Slight mass gain of Karakoram glaciers in the early 21st century. *Nature Geoscience*. 5, 322–325, 2012.
- Gardelle, J., Berthier, E. and Arnaud, Y.: Impact of resolution and radar penetration on glacier elevation changes computed from multi-temporal DEMs, *Journal of Glaciology*, 58(208), 419–422, 2012.
- Gardelle, J., Berthier, E., Arnaud, Y. and Kääb, A. Region-wide glacier mass balances over the Pamir-Karakoram-Himalaya during 1999–2011. *The Cryosphere*. 7, 1263–1286, 2013
- Gardner, A.S., Moholdt, G., Cogley, J.G., Wouters, B., Arendt, A.A., Wahr, J., Berthier, E., Hock, R., Pfeffer, W.T., Kaser, G., Ligtenberg, S.R.M., Bolch, T., Sharp, M.J., Hagen, J.O., van den Broeke, M.R. and Paul, F. A reconciled estimate of glacier contributions to sea level rise: 2003 to 2009. *Science*. 340(6134), 852–857, 2013.
- Hewitt, K. Glacier change, concentration, and elevation effects in the Karakoram Himalaya, Upper Indus Basin. *Mountain Research and Development*. 31(3), 188–200, 2011.
- Huss, M. Density assumptions for converting geodetic glacier volume change to mass change. *The Cryosphere*. 7(3), 877–887, 2011.
- Immerzeel, W.W., Pellicciotti, F. and Shrestha, A.B. Glaciers as a proxy to quantify the spatial distribution of precipitation in the Hunza basin. *Mountain Research and Development*. 32(1), 30–38, 2012.
- Jacob, T., Wahr, J., Pfeffer, W.T. and Swenson, S. Recent contributions of glaciers and ice caps to sea level rise. *Nature*. 482, 514–518, 2012.
- Kääb, A., Berthier, E., Nuth, C., Gardelle, J. and Arnaud, Y. Contrasting patterns of early twenty-first-century glacier mass change in the Himalayas. *Nature*. 488(7412), 495–498, 2012.
- Kapnick, S. B., Delworth, T. L., Ashfaq, M., Malyshev, S., and Milly, P. C. D.: Snowfall less sensitive to warming in Karakoram than in Himalayas due to a unique seasonal cycle, 7, 834–840, 10.1038/ngeo2269, 2014

- 391 Langley, K., Hamran, S. E., Hogda, K. A., Storvold, R., Brandt, O., Hagen, J. O., and Kohler,
 392 J.: Use of C-band ground penetrating radar to determine backscatter sources within
 393 glaciers, *Ieee T Geosci Remote*, 45, 1236-1246, Doi 10.1109/Tgrs.2007.892600, 2007.
- 394 Loibl, D., Lehmkuhl, F., and Griessinger, J.: Reconstructing glacier retreat since the Little Ice
 395 Age in SE Tibet by glacier mapping and equilibrium line altitude calculation,
 396 *Geomorphology*, 214, 22-39, DOI 10.1016/j.geomorph.2014.03.018, 2014.
- 397 Maussion, F., Scherer, D., Mölg, T., Collier, E., Curio, J., and Finkelnburg, R.: Precipitation
 398 seasonality and variability over the Tibetan Plateau as resolved by the High Asia
 399 Reanalysis, *J Climate*, 27, 1910-1927, Doi 10.1175/Jcli-D-13-00282.1, 2014.
- 400 Müller, K.: Microwave penetration in polar snow and ice: implications for GPR and SAR,
 401 Dissertation, Faculty of Mathematics and Natural Sciences. University of Oslo,
 402 available from <http://www.duo.uio.no/>, 2011.
- 403 Neckel, N., Kropacek, J., Bolch, T. and Hochschild, V. Glacier mass changes on the Tibetan
 404 Plateau 2003-2009 derived from ICESat laser altimetry measurements. *Environmental*
 405 *Research Letters*. 9(1), 2014
- 406 Nuimura, T., Sakai, A., Taniguchi, K., Nagai, H., Lamsal, D., Tsutaki, S., Kozawa, A.,
 407 Hoshina, Y., Takenaka, S., Omiya, S., Tsunematsu, K., Tshering, P., and Fujita, K.:
 408 The GAMDAM Glacier Inventory: a quality controlled inventory of Asian glaciers,
 409 *The Cryosphere*, 8, 2799-2829, 10.5194/tcd-8-2799-2014, 2014.
- 410 Pelto, M. S.: Forecasting temperate alpine glacier survival from accumulation zone
 411 observations, *The Cryosphere*, 4, 67-75, 2010.
- 412 Pfeffer, W. T., Arendt, A. A., Bliss, A., Bolch, T., Cogley, J. G., Gardner, A. S., Hagen, J. O.,
 413 Hock, R., Kaser, G., Kienholz, C., Miles, E. S., Moholdt, G., Molg, N., Paul, F.,
 414 Radic, V., Rastner, P., Raup, B. H., Rich, J., Sharp, M. J., Andeassen, L. M.,
 415 Bajracharya, S., Barrand, N. E., Beedle, M. J., Berthier, E., Bhambri, R., Brown, I.,
 416 Burgess, D. O., Burgess, E. W., Cawkwell, F., Chinn, T., Copland, L., Cullen, N. J.,
 417 Davies, B., De Angelis, H., Fountain, A. G., Frey, H., Giffen, B. A., Glasser, N. F.,
 418 Gurney, S. D., Hagg, W., Hall, D. K., Haritashya, U. K., Hartmann, G., Herreid, S.,
 419 Howat, I., Jiskoot, H., Khromova, T. E., Klein, A., Kohler, J., Konig, M., Kriegel, D.,
 420 Kutuzov, S., Lavrentiev, I., Le Bris, R., Li, X., Manley, W. F., Mayer, C., Menounos,
 421 B., Mercer, A., Mool, P., Negrete, A., Nosenko, G., Nuth, C., Osmonov, A.,
 422 Pettersson, R., Racoviteanu, A., Ranzi, R., Sarikaya, M. A., Schneider, C., Sigurdsson,
 423 O., Sirguey, P., Stokes, C. R., Wheate, R., Wolken, G. J., Wu, L. Z., Wyatt, F. R., and
 424 Consortium, R.: The Randolph Glacier Inventory: a globally complete inventory of
 425 glaciers, *J Glaciol*, 60, 537-552, Doi 10.3189/2014jog13j176, 2014.
- 426 Reggiani P. and T. H. M. Rientjes (2014). A reflection on the long-term water balance of the
 427 Upper Indus Basin. *Hydrology Research*. In press.
- 428 Rignot, E., Echelmeyer, K. and Krabill, W. Penetration depth of interferometric synthetic-
 429 aperture radar signals in snow and ice. *Geophysical Research Letters*. 28(18), 3501-
 430 3504, 2001.
- 431 Rodell, M., Velicogna, I. and Famiglietti, J.S. Satellite-based estimates of groundwater
 432 depletion in India. *Nature*. 460(7258), 999-1002, 2009.
- 433 Sakai, A., Nuimura, T., Fujita, K., Takenaka, S., Nagai, H., and Lamsal, D.: Climate regime
 434 of Asian glaciers revealed by GAMDAM Glacier Inventory, *The Cryosphere*, 8, 3629-
 435 3663, 10.5194/tcd-8-3629-2014, 2014.
- 436 Sarikaya, M.A., Bishop, M.P., Shroder, J.F. and Olsenholler, J.A. Space-based observations
 437 of Eastern Hindu Kush glaciers between 1976 and 2007, Afghanistan and Pakistan.
 438 *Remote Sensing Letters*. 3(1), 77-84, 2012.
- 439 Savoskul, O. S., and Smakhtin, V.: Glacier systems and seasonal snow cover in six major
 440 Asian river basins: hydrological role under changing climate, *International Water*
 441 *Management Institute (IWMI), Colombo, Sri Lanka*. 150, 2013.

- 442 Yao, T.D., Thompson, L., Yang, W., Yu, W.S., Gao, Y., Guo, X.J., Yang, X.X., Duan, K.Q.,
443 Zhao, H.B., Xu, B.Q., Pu, J.C., Lu, A.X., Xiang, Y., Kattel, D.B. and Joswiak, D.
444 Different glacier status with atmospheric circulations in Tibetan Plateau and
445 surroundings. *Nature Climate Change*. 2(9), 663-667, 2012.
- 446 Yi, S. and Sun, W.: Evaluation of glacier changes in high-mountain Asia based on 10 year
447 GRACE RL05 models, *Journal of Geophysical Research: Solid Earth*, 119(3),
448 2013JB010860, 2013.
- 449 Tao, H., Borth, H., Fraedrich, K., Su, B., & Zhu, X. Drought and wetness variability in the
450 Tarim River Basin and connection to large-scale atmospheric circulation. *International*
451 *Journal of Climatology*, 34: 2678 – 2684, 2014.
- 452 Zhang, G.Q., Yao, T.D., Xie, H.J., Kang, S.C. and Lei, Y.B. Increased mass over the Tibetan
453 Plateau: From lakes or glaciers? *Geophysical Research Letters*. 40(10), 2125-2130,
454 2013.
- 455

456

457