

Interactive comment on “Representativeness and seasonality of major ion records derived from NEEM firn cores” by G. Gfeller et al.

Anonymous Referee #3

Received and published: 18 July 2014

Gfeller et al. have assessed the spatial “representativeness” of several major ions commonly found in snow using a suite of snowpits and firn/ice cores from northwestern Greenland. Using complex statistical analysis, the authors identify which species are most likely to reflect the original atmospheric loading/source strength signal on seasonal to interannual timescales when measured in a single record, and which species would require several spatially distributed cores in order to overcome localized post-depositional alteration, most likely wind scouring in this case. The conclusions are specific to the region of study, though it is suggested that sites with high snow accumulation and calm winds are more likely to preserve representative signals and may not require extensive replicate coring.

This work represents a critical, though all too often overlooked, step in interpreting ma-

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



jor ion chemistry in ice cores. The analytical methods appear sound and there is careful consideration given to error and the validity of the statistical techniques employed. Importantly, authors consider two possible accumulation scenarios. The manuscript is appropriate for The Cryosphere and written in a clear and concise manner. I recommend publication largely as is and offer several minor comments and suggestions below.

Comments:

It is very interesting that nitrate is one of the most representative ions. As you note, nitrate is vulnerable to post-depositional loss, especially at low accumulation sites, though it may largely be preserved under higher accumulation regimes such as at Summit. Still, it is interesting that conservative species such as calcium and sodium are less representative, at least on the interannual scale. Do you feel comfortable speculating about what accounts for this difference? Calcium and sodium are largely deposited in the colder months, i.e., when wind speed and reworking is greater (you note that the AWS observations are scarce, but you could refer to Section 3.5 in Steffen and Box (2001), Surface climatology of the Greenland ice sheet: Greenland Climate Network 1995-1999, JGR, D24. Conductivity is more representative but also peaks in the colder months, though it may be different since it is a more integrated signal.

2531(10) and 2552(14): I don't believe decades contain apostrophes (e.g., 1970s) since they are not possessive.

Be consistent in your choice of time scale vs. time-scale, inter-annual vs. inter annual, and scenario 2 vs. scenario two (e.g., p2531 and p2545).

2549(8): worthwhile?

2551(22): Can you clarify "completely obliterated" in this sentence? I read this as the pre-Industrial summer nitrate peak being obscured by the higher, present-day concentrations and distinct seasonality. Is that correct?

Figure 7: The colored (dice 5) and dark colored lines (F2/F3) are very difficult to distinguish. Also, what are the colored stepped lines to the right? Are these the snowpit data? A legend would be useful in addition to the caption descriptions, if you can find room for it.

Figure 8: The step line plots for the modern seasonality are difficult to see against the similarly-colored variability. Can you darken these or otherwise help them stand out?

Interactive comment on The Cryosphere Discuss., 8, 2529, 2014.

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

[Discussion Paper](#)

