

Editor comments on “Precipitation measurement intercomparison in the Qilian Mountains, Northeastern Tibetan Plateau” by R. Chen et al.

General comments from the Editor (October 8, 2015)

Minor revision

DETAILED COMMENTS

The detailed comments are derived from the editor's marked PDF document by authors.

1. Page 1 Line 1:

Editor's comments: the old title is better, use that one!

Authors' response: The new title is replaced by the old title.

Author's changes in manuscript: Precipitation measurement intercomparison in the Qilian Mountains, Northeastern Tibetan Plateau

2. Page 1 Line 26:

Editor's comments: delete" independently of the local environmental conditions."

Authors' response: It has been deleted.

Author's changes in manuscript: The "independently of the local environmental conditions." in the abstract and text has been deleted

3. Page 2 Line 4:

Editor's comments: "simulate" should be revised as "define or quantify"

Authors' response: It has been replaced by "quantify" in two places in the text.

Author's changes in manuscript: used to ~~quantify~~simulate the

4. Page 3 Line 13:

Editor's comments: "huge volume" should be revised as "large volume"

Authors' response: It has been replaced by "large volume" in two places in the text.

Author's changes in manuscript: on the ~~huge-large~~ volume of

5. Page 4 Line 28~Page 5 Line 4:

Editor's comments: The marked sentence is not described clearly. "ratios" should be deleted. "shield" should be "gauge".

Authors' response: The paragraph has been revised.

Author's changes in manuscript: Therefore, an unshielded CSPG, a single Alter shield CSPG (SA), a DFIR with a Tretyakov-shielded CSPG and a CSPG in a pit were selected as the field experiment of wind-induced bias study. This paper presents the intercomparison experiments and their relevant data, introduces the adjustment methods, discusses wind-induced bias in precipitation measurements by CSPGs for different precipitation phases, analyses the correlations between shielded and unshielded CSPGs and quantifies the relationships between catch ratio and wind speed. The results of the present study are also compared with other studies. In addition, the pit gauge is evaluated for solid precipitation under these climatic conditions. The limitations of the present study are then discussed.

6. Page 5 Line 12:

Editor's comments: "criteria" should be revised as "standard"

Authors' response: Total 4 "criteria" are replaced by "standard" in the whole text.

Author's changes in manuscript: CMA's ~~eriteria-standard~~

7. Page 8 Line 27:

Editor's comments: 3.2 Comparisons of wind-induced bias: use one gauge as the reference.

Authors' response: In the Table 3, the CSPG_{PIT} was used as reference to calculate catch ratios before October 2012 when the CSPG_{DFIR} was not installed. It is not appropriate.

Author's changes in manuscript: The relevant catch ratio in the Table 3 are deleted.

8. Page 8 Line 28:

Editor's comments: Title "3.2.1 Rainfall", "3.2.2 mixed precipitation" and "3.2.3 snowfall" should be deleted

Authors' response: After delete the subtitle, the text is more appropriate.

Author's changes in manuscript: The three subtitles are deleted and the relevant paragraphs are combined.

9. Page 9 Line 24:

Editor's comments: What is your standard for a ref? It should define the limits.

Authors' response: Several words are added in the end of the sentence.

Author's changes in manuscript: at the ~~experimental~~ site with shallow snow cover and rare blowing snow event.

10. Page 12 Line 11:

Editor's comments: "no" should be revised as "not"

Authors' response:

Author's changes in manuscript: are ~~no~~ not

11. Page 12 Line 26:

Editor's comments: "reference" should be revised as "study"

Authors' response:

Author's changes in manuscript: Tianshan ~~reference~~ study

12. Page 13 Line 1 and Line 4:

Editor's comments: "largest" should be revised as "highest", "compared to" should be revised as "than"

Authors' response:

Author's changes in manuscript: the ~~largest-highest~~ wind : snowfall ~~compared to~~than the CSPG_{UN}.

13. Page 13 Line 18:

Editor's comments: modeling of wind field? useful too?

Authors' response: Yes. Modeling of wind field is very useful for knowing about wind-induced errors in precipitation measurements. A new reference paper is added:

Yang, D., and Simonenko, A.: Comparison of winter precipitation measurements by six Tretyakov gauges at the Valdai experimental site, Atmosphere-Ocean, 52:1, 39-53, doi: 10.1080/07055900.2013.865156, 2014.⁴

Author's changes in manuscript:

measurements for different mountain watersheds, field experiments need to be carried out continuously. Further investigation is also necessary to consider the influence of micrometeorology on gauge observations, particularly wind distribution and turbulence across this site (Yang and Simonenko, 2014).⁴

14. Page 14 Line 4:

Editor's comments: "snowfalls are" should be revised as "snowfall is "

Authors' response:

Author's changes in manuscript: snowfalls-are is

15. Page 15 Line 7:

Editor's comments: "The present experimental field" should be revised as "This "

Authors' response:

Author's changes in manuscript:~~The present experimental field~~This

16. Page 19 Line 8:

Editor's comments: This reference paper (Ye et al., 2007) is similar to the next one (Ye et al., 2004).

Authors' response: The reference paper(Ye et al., 2007) is deleted.

Author's changes in manuscript:⁴ | Ye, B., Yang, D., Ding, Y., and Han, T.: A bias-corrected precipitation climatology for China. Acta Geogr. Sin., 62, 3-13, 2007.⁴

In addition, the new reference is added:

Yang, D., and Simonenko, A.: Comparison of winter precipitation measurements by six Tretyakov gauges at the Valdai experimental site, Atmosphere-Ocean, 52:1, 39-53, doi: 10.1080/07055900.2013.865156, 2014.⁴

and some sentences in the new reference paper (Yang and Simonenko, 2014) are used in the text:

- 1) From the above mentioned relationships of $CR_{UN/DFIR}$ and $CR_{SA/DFIR}$ vs. wind speed, the following points can be drawn for our understanding. For daily rain and mixed precipitation, the relationships are not statistically significant. The use of daily mean wind speed may lead to uncertainties in gauge comparisons. Data collections and analyses on shorter time scales, such as hourly or 6-hourly, are expected to produce more reliable results, because wind speed may vary throughout the day and daily mean wind speeds may not be representative of the wind conditions over the precipitation period (Yang and Simonenko, 2014). Daily maximum and minimum
- 2) precipitation period, but the $CSPG_{UN}$, $CSPG_{SA}$, $CSPG_{PIT}$ and $CSPG_{DFIR}$ were observed only twice per day. In this field experiment, the precipitation phases were also distinguished by observers. This method is somewhat imprecise although this has remained the traditional method since the 1950s at the CMA stations (CMA, 2007b). Automatic sensors will also be important to detect precipitation types at operational and research networks (Yang and Simonenko, 2014).⁴

1 Precipitation measurement intercomparison in the Qilian Mountains,

2 Northeastern Tibetan Plateau

3 ~~Experimental wind-induced bias in precipitation measurements in a mountain-~~
4 ~~watershed on the north-eastern Tibetan Plateau~~

5 R. Chen^{*}, J. Liu, E. Kang, Y. Yang, C. Han, Z. Liu, Y. Song, W. Qing, P. Zhu

6 Qilian Alpine Ecology and Hydrology Research Station, Key Laboratory of Inland River Ecohydrology, Cold and Arid Regions
7 Environmental and Engineering Research Institute, Chinese Academy of Sciences, Lanzhou 730000, China

8

9 **Abstract:** An experimental field study of wind-induced bias in precipitation measurements was conducted from
10 September 2010 to April 2015 at a grassland site (99°52.9', 38°16.1', 2980 m) in the Hulu watershed in the Qilian
11 Mountains, on the north-eastern Tibetan Plateau, in China. The experiment included (1) an unshielded Chinese
12 standard precipitation gauge (CSPG_{UN}; orifice diameter=20 cm, height=70 cm), (2) a single Alter shield around a
13 CSPG (CSPG_{SA}), (3) a CSPG in a pit (CSPG_{PIT}) and (4) a Double-Fence International Reference (DFIR) ~~shield~~
14 with a Tretyakov-shielded CSPG (CSPG_{DFIR}). The catch ratio (CR) used the CSPG_{DFIR} as a reference
15 ($CR = \text{CSPG}_X / \text{CSPG}_{DFIR}$, %; X denotes UN, SA or PIT). The results show that the CSPG_{SA}, CSPG_{PIT} and
16 CSPG_{DFIR} caught 0.9%, 4.5% and 3.4% more rainfall, 7.7%, 15.6% and 14.2% more mixed precipitation (snow
17 with rain, rain with snow), 11.1%, 16.0% and 20.6% more snowfall, and 2.0%, 6.0% and 5.3% more precipitation
18 (of all types), respectively, than the CSPG_{UN} from September 2012 to April 2015. The CSPG_{PIT} and CSPG_{DFIR}
19 caught 3.6% and 2.5% more rainfall, 7.3% and 6.0% more mixed precipitation, 4.4% and 8.5% more snowfall and
20 3.9% and 3.2% more total precipitation, respectively, than the CSPG_{SA}. However, the CSPG_{DFIR} caught 1.0% less
21 rainfall, 1.2% less mixed precipitation, 3.9% more snowfall and 0.6% less total precipitation than the CSPG_{PIT}.
22 From most to least precipitation measured, the instruments ranked as follows: for rain and mixed precipitation,
23 $\text{CSPG}_{PIT} > \text{CSPG}_{DFIR} > \text{CSPG}_{SA} > \text{CSPG}_{UN}$; for snowfall, $\text{CSPG}_{DFIR} > \text{CSPG}_{PIT} > \text{CSPG}_{SA} > \text{CSPG}_{UN}$. The CR vs.
24 10 m wind speed for the period of precipitation indicated that with increasing wind speed from 0 to 8.0m/s, the
25 $\text{CR}_{UN/DFIR}$ and $\text{CR}_{SA/DFIR}$ for rainfall decreased slightly. For mixed precipitation, the wind speed showed no
26 significant effect on $\text{CR}_{UN/DFIR}$ and $\text{CR}_{SA/DFIR}$ below 3.5m/s. For snowfall, the $\text{CR}_{UN/DFIR}$ and $\text{CR}_{SA/DFIR}$ vs. wind

*Corresponding author. E-mail address: crs2008@lzb.ac.cn (R. Chen)

1 speed showed that CR decreased with increasing wind speed. The precipitation measured by the shielded gauges
2 increased linearly relative to that of the unshielded gauges ~~independently of the local environmental conditions~~.
3 However, the increase in the ratio of the linear correlation should depend on specific environmental conditions. A
4 comparison of the wind-induced bias indicates that the $CSPG_{PIT}$ could be used as a reference gauge for rain,
5 mixed and snow precipitation events at the experimental site. As both the PIT and DFIR effectively prevented
6 wind from influencing the catch of the precipitation gauge, the $CR_{PIT/DFIR}$ had no relationship with wind speed.
7 Cubic polynomials and exponential functions were used to ~~quantify~~~~simulate~~ the relationship between catch ratio
8 and wind speed. For snow, for both event and daily scales, the $CR_{UN/DFIR}$ and $CR_{SA/DFIR}$ were significantly related
9 to wind speed; while for rain and mixed precipitation, only the event scale showed a significant relationship.

10 **Keywords:** Precipitation, Gauge catch ratio, Wind-induced undercatch, Field observation, Qilian Mountains

11

12 **1 Introduction**

13 In western China, mountainous watersheds are the source areas of runoff generation and water resources, and
14 accurate precipitation measurements are extremely important for calculating the water balance and understanding
15 the water cycle processes in these high mountains. It is widely recognised that precipitation gauge measurements
16 contain systematic errors caused mainly by wetting, evaporation loss and wind-induced undercatch, and that
17 snowfall observation errors are very large under high wind (Sugiura et al., 2003). These errors affect the
18 evaluation of available water in a large number of economic and environmental applications (Tian et al., 2007; Ye
19 et al., 2012).

20 For decades, all knowledge of precipitation measurement errors has relied on field experiments. Back in 1955,
21 the World Meteorological Organization (WMO) conducted the first precipitation measurement intercomparisons
22 (Rodda, 1973). The reference standard was a British Meteorological Office gauge of the Snowdon type (Mk2)
23 elevated 1 m above the ground and equipped with the Alter wind shield, which did not accurately reflect the
24 precipitation level (Struzer, 1971). Rodda (1967) compared the catch of a UK 5" manual gauge, exposed
25 normally at the standard height of 30.5 cm above ground, with a Koschmieder-type gauge exposed in a pit. The
26 gauge in the pit caught 6% more precipitation than the normally exposed gauge. In the second WMO precipitation
27 measurement intercomparison (Rain, 1972–1976), a pit with an anti-splash grid was designated the reference
28 standard shield for rain gauges (Sevruk and Hamon, 1984). In the third WMO precipitation measurement
29 intercomparison (Snow, 1986–1993), the Double Fence International Reference (DFIR) ~~shield~~ with a Tretyakov

1 shield was designated the reference standard snow gauge configuration (Goodison et al., 1998). In the fourth
2 WMO precipitation measurement intercomparison (Rain Intensity, 2004–2008), different principles were tested to
3 measure rainfall intensity and define a standardised adjustment procedure (Lanza et al., 2005). Because
4 automation of precipitation measurements was widespread, the WMO Commission for Instruments and Methods
5 of Observation (CIMO) organised the WMO Solid Precipitation Intercomparison Experiment (WMO-SPICE;
6 Wolff et al., 2014) to define and validate automatic field instruments as references for gauge intercomparison, and
7 to assess the automatic systems and operational networks for precipitation observations. The experiments and
8 investigations are ongoing, and the WMO-SPICE project confirms the DFIR shield to be a part of the reference
9 configurations.

10 The DFIR shield has been operated at 25 stations in 13 countries around the world (Golubev, 1985; Sevruk et
11 al., 2009), but deviations from the DFIR measurements vary by gauge type and precipitation type (Goodison et al.,
12 1998). In China, the Chinese standard precipitation gauge (CSPG) and the Hellmann gauge were first compared
13 using the DFIR shield as a reference configuration at the Tianshan site (43°7' N, 86°49' E, 3720 m), during the
14 third WMO precipitation measurement intercomparison experiment from 1985 to 1987 (Yang, 1988; Yang et al.,
15 1991). The wetting loss, evaporation loss, wind-induced undercatch and trace precipitation of the CSPGs were
16 well quantified based on the ~~huge-large~~ volume of observation data at the Tianshan site (Yang et al., 1991). For
17 wind-induced undercatch, the derived CSPG catch ratio equations were based on the 10 m height wind speed at
18 the Daxigou Meteorological Station (43.06°, 86.5°E, 3540 m) and at several other standard meteorological
19 stations near the measurement site (Yang, 1988; Yang et al., 1991). This intensive experimental field study created
20 a basis for later work on the correction of systematic bias in precipitation measurements in China. From 1992 to
21 1998, Ren and Li (2007) conducted an intercomparison experiment at 30 sites (the altitude ranged from about 4.8
22 to 3837 m) using the pit as a reference across China, and a total of 29, 276 precipitation events were observed.
23 Yang et al. (1999) emphasised that among all known systematic errors in precipitation observation, wind-induced
24 gauge undercatch was the greatest source of bias, particularly in cold regions, and recommended testing for the
25 application of adjustment techniques in regional observation networks. In the mountainous watersheds of western
26 China, the complex high mountain topography and underlying surfaces with inhomogeneous glaciers, permafrost
27 and alpine vegetation make the wind vector field in the lower boundary layer extremely complex, causing equally
28 complex wind field deformations over the gauge orifice. At present, our investigation of wind-induced error in
29 precipitation measurements is based on the horizontal time-averaged wind speed. Thus it is reasonable to
30 investigate the regional average characteristics of wind fields and the interaction between wind fields and the

1 precipitation gauges at our present research level. In addition to Yang's experimental field work on systematic
2 error adjustments for precipitation measurements in eastern Tianshan from 1985 to 1987 (Yang, 1988), it is ~~very~~
3 necessary to carry out field experiments on precipitation measurement in the other mountainous regions of
4 western China.

5 Adjustment procedures and reference measurements were developed during several WMO international
6 precipitation measurement intercomparisons (Goodison et al., 1998; Sevruk et al., 2009; Yang, 2014). The
7 application of all of these adjustment procedures and methods depends on both environmental factors and
8 precipitation features, and among the factors considered, wind speed and temperature have been found to have the
9 most important effect on gauge catch (Yang et al., 1999). Ye et al. (2004,~~2007~~) developed a bias-error adjustment
10 method for CSPGs based on observation data from 1985 to 1997 at the Tianshan site (Yang et al., 1991), and
11 found a new precipitation trend in the adjusted precipitation data for the past 50 years in China (Ding et al., 2007).
12 The new precipitation adjustment has improved the precipitation estimation in water balance computation for
13 many basins in China (Ye et al., 2004; Tian et al., 2007; Ye et al., 2012). Ma et al. (2014) used the adjusted
14 equations from neighbouring countries in addition to the experimental results from eastern Tianshan in China
15 (Yang et al., 1991) to correct for wind-induced errors on the Tibetan Plateau. However, the precipitation gauges
16 used in the neighbouring countries were the Tretyakov, MK2, Nepal203, Indian standard and US 8". As the
17 world's third polar region, the Tibetan Plateau and its surrounding mountain ranges are ecologically fragile and
18 the source of several large rivers in China and neighbouring countries, and accurate precipitation data are urgently
19 needed for water resource exploitation and environmental protection. The problem is how to apply and test the
20 already established principal adjustment procedures and methods to correct for precipitation measurement errors
21 in the vast plateau and high mountains of western China, where climatic and environmental conditions are highly
22 complex and variable, both spatially and temporally. To quantify and understand the specific influences of
23 climatic and environmental factors on wind-induced bias in precipitation measurements in a mountain watershed,
24 and then test and parameterise the adjustment equations, an intercomparison experiment was carried out for nearly
25 five years on both unshielded and shielded CSPGs in a watershed in the Qilian Mountains on the north-eastern
26 Tibetan Plateau in China.

27 The CSPG is the standard manual precipitation gauge that has been used by the China Meteorological
28 Administration (CMA) in more than 700 stations since the 1950s. ~~The present experiment is to investigate the~~
29 ~~wind-induced bias of the CSPG in the high mountain environment.~~ The Alter shield (Struzer, 1971) was used by
30 the CMA to enhance catch ~~ratios~~ of automatic gauges (Yang, 2014), and the pit and DFIR ~~was~~ were used to

1 provide true rainfall and snowfall values for the WMO intercomparison project, respectively (Yang et al., 1999).
2 Therefore, an unshielded CSPG, a single Alter shield CSPG (SA), a DFIR with a Tretyakov-shielded CSPG and a
3 CSPG in a pit were selected as the field experiment of wind-induced bias study.~~a single Alter shield (SA) (Struzer,~~
4 ~~1971), a Double Fence International Reference shield with a Tretyakov-shielded (DFIR) and a pit were selected to~~
5 ~~shield the CSPGs, which were distributed by an unshielded CSPG. The SA shield was used by the CMA to~~
6 ~~enhance the catch ratios of automatic gauges (Yang, 2014), and the DFIR was used to provide true snowfall~~
7 ~~values for the WMO intercomparison project (Yang et al., 1999).~~ This paper presents the intercomparison
8 experiments and their relevant data, introduces the adjustment methods, discusses wind-induced bias in
9 precipitation measurements by CSPGs for different precipitation phases, analyses the correlations between
10 shielded and unshielded CSPGs and specifies-quantifies the relationships between catch ratio and wind speed. The
11 results of the present study are also compared with other studies. In addition, the pit shield-gauge is evaluated for
12 solid precipitation under these climatic conditions. The limitations of the present study are then discussed.

13 **2 Experiments and methods**

14 **2.1 Intercomparisons and data**

15 Precipitation intercomparison experiments (Fig.1, Table 1) were conducted at a grassland site (99°52.9',
16 38°16.1', 2980 m) in the Hulu watershed in the Qilian Mountains, on the north-eastern edge of the Tibetan Plateau,
17 in China. A meteorological cryosphere-hydrology observation system (Chen et al., 2014) was established in 2008
18 in the Hulu watershed. The mean annual precipitation was 447.2 mm during 2010-2012 and was concentrated
19 during the warm season from May to September. The annual mean temperature was 1.1 °C, with a July mean
20 (T_{mean}) of 12.5 °C and a January mean of -12.4°C over the years (Table 1). The annual potential evaporation (E_0)
21 was 1102 mm (Table 1).

22 The intercomparative experiments included (1) an unshielded CSPG (CSPG_{UN}; orifice diameter=20 cm,
23 height=70 cm), (2) a single Alter shield around a CSPG (CSPG_{SA}), (3) a CSPG in a pit (CSPG_{PIT}), and (4) a DFIR
24 ~~shield~~-with a Tretyakov-shielded CSPG (CSPG_{DFIR}) (Fig.1, Table 2). The CSPG_{UN}, CSPG_{SA} and CSPG_{PIT} were
25 installed before September 2010, whereas the CSPG_{DFIR} was installed in September 2012 (Table 2). In the cold
26 season (October to April), snowfalls dominated the precipitation events, and in the warm season (May to
27 September), rainfall was dominate. The precipitation was measured manually twice a day at 08:00 and 20:00 local
28 time (Beijing time) according to the CMA's criteria-standard (CMA, 2007a). In the warm season, precipitation
29 was measured by volume. Whereas in the cold season, the funnel and glass bottle were removed from the CSPG

1 and precipitation was weighed under a windproof box. Any frost on the outside surface of the collector was wiped
2 off using a dry hand towel. In rare cases where snow had accumulated on the rim of the collector, this was
3 removed before weighing.

4 The precipitation phases (snow, rain and mixed) were distinguished using the CMA's ~~criteria-standard~~ (CMA,
5 2007b). Meteorological elements, including maximum air temperature T_{max} and minimum T_{min} , have been
6 measured in conformation with the meteorological observation manual at the site since June, 2009. A
7 meteorological tower was used to measure wind speed (Lisa/Rita, SG GmbH; W_s), air temperature (HMP45D,
8 Vaisala) and relative humidity (HMP45D, Vaisala) at 1.5m and 2.5m heights in association with precipitation
9 measurements (Chen et al., 2014). The time step of the observations of the tower was 30 seconds and half-hourly
10 values were obtained. The specific meteorological conditions at the site are summarised in Table 1.

11
12 **Fig.1 about here**

13 **Table 1 and Table 2 about here**

14 **2.2 Adjustment methods**

15 This field experiment focused on two key aspects. One was a comparison of the CSPG_{UN}, CSPG_{SA}, CSPG_{PIT}
16 and CSPG_{DFIR} gauges. The other was the establishment of adjustment equations for the CSPG_{UN} and CSPG_{SA}
17 using the CSPG_{DFIR} as a reference. To adjust gauge-measured precipitation, Sevruk and Hamon (1984) provided
18 the general formula as:

$$19 \quad P_c = KP_g + \Delta P_w + \Delta P_e + \Delta P_t = P_{DFIR} + \Delta P_w + \Delta P_e + \Delta P_t \quad (1)$$

20 where P_c is the adjusted precipitation, K is the wind-induced coefficient, P_g is the gauge-measured precipitation.
21 P_w is the wetting loss, P_e is the evaporation loss, P_t is trace precipitation and P_{DFIR} is the DFIR-shielded
22 precipitation. For loss by the CSPG per observation, P_w is 0.23 mm for rainfall measurements, 0.30 mm for snow
23 and 0.29 mm for mixed precipitation (snow with rain, rain with snow), based on the measurements at the Tianshan
24 site (Yang, 1988; Yang et al., 1991). Ren and Li (2007) reported a mean P_w of about 0.19 mm for the total
25 precipitation over eastern China. The CSPG design reduces P_e to a near-zero value smaller than other losses in the
26 warm, rainy season (Ye et al., 2004; Ren and Li, 2007). In winter, P_e is already small (0.10–0.20 mm/day)
27 according to the results from Finland (Aaltonen et al., 1993) and Mongolia (Zhang et al., 2004). To prevent
28 evaporation loss in Chinese operational observations on particular days, e.g., hot, dry days or days of snow,
29 precipitation is measured as soon as the precipitation event stops (CMA, 2007a; Ren and Li, 2007). A
30 precipitation event of less than 0.10 mm is beyond the resolution of the CSPG and is recorded as trace

1 precipitation (P_t). Ye et al. (2004) recommended assigning a value of 0.1 mm, regardless of the number of trace
 2 observations per day. The present study focused on wind-induced bias in precipitation measurement by CSPGs,
 3 specifically in high mountain environments, therefore the above mentioned P_w , P_e and P_t values were assumed to
 4 be constant in the computation equations.

5 The WMO proposed Eqs.(2)–(4) to compute the catch ratio of unshielded over shielded Tretyakov gauges on a
 6 daily time step for three precipitation types, and the independent variables were wind speed (W_s , ms^{-1}) at the
 7 gauge height and the daily maximum and minimum temperatures (T_{\max} , T_{\min} , $^{\circ}\text{C}$) (Yang et al., 1995; Goodison et
 8 al., 1998). These equations are used over a great range of environmental conditions (Goodison et al., 1998).

$$CR_{\text{snow}} = 103.1 - 8.67W_s + 0.3T_{\max} \quad (2)$$

$$9 \quad CR_{\text{mix}} = 96.99 - 4.46W_s + 0.88T_{\max} + 0.22T_{\min} \quad (3)$$

$$CR_{\text{rain}} = 100.0 - 4.77W_s^{0.56} \quad (4)$$

10

11 where CR_{snow} (%), CR_{mix} (%) and CR_{rain} (%) are the catch ratios for snow, mixed precipitation and rain,
 12 respectively.

13 As the CMA stations usually observe wind speed at a height of 10m, Eqs.(5)–(7) were used for the CSPG catch
 14 ratio versus the daily mean wind speed W_s (ms^{-1}) at 10m (Yang et al., 1991). These equations are based on the
 15 huge-large volume of experimental precipitation gauge intercomparison data at the Tianshan site and the wind
 16 speed data at the Daxigou station:

$$CR_{\text{snow}} = 100 \exp(-0.056W_{s10}) \quad (0 < W_s < 6.2) \quad (5)$$

$$17 \quad CR_{\text{rain}} = 100 \exp(-0.04W_{s10}) \quad (0 < W_s < 7.3) \quad (6)$$

$$CR_{\text{mix}} = CR_{\text{snow}} - (CR_{\text{snow}} - CR_{\text{rain}})(T_{\text{mean}} + 2) / 4 \quad (7)$$

18 where T_{mean} is the daily mean air temperature ($^{\circ}\text{C}$).

19 Referring to Eqs.(2)–(7), two types of equation were used. One is for easy application using the 10m height
 20 wind speed during the period of precipitation in China. These are similar to a revised version of Eqs.(5)–(7). The
 21 other type is similar to Eqs.(2)–(4), which use the daily mean wind speed at gauge height. For the CSPGs, the
 22 gauge height was 70cm (Table 2). The catch ratio uses $\text{CSPG}_{\text{DFIR}}$ as the reference ($\text{CR} = \text{CSPG}_X / \text{CSPG}_{\text{DFIR}}$, %; X
 23 denotes UN, SA or PIT). The equations were fitted using SPSS software version 19.0 (IBM, 2010) and Microsoft
 24 Excel 2007 based on the mathematical least squares method (Charnes et al., 1976). The significance of the
 25 equations was evaluated using the F-test method (Snedecor and Cochran, 1989). If the significance level (α) of the
 26 F-test is below 0.05, the fitted equation is significant. The lower the α value, the greater the significance.

27 Wind speeds at gauge height ($W_{s0.7}$) and at the 10 m height (W_{s10}) were calculated using half-hourly wind speed

1 | data at 1.5 m ($W_{s1.5}$) and 2.5 m ($W_{s2.5}$) according to the Monin-Obukhov theory and the gradient method (Bagnold,
2 | 1941; Dyer and Bradley, 1982):

$$3 \quad W_{sZ} = \frac{\ln Z - \ln Z_0}{\ln 1.5 - \ln Z_0} W_{s1.5} \quad (8)$$

$$4 \quad \ln Z_0 = \frac{W_{s2.5} \ln 1.5 - W_{s1.5} \ln 2.5}{W_{s2.5} - W_{s1.5}} \quad (9)$$

5 | where Z denotes the height referred to.

6 | **3 Results**

7 | From September 2010 to April 2015, a total of 608 precipitation events were recorded at the intercomparison
8 | site for CSPG_{UN}, CSPG_{SA} and CSPG_{PIT}, respectively (Table 3). Snow occurred 84 times, mixed precipitation 44
9 | times and rain 480 times during this period. From September 2012 to April 2015, a subset of 283 precipitation
10 | events was recorded for the CSPG_{UN}, CSPG_{SA}, CSPG_{PIT}, and CSPG_{DFIR} gauges, respectively (Table 3). During
11 | this period, snow occurred 43 times, mixed precipitation 29 times and rainfall 211 times.

13 | **Table 3 about here**

15 | **3.1 Linear correlation of gauge precipitation**

16 | At the 14 WMO intercomparison sites, a strong linear relationship was found between Alter-shielded and
17 | unshielded Belfort gauges, Alter-shielded and unshielded NWS 8-inch gauges, and shielded and unshielded
18 | Tretyakov gauges for all types of precipitation, with a higher correlation for rain than for snow (Yang et al., 1999).
19 | In the present study in the Qilian Mountains, which experiences different environmental conditions compared to
20 | the other 14 sites, the same strong linear correlation was found among the four CSPG instalments for rainfall,
21 | mixed precipitation and snowfall, with a higher correlation for rain than for mixed precipitation, successively
22 | more than for snow (Figures 2–4). It is therefore considered that in general the precipitation measured by shielded
23 | gauges increases linearly with that of unshielded gauges, ~~independently of local environmental conditions.~~
24 | However, the relative increase in linear correlation should depend on the specific environmental conditions. For
25 | solid precipitation, some non-linear factors interfered with the linear relationship to reduce the correlation
26 | coefficient.

1 **Fig.2 about here**

2 **Fig.3 about here**

3 **Fig.4 about here**

4
5 **3.2 Comparisons of wind-induced bias**

6 **3.2.1 Rainfall**

7 From September 2010 to April 2015, the $CSPG_{PIT}$ caught 4.7% and 3.4% more rainfall than the $CSPG_{UN}$ and
8 the $CSPG_{SA}$ respectively ($(CSPG_{PIT}-CSPG_{UN})/CSPG_{UN}*100$; similarly hereinafter). The $CSPG_{SA}$ caught 1.3%
9 more rainfall than the $CSPG_{UN}$ (Table 3). During the period from September 2012 to April 2015, the $CSPG_{SA}$,
10 $CSPG_{PIT}$ and $CSPG_{DFIR}$ caught 0.9%, 4.5% and 3.4% more rainfall, respectively, than the $CSPG_{UN}$, and the
11 $CSPG_{PIT}$ and $CSPG_{DFIR}$ caught 3.6% and 2.5% more rainfall, respectively, than the $CSPG_{SA}$. However, the
12 $CSPG_{DFIR}$ caught 1.0% less rainfall than the $CSPG_{PIT}$ (Table 3, Fig.2). These comparative results indicate that the
13 $CSPG_{PIT}$ caught more rainfall and total precipitation compared to the $CSPG_{DFIR}$ and other gauges at the
14 experimental site (Table 3, Fig.2).

15 **3.2.2 Mixed precipitation**

16 A total of 29 mixed precipitation events were observed from September 2012 to April 2015. As shown in Table
17 3, the $CSPG_{PIT}$ caught the most mixed precipitation among the gauges, capturing 82.2 mm of mixed precipitation
18 in 29 events, but only 1.1 mm more than the $CSPG_{DFIR}$. The linear relationship between the $CSPG_{PIT}$ and
19 $CSPG_{DFIR}$ is statistically significant with an R^2 value of about 0.98 (Fig.3f). Thus for mixed precipitation, in
20 addition to the $CSPG_{DFIR}$, the $CSPG_{PIT}$ could also be selected as a reference gauge for the $CSPG_{UN}$ and $CSPG_{SA}$ at
21 the experimental site.

22 **3.2.3 Snowfall**

23 From September 2012 to April 2015, the $CSPG_{SA}$, $CSPG_{PIT}$ and ~~$CSPG_{DFIR}-CSPG_{DFIR}$~~ caught 11.1%, 16.0% and
24 20.6% more snowfall, respectively, than the $CSPG_{UN}$, and the $CSPG_{PIT}$ and $CSPG_{DFIR}$ caught 4.4% and 8.5%
25 more snowfall, respectively, than the $CSPG_{SA}$ (Table 3). Although the $CSPG_{DFIR}$ caught 3.9% more snowfall
26 compared to the $CSPG_{PIT}$ (Table 3), the difference in total snowfall (43 events) between the $CSPG_{DFIR}$ and
27 $CSPG_{PIT}$ was only about 3.4 mm (Table 3). Their linear correlation was highly significant with an R^2 value of
28 0.994 (Fig.4f). Blowing snow and thick snow cover have traditionally limited the pit's use as a reference ~~shield~~

1 for snowfall and mixed precipitation. At the experimental site, blowing snow was rarely observed and the snow
2 cover was usually shallow. This suggests that the $CSPG_{PIT}$ could be used as a reference gauge for snow
3 precipitation events at the ~~experimental site~~ with shallow snow cover and rare blowing snow event.

4 To sum up the comparisons of wind-induced bias, from most to least rainfall and mixed precipitation measured,
5 the instruments ranked as follows: $CSPG_{PIT} > CSPG_{DFIR} > CSPG_{SA} > CSPG_{UN}$, while for snowfall their ranking was
6 $CSPG_{DFIR} > CSPG_{PIT} > CSPG_{SA} > CSPG_{UN}$.

7 **3.3 Catch ratio vs. wind speed**

8 Previous studies have shown that wind speed during the precipitation period is the most significant variable
9 affecting gauge catch efficiency (Metcalf and Goodison, 1993; Yang et al., 1995; Goodison et al., 1998). Because
10 the CMA stations observe wind speeds at the 10m height, the $CSPG_{UN}$ and $CSPG_{SA}$ adjustment equations for a
11 single precipitation event were obtained for 10m height wind speeds. On the daily scale, adjustment equations
12 similar to Eqs.(2)–(4) were also obtained, based on the daily mean wind speed converted to gauge height (0.7m
13 for the CSPGs) and air temperature.

14 To minimise ratio scatter for the different gauges, precipitation events greater than 3.0 mm are normally
15 selected for the CR vs. wind analysis (Yang et al. 1995; Yang et al., 2014). However, in the Hulu watershed, most
16 snowfall and mixed precipitation events were less than 3.0 mm, thus the limit was reduced and single or daily
17 snowfall and mixed precipitation events greater than 1.0 mm were selected, while rainfall events greater than 3.0
18 mm were selected. The numbers of selected precipitation events are shown in Table 4. The CR vs. wind speed
19 relationships for different precipitation types were ~~simulated~~ determined using cubic polynomials and
20 exponential functions and were summarised in Table 4. The $CR_{UN/DFIR}$ and $CR_{SA/DFIR}$ vs. wind speed relationships
21 are statistically significant, but the $CR_{PIT/DFIR}$ vs. $W_{s0.7}$ or W_{s10} relationships do not pass the F-test with $\alpha=0.10$.
22 This phenomenon indicates that both PIT and DFIR are effective in preventing wind from influencing the gauge
23 catch of precipitation, therefore the $CR_{PIT/DFIR}$ is not related to wind speed.

24

25

26

27 Fig.5 presents scatter plots for the $CR_{UN/DFIR}$ and $CR_{SA/DFIR}$ vs. wind speed for rainfall. The CRs vary from 80%
28 to 110%. With increasing wind speed, the CRs decrease slightly. Only Eq. (10) shown in Fig.5 and Table 4 could
29 be used to adjust the rainfall event data from the $CSPG_{SA}$. It is significant at 0.03 level (Table 4). As described in

1 section 2.2, Eq.(10) was fitted using the NONLINEAR function in SPSS software
 2 (Analyze\Regression\Nonlinear). The F-value was then calculated using regression and the residual sum of
 3 squares from SPSS (Snedecor and Cochran, 1989). Based on the F-value and the degrees of freedom (Df), the
 4 significance level (α) was obtained using the FDIST function in Microsoft Excel. Other forms such as the
 5 exponential expression were treated in a similar way.

$$6 \quad CR_{SA/DFIR,Rain} = 0.188W_{s10}^3 - 0.719W_{s10}^2 + 0.551W_{s10} + 100 \quad 0 < W_{s10} < 7.4 \quad (10)$$

7 where $CR_{SA/DFIR,Rain}$ is the rainfall catch ratio (%) per observation of the CSPG_{SA} and W_{s10} is the wind speed at
 8 10m during the rainfall period ($m s^{-1}$).

9
 10 **Fig.5 about here**

11
 12 On the daily scale, the relationships between rainfall CR and wind speed at gauge height ($W_{s0.7}$) are also cubic
 13 functions, but they do not pass the F-test with $\alpha=0.25$ (Table 4).

14 For the mixed precipitation events, the CR vs. W_{s10} relationships are exponential (Table 4, Fig.6). The CRs vary
 15 greatly from about 60% to 120%. For the CSPG_{UN}, the exponential relationship Eq. (11) passes the F-test with
 16 $\alpha=0.07$, whereas for the CSPG_{SA}, the Eq.(12) α value is about 0.16 (Table 4).

17
 18 **Fig.6 about here**

$$19 \quad CR_{UN/DFIR,Mixed} = 100e^{-0.06W_{s10}} \quad 0 < W_{s10} < 5.9 \quad (11)$$

$$20 \quad CR_{SA/DFIR,Mixed} = 100e^{-0.04W_{s10}} \quad 0 < W_{s10} < 5.9 \quad (12)$$

21
 22 On the daily scale, the relationships between mixed precipitation CR and wind speed at gauge height ($W_{s0.7}$) are
 23 also exponential expressions (Table 4, Fig.6). Similarly, for the CSPG_{UN}, Eq. (13) passes the F-test with $\alpha < 0.10$,
 24 whereas Eq.(14) with an α value of about 0.18 doesn't (Table 4).

$$25 \quad CR_{UN/DFIR,Mixed} = 100e^{-0.12W_{s0.7}} \quad 0 < W_{s0.7} < 2.9 \quad (13)$$

$$26 \quad CR_{SA/DFIR,Mixed} = 100e^{-0.07W_{s0.7}} \quad 0 < W_{s0.7} < 2.9 \quad (14)$$

27 From Eq. (3), air temperature may also affect the mixed precipitation CRs on the daily scale. Eqs. (15)–(16) are
 28 obtained as follows. However, these two new equations do not pass the F-test with $\alpha < 0.20$.

$$29 \quad CR_{UN/DFIR,Mixed} = 13.83W_{s0.7}^{-4.91} + 1.25T_{max} - 0.88T_{min} + 62.21 \quad \alpha=0.20 \quad (15)$$

$$1 \quad CR_{SA/DFIR,Mixed} = 10.74W_{s0.7}^{-4.74} + 0.85T_{\max} - 0.18T_{\min} + 76.20 \quad \alpha=0.29 \quad (16)$$

2 where T_{\max} and T_{\min} are the daily maximum and minimum air temperature ($^{\circ}\text{C}$), respectively.

3 For the snowfall events, the $CR_{UN/DFIR,Snow}$ and $CR_{SA/DFIR,Snow}$ vs. W_{s10} relationships are significant (Table 4,
4 Fig.7). For the CSPG_{UN}, the exponential relationship Eq.(17) passes the F-test with $\alpha<0.001$. Eq.(17) is similar to
5 Eq.(5) suggested by Yang et al. (1991). For the CSPG_{SA}, its exponential expression in Eq.(18) passes the F-test at
6 $\alpha=0.07$ (Table 4).

7
8 **Fig.7 about here**

$$11 \quad CR_{UN/DFIR,Snow} = 100e^{-0.08W_{s10}} \quad 0 < W_{s10} < 4.8 \quad (17)$$

$$12 \quad CR_{SA/DFIR,Snow} = 100e^{-0.02W_{s10}} \quad 0 < W_{s10} < 4.8 \quad (18)$$

13 On the daily scale, the relationships between snowfall CRs and wind speed at gauge height ($W_{s0.7}$) are also
14 exponential expressions (Table 4, Fig.7). For the CSPG_{UN} and CSPG_{SA}, the Eqs.(19)–(20) pass the F-test with
15 $\alpha<0.001$ and $\alpha=0.14$, respectively (Table 4). Eqs. (17)–(19) could therefore be directly used to calibrate the
16 wind-induced snowfall measurement errors for the CSPG_{UN} and CSPG_{SA}.

$$17 \quad CR_{UN/DFIR,Snow} = 100e^{-0.11W_{s0.7}} \quad 0 < W_{s0.7} < 3.1 \quad (19)$$

$$18 \quad CR_{SA/DFIR,Snow} = 100e^{-0.03W_{s0.7}} \quad 0 < W_{s0.7} < 3.1 \quad (20)$$

19 Air temperature may also affect the snowfall CR on the daily scale as shown in Eq.(2). Eqs. (21)–(22) are the
20 new equations associated with daily maximum air temperature. However, these two new equations are ~~no~~not
21 better than Eqs. (19) –(20) according to their F-test α values.

$$22 \quad CR_{UN/DFIR,Snow} = 42.29W_{s0.7}^{-1.06} - 1.06T_{\max} + 55.91 \quad \alpha=4.2E-5 \quad (21)$$

$$23 \quad CR_{SA/DFIR,Snow} = -9.46\ln(W_{s0.7}) - 0.31T_{\max} + 98.76 \quad \alpha=0.17 \quad (22)$$

24 From the above mentioned relationships of $CR_{UN/DFIR}$ and $CR_{SA/DFIR}$ vs. wind speed, the following points can be
25 drawn for our understanding. For daily rain and mixed precipitation, the relationships are not statistically
26 significant. The use of daily mean wind speed may lead to uncertainties in gauge comparisons. Data collections
27 and analyses on shorter time scales, such as hourly or 6-hourly, are expected to produce more reliable results,
28 because wind speed may vary throughout the day and daily mean wind speeds may not be representative of the
29 wind conditions over the precipitation period (Yang and Simonenko, 2014). Daily maximum and minimum

1 temperatures should reflect the atmospheric conditions of radiation and convection to some degree, and their
2 function in the CR vs. wind speed relationship needs further investigation in a mountain environment.

3 **4 Discussion**

4 **4.1 Comparison with other studies**

5 Yang et al. (1991) carried out a precipitation intercomparison experiment from 1985 to 1987 at the Tianshan
6 site. Their results indicated that the $CSPG_{DFIR}/CSPG_{UN}$ ratios for snowfall and mixed precipitation were 1.222 and
7 1.160, respectively. In the Hulu watershed, these ratios were 1.165 (Fig.4c) and 1.072 (Fig.3c), while those for
8 $CSPG_{PIT}/CSPG_{UN}$ were 1.162 (Fig.4b) and 1.082 (Fig.3b), respectively. Similar topographic features and shading
9 induced similar lower wind speeds and led to similar catch ratios at both sites. For the Tianshan ~~reference-study~~
10 site, wind speed (W_{s10}) on rainfall or snowfall days never exceeded 6 m s^{-1} , and 88% of the total annual
11 precipitation took place with wind speeds below 3 m s^{-1} . At the Hulu watershed site, daily mean wind speeds (W_{s10})
12 on precipitation days never exceeded 6.4 m s^{-1} , and over 55.2% of the precipitation events occurred with daily
13 mean wind speeds below 3 m s^{-1} . During the periods of precipitation, the ~~largest-highest~~ wind speed at the 10m
14 height was about 8.8 m s^{-1} , and over 54.2% of the precipitation events occurred with wind speeds below 3 m s^{-1} .

15 As Ren et al. (2003) reported, across 30 comparison stations in China, the $CSPG_{PIT}$ caught 3.2% (1.1~7.9%)
16 more rainfall and 11.0% (2.2~24.8%) more snowfall ~~compared-to~~ than the $CSPG_{UN}$. Large wind-induced
17 differences were often observed at the mountainous western stations and in north-eastern China. At the Gangcha
18 station ($100^{\circ}08'$, $37^{\circ}20'$, 3015 m), which also lies in the Qilian Mountains at a similar elevation about 200 km
19 from the Hulu watershed site, the $CSPG_{PIT}$ caught 7.9% more rainfall and 16.8% more snowfall than the $CSPG_{UN}$
20 from 1992 to 1998. In our study, the $CSPG_{PIT}$ captured 4.7% more rainfall, 21.0% more snowfall and 12.1% more
21 mixed precipitation than the $CSPG_{UN}$ from September 2010 to April 2015 (Table 3). The outcome presented in
22 this study is somewhat different from that reported by Ren et al. (2003) due to differences in the wind regime. At
23 the Gangcha station, daily mean wind speeds (W_{s10}) on precipitation days during the experimental period from
24 1992 to 1998 never exceeded 8.5 m s^{-1} , and over 35.1% of the precipitation events occurred with daily mean wind
25 speeds below 3 m s^{-1} . The average daily mean W_{s10} was about 3.4 m s^{-1} on precipitation days from 1992 to 1998 at
26 the Gangcha station, whereas at the Hulu watershed site from 2010 to 2015, the average value was about 2.9 m s^{-1}
27 on precipitation days.

28 It is recognised that in western China, climatic and environmental conditions in the mountains vary both
29 spatially and temporally. To understand the similarities and differences in wind-induced bias in precipitation

1 measurements for different mountain watersheds, field experiments need to be carried out continuously. Further
2 investigation is also necessary to consider the influence of micrometeorology on gauge observations, particularly
3 wind distribution and turbulence across this site (Yang and Simonenko, 2014).

4 **4.2 CSPG_{PIT} as a reference for solid precipitation**

5 The pit is the WMO reference configuration for liquid precipitation measurements and the DFIR is the
6 reference configuration for solid precipitation measurements (Sevruk et al., 2009). In this study, the CSPG_{PIT}
7 measured more rainfall and mixed precipitation than the CSPG_{DFIR}. For snowfall, the catch ratio for CSPG_{PIT} was
8 0.96, close to that of the CSPG_{DFIR} measurement. The difference in total snowfall (43 events) between the
9 CSPG_{PIT} and CSPG_{DFIR} was only about 3.4 mm from September 2012 to April 2015 at the Hulu watershed site.
10 The snowfall for autumn and spring was greater than for winter during the observation period at the
11 intercomparison site (Fig.8). The snowfall is wetter in autumn and spring than in winter, and wetter snowfall
12 means less blowing or drifting snow. Thus the CSPG_{PIT} could serve as a reference for liquid and solid
13 precipitation in environments similar to that of the Hulu watershed site. Precipitation collected by the CSPG_{PIT}
14 would be most affected by blowing or drifting snow, inducing a faulty precipitation value (Goodison et al., 1998;
15 Ren and Li, 2007). Previous studies have indicated, however, that for most of China the maximum snow depth in
16 the past 30 years has been less than 20 cm (Li, 1999), with average snow depths below 3 cm (Li et al., 2008; Che
17 et al., 2008). Fig.9 shows annual snowfall amounts and annual snowfall proportion distributions for 644
18 meteorological stations in China from 1960 to 1979, indicating that snowfalls ~~s-are~~ is concentrated in the middle
19 and south-western Tibetan Plateau, northern Xinjiang province and north-eastern China. Statistical analysis
20 indicates that for more than 94% of stations, solid precipitation comprises less than 15% of the annual
21 precipitation. Ren et al. (2003) reported, that among the 2286 snowfall events, only 54 were blowing or drifting
22 snow events accounting for about 2.4% for 26 stations across China. Based on the regionalisation of snow drift in
23 China, blowing or drifting snow events occur mostly on the central and south-western Tibetan Plateau, in the
24 northern Xinjiang province and in north-eastern China (Wang and Zhang, 1999). In these regions, the CSPG_{DFIR}
25 should be used as a reference gauge. In other regions, the CSPG_{PIT} may be applicable. Based on the CMA
26 snowfall and snow depth data, and the regionalisation of snow drift in China, the applicable regions for the
27 CSPG_{PIT} and CSPG_{DFIR} as reference gauges are shown in Fig.10.

28 **Fig.8 about here**

29 **Fig.9 about here**

1 **Fig.10 about here**

2 **4.3 Limitations of this experiment**

3 Although the measurement procedures were based on the CMA's ~~criteria~~standard, manual observations were
4 infrequent, and as a result, some precipitation events were summarised as single events, especially in the evenings.
5 The automatic meteorological tower could observe precipitation and wind speeds half-hourly during the
6 precipitation period, but the $CSPG_{UN}$, $CSPG_{SA}$, $CSPG_{PIT}$ and $CSPG_{DFIR}$ were observed only twice per day. In this
7 field experiment, the precipitation phases were also distinguished by observers. This method is somewhat
8 imprecise although this has remained the traditional method since the 1950s at the CMA stations (CMA, 2007b).

9 Automatic sensors will also be important to detect precipitation types at operational and research networks (Yang
10 and Simonenko, 2014).

11 The wind speeds at gauge height and the 10 m height were not observed directly but rather calculated from the
12 observed data at 1.5 m and 2.5m heights according to the Monin-Obukhov theory and the gradient method
13 (Eq.(8)). Although this method is widely used, it is effective only under neutral atmospheric conditions. For the
14 precipitation period from September 2012 to April 2015, the Z_0 was calculated using Eq. (9). The results showed
15 the Z_0 to be about 0.06m on average but it varied from nearly zero to 0.67m. As shown in Fig.11, in about 68.9%
16 and 95.1% of instances, the Z_0 was lower than 0.05 m and 0.25 m, respectively. In rare cases when the Z_0 was very
17 large, as shown in Fig.11, the Z_0 was arbitrarily assigned 1/2 of the grass height (h) at the site based on the
18 equation $Z_0=0.5hL_e$ provided by Lettau (1969). The very large Z_0 values usually appeared in late August and early
19 September when the vegetation coverage (L_e) was close to 100% at the Hulu watershed site.

20
21 **Fig. 11 about here**

22 **5 Conclusions**

23 ~~The present experimental field~~This study focused on wind-induced bias in precipitation measurements by
24 CSPGs specifically in a high mountain environment. The precipitation intercomparison experiment in the Hulu
25 watershed of the Qilian Mountains indicated that the $CSPG_{PIT}$ caught more rainfall, mixed precipitation and total
26 precipitation but less snowfall than the $CSPG_{DFIR}$. From most to least rainfall and mixed precipitation measured,
27 their ranking was $CSPG_{PIT} > CSPG_{DFIR} > CSPG_{SA} > CSPG_{UN}$, whereas in the snowy season, better wind shielding
28 increased the snow catch, leading to $CSPG_{DFIR} > CSPG_{PIT} > CSPG_{SA} > CSPG_{UN}$. The measured daily precipitation
29 by shielded gauges increases linearly with that of unshielded gauges. ~~and is independent of local environmental~~

1 ~~conditions. However, an increase in the ratio of the linear correlation should depend on specific environmental~~
2 ~~conditions.~~ For solid precipitation, some non-linear factors interfere with the linear relationship to reduce the
3 linear correlation coefficient.

4 In regions with lower snowfall, such as the southern and central parts of China (Zhang and Zhong, 2014), and
5 in regions with a similar climate and environment to that of the Hulu watershed site, the CSPG_{PIT} could be used as
6 a reference gauge because of its high catch ratio, simplicity and lower maintenance requirements. In north-eastern
7 China, northern Xinjiang province and the central and south-western Tibetan Plateau where snowfalls often occurs,
8 the best choice of reference gauge would be the CSPG_{PIT} for rainfall and the CSPG_{DFIR} for snowfall observations.

9 The catch ratio vs. wind speed relationship for different precipitation types is simulated by cubic polynomials
10 and exponential functions. The CR_{PIT/DFIR} does not have a significant relationship to wind speed, indicating that
11 both PIT and DFIR are effective in preventing wind from influencing the precipitation gauge catch. For daily rain
12 and mixed precipitation, the relationships are not statistically significant. Daily maximum and minimum
13 temperatures should reflect the atmospheric conditions of radiation and convection to some degree, and their
14 function in the CR vs. wind speed relationship needs further investigation in mountain environments. It is
15 recognised that in western China, the climatic and environmental conditions in the mountains vary both spatially
16 and temporally. To understand the similarities and differences among wind-induced biases in precipitation
17 measurements for the different mountain watersheds in western China, field experiments and modelling of wind
18 field need to be carried out continuously.

19 Acknowledgments

20 This ~~paper study~~ was supported primarily by the National Basic Research Program of China (2013CBA01806)
21 and the National Natural Sciences Foundation of China (91025011, 41222001, 91225302 and 41401078).

22 References

- 23 Aaltonen, A., Elomaa, E., Tuominen, A., and Valkovuori, P.: Measurement of precipitation, in: Proceedings of the
24 Symposium on Precipitation and Evaporation, edited by: Sevruck, B. and Lapin, M., Slovak
25 Hydrometeorological Institute and Swiss Federal Institute of Technology, Bratislava, Slovakia, 42–46, 1993.
- 26 Bagnold, R. A.: The Physics of Blown Sand and Desertdunes, Methuen, New York, 85-95, 1941.
- 27 Charnes, A., Frome, E. L., and Yu, P. L.: The equivalence of generalized least squares and maximum likelihood
28 estimates in the exponential family, Journal of the American Statistical Association, 71: 169, 1976.

1 Che, T., Li, X., Jin, R., Armstrong, R., and Zhang T.: Snow depth derived from passive microwave remote-sensing
2 data in China, *Ann. Glaciol.*, 49, 145-154, 2008.

3 Chen, R., Song, Y., Kang, E., Han, C., Liu, J., Yang, Y., Qing, W., and Liu, Z.: A Cryosphere-Hydrology
4 observation system in a small alpine watershed in the Qilian Mountains of China and its meteorological
5 gradient, *Arct. Antarct. Alp. Res.*, 46(2): 505-523. doi: <http://dx.doi.org/10.1657/1938-4246-46.2.505>, 2014.

6 China Meteorological Administration (CMA): Specifications for surface meteorological observation Part 8:
7 Measurement of precipitation (QX/T 52-2007), China Meteorological Press, Beijing, 2007a.

8 China Meteorological Administration (CMA): Specifications for surface meteorological observation Part 4:
9 Observation of weather phenomenon (QX/T 48-2007), China Meteorological Press, Beijing, 2007b.

10 Ding, Y., Yang, D., Ye, B., and Wang, N.: Effects of bias correction on precipitation trend over China, *J. Geophys.*
11 *Res.*, 112, D13116, doi:10.1029/2006JD007938, 2007.

12 Dyer, A. J., and Bradley, E. F.: An alternative analysis of flux-gradient relationships at the 1976 ITCE, *Bound.-*
13 *Lay. Meteorol.*, 22, 3–19, doi: 10.1007/BF00128053, 1982.

14 Golubev, V. S.: On the problem of actual precipitation measurements at the observations site, in: *Proceeding of the*
15 *International Workshop on the Correction of Precipitation Measurements WMO/TD 104*, World
16 *Meteorological Organization*, Geneva, Switzerland, 61–64, 1985.

17 Goodison, B. E., Louie, B. P. Y. T., and Yang, D.: WMO solid precipitation measurement intercomparison: Final
18 report, *Instrum. and Obs. Methods Rep. 67/Tech. Doc. 872*, World Meteorol. Organ., Geneva, Switzerland,
19 1998.

20 IBM Corp: IBM SPSS Statistics for Windows, Version 19.0, IBM Corp, Armonk, NY, USA, 2010.

21 Lanza, L. G., Leroy, M., Alexandropoulos, C., Stagi, L., and Wauben, U.: WMO Laboratory Intercomparison of
22 Rainfall Intensity Gauges - Final Report, IOM Report No. 84, WMO/TD No. 1304, WMO, Geneva,
23 Switzerland, 2005.

24 Lettau, H.: Note on aerodynamic roughness-parameter estimation on the basis of roughness element description,
25 *Journal of Applied Aerology*, 8, 828-832, 1969.

26 Li, P.: Variation of snow water resources in northwestern China, 1951-1997, *Sci. China Ser. D*, 42, 73-79, 1999.

27 Li, X., Cheng, G., Jin, H., Kang, E., Che, T., Jin, R., Wu, L., Nan, Z., Wang, J., and Shen, Y.: Cryospheric change

- 1 in China, *Global Planet. Change*, 62, 210-218, doi:10.1016/j.gloplacha.2008.02.001, 2008.
- 2 Ma, Y., Zhang, Y., Yang, D., and Farhan, S.: Precipitation bias variability versus various gauges under different
3 climatic conditions over the Third Pole Environment (TPE) region, *Int. J. Climatol.*, doi: 10.1002/joc.4045,
4 2014.
- 5 Metcalfe, J. R., and Goodison, B. E.: Correction of Canadian winter precipitation data. Preprints, in: Eighth Symp.
6 on Meteorological Observations and Instrumentation, Anaheim, CA, Am. Meteorol. Soc., 338–343, 1993.
- 7 Ren, Z., and Li, M.: Errors and correction of precipitation measurements in China, *Adv. Atmos. Sci.*, 24, 449–458,
8 doi: 10.1007/s00376-007-0449-3, 2007.
- 9 Ren, Z., Wang, G., Zou, F., and Zhang, H.: The research of precipitation measurement errors in China, *Acta*
10 *Meteorol. Sin.*, 61, 621-627, 2003.
- 11 Rodda, J. C.: Annotated Bibliography on Precipitation Measurement Instruments, WMO-No. 343, World Meteorol.
12 Org, Geneva., Switzerland, 1973.
- 13 Rodda, J. C.: The rainfall measurement problem, in: Proceedings of IAHS, General Assembly, Bern 1967,
14 Publication No. 78: 215-231, 1967.
- 15 Sevruk, B., and Hamon, W. R.: International comparison of national precipitation gauges with a reference pit gauge,
16 instruments and observing methods Rep., 17, 135, World Meteorol. Org., Geneva, 1984.
- 17 Sevruk, B., Ondras, M., and Chvıla, B.: The WMO precipitation measurement intercomparisons, *Atmos. Res.*, 92,
18 376–380, doi:10.1016/j.atmosres.2009.01.016, 2009.
- 19 Snedecor, G., and Cochran, W.: Statistical methods. Iowa State University Press, Iowa, 1989.
- 20 Struzer, L. R.: Practicability analysis of rain gauge international comparison test results (in Russian). *Trans.*
21 *Voyeykov Main Geophys. Observ*, 260, 77-94, 1971.
- 22 Sugiura, K., Yang, D., and Ohata, T.: Systematic error aspects of gauge-measured solid precipitation in the Arctic,
23 Barrow, Alaska, *Geophys. Res. Lett.*, 30, 1192, doi:10.1029/2002GL015547, 2003.
- 24 Tian, X., Dai, A., Yang, D., and Xie, Z.: Effects of precipitation-bias corrections on surface hydrology over
25 northern latitudes, *J. Geophys. Res.*, 112, D14101, doi:10.1029/2007JD008420, 2007.
- 26 Wang, Z., and Zhang, Z.: Regionalization of snow drift in China, *J. Mt. Sci.*, 17, 312-317, 1999.
- 27 Wolff, A. M., Nitu, R., Earle, M., Joe, P., Kochendorfer, J., Rasmussen, R., Reverdin, A., Sminth, C., Yang, D.,

- 1 and the SPICE-TEAM: WMO Solid Precipitation Intercomparison Experiment (SPICE): Report on the SPICE
2 Field Working Reference System for precipitation amount, WMO, IOM No. 116, TECO-2014, World
3 Meteorological Organization, Geneva, Switzerland, 2014.
- 4 Yang, D., Goodison, B. E., Metcalfe, J. R., Golubev, V. S., Elomaa, E., Gunther, T. H., Bates, R., Pangburn, T.,
5 Hanson, C. L., Emerson, D., Copaciu, V., and Milkovic, J.: Accuracy of Tretyakov precipitation gauge: Results
6 of WMO intercomparison, *Hydrol. Process.*, 9, 877– 895, doi:10.1002/hyp.3360090805, 1995.
- 7 Yang, D., Goodison, B. E., Metcalfe, J. R., Louie, P., Leavesley, G., Emerson, D., Hanson, C. L., Golubev, S. S.,
8 Elomaa, E., Gunthter, T., Pangburn, T., Kang, E., and Milkovic, J.: Quantification of precipitation
9 measurement discontinuity induced by wind shields on national gauges, *Water Resources Research*, 35, 491 –
10 508, doi: 10.1029/1998WR900042, 1999.
- 11 Yang, D., Metcalfe, J. R., Goodison, B. E., and Mekis, E.: An evaluation of Double-Fence Intercomparison
12 Reference (DFIR) gauge, in: *Proceedings of Eastern Snow Conference, 50th Meeting, Quebec, City*, 105–111,
13 1993.
- 14 Yang, D., Shi, Y., Kang, E., Zhang, Y., and Yang, X.: Results of solid precipitation measurement intercomparison
15 in the Alpine area of Urumqi River basin, *Chin. Sci. Bull.*, 36, 1105–1109, 1991.
- 16 [Yang, D., and Simonenko, A.: Comparison of winter precipitation measurements by six Tretyakov gauges at the](#)
17 [Valdai experimental site, *Atmosphere-Ocean*, 52, 1, 39-53, doi: 10.1080/07055900.2013.865156, 2014.](#)
- 18 Yang, D.: Double-Fence Intercomparison Reference (DFIR) vs. Bush Gauge for “true” snowfall measurement, *J.*
19 *Hydrol.*, 509, 94–100, doi:10.1016/j.jhydrol.2013.08.052, 2014.
- 20 Yang, D.: Research on analysis and correction of systematic errors in precipitation measurement in Urumqi River
21 basin, Tianshan, PhD thesis, Lanzhou Institute of Glaciology and Geocryology, Chinese Academy of Sciences,
22 Lanzhou, China, 169 pp., 1988.
- 23 Ye, B., Yang, D., and Ma, L.: Effect of precipitation bias correction on water budget calculation in Upper
24 YellowRiver, China, *Environ. Res. Lett.*, 7, 025201, doi:10.1088/1748-9326/7/2/025201, 2012.
- 25 ~~Ye, B., Yang, D., Ding, Y., and Han, T.: A bias-corrected precipitation climatology for China. *Acta Geogr. Sin.*, 62,~~
26 ~~3-13, 2007.~~
- 27 Ye, B., Yang, D., Ding, Y., Han, T., and Koike, T.: A bias-corrected precipitation climatology for China, *J.*
28 *Hydrometeorol.*, 5, 1147–1160, doi: http://dx.doi.org/10.1175/JHM-366.1, 2004.

1 Zhang, T., and Zhong, X.: Classification and regionalization of the seasonal snow cover across the Eurasian
2 Continent, *J. Glaciol. Geocryol.*, 36, 481-490, 2014.

3 Zhang, Y., Ohata, T., Yang, D., and Davaa, G.: Bias correction of daily precipitation measurements for Mongolia,
4 *Hydrol. Process.*, 18, 2991–3005, doi: 10.1002/hyp.5745, 2004.

5

6

7

1
2
3
4
5
6
7
8
9
10
11
12
13
14

Table 1. Monthly climate values at the experimental site (2010-2012).

Element	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Yearly
Monthly precipitation (mm)	3.5	2.5	11.0	8.8	67.7	69.6	87.1	111.6	57.7	24.0	2.7	1.0	447.2
Monthly mean air temperature (°C)	-12.4	-7.7	-4.4	2.2	7.0	11.2	12.5	12.1	8.0	1.4	-5.6	-11.3	1.1
Monthly mean daily maximum air temperature (°C)	-4.0	0.7	3.5	10.3	14.3	18.2	19.5	19.7	15.4	10.2	3.6	-1.9	9.1
Monthly mean daily minimum air temperature (°C)	-19.0	-14.8	-11.6	-5.2	0.6	4.9	6.8	5.8	1.8	-5.5	-12.7	-18.2	-5.6
Monthly mean wind speed at the 1.5m height (m s ⁻¹)	1.79	1.96	2.30	2.55	2.42	1.98	1.82	1.81	1.93	1.81	2.08	1.96	2.03
Monthly mean wind speed at the 2.5m height (m s ⁻¹)	1.79	2.02	2.43	2.77	2.65	2.16	2.04	2.02	2.16	1.99	2.19	2.01	2.18
Monthly potential evaporation (mm)	31.6	47.0	79.4	124.4	140.9	155.0	141.7	127.0	101.6	75.2	47.3	31.0	1102.2

Table 2. The precipitation measurement intercomparison experiment in the Qilian Mountains.

Gauge	Abbreviation	Size(φ denotes orifice diameter and h is observation height)	Start date	End date	Observation time
Unshielded China standard precipitation gauge (CMA, 2007a)	CSPG _{UN}	$\varphi=20\text{cm}, h=70\text{cm}$	Jun 2009	Apr, 2015	20:00 and 08:00, Local time
Single Alter shield (Struzer, 1971) around a CSPG	CSPG _{SA}	$\varphi=20\text{cm}, h=70\text{cm}$	Jun 2009	Apr, 2015	20:00 and 08:00, Local time
A CSPG in a Pit (Sevruk and Hamon, 1984)	CSPG _{PIT}	$\varphi=20\text{cm}, h=0\text{cm}$	Sep 2010	Apr, 2015	20:00 and 08:00, Local time
DFIR shield(Goodison et al., 1998) around a CSPG	CSPG _{DFIR}	$\varphi=20\text{cm}, h=3.0\text{m}$	Sep 2012	Apr, 2015	20:00 and 08:00, Local time

1

2

3

Table 3. Summary of precipitation observations at the Hulu watershed intercomparison site, 2010-2015.

Date	Phase	No. of events	Total precipitation and catch ratio (CR, %)													
			CSPG _{UN} (mm)	CR	$100 \left(\frac{\text{CSPG}_{\text{SA}}}{\text{CSPG}_{\text{UN}}} - 1 \right)$	$100 \left(\frac{\text{CSPG}_{\text{PIT}}}{\text{CSPG}_{\text{UN}}} - 1 \right)$	$100 \left(\frac{\text{CSPG}_{\text{DFIR}}}{\text{CSPG}_{\text{UN}}} - 1 \right)$	CSPG _{SA} (mm)	CR	$100 \left(\frac{\text{CSPG}_{\text{PIT}}}{\text{CSPG}_{\text{SA}}} - 1 \right)$	$100 \left(\frac{\text{CSPG}_{\text{DFIR}}}{\text{CSPG}_{\text{SA}}} - 1 \right)$	CSPG _{PIT} (mm)	CR	$100 \left(\frac{\text{CSPG}_{\text{DFIR}}}{\text{CSPG}_{\text{PIT}}} - 1 \right)$	CSPG _{DFIR} (mm)	CR
Sep 2010-	All	608	1986.8	93.9	2.6	6.5		2038.1	96.4	3.8		2115.1	100			
	rain	480	1700.7	95.5	1.3	4.7		1723.4	96.7	3.4		1781.4	100			
Apr 2015	mixed	44	139.9	89.2	6.1	12.1		148.5	94.7	5.6		156.8	100			
	snow	84	146.2	82.6	13.7	21.0		166.2	94.0	6.4		176.9	100			
Sep 2012-	All	283	1066.7	94.9	2.0	6.0	5.3	1088.4	96.9	3.9	3.2	1130.9	100.6	-0.6	1123.7	100
	rain	211	920.7	96.7	0.9	4.5	3.4	928.6	97.5	3.6	2.5	961.8	101.0	-1.0	952.2	100
Apr 2015	mixed	29	71.1	87.6	7.7	15.6	14.2	76.6	94.3	7.3	6.0	82.2	101.2	-1.2	81.2	100
	snow	43	74.9	82.9	11.1	16.0	20.6	83.2	92.1	4.4	8.5	86.9	96.2	3.9	90.3	100

4

5

1
2
3
4
5
6
7
8
9
10
11
12

Table 4. Catch ratio (CR) vs. wind speed relationships at the Hulu watershed intercomparison site, 2012-2015.

Temporal scale	Phase	Gauges	Catch ratio (CR) vs. wind speed relationships*	<i>P</i> (mm)	No. of events	F-test
Precipitation event	Rain	CSPG _{UN}	$CR_{UN/DFIR,Rain} = 0.181W_{s10}^3 - 0.256W_{s10}^2 - 0.795W_{s10} + 100$ $R^2=0.042$	<i>P</i> >3.0	103	$\alpha=0.23$
		CSPG _{SA}	$CR_{SA/DFIR,Rain} = 0.188W_{s10}^3 - 0.719W_{s10}^2 + 0.551W_{s10} + 100$ $R^2=0.083$			$\alpha=0.03$
		CSPG _{PIT}	$CR_{PIT/DFIR,Rain} = 0.150W_{s10}^3 - 0.425W_{s10}^2 + 1.119W_{s10} + 100$ $R^2=0.008$			$\alpha=0.83$
	Mixed	CSPG _{UN}	$CR_{UN/DFIR,Mixed} = 100e^{-0.06W_{s10}}$ $R^2=0.194$	<i>P</i> >1.0	24	$\alpha=0.07$
		CSPG _{SA}	$CR_{SA/DFIR,Mixed} = 100e^{-0.04W_{s10}}$ $R^2=0.100$			$\alpha=0.16$
		CSPG _{PIT}	$CR_{PIT/DFIR,Mixed} = 100e^{-7E-0W_{s10}}$ $R^2=0.000$			$\alpha=no\ data$
	Snow	CSPG _{UN}	$CR_{UN/DFIR,Snow} = 100e^{-0.08W_{s10}}$ $R^2=0.412$	<i>P</i> >1.0	34	$\alpha=6.4E-05$
		CSPG _{SA}	$CR_{SA/DFIR,Snow} = 100W_{s10}^{-0.02}$ $R^2=0.090$			$\alpha=0.07$
		CSPG _{PIT}	$CR_{PIT/DFIR,Snow} = 100e^{-0.01W_{s10}}$ $R^2=0.024$			$\alpha=0.35$
Daily precipitation	Rain	CSPG _{UN}	$CR_{UN/DFIR,Rain} = -1.400W_{s0.7}^3 + 2.987W_{s0.7}^2 - 6.116W_{s0.7} + 100$ $R^2=0.032$	<i>P</i> >3.0	90	$\alpha=0.37$
		CSPG _{SA}	$CR_{SA/DFIR,Rain} = -0.924W_{s0.7}^3 + 1.158W_{s0.7}^2 - 3.338W_{s0.7} + 100$ $R^2=0.021$			$\alpha=0.55$
		CSPG _{PIT}	$CR_{PIT/DFIR,Rain} = -0.952W_{s0.7}^3 - 1.503W_{s0.7}^2 + 2.237W_{s0.7} + 100$ $R^2=-0.00$			$\alpha=no\ data$
	Mixed	CSPG _{UN}	$CR_{UN/DFIR,Mixed} = 100e^{-0.12W_{s0.7}}$ $R^2=0.144$	<i>P</i> >1.0	21	$\alpha=0.09$
		CSPG _{SA}	$CR_{SA/DFIR,Mixed} = 100e^{-0.07W_{s0.7}}$ $R^2=0.094$			$\alpha=0.18$
		CSPG _{PIT}	$CR_{PIT/DFIR,Mixed} = 100e^{-0.001W_{s0.7}}$ $R^2=0.003$			$\alpha=no\ data$
	Snow	CSPG _{UN}	$CR_{UN/DFIR,Snow} = 100e^{-0.11W_{s0.7}}$ $R^2=0.477$	<i>P</i> >1.0	27	$\alpha=1.8E-04$
		CSPG _{SA}	$CR_{SA/DFIR,Snow} = 100e^{-0.03W_{s0.7}}$ $R^2=0.087$			$\alpha=0.14$
		CSPG _{PIT}	$CR_{PIT/DFIR,Snow} = 100e^{-0.01W_{s0.7}}$ $R^2=-0.00$			$\alpha=no\ data$

*: W_{s10} -Wind speed during period of precipitation at 10 m height; $W_{s0.7}$ -Daily mean wind speed at gauge height (0.7 m for CSPG).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36





Figure 1. Precipitation gauge intercomparison experiment in the Qilian Mountains, Tibetan Plateau.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27



28 **Figure 2.** Intercomparison plots among CSPG_{UN}, CSPG_{SA}, CSPG_{PIT} and CSPG_{DFIR} for the rainfall events from
29 September 2010 (a, b and d) and September 2012 (c, e and f) to April 2015.

30

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25





Figure 3. Intercomparison plots among CSPG_{UN}, CSPG_{SA}, CSPG_{PIT} and CSPG_{DFIR} for the mixed precipitation events from September 2010 (a, b and d) and September 2012 (c, e and f) to April 2015.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22



23 **Figure 4.** Intercomparison plots among CSPG_{UN}, CSPG_{SA}, CSPG_{PIT} and CSPG_{DFIR} for the snowfall events from
24 September 2010 (a, b and d) and September 2012 (c, e and f) to April 2015.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25




Figure 5. Catch ratios (CRs) vs. wind speed for rainfall events (a and b) and daily rainfall (c and d) greater than 3.0 mm.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30




Figure 6. Catch ratios (CRs) vs. wind speed for mixed precipitation events (a and b) and daily mixed precipitation (c and d) greater than 1.0 mm.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30




Figure 7. Catch ratios (CRs) vs. wind speed for the snowfall event (a and b) and the daily (c and d) snowfall greater than 1.0 mm.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30




Figure 8. Seasonal snowfall and its percentage from September 2010 to April 2015 at the Hulu watershed site.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25




Figure 9. (a) Annual snowfall (mm) and **(b)** ratio of annual snowfall to total precipitation in China.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25




Figure 10. Applicable regions in China for the $CSPG_{PIT}$ and $CSPG_{DFIR}$ as reference gauges.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21




Figure 11. Surface roughness during the precipitation period from September 2012 to April 2015.