



17 **Abstract**

18 This study quantifies the inconsistency in gauge precipitation observations across the border of Alaska  
19 and Yukon. It analyses the precipitation measurements by the national standard gauges (NWS 8-in gauge  
20 and Nipher gauge), and the bias-corrected data to account for wind effect on the gauge catch, wetting loss  
21 and trace events. The bias corrections show a significant amount of errors in the gauge records due to the  
22 windy and cold environment in the northern areas of Alaska and Yukon. Monthly corrections increase  
23 solid precipitation by 136% in January, 20% for July at the Barter Island in Alaska, and about 31% for  
24 January and 4% for July at the Yukon stations. Regression analyses of the monthly precipitation data  
25 show a stronger correlation for the warm months (mainly rainfall) than for cold month (mainly snowfall)  
26 between the station pairs, and small changes in the precipitation relationship due to the bias corrections.  
27 Double mass curves also indicate changes in the cumulative precipitation over the study periods. This  
28 change leads to a smaller and inverted precipitation gradient across the border, representing a significant  
29 modification in the precipitation pattern over the northern region. Overall, this study discovers significant  
30 inconsistency in the precipitation measurements across the US and Canada border. This discontinuity is  
31 greater for snowfall than for rainfall, as gauge snowfall observations have large errors in windy and cold  
32 conditions. This result will certainly impact regional, particularly cross borders, climate and hydrology  
33 investigations.

34

35 **Key words:** snowfall, national precipitation gauge, measurement errors, bias correction,  
36 precipitation gradient and distribution.

37

38 **1. Introduction**

39 It is known that discontinuities in precipitation measurements may exist across the national boundaries  
40 because of the different instruments and observation methods used (Nitu and Wong, 2010; Sanderson,  
41 1975; Sevruk and Klemm, 1989; Yang et al., 2001). For instance, the National Weather Service (NWS) 8-  
42 inch gauge is used for precipitation measurements in the United States (U.S.), and the Nipher snow gauge  
43 has been used in Canada for decades. Different instruments have also been used in various observational  
44 networks within the same country. In the synoptic network, the Type-B rain gauge and Nipher gauge are  
45 the standard manual instruments for rain and snow observations in Canada (Mekis and Vincent, 2011;  
46 Metcalfe and Goodison, 1993), and recently the Geonor automatic gauges have been installed

47 Instruments also change over time at most operational networks, resulting in significant breaks in data  
48 records. It has been realized that combination of regional precipitation records from different sources may  
49 result in inhomogeneous precipitation time series and can lead to incorrect spatial interpretations (Yang et  
50 al., 2005). Efforts have been reported to examine the precipitation discontinuity within a country  
51 (Groisman and Easterling, 1994; Sanderson, 1975). Leeper et al., (2015) found that the US COOP stations  
52 reported slightly more precipitation overall (1.5%) with network differences varying seasonally. The  
53 COOP gauges were sensitive to wind biases, particularly over winter when COOP observed (10%) less  
54 precipitation than the U.S. Climate Reference Network (USCRN). Conversely, wetting and evaporation  
55 losses, which dominate in summer, were sources of bias for USCRN. Mekis and Brown, (2010)  
56 developed adjustment method to link the Nipher gauge and ruler snowfall measurements over Canada  
57 Yang and Simonenko, (2013) compared the measurements among 6 Russian Tretyakov gauges at the  
58 Valdai experimental station, and reported the differences of less than 5-6% for the study period. These  
59 results are useful to determine the homogeneity of precipitation data collected by a standard gauge within  
60 the national and regional networks.

61 Many studies show that the national standard gauges, including the Canadian Nipher, and US 8-inch  
62 gauges, under measure precipitation especially for snowfall (Goodison, 1981; Goodison et al., 1998;  
63 Yang et al., 1995, 1998a, 1999). Compatibility analysis of precipitation measurements by various national  
64 gauges suggests little difference (less than 5%) for rainfall observations, but a significant discrepancy (up  
65 to 110%) for snowfall measurements (Yang et al., 2001). For instance, the experimental data from Valdai  
66 show that the U.S. 8-inch gauge at Valdai systematically measured 30-50% less snow and mixed  
67 precipitation than the Canadian Nipher gauge (Yang et al., 2001). This difference in national gauge catch  
68 has introduced a significant discontinuity in precipitation records between the U.S. and Canada borders,  
69 particularly in windy and cold regions. Differences in the snow measurements across the US and Canada  
70 border has also been noticed in other studies as a problem to produce gridded products and to develop  
71 precipitation input for basin hydrological investigations (Šeparović et al., 2013; Zhao et al., 2010).

72 Although Yang et al. (2001) compared the relative catch of many national standard gauges, little has been  
73 done to address the inconsistency of precipitation records across the national borders. This is an important  
74 issue, since most regional precipitation data and products have been compiled and derived from the  
75 combination of various data sources, assuming these data and observations were compatible across the  
76 borders and among the national observational networks. Simpson et al., (2005) studied temperature and  
77 precipitation distributions over the State of Alaska and west Yukon, and documented precipitation  
78 increase from north to south. They also report differences in mean monthly precipitation across the  
79 Alaska-Yukon border, i.e. about 5-15 mm in central-east Alaska and 15-40 mm in central-west Yukon.  
80 (Jones and Fahl, 1994) found a weak gradient in annual precipitation across the AK-YK border, including  
81 the headwaters of the Yukon River. Other studies also discuss precipitation distribution and changes over  
82 the arctic regions (Legates and Willmott, 1990; Serreze and Hurst, 2000; Yang et al., 2005).

83 The objective of this work is to examine the inconsistency in precipitation measurements across the  
84 border between Alaska and Yukon. We analyze both gauge-measured and bias-corrected monthly  
85 precipitation data at several climate stations across the border, and quantify the changes in precipitation

86 amounts and patterns due to the bias corrections. We also calculate the precipitation gradients across the  
87 border, and discuss precipitation distribution for the warm and cold seasons. The methods and results of  
88 this study are useful for cold region climate and hydrology investigations and applications.

## 89 **2. Study Area, Data and Methods**

90 The study areas include the northern and central regions of Alaska (AK) and Yukon (YK). We choose 5  
91 climate stations across the Yukon and Alaska border, which use the national standard gauges (NWS 8 in  
92 gauge and the Canadian Nipher gauge) for precipitation observations (Figure 1). These stations can be  
93 classified into 2 groups. The first group, 3 stations about 150 km apart, is the northern region along the  
94 coast of the Beaufort Sea; with the Barter Island station in Alaska and Komakuk and Shingle Point  
95 stations in Yukon. The second group is in the central part of the region; the Eagle station in Alaska and  
96 Dawson station in Yukon, about 130 km apart.

97 The three northern stations selected for this study are located north of the Brooks Range. The approximate  
98 distances to the mountain edge are 100 km for the Barter Island station, 90 km for Shingle Point station,  
99 and 150 km for the Komakuk station. Both stations in Yukon are along the shore line and the station in  
100 Alaska is an island site, very close to the coast line. The altitudes of the stations range from 7 to 49 m  
101 a.s.l. According to Manson and Solomon, (2007), the summer storm tracks are usually from the northwest  
102 coming from the open water in the Beaufort Sea and are the greatest contributor to annual precipitation.  
103 The storms are obstructed by the Brooks Range once moving inland. The weather patterns in the  
104 surrounding of the stations might be affected by the mountains, but the stations are not separated by the  
105 Brooks Range. Given this setting, it is expected to see little impact of mountain range on the precipitation  
106 process and distribution along the relatively flat coast line.

107 These stations have been operated by the NWS and Environment Canada (EC) since the early 1970's. The  
108 observations have been done according to the national standards of US and Canada. The detail  
109 information for these stations are given in Table 1, such as the location, period of measurement used for

110 this work, instrument types for precipitation observations, and a climate summary for yearly temperature,  
111 precipitation , and wind speed.

112 Yang et al. (2005) have developed a bias corrected daily precipitation dataset for the northern regions  
113 above 45°N. The source data are acquired from the National Centers for Environmental Information  
114 (NCEI), i.e. a global daily surface data archive for over 8,000 stations around the world  
115 (<https://www.ncdc.noaa.gov/data-access/quick-links#ghcn>). To focus on the high latitude regions, a  
116 subset of the global daily data, about 4,000 stations located north of 45°N with data records longer-than  
117 20 years during 1973-2003 has been created. Yang et al. (2005) applied a consistent procedure derived  
118 from the WMO Solid Precipitation Intercomparison (Goodison et al., 1998), using wind speed,  
119 temperature, and the precipitation as inputs (Yang et al., 1998b, 2005). They quantify the precipitation  
120 gauge measurement biases for the wind-induced undercatch, wetting losses, and trace amount of  
121 precipitation. For the US stations, wind data from the standard height was reduced to the gauge level of  
122 the NWS 8-in gauge (standard height is 1 m). Wind speeds and directions were measured at the Canadian  
123 climatic network; the same approach was applied to estimate the wind speed at the gauge height (standard  
124 height is 2 m) on precipitation days. The corrections were done only for those stations with wind  
125 observations. Unfortunately there are many stations in the US without wind information and this is a  
126 challenge to gauge bias corrections.

127 This study uses the updated (until 2013) monthly precipitation, temperature and wind speed data from  
128 Yang et al. (2005) for the selected AK and YK stations (Table 1). The selected data periods range from 7  
129 to 10 years for the stations that are considered long enough to examine precipitation patterns in these  
130 regions. Missing records affect regional climate data analyses. In this study, a threshold of 0°C of  
131 monthly temperature has been used to determine the cold and warm months for snow and rain. Mixed  
132 precipitation has not been classified separately. The frequency of missing values was calculated when the  
133 bias correction was made in Yang et al., (2005). For any month with less than 20 days (~30%) of  
134 measurements, it is excluded from data analysis. Statistical methods to compare the measured and

135 corrected monthly and yearly precipitation data across the selected border station pairs is used to analyze  
136 these data. It also carries out regression analysis on monthly precipitation records, and calculates the  
137 cumulative precipitation amounts to derive the Double Mass Curves (DMC) over the study period. The  
138 double mass curve (DMC) is a useful tool to evaluate the consistency of observation records over space  
139 and time (Searcy and Hardison, 1960). Some typical issues of observations that DMC can identify,  
140 include changes in the station location, and instruments or sensors. A reference station is needed for DMC  
141 analyses. In this study, the DMC has been applied without a reference station to mainly detect any shifts  
142 between the observed and corrected precipitation. Through the data analyses and comparisons with other  
143 studies, we document the spatial and temporal variations of bias corrections across the border stations.  
144 We also determine the precipitation gradients across the border, and examine the changes, due to the bias-  
145 corrections of the US and Canadian gauge data, in precipitation distributions on both seasonal and yearly  
146 time scales.

147

### 148 **3. Results**

149 Based on the analyses of the measured precipitation ( $P_m$ ) and corrected precipitation ( $P_c$ ) data, this section  
150 presents the results on the bias corrections of monthly and yearly precipitation for each station, regression  
151 and correlation of monthly precipitation data between the stations, and cumulative precipitation via the  
152 double mass curves for the warm (monthly temperature  $> 0^\circ\text{C}$ ) and cold seasons (monthly temperature  $<$   
153  $0^\circ\text{C}$ ).

#### 154 **3.1. Monthly data and corrections**

155 The monthly mean precipitation and bias corrections are illustrated in Figure 2 for the northern group  
156 during the corresponding observation period (Table 1). In Figure 2, the missing data percentages are also  
157 presented for each month. Barter Island had the lowest percentages of missing data, about 2% as a  
158 maximum monthly mean in December. The mean missing percentages for the Komakuk station was about

159 5% (in May), with the maximum month in July 1984 (16%). For Shingle Point, the mean missing values  
160 were 11% for both April and May, with the maximum (26%) in April 1979. Given the small percentages  
161 of missing records, its impact is insignificant on monthly mean and yearly precipitation calculations.  
162 Figure 2 shows that annual precipitation cycle was centered on August, with an approximate maximum  
163  $P_m$  around 40 to 80 mm between August and September. This maximum was coincident with the monthly  
164 mean maximum temperature in the area (around 10°C).

165 For the Barter Island station in AK, the corrections were variable through the months. The monthly  
166 corrections increased the  $P_m$  amount by 3-31 mm for snow to 4-9 mm for rain. The relative increases were  
167 59-136% for snow and 20-41% for rain, with a monthly mean of 9 mm (or 76%). The relative changes  
168 were usually large for months with low  $P_m$  and small for months with high precipitation. In other words,  
169 the monthly correction amounts do not always matched with the percentage changes, i.e. a small  
170 correction in a dry month can have a large percentage change.

171 It is important to note that gauge measurements at Barter showed the maximum precipitation in August,  
172 but the peak shifted to October due to the corrections; i.e. the mean monthly  $P_c$  in October were 98%  
173 (about 32mm) more than the  $P_m$  (Figure 2). Closer examination of the monthly precipitation time series  
174 for Barter Island (Figure 3) indicated that, for most of the years, October was the most significant  
175 contributor to the total annual (23% for  $P_m$  and 22% for  $P_c$ ). However, there were some years in the study  
176 period with the maximum  $P_m$  in other months; for example, the highest  $P_m$  in 1982 was in September, as  
177 documented by Yang et al., (1998b). Climate data and analyses showed the highest wind speed (4.5 m/s)  
178 and cold temperature (about -9°C) for October, indicating higher undercatch by the US standard gauge for  
179 snowfall. On the other hand, the wind speed showed the minimum values in July and August (3.3 m/s),  
180 coincident with the highest temperatures (4.6 and 4 °C) (Figure 2). Due to the combination of warm  
181 temperatures and low wind speeds, the corrections for summer months were the lowest at this station (20-  
182 27%).

183 For the Komakuk Beach station in Yukon, the corrections increased the precipitation by 0.7-5.5 mm (or  
184 14%-34%) for snow and 1-2.6 mm (4%-10%) for rain, with a total monthly mean change of 2.6 mm  
185 (14%) (Figure 2). The monthly maximum precipitation was in August, i.e. 48mm and 50mm,  
186 respectively, for the  $P_m$  and  $P_c$ . The monthly minimum precipitation was in March, i.e.  $P_m = 4.2$  mm and  
187  $P_c = 5$  mm. For this station, the extremes remained in the same month after the bias corrections. The wind  
188 speed had the minimum value in Aug. (3.1 m/s) and Sept. (3.2 m/s), and max in Dec. (4.3m/s) and Jan  
189 (4.7m/s). The temperatures were highest in July (6.9°C) and Aug. (5.8°C), and lowest in Feb and Mar (-25  
190 °C). Given this climate condition, the corrections were lower in the summer months (mean of 6%) and  
191 higher in winter (mean of 23%).

192 The monthly corrections for the Shingle Point station in Yukon ranged from 1-7.6 mm (3%-15%) for rain  
193 to 1-8.2 mm (14%-28%) for snow, with the monthly mean correction of 4.2 mm (14%). The maximum  
194 precipitation was in Aug., about 73-76 mm (or 20% of the annual total) (Figure 2). The minimum  
195 precipitation was in March with 9.8 mm for  $P_m$ ; and 11 mm for  $P_c$ . The monthly wind speeds were  
196 generally higher in winter and lower in summer, with the maximum in Feb. (4 m/s) and minimum in May  
197 (2.7 m/s). The temperatures had a common annual cycle with the maximum in July (11°C) and the  
198 minimum in Feb. (-24.3°C). Because of the higher wind speeds and cold temperatures in the cold months,  
199 the corrections were greater for the winter season.

200 It was necessary to compare the correction result across the border in order to quantify the effect of biases  
201 in gauge observations on precipitation analyses, such as precipitation distribution and seasonal patterns.  
202 The mean snowfall corrections were about 96% for Barter Island in Alaska and around 22% for both  
203 Shingle Point and Komakuk stations in Yukon; while the rainfall corrections were approximately 32% for  
204 Barter and 7% for the two Yukon stations. Bias corrections also demonstrated a clear shift in the  
205 maximum precipitation timing for the Barter Island, but no change for the Yukon stations. This  
206 remarkable contrast across the border was caused mainly by the difference in gauge types and their catch  
207 efficiency. Many experimental studies have shown that the Canadian Nipher snow gauge catches more

208 snowfall relative to the US gauge (Goodison et al., 1998; Yang et al., 1998b). For instance, the mean  
209 catch ratios for snowfall were about 40% and 85% for 4 m/s wind speed, respectively, for the NWS 8-in  
210 unshielded and Nipher gauges (Figure 4) (Yang et al., 1998b).

211 For the central group, the maximum and minimum  $P_m$  were in July and March for the Eagle station  
212 (Figure 5). The corrections did not modify the timings of maximum and minimum amounts; they  
213 remained in July for the maximum ( $P_m=67$  mm and  $P_c=70$  mm), and in March for the minimum ( $P_m=3$   
214 mm and  $P_c=4$  mm) precipitation. The correction increased the precipitation by 0.6-1.8 mm (8%-22%) for  
215 snow and 1-3 mm (5%-10%) for rain, with a monthly mean correction of 1.7 mm (12%). The annual  
216 temperature cycle for Eagle showed warmer temperatures relative to the northern station, with the  
217 maximum of 16.2°C and above 0°C during April to mid-October. Eagle had lower wind speeds around 1  
218 m/s (Figure 5).

219 For Dawson station, precipitation was more homogeneous throughout months; varying from 10 mm to 50  
220 mm in October and June, respectively. Another relative maximum occurs in January with  $P_m=38$  mm  
221 (Figure 5). The precipitation correction was small and fluctuated from 0.3 to 1 mm (or 2%-4%) for snow  
222 and 0.4-1.3 mm (3%-4%) for rain. This small correction was due to the lower undercatch correction for  
223 the Nipher gauge, besides the warmer temperatures and lighter winds. The temperature annual amplitude  
224 was between 16°C in July and -25°C in January, with temperatures above 0°C from April to September.  
225 Wind speeds showed a clear annual cycle with the maximum in May (1.6 m/s) and lighter winds in winter  
226 months, with the minimum in January (0.4 m/s).

227 The temperature and wind conditions were similar between the Eagle and Dawson stations, with mean  
228 temperature around 1°C and wind speed of 1m/s. The missing data percentages were also similar for  
229 Eagle and Dawson stations; less than 3% for most months, with the maximum of 10% in May 2006 for  
230 Eagle and 20% in September 2009 for Dawson. The bias corrections were quite different, with the mean  
231 corrections of 16% for snow and 7% for rain at Eagle, and about 2 % and 3% for both rain and snow at

232 Dawson. Overall, the correction was four times greater at Eagle than that at Dawson. This discrepancy  
233 reflects again the catch difference between the US and Canadian standard gauges.

234 In order to understand the effect of precipitation bias corrections on regional climate around the AK-YK  
235 border, it was useful to examine and compare the temperature and precipitation features between the  
236 northern and central regions. The monthly mean temperature threshold of 0°C did not occur exactly at the  
237 same time among the 2 groups; the warm months (above 0°C) were between June and September in the  
238 north group and between April and September in the central group. Although both regions had similar  
239 mean minimum temperatures, around -24°C and -27°C, the maximum temperature was considerably  
240 lower in the north part, with the average of 8°C in the north group vs. 16°C for the central region.  
241 Additionally the monthly mean wind speed was higher for the northern region, 4 m/s vs. 1 m/s.  
242 Therefore, because of the colder temperatures and higher winds in the northern region, the bias  
243 corrections were higher in the north relative to the central region.

### 244 **3.2. Yearly data and corrections**

245 The annual  $P_m$  and  $P_c$  time series for 11 years during 1978-1988 in the northern group is presented in  
246 Figure 6. There was almost no missing data for the whole period, except 3% for 1978. At the Barter  
247 Island station in Alaska, the yearly  $P_m$  ranged from 114 mm to 211 mm, with the long-term mean of 155  
248 mm. The mean annual corrections ranged from 67 to 138 mm, with a long-term mean of 101 mm (or  
249 65%). The  $P_c$  records varied from 181mm to 343 mm. The maximum precipitation was in 1985 for both  
250  $P_m$  and  $P_c$  (211 mm and 343 mm, respectively). The minimum precipitation was in 1983 for the  $P_m$  and  $P_c$   
251 (114 mm and 181 mm, respectively).

252 For Komakuk Beach station in Yukon, the  $P_m$  ranged from 103 mm to 306 mm, with the missing data  
253 between 0 and to 7% among the years. The bias corrections increased the precipitation by 13 mm to 45  
254 mm (or 8-19%). The long-term mean was about 194 mm for  $P_m$  and 220 mm with the corrections. The

255 maximum precipitation occurred in 1981, 306 mm and 347 mm for  $P_m$  and  $P_c$ , respectively. The minimum  
256 precipitation was in 1988 for both the  $P_m$  and  $P_c$ , 103 mm and 123 mm, respectively.

257 For Shingle Point station in Yukon, yearly  $P_m$  varied from 126 mm to 551 mm and the  $P_c$  ranges from 138  
258 to 638 mm. The mean annual total precipitation was about 302 mm for  $P_m$  and 341 mm after the  
259 corrections (change of 13%). The high and low extreme years were 1981 ( $P_m = 551$  mm,  $P_c = 638$  mm),  
260 and 1988 ( $P_m = 126$  mm,  $P_c = 138$  mm). Shingle station had missing data from 2% in 1983 to 10% in 1979.

261 Figure 7 displays the mean annual precipitation in cold and warm seasons for the northern group. The  
262 gauge measurements showed annual values from 155 mm at Barter Island, 194 mm at Komakuk to 302  
263 mm at Shingle Point, i.e. a strong precipitation increased from the west to the east, particularly between  
264 Komakuk Beach and Shingle Point. However, the corrected data ( $P_c$ ) showed a different pattern (Figure  
265 7), i.e. higher precipitation at Barter than Komakuk, so the gradient across the border changed the sign  
266 and magnitude. This change was caused mainly by the high correction at the Barter station, particularly  
267 for snowfall data during the cold months (Fig. 2).

268 For the central group, the annual results are shown for 8 years (2006-2013) in Figure 8. The  $P_m$  ranged  
269 from 66 to 391 mm at the Eagle, and the bias corrections were 5-27 mm, correspondingly, which on  
270 average increase the total precipitation by 7%. While at Dawson, the  $P_m$  ranged from 158 to 333 mm, and  
271 the adjustments were from 4 mm to 10 mm, with an average increase in yearly precipitation by 3%. The  
272 gauge data showed a slight increase (12 mm) of mean precipitation from west to the east, i.e. slightly  
273 higher  $P$  in Yukon relative to Alaska. This result is consistent with other studies (Simpson et al., 2002,  
274 2005). The corrected data, on the other hand, suggest a smaller gradient (1 mm) across the border (Figure  
275 9). This change was mainly due to the higher corrections for the US 8-inch gauge at Eagle.

276 Similar to the monthly results, the northern stations exhibited higher yearly corrections for snowfall and  
277 rainfall measurements relative to the central group. This was because of higher winds in the northern

278 stations, i.e. yearly mean wind speeds of 3.8 m/s in the north group and 1 m/s in the central group. This  
279 windy and snowy environment in the north produced higher wind-loss for the snowfall measurements by  
280 the gauges, which was the largest errors in precipitation records in the high latitudes (Benning and Yang,  
281 2005; Yang and Ohata, 2001; Yang et al., 1998b). It is important to note that gauge measured and bias  
282 corrected data showed different pattern in seasonal and yearly precipitation in the northern region. In  
283 other words, bias corrections of gauge measurements alter the precipitation gradient in the northern areas;  
284 this change was mainly due to the difference in the catch efficiency between the US and Canadian  
285 standard gauges. The corrections for the US gauge snow measurements were much higher than the  
286 Canadian gauge, particularly in the cold and windy coastal regions.

### 287 **3.3. Regression analysis of monthly data**

288 The scatter plots of corresponding monthly precipitation for the two stations across the border and  
289 between the two Yukon stations in Canada are illustrated in Figure 10. For the cold season (Figure 10.A),  
290 the gauge data showed more snowfall at Barter for most years. Regression analysis suggested a weak  
291 relationship, with  $R^2=0.34$ . The corrected data showed a similar relationship, but a shift in the regression  
292 line, indicating a greater precipitation difference over the cold season across the border. For the warm  
293 season (Figure 10.B), the gauge data showed higher precipitation at the Komakuk station, and the  
294 regression suggested a much stronger relationship. The corrected data revealed a closer relationship  
295 between these two stations, proposing a smaller gradient for the warm months.

296 The scatter plot between the two stations in the Yukon Territory showed higher precipitation at Shingle  
297 point for both cold and warm seasons. It also gave another point of view about the effect of the correction  
298 in this area. Relative to the cold months (Figure 10.C), the corrections were smaller for the warm months  
299 (Figure 10.D), and correlation improved ( $R^2=0.72-0.76$ ). However, the relationship did not change much  
300 in both cases between the measured and corrected data. This was because very small amount of  
301 corrections for the lower wind conditions and higher catch efficiency of the Canadian Nipher gauge.

302 For the central group, the scatter plot between Eagle and Dawson stations illustrated a clear difference in  
303 precipitation amount for the cold and warm months (Figure 10.E-F). The cold months showed more  
304 precipitation at Dawson, particularly for the wettest events, while Eagle did not show any comparable  
305 amount. The correlation was weak, and insignificant ( $R^2 = 0.13$ ). The shift in the fit line between  
306 measured and corrected data was also very small. The warm months showed low precipitation at Dawson;  
307 a different pattern from the cold months. The regression was better,  $R^2 = 0.59$  with a smaller shift due to  
308 the corrections.

309 Overall, we obtained consistent results among the Alaska and Yukon stations. The correlations were  
310 higher in warm months ( $R^2 = 0.58$  to  $0.76$ ) and lower for the cold season ( $R^2$  between  $0.13$  and  $0.52$ ). This  
311 result may suggest that the rainfall was more homogeneous over the regions in summer, and greater  
312 difficulty and errors in snowfall measurements during the cold months.

#### 313 **3.4. Cumulative precipitation via double mass curves (DMC)**

314 The DMC plot for Barter Island and Komakuk Beach showed more  $P_m$  at Komakuk than Barter (Figure  
315 11.A). The bias corrections led to a shift of the relationship with a significant increase in the total  
316 precipitation amount at Barter. Relatively, the total cumulative precipitation for Barter Island increased by  
317 65% after the correction and by 14% at Komakuk. The difference between the two stations at the last  
318 cumulative point (December 1988) is 426 mm for  $P_m$ , and 393 mm for  $P_c$ . This shift represented a  
319 modification in the precipitation difference between these stations, i.e. a change in the gradient's direction  
320 (Figure 7).

321 The comparison of cumulative precipitation values between Shingle Point and Komakuk, both in Yukon,  
322 is illustrated in Figure 11.B. Shingle Point showed more cumulative precipitation at the end of the period  
323 ( $P_m = 3322$  mm vs.  $P_m = 2115$  mm for Komakuk). Although the relationship was more homogeneous  
324 between these stations, there was a break in the records around 1300 mm for Komakuk, maybe associated  
325 with changes in instruments or sensors. Examination of the station history and information revealed an

326 anemometer issue around the critical time that was fixed by August 1980. This may affect wind data and  
327 thus the corrected precipitation values. Both stations showed increases in total cumulative precipitation by  
328 13%.

329 The central stations showed a greater amount of  $P_m$  in Dawson (2065 mm) than in Eagle (1973 mm) over  
330 the study period. Bias corrections changed the total precipitation by 3% and 7% for Dawson and Eagle,  
331 respectively, resulting in a shift in the DMC (Figure 11.C), particularly for the last period of time, to 2123  
332 mm in Dawson and to 2116 mm in Eagle. This shift also represented a slightly smaller precipitation  
333 difference between the two stations. During the 8 years, the cumulative difference decreased from 92 mm  
334 to 7.3 mm.

335 In summary, the DMC for measured and corrected precipitation showed that the main change was due to  
336 the difference in their corrections (Figure 11); the north stations showed a greater change compared with  
337 the central group. The  $P_c$  showed in all the cases a smaller precipitation difference between the two  
338 countries. This smaller difference led to a decrease in the precipitation gradient across the border. This  
339 result implies that existing precipitation climate maps and information derived from gauge measurement  
340 without bias corrections may over-estimate the precipitation gradient in these regions. This  
341 overestimation will affect regional climate and hydrology analyses.

342

#### 343 **4. Summary and Discussion**

344 This study documents and quantifies the inconsistency in precipitation measurements in the northern and  
345 central regions of Alaska/Yukon, with a focus on station pairs across US-Canada border. The monthly  
346 bias corrections show large errors in the gauge records due to the windy and cold environment in the  
347 northern areas of Alaska and Yukon. The corrections for gauge undercatch increase the snowfall by 136%  
348 in January for Barter Island station in Alaska. For the Yukon stations, the increase is about 31% in  
349 January and 4% in July. These represent an annual mean loss of 81 mm (101%) in snowfall and 20 mm

350 (29%) of rain at Barter, while at Shingle Point and Komakuk Beach in Yukon the corrections are, on  
351 average, about 25 mm (21%) for snow and 8 mm (6%) for rain. For Eagle (AK) and Dawson (YK)  
352 stations in the central region, the bias corrections are small. The monthly corrections range from 2% to  
353 22% in winter and from 3% to 10% on summer months.

354 On the annual scale, Barter Island station in AK shows a yearly mean correction around 65%, five times  
355 greater than the correction at Shingle Point and Komakuk Beach (13% and 14%) in Canada. In the central  
356 region, Eagle station shows an increase by 7%, meanwhile for Dawson the increase is only 3%. Thus, the  
357 bias correction is twice for Alaska compared to the Yukon stations. Relative to the northern region, these  
358 corrections are small mainly due to warmer temperatures and lower winds in the central region. These  
359 results clearly demonstrate that bias corrections may affect the spatial distribution of precipitation across  
360 the border.

361 Regression analyses of the monthly data show small changes in the relationship due to the bias  
362 corrections. The most evident change in the regression is between Barter Island and Komakuk Beach for  
363 both warm and cold seasons. The rest of the scatter plots, for the Komakuk Beach-Shingle Point and  
364 Eagle-Dawson, do not show any appreciable change as the result of the bias corrections. There is a  
365 stronger precipitation correlation for the warm months (mainly rainfall) than for the cold month (mainly  
366 snowfall) for all the station pairs. The cold months seem to have greater precipitation variability across  
367 the regions.

368 The double mass curve analyses demonstrate a significant change in the precipitation accumulation and  
369 difference between the two stations across the AK-YK border for the northern region, little changes for  
370 the two stations in Yukon, and a smaller change in the central group. These changes, caused by gauge  
371 catch efficiency, alters the precipitation difference, resulting in a smaller and inverted precipitation  
372 gradient across the border in the northern region. The double mass curve (DMC) is a useful tool to  
373 evaluate the consistency of observation records over space and time (Searcy and Hardison, 1960).

374 Although in this work the DMC has not been constructed against a reference station, the results clearly  
375 show some breaks on the slope and gaps in the curves, indicating changes in precipitation relationship  
376 across the border that could be caused by any of the two stations. This information provides the timing  
377 when significant changes occurred in the precipitation regime. Detail metadata and information for the  
378 stations/networks are necessary to understand the changes in precipitation observations and to improve  
379 the homogenization of the precipitation records over the high latitudes.

380 This study shows similar monthly  $P_m$  across the north border region and higher  $P_m$  in Yukon than Alaska  
381 over the central region. This result is similar to other studies (Serreze and Hurst, 2000; Simpson et al.,  
382 2005). After the bias corrections, precipitation patterns across the border changed, i.e. higher precipitation  
383 in Barter than Komakuk, in other words, an inverted gradient across the borderline. Over the central  
384 region, the measured mean annual precipitation is slightly higher in Yukon than Alaska, which is also  
385 consistent with Simpson et al., (2002) and (2005). Our results suggest that the gradient between the  
386 central pair of stations becomes smaller after the bias correction. This discrepancy should be taken into  
387 account when using the precipitation data across the national borders for regional climate and hydrology  
388 investigations.

389 Missing data may affect regional precipitation analyses. In this study, we calculated the missing data  
390 percentages for all stations during the corresponding study periods, and set up a threshold of 30% to  
391 exclude those months with higher missing values from monthly precipitation calculations. We compared  
392 the precipitation amounts with and without the application of the threshold. The results do not show any  
393 significant changes in the differences of gauge measured annual mean precipitation across the border,  
394 although this filter affected annual precipitation in certain years. For instance, the northern station pair  
395 (Barter and Komakuk stations) has missing value of 32% on July 1987. Calculations of yearly  
396 precipitation for 1987 with and without this month show 16% and 10% difference at Komakuk and Barter  
397 Island stations, respectively. Over the study period of 11 years, the annual mean bias correction  
398 percentages remain the same (65% in Barter and 13% in Komakuk) with or without the missing months.

399 The mean annual decrease in bias correction amounts after the consideration of missing data is about 1-  
400 3% in the northern region. This analysis suggests that the effect of missing data for our study is not  
401 significant, particularly with the application of 30% missing threshold. More efforts are needed to further  
402 examine the issues of missing records in climate analyses.

403 Classification of precipitation types is the first step for the bias corrections of gauge records. It is also  
404 important for climate change analyses over the cold regions. Leeper et al., (2015), in comparison of US  
405 CRN with the CO-OP station network precipitation measurements, averaged the USCRN hourly  
406 temperatures data during precipitation periods into an event mean and used it to group precipitation  
407 events into warm (mean temperature  $> 5^{\circ}\text{C}$ ), near-freezing (mean temperature between  $0^{\circ}\text{C}$  and  $5^{\circ}\text{C}$ ), and  
408 freezing (mean temperature  $< 0^{\circ}\text{C}$ ) conditions. Yang et al., (2005) used the daily mean air temperature to  
409 estimate precipitation types (snow, mixed, and rain) when this information is not available for the  
410 northern regions. In this study, monthly mean temperatures have been used to determine the warm months  
411 (mainly for rain) and cold months (mainly for snow). Mixed precipitation has not been classified  
412 separately. This approach is reasonable for our analysis to focus on the inconsistency in the monthly and  
413 yearly  $P_m$  records across the border. Data collections and analyses on shorter timescales, such as daily or  
414 hourly steps, are expected to produce better results, since temperatures vary throughout the days in a  
415 month, particularly in the spring and fall seasons. Automatic sensors will also be important to decide  
416 precipitation types at the operational and research networks.

417 The bias-corrected precipitation dataset developed by Yang et al., (2005) has been used for this analysis.  
418 The corrections have been done systematically on a daily time scale that affects the daily  $P_m$  time series.  
419 This analysis focuses on the results of monthly and yearly precipitation data and quantifies the changes in  
420 precipitation pattern across the AK-YK border. Careful analyses of available daily measured  $P_m$  and  
421 corrected  $P_c$  data are necessary, since in the northern regions with low precipitation in winter, the bias  
422 corrections can easily increase the daily  $P_m$  by a factor of up to 4-5 (Benning and Yang, 2005; Kane and

423 Stuefer, 2015; Yang et al., 1998b, 2005). This means that extreme precipitation events have been very  
424 likely and seriously underestimated by using the gauge records without any bias corrections. The  
425 consequence is certainly significant for climate regime and change investigations. To fill this knowledge  
426 gap, our efforts are underway to examine the daily corrections, particularly on the windy and heavy  
427 precipitation days, and to document the possible underestimation of precipitation extremes over the large  
428 northern regions.

429 Automation of the meteorological observation networks and instruments has been a trend over the past  
430 few decades around the world, including both the developed and developing nations. There is a large  
431 variety of automatic gauges currently used for precipitation measurements at the national networks (Nitu  
432 and Wong, 2010). These gauges differ in the measuring system, orifice area, capacity, sensitivity, and  
433 configuration. The variation in automatic gauges is much greater relative to the manual standard gauges  
434 (Goodison et al., 1998; Sevruk and Klemm, 1989). As demonstrated by (Yang et al., 2001) and this study,  
435 the use of different instruments and configurations significantly affect the accuracy and consistency of  
436 regional precipitation data. Fortunately, the Geonor gauge has recently been chosen and used at both the  
437 US Climate Reference Network (USCRN) and the Surface Weather and Climate Network (SWCN) in  
438 Canada. This may reduce the inconsistency in precipitation measurements across US and Canada borders,  
439 although the double and single Alter wind shields have been installed with the Geonor gauges in US and  
440 Canada, respectively.

441 Finally, it is important to emphasize that automatic gauges also significantly under catch snowfall (Wolff  
442 et al., 2015) and bias corrections are necessary in order to obtain reliable precipitation data for the cold  
443 regions and seasons. The WMO SPICE project aims to examine the performance of automatic gauges and  
444 instruments for snowfall observations in various climate conditions. It has tested many different automatic  
445 gauges, including the Geonor gauge, at more than 20 field sites around the globe (Nitu et al., 2012;  
446 Rasmussen et al., 2012; Wolff et al., 2015). The results of this project will be very useful to improve

447 precipitation data quality and regional climate analyses, including the border regions between US and  
448 Canada.

449

450       **Acknowledgments**

451       The authors gratefully acknowledge the support from the Global Institute of Water Security at University  
452       of Saskatchewan and Environment Canada.

453

454       **References**

455       Benning, J. and Yang, D.: Adjustment of Daily Precipitation Data at Barrow and Nome Alaska for 1995–  
456       2001,       Arctic,       Antarct.       Alp.       Res.,       37(3),       276–283,       doi:10.1657/1523-  
457       0430(2005)037[0276:AODPDA]2.0.CO;2, 2005.

458       Goodison, B. E.: Compatibility of Canadian snowfall and snow cover data, *Water Resour. Res.*, 17(4),  
459       893, doi:10.1029/WR017i004p00893, 1981.

460       Goodison, B. E., Louie, P. Y. T. and Yang, D.: Report No . 67 WMO SOLID PRECIPITATION  
461       MEASUREMENT INTERCOMPARISON., 1998.

462       Groisman, P. Y. and Easterling, D. R.: Variability and Trends of Total Precipitation and Snowfall over  
463       the       United       States       and       Canada,       *J.       Clim.*,       7,       184–205,       doi:10.1175/1520-  
464       0442(1994)007<0184:VATOTP>2.0.CO;2, 1994.

465       Jones, S. H. and Fahl, C. B.: Magnitude and Frequency of Floods in Alaska and Conterminous Basins of  
466       Canada, Anchorage, Alaska., 1994.

467       Kane, D. L. and Stuefer, S. L.: Reflecting on the status of precipitation data collection in Alaska: a case  
468       study, *Hydrol. Res.*, 46(4), 478, doi:10.2166/nh.2014.023, 2015.

469 Leeper, R. D., Rennie, J. and Palecki, M. A.: Observational Perspectives from U.S. Climate Reference  
470 Network (USCRN) and Cooperative Observer Program (COOP) Network: Temperature and Precipitation  
471 Comparison., *J. Atmos. Ocean. Technol.*, 32(4), 703–721, 2015.

472 Legates, D. R. and Willmott, C. J.: Mean seasonal and spatial variability in gauge-corrected, global  
473 precipitation, *Int. J. Climatol.*, 10(2), 111–127, doi:10.1002/joc.3370100202, 1990.

474 Manson, G. K. and Solomon, S. M.: Past and future forcing of Beaufort Sea coastal change, *Atmosphere-*  
475 *Ocean*, 45(2), 107–122, doi:10.3137/ao.450204, 2007.

476 Mekis, É. and Vincent, L. a.: An Overview of the Second Generation Adjusted Daily Precipitation  
477 Dataset for Trend Analysis in Canada, *Atmosphere-Ocean*, 49(2), 163–177,  
478 doi:10.1080/07055900.2011.583910, 2011.

479 Metcalfe, J. R. and Goodison, B. E.: Correction of Canadian winter precipitation data, in *Proc. Eighth*  
480 *Symp. on Meteorological Observations and Instrumentation*, pp. 338–343, Amer. Meteor. Soc., Anaheim,  
481 CA., 1993.

482 Nitu, R. and Wong, K.: CIMO Survey on National Summaries of Methods and Instruments Related to  
483 Solid Precipitation Measurement at Automatic Weather Stations-Preliminary., 2010.

484 Nitu, R., Rasmussen, R., Baker, B., Lanzinger, E., Joe, P., Yang, D., Smith, C., Roulet, Y. A., Goodison,  
485 B., Liang, H., Sabatini, F., Kochendorfer, J., Wolff, M., Hendrikx, J., Vuerich, E., Lanza, L., Aulamo, O.  
486 and Vuglinsky, V.: WMP intercomparison of instruments and methods for the measurement of solid  
487 precipitation and snow on the ground : Organization of the Experiment., 2012.

488 Rasmussen, R., Baker, B., Kochendorfer, J., Meyers, T., Landolt, S., Fischer, A. P., Black, J., Thériault, J.  
489 M., Kucera, P., Gochis, D., Smith, C., Nitu, R., Hall, M., Ikeda, K. and Gutmann, E.: How Well Are We

490 Measuring Snow: The NOAA/FAA/NCAR Winter Precipitation Test Bed, *Bull. Am. Meteorol. Soc.*,  
491 93(6), 811–829, doi:10.1175/BAMS-D-11-00052.1, 2012.

492 Sanderson, M.: NOTES AND CORRESPONDENCE: A comparison of Canadian and United States  
493 Standard Methods of Measuring Precipitation, *J. Appl. Meteorol.*, 14, 1197–1199, 1975.

494 Searcy, J. and Hardison, C.: Double-Mass Curves, United States Department of the Interior, Washington  
495 DC., 1960.

496 Šeparović, L., Alexandru, A., Laprise, R., Martynov, A., Sushama, L., Winger, K., Tete, K. and Valin,  
497 M.: Present climate and climate change over North America as simulated by the fifth-generation  
498 Canadian regional climate model., 2013.

499 Serreze, M. C. and Hurst, C. M.: Representation of mean arctic precipitation from NCEP-NCAR and  
500 ERA reanalyses, *J. Clim.*, 13(1), 182–201, doi:10.1175/1520-0442(2000)013<0182:ROMAPF>2.0.CO;2,  
501 2000.

502 Sevruk, B. and Klemm, S.: Types of standard precipitation gauges, in *Proceedings of International*  
503 *Workshop on Precipitation Measurement, WMO/IAHS/ETH*, vol. 227236, St. Moritz, Switzerland., 1989.

504 Simpson, J. J., Hufford, G. L., Fleming, M. D., Berg, J. S. and Ashton, J. B.: Long-term climate patterns  
505 in Alaskan surface temperature and precipitation and their biological consequences, *IEEE Trans. Geosci.*  
506 *Remote Sens.*, 40(5), 1164–1184, doi:10.1109/TGRS.2002.1010902, 2002.

507 Simpson, J. J., Hufford, G. L., Daly, C., Berg, J. S. and Fleming, M. D.: Comparing maps of mean  
508 monthly surface temperature and precipitation for Alaska and adjacent areas of Canada produced by two  
509 different methods, *Arctic*, 58(2), 137–161, 2005.

510 Wolff, M. A., Isaksen, K., Petersen-Øverleir, A., Ødemark, K., Reitan, T. and Brækkan, R.: Derivation of  
511 a new continuous adjustment function for correcting wind-induced loss of solid precipitation: results of a  
512 Norwegian field study, *Hydrol. Earth Syst. Sci.*, 19(2), 951–967, doi:10.5194/hess-19-951-2015, 2015.

513 Yang, D. and Ohata, T.: A Bias-Corrected Siberian Regional Precipitation Climatology, *J.*  
514 *Hydrometeorol.*, 2(2), 122–139, doi:10.1175/1525-7541(2001)002<0122:ABCSRP>2.0.CO;2, 2001.

515 Yang, D. and Simonenko, A.: Comparison of Winter Precipitation Measurements by Six Tretyakov  
516 Gauges at the Valdai Experimental Site, *Atmosphere-Ocean*, 52(1), 39–53,  
517 doi:10.1080/07055900.2013.865156, 2013.

518 Yang, D., Goodison, B. E., Metcalfe, J. R., Golubev, V. S., Elomaa, E., Gunther, T., Bates, R., Pangburn,  
519 T., Hanson, C. L., Emerson, D., Copaciu, V. and Milkovic, J.: Accuracy of tretyakov precipitation gauge:  
520 Result of wmo intercomparison, *Hydrol. Process.*, 9(8), 877–895, doi:10.1002/hyp.3360090805, 1995.

521 Yang, D., Goodison, B. E., Metcalfe, J. R., Golubev, V. S., Bates, R., Pangburn, T. and Hanson, C. L.:  
522 Accuracy of NWS 8" standard nonrecording precipitation gauge: Results and application of WMO  
523 intercomparison, *J. Atmos. Ocean. Technol.*, 15(1), 54–68, doi:10.1175/1520-  
524 0426(1998)015<0054:AONSNP>2.0.CO;2, 1998a.

525 Yang, D., Goodison, B. E., Ishida, S. and Benson, C. S.: Adjustment of daily precipitation data at 10  
526 climate stations in Alaska: Application of World Meteorological Organization intercomparison results,  
527 *Water Resour. Res.*, 34(2), 241–256, doi:10.1029/97WR02681, 1998b.

528 Yang, D., Ishida, S., Goodison, B. E. and Gunther, T.: Bias correction of daily precipitation  
529 measurements for Greenland, *J. Geophys. Res.*, 104(D6), 6171, doi:10.1029/1998JD200110, 1999.

530 Yang, D., Goodison, B., Metcalfe, J., Louie, P., Elomaa, E., Hanson, C., Golubev, V., Gunther, T.,  
531 Milkovic, J. and Lapin, M.: Compatibility evaluation of national precipitation gage measurements, J.  
532 Geophys. Res., 106(D2), 1481, doi:10.1029/2000JD900612, 2001.

533 Yang, D., Kane, D., Zhang, Z., Legates, D. and Goodison, B.: Bias corrections of long-term (1973–2004)  
534 daily precipitation data over the northern regions, Geophys. Res. Lett., 32(19), L19501,  
535 doi:10.1029/2005GL024057, 2005.

536 Zhao, K., Stadnyk, T., Koenig, K. and Crawford, J.: Better Precipitation Product over the Red River  
537 Basin., 2010.

538


539  
540

Table 1: Station information and climate summary

ID	Country	Station Name	Location			Data Period		Measurement Device	Annual Means				
			Latitude (°)	Longitude (°)	Altitude (m)	Start	End	Precipitation gauge	Precipitation (mm)	Missing Precipitation data %	Minimum Temperature (°C)	Maximum Temperature (°C)	Wind Speed (m/s)
700860	US	BARTER IS WSO AP	70.13	-143.63	11	1978	1988	US-8 inch Unshielded	155	0.3	-27.1	4.6	4.0
719690	CA	KOMAKUK BEACH ARPT	69.58	-140.18	7	1978	1988	Nipher Type B gauge	191.8	2.9	-27.5	7.4	3.9
719680	CA	SHINGLE POINT ARPT	68.95	-137.21	49	1978	1988	Nipher Type B gauge	302	6	-26.6	10.6	3.4
701975	US	EAGLE	64.78	-141.16	268	2006	2013	US-8 inch Unshielded	247	0.2	-22.7	15.5	0.9
719660	CA	DAWSON AIRPORT	64.05	-139.13	369	2006	2013	Nipher Type B gauge	258	0.6	-25.8	15.9	1

541

542









543

544

545 Figure 1: Study areas and locations of selected climate stations, and photos of the national standard gauges, NWS 8 in gauge (left) and the Nipher snow  
546 gauge (right), respectively, for USA and Canada.

547



548

549

550 Figure 2: Monthly mean precipitation at 3 stations during 1977 - 1988 (upper panels) and corresponding monthly mean wind speed and air temperature

551 (bottom panels). Shadows represent the 95% confidence interval for the temperature and wind speed. The percentages above the bars represent the


552 missing data for the corresponding time step. The bold percentage is the monthly mean and the one in the parenthesis is the maximum missing value in

553

the study period.

554

555




556

557





558 Figure 3: Monthly precipitation records at the Barter station during 1978-1988. The months with more than 50 mm (black line) are labeled.

559



560  
 561  
 562  
 563  
 564

Figure 4: Comparison of the catch ratio of snowfall as a function of wind speed at gauge height for the Alter-shielded or unshielded NWS 8-inc standard gauge and the Canadian Nipher snow gauge. DFIR is the Double Fence Intercomparison Reference (Yang et al. 1998)



565




566

567

568

569


Figure 5: Monthly mean precipitation at 2 stations during 2006 - 2013 (upper panels) and corresponding monthly mean wind speed and air temperature (bottom panels). Shadows represent the 95% confidence interval for the temperature and wind speed. The percentages above the bars represent the missing data for the corresponding time step. The bold percentage is the monthly mean and the one in the parenthesis is the maximum missing value in the study period.



570

571 Figure 6: Annual precipitations during 1978-1988 for the 3 stations in the northern group across the

572 border. The percentages above the bars represent the missing data for the corresponding year.



573

574

Figure 7: Mean Annual (1978-1988) measured and corrected precipitation for cold ( $T < 0^{\circ}\text{C}$ ) and warm



575

( $T > 0^{\circ}\text{C}$ ) months. The percentages are the changes from measured to corrected precipitation. The

576

approximate horizontal distance between the stations is displayed at the bottom.

577




578  
579

Figure 8: Annual precipitations during 2006-2013 for two stations in the central part of the AK/YK border. The percentages above the bars represent the missing data for the corresponding year.

580


581



583

584 Figure 9: Mean Annual (2006-2013) measured and corrected precipitation for cold ( $T < 0^{\circ}\text{C}$ ) and warm  
 585 ( $T > 0^{\circ}\text{C}$ ) months. The percentages are the change from measured to corrected precipitation. The  
 586 approximate horizontal distance between the stations is displayed at the bottom.

587



588




589 Figure 10: Scatter plots between station pairs for the measured and corrected precipitation (mm). The red

590 color shows warm months and the blue represents the cold months. A and B - Barter and Komakuk

591 comparison across the border, the highest corrected values for Barter (AK) are labeled with the date to

592 compare with Figure 4. C and D - Komakuk and Shingle Point comparison within Canada. E and F-

593 vs. Dawson across the border for the central group.



594

595

596

597

Figure 11: Double mass curves between station pairs. The red color shows the warm months and blue represents the cold months. The top and the central plots compare the stations for the northern group and the bottom one is the central station comparison across the border.