Reply to the reviews of our manuscript "Brief Communication: Global glacier mass loss reconstructions during the 20th century are consistent"

We would like to thank Hilmar Gudmundsson for obtaining the reviews, and we would like to thank both reviewers for providing their constructive and very helpful comments on our manuscript. We were able to address all their points (as detailed below).

Response to reviewer 1:

Specific comment:

1. Comment: Section 3.2 should be expanded from a short paragraph to ~ 3 paragraphs. The authors should start by pointing out the regions where the match between different methods is good (and has been previously), explaining why. They should then discuss locations where mismatches have now been resolved (this is the current focus of the section and it should be retained). They should finish by explaining where the match remains poor, and why. It would be helpful to make suggestions here (e.g., we don't know why, or more/better mass balance measurements or further updates of RGI, are required in region x, climate data needs more evaluation in region y, satellite data have limited usefulness in region z providing an indication of the greatest needs and priorities). What are the other sources of uncertainty and mismatch? Dealing with surface debris cover? Iceberg calving? Some of the discussion that I propose here may be speculative and the authors can make this clear ("we speculate..") but providing this short "road map" will help the field to move forward. I am confident that the authors have the knowledge to write this short section.

Reply: We expanded Sect. 3.2 significantly along the lines suggested by the reviewer.

Technical Corrections:

1. **Comment:** Title, abstract and paper body. The authors refer to Global glacier mass loss. Loss should be replaced with change. Even though the overall pattern is of glacier mass loss, the rate of loss has changed through time (and regionally, there have been periods of mass gain).

Reply: Done – except for those occasions where we really mean a mass loss.

2. Comment: Pg 3808: Line 1. Use "Recent estimates..." and remove "that were published in recent years".

Reply: Done.

3. Comment: Line 19. "However" and "this" shouldn't be used at the start of a new paragraph. This sentence (lines 19-21) needs to be rewritten. Reply: Done.

4. Comment: Pg 3809: Lines 4-6. Avoid starting a sentence with "But", and clarify the meaning of this sentence. It is confusing.

Reply: Done.

5. Comment: Lines 26-29. Please split into two or more sentences.

Reply: Done.

6. Comment: Pg3811: Line 8 (and elsewhere in paper): "Pentadal" is not in common usage in the English language. It would be straightforward to replace this with "5-year" and it wouldn't take up much more space.

Reply: Done.

7. Comment: Pg3811: Line 18. "Note that the results presented in this paper as Leclercq". Rewrite in active voice. "Note that the results that we present as Leclercq 2011..."

Reply: Done.

Response to reviewer 2:

General comments:

1. Comment: No numbers/results are given in the text. Although everything is put together in Table 1 and Figure 1, I have the impression that it would facilitate the reading of the paper, if some key results / some key comparisons between previous and updated estimates were given in the Results section. E.g. by how how did the estimates change with the reassessment?

Reply: We agree that readers will be interested in individual differences resulting from the updates. It is, however, hard to foresee what numbers will be most interesting, and picking examples seems arbitrary to us. For this reason (but also because of its more general usefulness) we added a data supplement, including the data shown in Figures 1 and 2, in both their old and updated versions.

2. Comment: In agreement with the first reviewer I also suggest that the analysis/the discussion of the regional differences should be extended. This might point out the regions on which to focus in future studies.

Reply: Please see our answer to reviewer 1.

3. Comment: The results indicated that the model yielded too high mass loss for certain regions due to an oversensitivity. It would be interesting to shortly discuss whether this pattern might also have affected the future projections performed by the same model (Marzeion et al., 2012).

Reply: We agree. There are small differences in the projections, but they are very small and negligible compared to the dominant uncertainty,

which for projections is the spread of the climate model ensemble used to force the glacier model. We added this to the discussion.

Detailed comments:

 Comment: page 3813, line 12: I do not fully agree with this statement. The RGIv4.0 still uses the ASTER GDEMv2.0 for the Russian Arctic and Svalbard, i.e. the same source of information as Marzeion et al. (2012). For Greenland, the GIMP DEM is used. I suggest that this somewhat fuzzy description is clarified here (also by using the reference to Arendt et al., 2014). Nevertheless, this does not explain why the maximum elevations turned out to be so different for Russian Arctic and Svalbard.

Reply: Below, we show an example of a problematic case from the Russian Arctic (Graham Bell Island in Franz Josef-Land). Obviously, there are great errors in both directions in the DEM. The too low elevations do not affect the elevation range, but the too high elevations do. While we did perform sanity checks based on the elevation distribution when we automatically extracted elevation information for Marzeion et al. (2012), we did not catch cases like this one, as the areas affected were too large to be considered errors by our routine (note also that the elevation histogram of this glacier does not look unreasonable for elevations greater than 600 m, while the true elevation maximum of the island is 509 m.). In RGIv4.0, spatial filters are applied to remove noise before the elevation distribution is determined (pers. comm. from Matthias Huss, who is calculating the elevation data for RGI). Because of the scatter of the elevation errors, the spatial filtering is better suited to remove them than the histogram-based filter we applied for RGIv1.0. We explicitly mention the spatial filtering in the revised manuscript.

2. Comment: Page 3813, line 16: Can the authors explain this 48% increase in area? I thought the same inventory (Rastner et al., 2012, TC) was included in both RGIv1.0 and RGIv4.0? However, the latter version of the RGI includes all connectivity levels, whereas the earlier versions did not include connectivity level II. If this is the case, this issue should be handled consistently in the re-assessment.

Reply: This was in fact due to the connectivity level 2 glaciers; we corrected this in the revised figures.

3. **Comment:** Page 3814, footnote: Just the glacier length feedback is mentioned here. It is, however, counteracted by the mass balance elevation feedback (which is not explicitly included in the model). Thus, part of the effect is offset. This should also be acknowledged here.

Reply: We moved this into the main text and expand a bit here, because this issue is also relevant for the reviewer's question regarding the projections. We added a statement regarding the mass balance elevation feedback, as suggested by the reviewer.

4. Comment: Table 1: I suggest to restructure Table 1 with a clear separation of previous results (Leclercq, Marzeion) and revised estimates. The



Figure 1: Example from Graham Bell Island, Franz Josef-Land. Upper panel: Black line: Glacier outline from RGIv1.0. Background shading: ASTER GDEMv2.0. White: missing data. Lower panel: distribution of elevations within the glacier outline from ASTER GDEMv2.0. The true elevation maximum of the island is 509 m.

best available estimates presently available (Cogley, Leclercq, Marzeion, Gardner) should be displayed together and not be mixed with the now obsolete numbers.

Reply: Done.

5. Comment: Figure 2: Wouldn't it be easier to read this figure if results were displayed in specific units (kg m-2 yr-1) instead of mm SLE yr-1? This would allow using the same y-scale for all panels and promote the comparison of the individual regions.

Reply: We see both advantages and disadvantages to displaying this in specific units. The reviewer mentions the most important advantage – the most important disadvantage we see is that it becomes harder to understand how the reduced mass loss of the Marzeion et al. model shown in Fig. 1 is regionally distributed. The differences of the glacier surface areas – both from the different RGI versions and the different temporal evolutions – further complicate the comparison based on specific units. For these reasons, we prefer to keep the Figure as it is.

Manuscript prepared for The Cryosphere Discuss. with version 2014/09/16 7.15 Copernicus papers of the LATEX class copernicus.cls. Date: 11 November 2015

Brief Communication: Global reconstructions of glacier mass change during the 20th century are consistent

B. Marzeion¹, P. W. Leclercq², J. G. Cogley³, and A. H. Jarosch⁴

¹Institute of Meteorology and Geophysics, University of Innsbruck, Innsbruck, Austria
 ²Department of Geosciences, University of Oslo, Oslo, Norway
 ³Department of Geography, Trent University, Peterborough, Canada
 ⁴Institute of Earth Sciences, University of Iceland, Reykjavik, Iceland

Correspondence to: B. Marzeion (ben.marzeion@uibk.ac.at)

5

10

15

Estimates Recent estimates of the contribution of glaciers to sea-level rise during the 20th century that were published in recent years are strongly divergent. Advances in data availability have allowed revisions of some of these published estimates. Here we show that outside of Antarctica, the global estimates of glacier mass loss change obtained from glacier-length-based reconstructions and from a glacier model driven by gridded climate observations are now consistent with each other, and also with an estimate for the years 2003–2009 that is mostly based on remotely sensed data. This consistency is found throughout the entire common periods of the respective data sets. Inconsistencies of reconstructions and observations persist in estimates on regional scales.

Introduction 1

Reconstructions of past glacier mass loss change are of interest for several reasons: they help constrain the budget of past sea-level change (e.g., Gregory et al., 2013); they can contribute to the understanding of the magnitude of natural (internal and forced) climate variability, and to isolating the anthropogenic signal in the climate system (Marzeion et al., 2014). Confidence in projections of future glacier change can will also be increased by the reproduction of if we can reproduce past glacier mass loss as a benchmark.

However, this requires an understanding of past glacier mass loss and the uncertaintiesinvolved in the reconstruction, and obtaining uncertainty estimates change.

- To do so, however, requires that we understand the past changes and their uncertainties. 20 Estimating uncertainties that are robust in time and space is very hard for each individual reconstruction method. If the reconstruction is based on observations (both direct and geodetic mass change observations, as well as glacier length change observations, e.g., Cogley, 2009; Leclercq et al., 2011) the uncertainty will be fundamentally governed by the sparse and probably unrepresentative sampling of observations from the entirety of 25
- Discussion Paper

the world's glaciers. In principle, it can be estimated whether the observed glaciers are

representative of the global mean for periods where global observations exist (i.e., for the satellite period; but note that remotely sensed data, e.g., Gardner et al., 2013, also have considerable uncertainty that can be hard to break down to the level of individual glaciers). But neither is the samplingThe sampling, however, is not constant in time, nor can we

- 5 . We also cannot assume that a sample of glaciers representative of the global mean now has been so in the past, or will be so in the future. If, in order to extend the period of observations, glacier length changes are used as a proxy for glacier volume change (Oerlemans et al., 2007; Leclercq et al., 2011), uncertainty in the relation between glacier length and volume is added to the sampling uncertainty.
- Reconstructions based on modeling glacier change as a response to past observed (or modeled) climate change (e.g., Marzeion et al., 2012) carry some of the uncertainty from direct glacier mass change measurements (of individual glaciers; if all of the world's glaciers are modeled individually, the potential sampling bias is not carried on). More importantly, the errors of the glacier model can only be determined at times (typically, the second half of the
- 15 20th century) and places where glacier mass change was measured (many observations from Europe, but few from heavily glacierized regions). The regions with above-average glaciological observations are also the regions with an above-average meteorological measurement density. This implies that generally speaking, the reconstructed climate used to force the glacier model is based on an above-average quality and density of atmospheric
- observations around those model uncertainty is evaluated on glaciers where the uncertainty estimate is obtained forcing is based on atmospheric observations of above-average density and guality. Since there is no practical way to estimate the impact of spatial and temporal deterioration of climate reconstructions on glacier mass balance error, it is reasonable to assume that even model validation techniques that estimate uncertainty outside of the calibration sample (such as cross validation) will underestimate the glacier model
 - uncertainty. The same argument can be made regarding the quality of glacier outlines and hypsometries (Marzeion et al., 2012).

These considerations show that it is not surprising that reconstructions of glacier mass loss change tend to agree better within the second half of the 20th century than in

earlier times (see Fig. 1), and they . They also illustrate the benefit of a comparison of reconstructions based on multiple methods: the better the agreement of the different methods within their uncertainties, the higher the confidence in their robustness – irrespective of the shortcomings of the individual uncertainty estimates.

Here we present revisions and updates of previously published estimates of past glacier mass losschange. We consider all glaciers outside of the Antarctic periphery, i.e. we include glaciers in the periphery of the Greenland ice sheet. We discuss the revisions and updates for each of the methods in Sect. 2 and show and discuss the results in Sect. 3.

2 Revisions and updates of reconstructions

10 2.1 Direct and geodetic observations

Cogley (2009) introduced a global mass-balance compilation that included geodetic as well as direct measurements. The compilation, available at http://people.trentu.ca/~gcogley/glaciology/glglmbal.htm, has been updated several times. Gardner et al. (2013, see also Vaughan et al., 2013) found inconsistencies between regional and global mass changes obtained with measurements from release 1202 of the compilation and those obtained by orbital altimetry and gravimetry. Substantial additions of geodetic and especially regional-scale geodetic measurements were made in release 1203 and carried over to release 1301. Marzeion et al. (2014) relied on release 1301 and found less negative balances than those obtained with release 1202. In particular, the inconsistency reported for 2003–2009 by Gardner et al. (2013) was somewhat reduced. It is further reduced in the present study by the exclusion of the Antarctic periphery, where measurements are few and in consequence the interpolation algorithm of Cogley (2009) yields unrealistically negative estimates of mass balance.

To illustrate the improvement of coverage in release 1301 relative to release 1202, over 1960–2012 the number of years of measured mass balance increased by 21% (from 14627 to 17673 balance years), with most of the increase accounted for by new geodetic measurements (increased by 28 %) rather than by new direct measurements (increased by 5 %).

Thus the density of temporal coverage is improved by the incorporation of geodetic massbalance measurements, which also improve spatial coverage. However because most geodetic measurements are multi-annual they tend to suppress interannual variability in regional and global estimates. This cost is offset to some extent by allowing explicitly for it in the calculation of uncertainties (as in Cogley, 2009), and also by calculating pentadal 5-year rather than annual averages.

2.2 Reconstruction based on glacier length changes

- Leclercq et al. (2011) extended the observation period of direct and geodetic observations (Cogley, 2009) with observations of glacier length changes to reconstruct the glacier mass loss change over the last two centuries. They used records of length change of 349 glaciers, distributed over 13 regions. To convert observed length change to global glacier mass change, the normalized glacier length changes were averaged to a global mean and then
- ¹⁵ scaled to get a normalized global volume change. This normalized global volume change was translated into glacier mass change by calibration against the global glacier mass loss change over the period 1950–2005 based on Cogley (2009). Note that the results presented in this paper that we present here as Leclercq et al. (2011) differ from the published results, as we here consider only glaciers outside of the Antarctic periphery. The conversion of the results for global glacier mass loss change to glacier mass loss change aveluding the
- the results for global glacier mass loss change to glacier mass loss change excluding the Antarctic periphery is straightforward as the mass loss change of the Antarctic periphery in Leclercq et al. (2011) was based on upscaling the estimate for the rest of the world.

Here, we make use of additional data on both glacier length changes (Leclercq et al., 2012, 2014) and direct and geodetic mass changes (Cogley, 2009, release 1301). The
reconstruction is now based on 456 glacier length records distributed over 15 regions. Compared to Leclercq et al. (2011) the number of records in the Arctic regions increased substantially, with over ninety new records in Novaya Zemlya, Alaska, and Greenland, such that the data set is more representative of the world's glaciers. Available length-change data

of glaciers in the Antarctic periphery are not considered here. The updates of the glacier length changes also made it possible to extend the reconstruction to 2010, instead of 2005, such that it now covers the period 1800–2010.

- The new reconstruction presented here shows considerably more glacier mass loss <u>change</u> than the reconstruction of Leclercq et al. (2011) does. The difference can partly be ascribed to the updated glacier length change data set. Using the updated data set the global normalised glacier change in the 1970s and 1980s (i.e. within the calibration period) is smaller than that in Leclercq et al. (2011), while it is similar before 1960. This leads to an increase of 10 mm SLE in the reconstructed 20th century glacier mass loss change. The cumulative mass loss change over the period 1950–2005 in the updated direct and geodetic observations is higher than in the earlier release used in Leclercq et al. (2011), which
- leads to an additional increase in the reconstructed glacier mass loss change of roughly 15 mm SLE. As discussed more extensively by Leclercq et al. (2011), the reconstruction is strongly sensitive and directly proportional to the mass loss change derived from direct and geodetic observations, and uncertainties related to the calibration using the direct and
- geodetic observations are the main cause of uncertainty in the reconstruction.

2.3 Modeled mass balances

Marzeion et al. (2012) modeled the response of each glacier contained in version 1.0 of the Randolph Glacier Inventory (RGIv1.0, Arendt et al., 2014) based on gridded climate
 observations (CRU CL 2.0 and CRU TS 3.0, New et al., 2002; Mitchell and Jones, 2005). Terminus elevation, mean elevation, and maximum elevation of the glaciers were derived from version 2 of the ASTER global digital elevation model (GDEMv2). Here, we update their reconstruction to RGIv4.0 and CRU TS 3.22 (Harris et al., 2014).

For the purpose of glacier modeling, the most significant difference between the CRU TS 3.0 and CRU TS 3.22 data sets is the extension of the covered time period from 1901–2009 to 1901–2013. While there are differences in the reconstructed glacier mass loss change also within the 20th century as a response to the updated climate forcing (globally

a reduced 4.6 mm SLE less mass loss), they are very small compared to the effects of updating from RGIv1.0 to RGIv4.0, and associated changes to the glacier model.

Along the northern boundary of the coverage of the GDEMv2, we discovered elevation errors of several hundred meters, often covering several square kilometers (probably cloud

- ⁵ tops misclassified as land surface), that impacted the terminus, mean, and maximum elevation calculations in Marzeion et al. (2012). These elevation errors led to overestimates of the elevation range of some glaciers, and therefore to overestimates of the solid precipitation, for which a lapse rate of 3 %/100 m elevation was assumed. This, through the calibration procedure, produced too high temperature sensitivities of the affected glaciers.
- The region affected most strongly by these errors was the Russian Arctic, with some significant effects also in Svalbard and the northern periphery of Greenland. RGI v4.0 contains hypsometry data for almost all glaciers, and avoids the GDEMv2 errors by either considering other topographic data sets, or by filteringand by applying a spatial filter before calculating the hypsometric data. For some regions, there were considerable changes to the selector outline at the methods are provided by a spatial filter before of the selector outlines.
- the glacier outlines themselves, most notably e.g. resulting in an increase in glacier area of 4822 % for the Greenland peripheryNorth Asia, and a decrease in glacier area of 53 % in the Low Latitudes. For detailed information on the differences between RGIv1.0 and RGIv4.0 see Arendt et al. (2014).

3 Results

20 3.1 Global scale

The lowered temperature sensitivities of the glaciers affected by the elevation error in Marzeion et al. (2012) lead to lower estimates of global glacier mass loss change in the reconstruction based on climate observations (Fig. 1a and Table 1). The differences are greatest in the first half of the 20th century¹.... This is a result of the negative

¹Note that the negative feedback between terminus elevation and mass balance becomes positive when going backwards in time: mass loss in one year implies a lower terminus in the

Discussion Paper

feedback between terminus elevation and mass balance, which becomes positive when going backwards in time: mass loss in one year implies a lower terminus in the preceding year, which leads to a more negative mass balance in the preceding year¹. This feedback was further erroneously enhanced in Marzeion et al. (2012) through the overestimated elevations ranges. I.e., for reconstructions, systematic mass balance errors were amplified further back in time. In forward model runs, the errors are accordingly dampened, which explains why the corrected elevation ranges affect the projections of Marzeion et al. (2012) far less. The resulting differences are negligible compared to the dominant uncertainty, which for projections is the spread of the climate model ensemble

¹⁰ used to force the glacier model (not shown).

The revised mass change reconstruction based on glacier length change shows higher mass loss during the 20th century than in Leclercq et al. (2011), leading to an agreement (within their respective uncertainties) of the reconstructions based on glacier length change and climate observations, throughout the entire length of their overlapping periods. There is also agreement of the pentadal5-year, global mean rates of glacier mass change (Fig. 1b) of the revised reconstructions, as well as for the 2003–2009 period (Fig. 1c), for which altimetric and gravimetric data give relatively tight uncertainty constraints for several strongly glaciated regions (Gardner et al., 2013). Even though the uncertainty ranges are still relatively large, this result indicates that the different reconstruction methods are converging as more and higher quality data are becoming available. This increases confidence in their viability.

Strictly speaking, the three reconstructions considered here are not independent over the entire time, because (i) the glacier length-based estimate of Leclercq et al. (2011) is calibrated globally using the direct and geodetic mass change observations of Cogley (2009), and (ii) the estimate based on climate observations of Marzeion et al. (2012) is

25

5

15

20

¹Note that in reality this feedback to some extent is balanced by the mass balance–surface elevation feedback. This feedback, however, is not included in the model.

preceding year, which leads to a more negative mass balance in the preceding year. I.e., for reconstructions, systematic mass balance errors are amplified further back in time.

calibrated using direct mass change observations on 255 glaciers that also enter Cogley (2009). The practical limitations caused by this dependence are minor, as can be seen in Fig. 102: even though the validation in Marzeion et al. (2012) does not indicate a model bias on those glaciers that enter Cogley (2009), the regional mean estimates of the two methods are strongly divergent in some regions, which would not be possible if the dependence was strong.

3.2 Regional scale

subdued.

5

While the agreement of the reconstructions on the global scale is clearly improved, the comparison of the climate observation-based reconstruction with the results of Gardner et al. (2013) still shows considerable differences on the 10 regional scale, and the improvement through the revision of the reconstruction is marginal. The changes are greatest in the for some of the regions. In regions where the spatial and temporal densitiv of glacier observations is high (e.g., Western Canada & US, Scandinavia, Central Europe, Caucasus & Middle East, see Gardr

there is good agreement between both the old and the revised model-based reconstructions 15 and observations. This reflects the abundance and generally high quality of glaciological and meteorological data in these regions.

The greatest improvement in the agreement between observations and reconstruction can be found in the Russian Arctic. There are also considerable changes to the reconstruction in Greenland and Svalbard (in those three cases, these cases, as with the 20 Russian Arctic, due to the corrected elevation errors), and in the Low Latitudes (due to the strongly reduced glacier area in RGIv4.0). However, since the changes mostly affect the period prior to the estimate of Gardner et al. (2013), it is unclear whether these changes improve the regional estimate (we can only conclude that the total sum of regional changes improved, based on the better agreement on global scale described in Sect. 3.1). These 25 regions, especially Greenland (Fig. 2), accounted for much of the rapid mass loss simulated reconstructed for 1930-1935 by Marzeion et al. (2012), and this excursion is now more

In all other regions, the effects of the revision are negligible, and the. The strong disagreement during the 2003-2009 period, particularly in Svalbard, the and Canadian Arctic, and is not resolved and is even increased in the Southern Andesis not potential explanations: (i) The model-based reconstruction is motivated by the generally 5 better availability and guality of meteorological observations compared to glaciological observations, particularly in the early period of the reconstruction. During the most recent years it is possible that the station-based meteorological observations reflect the spatial and temporal atmospheric variability less well than remotely sensed data of glaciers. (ii) The glacier model's calibration routine (Marzeion et al., 2012) relies on the assumption that the 10 sampling of direct glacier observations is dense enough to reflect the spatial scale of climate anomaly patterns. In sparsely sampled regions, and in regions with small scale variability, this assumption will limit the success of glacier mass balance reconstructions. (iii) Calving is not explicitly considered in the model, but is responsible for a considerable fraction of the mass budget of many high-latitude glaciers. The implicit treatment of calving will limit 15 the model's ability to reflect the full temporal variability of the glacier mass balance, and in particular the partial decoupling of glacier mass change and climate forcing that may occur for calving glaciers. (iv) The sampling of glacier observations may not be representative for entire regions if based on direct observations of individual glaciers, and may be affected by methodological uncertainties (e.g., in the conversion of volume change to mass change 20 for geodetic estimates, or by leakage of mass change from other components of the Earth system and/or other regions for gravimetric data).

Conclusions 4

Additional glacier length data (Leclercq et al., 2012, 2014), updates of the RGI (Arendt et al., 2014) and associated corrections of errors in glacier elevation data, additional more 25 extensive geodetic measurements of glacier mass change, and extensions of gridded climate observations (Harris et al., 2014) encouraged us to revise reconstructions of 20th

Discussion Paper

century glacier mass change. These revisions lead to results that are consistent with each other on the global scale, and on all common time scales. Inconsistencies remain in the recent past (2003–2009) on the regional scale in some regions between our reconstructions and a consensus estimate that relies strongly on altimetric and gravimetric data (Gardner et al., 2013), particularly in Arctic regions.

The newly-achieved consistency between the two reconstructions may simply mean that they are consistently wrong, such that future improvements in observations of glacier length change, glacier mass change, glacier geometries included in the RGI, as well as model formulation may lead to different estimates of 20th century glacier mass loss change. But the

¹⁰ strongest evidence for this argument, namely the discrepancy with altimetric and gravimetric estimates during 2003–2009, is now less strong, as seen in Fig. 1c and Table 1.

Supplementary material related to this article is available online at ...

Acknowledgements. This work was funded by the Austrian Science Fund (FWF): P22443-N21 and P25362-N26, and supported by the Austrian Ministry of Science BMWF as part of the UniInfrastrukturprogramm of the Focal Point Scientific Computing at the University of Innsbruck. P. W. Leclercq acknowledges funding by the European Research Council under the European Union's Seventh Framework Programme (FP/2007-2013)/ERC grant agreement no. 320816. We thank two anonymous reviewers for their comments that helped to improve the manuscript.

References

5

15

- Arendt, A., Bliss, A., Bolch, T., et al.: Randolph Glacier Inventory A Dataset of Global Glacier Outlines: Version 4.0, Global Land Ice Measurements from Space, Boulder Colorado, USA, Digital Media, 2014.
 - Cogley, J. G.: Geodetic and direct mass-balance measurements: comparison and joint analysis, Ann. Glaciol., 50(50), 96–100, doi:10.3189/172756409787769744, 2009.
- ²⁵ Gardner, A. S., Moholdt, G., Cogley, J. G., Wouters, B., Arendt, A. A., Wahr, J., Berthier, E., Hock, R., Pfeffer, W. T., Kaser, G., Ligtenberg, S. R. M., Bolch, T., Sharp, M. J., Hagen, J. O., van den

Broeke, M. R., and Paul, F.: A reconciled estimate of glacier contributions to sea level rise: 2003 to 2009, Science, 340, 852–857, doi:10.1126/science.1234532, 2013.

Gregory, J. M., White, N. J., Church, J. A., Bierkens, M. F. P., Box, J. E., van den Broeke, M. R., Cogley, J. G., Fettweis, X., Hanna, E., Huybrechts, P., Konikow, L. F., Leclercq, P. W., Marzeion, B.,

Oerlemans, J., Tamisiea, M. E., Wada, Y., Wake, L. M., and van de Wal, R. S.: Twentieth-century global-mean sea-level rise: is the whole greater than the sum of the parts?, J. Climate, 26, 4476– 4499, doi:10.1175/JCLI-D-12-00319.1, 2013.

Harris, I., Jones, P., Osborn, T., and Lister, D.: Updated high-resolution grids of monthly climatic observations – the CRU TS3.10 Dataset, Int. J. Climatol., 34, 623–642, doi:10.1002/joc.3711, 2014.

Leclercq, P. W., Oerlemans, J., and Cogley, J. G.: Estimating the glacier contribution to sea-level rise for the period 1800–2005, Surv. Geophys., 32, 519–535, doi:10.1007/s10712-011-9121-7, 2011.
Leclercq, P. W., Weidick, A., Paul, F., Bolch, T., Citterio, M., and Oerlemans, J.: Brief communication "Historical glacier length changes in West Greenland", The Cryosphere, 6, 1339–1343,

doi:10.5194/tc-6-1339-2012, 2012.

10

15

20

Leclercq, P. W., Oerlemans, J., Basagic, H. J., Bushueva, I., Cook, A. J., and Le Bris, R.: A data set of worldwide glacier length fluctuations, The Cryosphere, 8, 659–672, doi:10.5194/tc-8-659-2014, 2014.

Marzeion, B., Jarosch, A. H., and Hofer, M.: Past and future sea-level change from the surface mass balance of glaciers, The Cryosphere, 6, 1295–1322, doi:10.5194/tc-6-1295-2012, 2012.

- Marzeion, B., Cogley, J. G., Richter, K., and Parkes, D.: Attribution of global glacier mass loss to anthropogenic and natural causes, Science, 345, 919–921, doi:10.1126/science.1254702, 2014.
 Mitchell, T. D. and Jones, P. D.: An improved method of constructing a database of monthly climate observations and associated high-resolution grids, Int. J. Climatol., 25, 693–712, doi:10.1002/ioo.1181.2005
- ²⁵ doi:10.1002/joc.1181, 2005.
 - New, M., Lister, D., Hulme, M., and Makin, I.: A high-resolution data set of surface climate over global land areas, Clim. Res., 21, 1–25, doi:10.3354/cr021001, 2002.

Oerlemans, J., Dyurgerov, M., and van de Wal, R. S. W.: Reconstructing the glacier contribution to sea-level rise back to 1850, The Cryosphere, 1, 59–65, doi:10.5194/tc-1-59-2007, 2007.

Vaughan, D., Comiso, J., Allison, I., Carrasco, J., Kaser, G., Kwok, R., Mote, P., Murray, T., Paul, F., Ren, J., Rignot, E., Solomina, O., Steffen, K., and Zhang, T.: Observations: cryosphere, in: Climate Change 2013: the Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, edited by: Stocker, T., Qin, D., Plattner, G.-K., Tignor, M., Allen, S., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P., Cambridge University Press, Cambridge, UK, New York, NY, USA, 317–382, 2013.

Table 1. Globally integrated glacier mass change reconstructions. Note that all reconstructions exclude glaciers in the Antarctic periphery.

	Mass change [mm SLE]		Mass change rate $[mm SLE yr^{-1}]$
	1961–2005	1902–2005	2003–2009
Leclercq et al. (2011)	19.2 ± 4.9	57.5 ± 14.4	_
Marzeion et al. (2012)	26.4 ± 3.1	92.3 ± 7.7	0.96 ± 0.12
Gardner et al. (2013)	_	_	0.70 ± 0.07
Cogley (2009, R1301)	23.0 ± 2.8	-	0.75 ± 0.07
Updated from Leclercq et al. (2011)	24.3 ± 7.1	80.4 ± 21.1	0.87 ± 0.64
Updated from Marzeion et al. (2012)	19.9 ± 3.4	63.2 ± 7.9	0.78 ± 0.15



Figure 1. Globally integrated glacier mass change; (a) accumulated in time, relative to the year 2000; (b) pentadal 5-year mean values of mass loss change rates; (c) mean mass loss change rates during 2003 to 2009. Shading/error bars indicate 90% confidence interval. Note that all reconstructions exclude glaciers in the Antarctic periphery.



Figure 2. Regional glacier mass change rates, colors as in Fig. 1. Shading indicates the 90% confidence interval of the pentadal 5-year means. Numbers given on top of each panel indicate glacier area in RGIv1.0 (green) and RGIv4.0 (blue). Crosses on the right of each panel give mass change rate and 90% confidence interval during 2003–2009.