

## Reply to reviewer 1

We would like to thank the reviewers for their constructive comments on our manuscript. Both reviewers are concerned about the novelty of the methods employed and the results obtained in this study. We therefore have to acknowledge that we were unable to explain the purpose of our study and to highlight the novelty of our results. Before addressing the reviewer's specific comments in detail, we will therefore briefly mention the goals and the novelty of this work:

As mentioned in the introduction the three goals of this study were:

- i) Modelling the annual mass balances of eight Scandinavian glaciers with long annual mass balance series using a suite of statistical models using seasonally averaged climate data as input variables. These models enable us to compare the relative importance of accumulation-season precipitation and ablation-season temperature on annual mass balances of glaciers.
- ii) Assessing temporal changes of relative importance of accumulation-season precipitation and ablation-season temperature. These temporal changes are then compared to large-scale oceanic and atmospheric modes, such as the Atlantic Multidecadal Oscillation (AMO) and the North Atlantic Oscillation (NAO).
- iii) In a last step we compare the climate sensitivities of ablation-season temperature and accumulation-season precipitation of statistical models to results from other modelling approaches, namely temperature index models and physically based models. We then used the mass balance models and climate projections for the years 2050 and 2100 to predict average annual mass balances for these years.

The reviewers are very concerned about point iii). The projections were only a side product of this work, and as the reviewers point out a rather poor one. We therefore removed this part of the manuscript. As the reviewers point out, the glaciers studied in this study have all been studied previously, most of them with more sophisticated models.

These studies have focused on two aspects of glacier mass balances:

- i) Many studies have focused on sensitivities of mass balances to changes in temperature and precipitation, i.e. on expected changes of net balances for given changes in temperature (in °C or K) and precipitation (in % of a reference level).

The different units of temperature and precipitation make a direct comparison of the climate sensitivities in term of relative importance difficult. For instance 1 K and 10% of the precipitation of a reference period do most probably not cover the same proportion of the range of precipitation and temperature data.

Considering data from Bergen for the period 1962 – 2010, the standard deviation (sd) of temperature from May to September (T MJJAS) is 0.856 °C. 1°C is then  $1.16 * sd(T \text{ MJJAS})$ . The average monthly precipitation from October to April (P ONDJFMA) is 200 mm, the standard deviation is 52 mm.  $1.16 * sd(P \text{ ONDJFMA})$  is then 61 mm which corresponds to a precipitation change of 30%. In this case, climate sensitivity to a 1°C change similar to a 30% precipitation change is indicative of equal relative importance of T MJJAS and P ONDJFMA. The standard deviation of summer temperature and winter precipitation is different for each station and for each point of a gridded data set.

In this study, we therefore want to estimate the relative importance of summer temperature and winter precipitation, most importantly these relative importances are directly comparable, which is not the case for climate sensitivities.

ii) Some studies have focused on the relative importance of summer (Bs) and winter balance (Bw) for the annual balance (Ba) (e.g. Nesje et al. 2000, Andreassen et al. 2005, Mernild et al. 2014). Nesje et al. (2000) used the correlation between Bs and Ba and Bw and Ba as estimators of relative importance, whereas Andreassen et al. (2005) used the more direct measures of ratios of standard deviations of Bs and Ba ( $sBs/sBa$ ) and Bw and BA ( $sBw/sBa$ ).

In this study, we use a combination of these two approaches: We want to estimate the relative importance of climate variables (summer temperature and winter precipitation) for the annual balance. Additionally, we are interested in changes of relative importance of summer temperature and winter precipitation for the annual mass balance, and we test if the relative importances change when only considering years characterised by certain states of climate indexes.

In this study, we propose a statistical framework in which we are able to automatically/ objectively assign relative importance to variations in summer temperature and variations in winter precipitation. Standardising all variables involved in a linear model (as outlined on page 389) enables the calculation of relative importance of summer temperature and winter precipitation for net balances. Most importantly these relative influences are then comparable. I.e. a higher absolute value of the relative importance of summer temperature (the standardized partial regression coefficient to use some jargon) than of winter precipitation means that summer temperature is more important.

We are interested in the relative importance of summer temperature and winter precipitation for the entire measurement period, but we are also interested in changes of the relative importance of summer temperature and winter precipitation through time. Therefore, we estimate the relative importance of summer temperature and winter precipitation in 25-year moving windows (changed from 30-year windows in the original manuscript).

As there is one main pattern of oceanic variability, the AMO and one main pattern of atmospheric variability, the NAO, over the North Atlantic, we are interested to see if there are changes in relative importance of summer temperature and winter precipitation for the net balance for the two states of the AMO and the two states of the NAO. In addition to 25-year running windows, we therefore estimated relative importance of summer temperature and winter precipitation in phases where we expect above and below normal summer temperature (as expressed by the AMO) and in phases where we expect above and below normal winter precipitation (as expressed by the NAO).

The results of the analyses in moving windows and the estimation of relative importance of summer temperature and winter precipitation for different states of the NAO and the AMO are the main novelty of this study. These results demonstrate changes of relative importance of summer temperature and winter precipitation depending on the calibration period, and their association with large scale oceanic and atmospheric patterns.

To our knowledge, such a thorough assessment of relative importance of summer temperature and winter precipitation on net balances has never been presented for Scandinavian glaciers before. Hence, we believe that the results presented in this study are indeed novel and interesting from a climatological viewpoint.

These results are especially interesting for palaeoclimatological studies, where at maximum seasonal precipitation and temperature, or possibly only one of them is available. This study cautions against the assumption of a constant relative importance of winter precipitation and summer temperature for net balances and against the use of mass balance reconstructions for reconstructing the NAO-index.

In comparison to methods used by Nesje et al. (2000) and Andreassen et al (2005) we only need annual mass balance, and not winter and summer balance to estimate relative importance of summer temperature and winter precipitation.

We rewrote large portions of the manuscript to improve the explanations of methods used and to emphasise the points made above. We also changed some analyses as follows:

We also replaced the analysis shown in Fig 3 (correlations of winter and summer balance with net balance) with a direct measure of the relative importance of winter (Bw) and summer (Bs) balance on the annual balance (Ba). Like Andreassen et al. (2005) we used the ratios of the standard deviations of winter balance (sBw) and annual balance (sBa) and summer balance (sBs) and net balance (sBn) and compared the entire measurement period with the two states of the NAO and if possible the two states of the AMO.

We also modified most of the figures according to the reviewers' suggestions.

### **Reply to reviewer 1:**

*Page 384, line 23: What about summer accumulation and winter melting? Throughout the entire paper the authors only talk about winter accumulation and summer ablation. Of course, these components are the most relevant ones, but for maritime glaciers considerable snow fall amounts can also occur during summer in the higher regions, and the glacier tongues can experience melting over the winter season. These problems are not discussed at all.*

The influence of summer snow-fall and winter melt are indeed not accounted for with the methods used in this study. The reviewer points out that these are minor effects compared to winter snow-fall and summer melt, and suggests that winter melt and summer accumulation are particularly important for maritime glaciers. Still, mass balances of maritime glaciers are very well modelled by the statistical models and models explain more of the variance of net balance for maritime glaciers than for continental glaciers. This clearly shows that winter melt and summer accumulation are (currently) not a major problem for modelled mass balances.

We added a short discussion related to this problem.

*Page 385, line 21: This sentence appears to be circular to me – or maybe too complicated to get its essence. It occurs in similar form several times in the paper.*

The sentence reads 'When modelling the joint influence of ablation-season temperature and accumulation-season precipitation on annual mass balance, statistical models allow for assessing the individual influence of ablation-season temperature and accumulation season precipitation on annual mass balance.'

This formulation was unclear and was changed to: When modelling the joint influence of summer temperature and winter precipitation on annual mass balance, statistical models enable us to compare the relative importance of summer temperature and winter precipitation on the annual mass balance.

*Page 386, line 13: The uncertainty in the mass balance data is not addressed. At least for some of the maritime glaciers, there is an indication that the glaciologically derived mass balances are significantly*

*more positive than mass balance based on long-term geodetic surveys. As the mass balance data are the backbone of the study, more effort could be invested to discuss the uncertainty in the input data and potential effects on the results.*

The uncertainty in the mass balance measurements is estimated to between 0.2 and 0.4 mwe per year (Andreassen et al. 2005). The effect of the uncertainties is determined by their nature. If the mass balance measurements are mainly biased, then the estimates of the relative importance of summer temperature and winter precipitation will change (if the measurements are biased in the way described by reviewer 1, then the relative importance of temperature will increase). If the uncertainties are mainly random, then they are affecting our models in a random way, i.e. estimates of relative importance may change either way, and variance explained will change as well.

On the one hand, the unexpected changes of relative importance of winter and summer balance for net balance as function of NAO-index for Aalfotbreen and Engabreen (decreased relative importance of winter balance for years with above median NAO-Index) might suggest problems with the mass balance data, on the other hand, arguing: as we do not get the expected results the data we used is poor is a very dangerous line of argument.

We also added a paragraph discussing possible uncertainties.

*Page 392, line 25: What is the objective of analysing positive and negative AMO phases separately? I.e. what do the authors want to find out?*

We hope that we could answer this question in the general part of this response. We are mainly interested in the relative importance of winter precipitation and summer temperature for the net balance. We suspect that the AMO has an influence on this relative importance, as the AMO expresses changes in sea-surface temperatures. That is why we tested the differences between models including years with exclusively AMO + and exclusively AMO-, respectively.

*Section 2.2.6.: The first paragraph should rather be in the data section*

This entire section was removed.

*Page 387, line 15: Here and elsewhere. Why not simply “winter” balance and “summer” balance? The use of “ablation-season mass balance” etc is relatively complicated and sometimes awkward.*

As suggested by reviewer 1, we now use winter and summer instead of accumulation season and ablation season

*Figure 3: What is on the y-axis? The label is tiny. Better use text. Furthermore, I found the point cloud not very intuitive. When printed in black-white the information is almost impossible to extract. Also the*

*experiments are difficult to understand from the caption. It might also be reasonable to reduce the number of experiments shown.*

We improved the readability following the reviewer's suggestion. We kept the number of experiments constant as we compare correlations for the entire measurement period and for years with above and below median NAO-index.

*- Figure 4: add (a), (b) etc to the panels. What is on the x- and y-axis? It would be much more intuitive when writing "Summer temperature" and "winter precipitation" –*

The labels of the y-axis are indeed unclear, these are the smooth terms of the additive model.

*Figure 5: add (a), (b) etc. Shouldn't this figure be flipped by 90 degrees?*

Yes, this figure was arranged in this way by the editorial team.

*- Figure 6: I really have troubles with this figure. The labels are tiny! Furthermore, even after re-reading the text, the figure is difficult to understand. The approach of evaluation, the presentation of the results and their interpretation (text) should be improved.*

In Fig. 6, the changes of relative importance of summer temperature and winter precipitation through time are indicated. Hence, this is one of the main figures of this manuscript. We tried to improve the explanation of this figure.

## **Reply to reviewer 2.**

### *Specific comments*

*P 385. The introduction needs to be rewritten. The authors use the term physically based mass balance models in contrast to statistical models. They here divide statistical models in i) temperature-index-models and (ii) models that use seasonal and monthly mean T and P. This description in the introduction is a bit unclear and division is not necessarily appropriate for the papers they refer to. There is a whole range of mass balance models from sophisticated energy balance models to simplified temperature-index models and regression models (e.g. see overview by Hock, 2005). Furthermore, mass balance models can be for a point, a profile or they can be spatially distributed, and they may or may not be coupled with a dynamical model. For example an energy balance model can be just for a point, whereas a degree day model can*

*be coupled to a dynamical model. Moreover, temperature-index models also need calibration.*

*C131P 385/386 Aims i-iii. What is the purpose of these aims? In particular iii) seems not to be a good approach based on current knowledge (e.g. a model needs to take into account the changes in ablation and accumulation season as well as dynamical changes). Furthermore, why are these years chosen and more importantly, what is the advantage of this approach compared to already published studies?*

We tried to address these points in the replies to the reviewers' general comments. But most importantly: we are interested in relative importance of summer temperature and winter precipitation for the annual balance and not in absolute importance (climate sensitivity).

We are interested in changes of relative importance through time and when only considering states of certain climate indexes.

*Could here or rather in the introduction mention other data sources, e.g. seNorge.no, reanalysis data etc. In the discussion could be interesting to have a comparison of the results using other data sources. e.g. that mass balance have been modelled for all of Norway using temperature-index models (Engelhardt et al., 2013)*

The direct comparison to results by Engelhardt et al. (2013) who used data from senorge.no is extremely difficult. For Nigardsbreen and Storbreen (distance of about 50km) the predicted net balances have a strong negative bias for Storbreen and a strong positive bias for Nigardsbreen. This might be caused by model bias, or by biases in the climate data. We assume these are biases caused by the model and not by the data, as opposite bias on a short distance of 50 km would be very surprising.

We also tested effects of using different station data. For instance, we used climate data from Lom between 20 and 30 km to the north-east of the continental glaciers in southern Norway. Model performance was reduced compared to models using meteo data from Bergen as input. Using data from Sogndal (to the southwest of the three continental glaciers), but much closer than Bergen had no systematic influence on the performance of statistical models (the performance remained unchanged for one glacier, was slightly decreased for one glacier and was slightly increased for one glacier).

Additionally, for our models (except when calculating climate sensitivities), it is only important if there is a linear relation between temperature and precipitation in Bergen and close to the glaciers, as the climate data are standardized.

*P 393. Line 9. Are the results then directly comparable when different (the most parsimonious) models are used for each glacier?*

In this study, we are first of all interested to compare the relative importance of summer temperature and winter precipitation for the mass balances of individual glaciers. Hence we are not interested to compare the models between glaciers. We are simply interested to see if for individual glaciers, the cost of additional complexity (individual weights for temperature and precipitation, and for additive models, the use of smooths instead of coefficients) is justified by increased variance explained. In this context, it is only important that we always use the same model for one glacier but not between glaciers.

*Chapter 3. Line 20->The cumulative mass balance records are not a result of this paper and does not strictly belong here.*

This is indeed true, however, not showing the data would make the presentation of results and their discussion extremely difficult.

*P 397 Line 5 This is the first time fig 5 is mentioned, but it ends abruptly with a new paragraph, rewrite. Please explain a bit what you mean and what is new here compared to this rather well known results of cumulative balances and mass surpluses for certain periods. Why just AMO and NAO, since there are other indices too that has been studied? The chapter 4 discussion has one subheading for the last ~third of the text, suggest to add 1-2 subheading(s) also for the first part.*

In the new manuscript Figs 2, 5 and 6 have been merged. The temporal changes of relative importance of summer temperature and winter precipitation (25-year windows) are now shown together the cumulative mass balances.

Using 25-year running windows, we find changes in relative importance of summer temperature and winter precipitation through time. Results are especially interesting for continental glaciers in southern Norway. For 25-year windows centred before 1978, relative importance of summer temperature and winter precipitation are equal, the relative importance of winter precipitation is higher in the following 25-year windows, and summer temperature gets more important than winter precipitation for 25-year windows centred after 1990.

With the method employed in this study, we are able to directly assign relative importance to summer temperature and winter precipitation in the 25-year moving windows. We are using the same mass balance data as in previous studies, hence we do not expect to find completely different results. For instance we do not expect to find increased relative importance of summer temperature in phases of positive mass balances.

Besides the AMO and the NAO, other indexes for instance the Arctic Oscillation (AO) index could have been used. The NAO and the AO are very similar concepts (Thompson and Wallace, 1998), the NAO-index is traditionally based on pressure measurements of two meteorological stations (on Island and the Azores or Lisbon), whereas the AO is the first PC of the surface level pressure field north of 20N. As NAO and AO are strongly related (Peings and Magnusdottir, 2014, do mention NAO / northern annual model simultaneously) it seems only reasonable to use either the NAO or the AO for comparison with glacier mass balances (as the use of both would be redundant). We decided to use the NAO as the NAO-index is only considering data from the North Atlantic region and not the entire area north of 20N. Additionally, Li et al. (2013) find the NAO to be the dominant mode of atmospheric variability over the North Atlantic.

We now use four subheadings in the results and discussion sections.

*P398 Line 18. Incorrect to write that glacier melt started in the early 2000s in Scandinavia. Some of the continental glaciers have in general lost mass throughout since measurements began in 1940s or 1960s, except for the period with transient mass surplus 1989-2005. That glaciers in the alps and Scandinavia correlate differently to the NAO-index is not new, here more references to previously studies should be*

added, e.g. Marzeion & Nesje (2012) and see also Rasmussen (2007) for more references. P400. Line 9. Might is a weak word here, it will. Line 24. Here you refer to previous studies, then use present tense. Then you refer to your own results, a bit abrupt, rewrite.

This is indeed true, we meant to say that the transient mass loss after the transient positive mass gain between 1989 and 1995 (not 2005, as stated by the reviewer) started in the early 2000s.

Regarding NAO and the Alps we added references to Marzeion and Nesje (2012) and Six et al. (2001).

P401. Line 4-5. I find this way of using references with your results confusing. Use 'as shown by' or similar. Conclusion: should be rewritten to focus on the new outcomes of your study.

The conclusions were rewritten.

Table 2. The first sentence is repeated in the next sentence. Giesen is written with one s.

Figure 1. The mid panel is cut in a strange way. It should be possible to show it in three full panels, using the same orientation and a different location map than the left panel. The source of the P and T data should be given, e.g. seNorge.no? It looks like three maps, not a location map with inserts (they are the same size), thus, rewrite text. Figure 2. This is the same as Fig 5a. As this is not a result by the authors I suggest to remove it here and add the data source to the figure text in 5a.

Figure 3. The first two sentences are nearly identical. Change Bn to Ba as Ba is standard term shortening for annual balance now in common use (Cogley et al., 2011). Bn is usually used as shortening for net balance.

Changed.

Figure 4. could specify the confidence bounds in figure text.

Figure 5. See comment to Figure 2. a)-d) are missing from the figure. c) Jones et al., 1997 are referred to, but data spans to 2010, could add an updated ref or write Jones et al., 1997 with updates.

Was changed.

Figure 6. Very small font compared to other figs.

Font was increased

Figure 7. Annual balance = Ba



*Changed.*

*Figure 8. Not sure if it is worthwhile to show this result due to the limitations of the model. See previous comments.*

This result was removed.

#### **References:**

Andreassen, L. M., Elvehøy, H., Kjollmoen, B., Engeset, R. V., and Haakensen, N.: Glacier mass-balance and length variation in Norway, *Ann. Glaciol.*, 42, 317–325, 2005. Engelhard et al. 2013.

Li, J., Sun, C., and Jin, F.-F.: NAO implicated as a predictor of Northern Hemisphere mean temperature multidecadal variability, *Geophys. Res. Lett.*, 40, 5497–5502, doi:10.1002/2013GL057877, 2013.

Marzeion, B. and Nesje, A.: Spatial patterns of North Atlantic Oscillation influence on mass balance variability of European glaciers, *The Cryosphere*, 6, 661-673, doi:10.5194/tc-6-661-2012, 2012.

Mernild, S.H., Hanna, E., Yde, J.C., Seidenkrantz, M.-S., Wilson, R. and Knudsen, N.T. Atmospheric and oceanic influence on mass balance of northern North Atlantic region land-terminating glaciers. *Geografiska Annaler: Series A, Physical Geography*, 96, 561–577. doi:10.1111/geoa.12053

Nesje A, Lie Ø and Dahl SO (2000) Is the North Atlantic Oscillation reflected in Scandinavian glacier mass balance records? *Journal of Quaternary Science* 15(6): 587–601.

Peings, Y. and Magnusdottir, G.: Forcing of the wintertime atmospheric circulation by the multidecadal fluctuations of the North Atlantic ocean, *Environ. Res. Lett.*, 9, 034018, doi:10.1088/1748-9326/9/3/034018, 2014.

Six, D. Reynaud L., Letreguilly A. (2001). Bilans de masse des glaciers alpins et scandinaves, leurs relations avec l'oscillation du climat de l'Atlantique nord. *Earth and Planetary Sciences* 333 (2001) 693–698.

### **Changes in the Manuscript:**

- We removed the part dealing with projections of future mass balances
- The number of Figures was reduced from eight to five.
- We removed the analysis comparing correlations between  $B_w$  and  $B_a$  and  $B_s$  and  $B_a$  and replaced this part by comparing standard deviation ratios ( $sB_w/sB_a$  and  $sB_s/sB_a$ ) as they are more direct and more useful measures of the relative importance of summer and winter balance for annual balances.
- The entire manuscript except the methods section was rewritten following the reviewers suggestions. We therefore do not attach a manuscript with track changes.
- Unfortunately, we could not include a discussion of the publication by Mernild et al. (2014). We were unable to reproduce their results and therefore think that they used methods differing from the methods described in their paper. We asked S. Mernild for clarification but, did not get answers containing more than an acknowledgment of receipt of our inquiry. As we do not know what Mernild et al. (2014) did in their paper, we are not able to discuss their results. (Please contact the editor for further clarifications of this issue).

# 1 Modelling annual mass balances of eight Scandinavian 2 glaciers using statistical models

3

4 **M. Trachsel<sup>1</sup> and A. Nesje<sup>2</sup>**

5 [1]{ Department of Biology, Bergen, Norway }

6 [2]{ Department of Earth Science and Uni Climate, Bergen, Norway }

7 Correspondence to: M. Trachsel (mathias.trachsel@uib.no)

8

## 9 **Abstract**

10 Glacier mass balances are mainly influenced by winter precipitation and summer temperature.  
11 We used simple statistical models to assess the relative importance of summer temperature  
12 and winter precipitation for annual balances of eight glaciers in Scandinavia. Winter  
13 precipitation is more important for maritime glaciers, whereas summer temperature is more  
14 important for annual balances of continental glaciers. Most importantly relative importances  
15 of summer temperature and winter precipitation were not stable in time. For instance, winter  
16 precipitation was more important than summer temperature for all glaciers in the 25-year  
17 period 1972 – 1996, whereas the relative importance of summer temperature was increasing  
18 towards the present. Between 1963 and 1996 the Atlantic Multidecadal Oscillation (AMO)  
19 index was consistently negative and the North Atlantic Oscillation (NAO) Index was  
20 consistently positive between 1987 and 1995, both being favourable for glacier growth.  
21 Winter precipitation was more important than summer temperature for annual balances when  
22 only considering subsets of years with high NAO-index and negative AMO-index,  
23 respectively, whereas the importance of summer temperature was increased analysing subsets  
24 of years with low NAO-index and positive AMO-index, respectively. Hence, the relative  
25 importance of precipitation and temperature for mass balances was probably influenced by the  
26 state of the AMO and the NAO, as these two indexes are associated with changes in summer  
27 temperature (AMO) and winter precipitation (NAO).

28

# 1 Introduction

Glaciers respond to climate change because their mass balance and extent are mainly a result of variations in winter accumulation and summer ablation. Over time, glacier changes exhibit some of the clearest evidence of variations in the earth's climate system. As a result, glaciers are a key indicator of global, regional and local climate change (IPCC, 2007, 2013). Glaciers integrate changes in accumulation as well as changes in ablation. Past (e.g. Nesje, 2009), present (e.g. Andreassen and Oerlemans, 2009) and future (e.g. Giesen and Oerlemans, 2011) of Scandinavian has been studied extensively. The accumulation on Scandinavian glaciers is mainly a result of winter precipitation (as snow) and wind redistribution of snow, whereas glacier ablation is more complex and depends on the total energy available for melt. Accumulation and ablation processes have been extensively studied by means of mass balance models of varying complexity (e.g. Andreassen et al., 2006; Andreassen and Oerlemans, 2009; Engelhardt et al. 2013; Giesen and Oerlemans, 2010; Hock et al. 2007; Laumann and Nesje, 2009a, 2009b, 2014; Oerlemans, 1992, 1997; Rasmussen and Conway 2005; Rasmussen et al. 2007; Schuler et al. 2005). Most of these studies have focused on calculating sensitivities of winter balances, summer balances and annual balances to changes in temperature and precipitation. Many studies provided projections of future mass balances based on climate projections (e.g. Giesen and Oerlemans, 2011). Climate sensitivities are absolute influences of temperature and precipitation changes on mass balances. They are, however, measured in different units and are therefore difficult to compare directly ( $\Delta m w.e$  for changes in K and in % of precipitation). It is possible to directly deduce from climate sensitivities that changes in temperature are more important for continental glaciers than for maritime glaciers in southern Norway, as a larger change in precipitation is needed to counterbalance a temperature change of 1 K. But it is not possible to directly assess if temperature or precipitation are more important for the annual balances of one glacier. Relative and thereby directly comparable sensitivities of annual balances to changes in temperature and precipitation are therefore not obtained from climate sensitivities.

Further studies have explicitly assessed the relative importance of winter balance and summer balance for annual balance by correlating the summer and winter balances with annual balance (Nesje et al. 2000). Nesje et al. (2000) showed that the correlation between winter balance and annual balance is higher than the correlation between summer balance and annual balance for maritime glaciers and *vice versa* for continental glaciers. Mernild et al. (2014)

1 replicated this analysis using data from 1970 to 2009. Andreassen et al. (2005) used ratios of  
2 standard deviations of winter balances (sBw) to standard deviations of annual balances (sBa,  
3 sBw/sBa) and standard deviations of summer balances (sBs) to standard deviations of annual  
4 balances (sBs/sBa) to assess the relative importance of summer and winter balance for the  
5 annual balance. These ratios are direct measures of the relative importance of summer balance  
6 and winter balance for annual balances. Hence absolute influences of temperature and  
7 precipitation on annual balances as well as relative influences of winter and summer balance  
8 on annual balances have been assessed. In this study, we combine these two approaches and  
9 focus on determining relative and thereby directly comparable importances of winter  
10 precipitation and summer temperature for annual balances of glaciers in Scandinavia.

11 Assessing the relative importance of seasonally averaged summer temperature and winter  
12 precipitation for annual balances and possible changes in time, is especially interesting in  
13 light of palaeoclimatological interpretation of glacier records. In palaeoclimatology, at best  
14 summer temperature, winter precipitation and annual balance reconstructions are available.  
15 Attempts have been made to reconstruct winter precipitation based on glacier reconstructions  
16 and independent summer temperature reconstructions (e.g. Bakke et al. 2005).

17 There are well-known transient phases of positive annual balances (e.g. 1987 – 1995, e.g.  
18 Nesje et al. 2000). It is therefore interesting to assess if the relative importance of summer  
19 temperature and winter precipitation for annual balance changes through time. Until now,  
20 attempts of quantifying temporal changes of summer balance and winter balance on annual  
21 balance have been constrained to calculating running means of summer and winter balances  
22 and comparing the absolute values of these running means (e.g. Engelhardt et al. 2013).  
23 However, a direct assessment of temporal changes of the relative importance of summer  
24 temperature and winter precipitation for annual balances is still missing. Cumulative annual  
25 balances show clear patterns of consistently positive mass balances and thereafter consistently  
26 negative mass balances (e.g. Nesje et al. 2000, Fig. 3). We therefore hypothesise that the  
27 relative importance of summer temperature and winter precipitation for annual balances is not  
28 stable in time and that there is a large-scale forcing mechanism causing these changes. These  
29 forcings could either be of atmospheric or oceanic origin. It is for instance well known that  
30 increased amounts of winter precipitation in Scandinavia are associated with stronger zonal  
31 moisture advection that is due to pressure differences between Iceland and the Azores (e.g.  
32 Wanner et al. 2001). These pressure differences are summarized by the North Atlantic

1 Oscillation (NAO) Index. In addition to the atmosphere, systematic changes in ocean  
2 temperatures may also influence the relative importance of summer temperature and winter  
3 precipitation for annual balances of glaciers in Scandinavia. The Atlantic Multidecadal  
4 Oscillation (AMO) is a pattern of changing sea-surface temperatures in the North Atlantic  
5 (e.g. Schlesinger and Ramankutty, 1994). Changing sea surface temperatures might result in  
6 changing temperatures over land and thereby also alter the relative importance of summer  
7 temperature and winter precipitation for annual balances.

8 In this study, we focus on assessing the relative importance of winter precipitation and  
9 summer temperature for annual mass balances, temporal changes of these influences and on  
10 possible influences of large scale atmospheric and oceanic patterns on these temporal  
11 changes. The aims of this study are therefore threefold: i) model the annual mass balances of  
12 eight Scandinavian glaciers with long annual mass balance series using a suite of statistical  
13 models using seasonally averaged climate data as input variables. These models enable us to  
14 compare the relative importance of winter precipitation and summer temperature for annual  
15 mass balances of glaciers; ii) assessing temporal changes of relative importances of winter  
16 precipitation and summer temperature. iii) Compare these temporal changes to large-scale  
17 oceanic and atmospheric modes, such as the Atlantic Multidecadal Oscillation (AMO) and the  
18 North Atlantic Oscillation (NAO).

## 20 **2 Data and Methods**

### 21 **2.1 Data**

22 We modelled the mass balances of eight glaciers in Scandinavia, Ålfotbreen (ALF),  
23 Rembesdalskåka (REM), Nigardsbreen (NIG), Storbreen (STO), Hellstugubreen (HEL),  
24 Gråsubreen (GR) in southern Norway and Engabreen (ENG) and Storglaciären (STORGL) in  
25 northern Norway and northern Sweden, respectively (Fig. 1). Storglaciären has the longest  
26 annual mass balance time series, beginning in 1946 and Engabreen has the shortest time  
27 series, initiated in 1970. For all glaciers, data until 2010 was considered. Glacier mass balance  
28 data are available at [www.nve.no/bre](http://www.nve.no/bre) (Kjøllmoen, 2011; Andreassen and Winsvold, 2012)  
29 and [bolin.su.se/data/tarfala](http://bolin.su.se/data/tarfala). For all glaciers, winter balances, summer balances and annual  
30 balances are available. **Uncertainties of mass balance measurements and their possible sources**

1 are thoroughly discussed in Andreassen et al. (2005) and are estimated to between  $\pm 0.2$  and  
2  $\pm 0.4$  m w.e. per year.

3 Cumulative mass balance changes are shown in Figure 3. The three maritime glaciers  
4 Ålfotbreen (ALF), Rembesdalsskåka (REM), and Nigardsbreen (NIG) in southern Norway  
5 and the maritime glacier Engabreen (ENG) in northern Norway show positive cumulative net  
6 balances between the initiation of the measurements and 2010 (Fig. 3). Mass balances are  
7 especially positive during the first half of the 1990s. The continental glaciers Storbreen  
8 (STO), Hellstugubreen (HEL), and Gråsubreen (GR) in southern Norway and the continental  
9 glacier Storglaciären (STORGL) in northern Sweden experienced negative cumulative mass  
10 balances between the start of the measurements and 2010. For these glaciers the mass balance  
11 loss was reduced in the first half of the 1990s.

12 We used meteorological data from the meteorological station Bergen-Florida to model mass  
13 balances in southern Norway. We decided to exclusively use precipitation data from Bergen-  
14 Florida for all glaciers in southern Norway since Bergen-Florida records the large synoptic  
15 weather systems and is not affected by local topographic effects that are affecting  
16 meteorological stations in the deep and narrow valleys closer to the glaciers studied (e.g.  
17 Nesje 2005). For glaciers in northern Scandinavia, we used meteorological data from the  
18 coastal station Glomfjord available from the beginning of the mass balance series. The  
19 temperature measurements are continuous, but the precipitation series ends in 2003. We  
20 extended the precipitation series with data from the nearby Bodø meteorological station. The  
21 precipitation data from Bodø was scaled to the data from Glomfjord in the period of overlap  
22 (1953 - 2003) of the two data series.

## 23 **2.2 Methods**

24 To directly quantify the relative importances of summer temperature and winter precipitation  
25 on annual balances, we used a suite of three statistical models with increasing complexity and  
26 number of parameters that needed to be estimated:

- 27 i) Linear models using a climate index as independent variable
- 28 ii) Linear models using summer temperature and winter precipitation as independent variables
- 29 iii) Additive models using summer temperature and winter precipitation as independent  
30 variables

1 If the variance explained by two models was not significantly different, we favoured the  
2 simpler model, as it was more parsimonious.

3 As glaciers are mainly sensitive to summer temperatures and winter precipitation, models  
4 were calculated using one summer temperature and one winter precipitation as independent  
5 variables. We tested the influences of two summer temperatures, namely temperatures from  
6 May to September (T MJJAS) and temperatures from June to August (T JJA), and two winter  
7 precipitation variables, precipitation October to April (P ONDJFMA) and precipitation from  
8 November to March (P NDJFM) on annual glacier mass balances. This resulted in a total of  
9 four possible combinations of input variables. We chose the combination that resulted in  
10 lowest Akaike information criterion (AIC).

11

## 12 **2.2.1 Climate indices**

13 The simplest way of modelling the influence of (winter) precipitation and (summer)  
14 temperature on glacier mass balances is to generate a climate index, where winter  
15 precipitation and summer temperature are equally weighted (Imhof et al., 2012; Nesje, 2005),  
16 i.e. they are assigned the same relative importance for the annual balance. This was achieved  
17 by standardising summer temperature and winter precipitation and subtracting standardised  
18 summer temperature from standardised winter precipitation, as the two variables have  
19 opposed influences.

$$20 \quad z = \left( \frac{P - \bar{P}}{s_P} \right) - \left( \frac{T - \bar{T}}{s_T} \right) \quad (1)$$

$$21 \quad y = a + b \cdot z \quad (2)$$

22 Where  $z$  is the climate index,  $P$  winter precipitation,  $T$  summer temperature,  $s$  are standard  
23 deviations, bars denote means,  $y$  is the annual mass balance and  $a$  and  $b$  are regression  
24 coefficients.

25

## 26 **2.2.2 Linear models**

27 Annual mass balances were modelled using linear models with one (summer) temperature and  
28 one (winter) precipitation variable as independent variables. In a first step, we tested



1 interactions between (summer) temperature and (winter) precipitation and quadratic terms for  
2 significance. F-tests indicated that neither interaction terms, nor quadratic terms were  
3 significant ( $p < 0.05$ ).

4 The linear regression equation

$$5 \quad y = a + b_1 \cdot x_1 + b_2 \cdot x_2 \quad (3)$$

6 is interpreted as follows: if  $x_2$  is kept constant and  $x_1$  is changed by one unit,  $y$  changes by  $b_1$   
7 units (e.g. Legendre and Legendre 2012). Hence the regression coefficients of unscaled  
8 variables are also the climate sensitivity of this variable. Usually,  $x_1$  and  $x_2$  are measured in  
9 different units hampering the comparison of the influence of the two variables on  $y$ . This  
10 problem is, however, solved by standardising all the variables. The effect of standardisation is  
11 two-fold:

12 i) The intercept of the regression model is zero, and more importantly

13 ii) The standard regression coefficients are now comparable and are a 'means of assessing the  
14 relative importance of each explanatory variable  $x_j$  included in the regression model: the  
15 variables with the highest standard regression coefficient (in absolute values) are those that  
16 contribute the most to the estimated  $\hat{y}$  values' (Legendre and Legendre, 2012). In our case,  
17 using standardised annual balances, standardised winter precipitation and standardised  
18 summer temperature, the standard regression coefficients for winter precipitation and summer  
19 temperature are directly comparable and indicate the relative importance of summer  
20 temperature and winter precipitation for the annual mass balance.

21 For standardized variables, calculus with

$$22 \quad B = (X' \cdot X)^{-1} \cdot (X' \cdot Y) \quad (4)$$

23 as starting point (Legendre and Legendre, 2012), where  $X$  is a matrix of independent  
24 variables,  $Y$  is the dependent variable and  $B$  is a vector of coefficients linking  $X$  and  $Y$  in the  
25 regression equation proves that the standard regression coefficients are estimated as:

$$26 \quad b_1 = \frac{r_{x_1y} - r_{x_1x_2} \cdot r_{x_2y}}{1 - r_{x_1x_2}^2} \quad (5)$$

$$27 \quad b_2 = \frac{r_{x_2y} - r_{x_1x_2} \cdot r_{x_1y}}{1 - r_{x_1x_2}^2} \quad (6)$$

1

2 Where  $b_1$  and  $b_2$  are the standard regression coefficients of the first and second independent  
3 variable, respectively,  $r_{x_1y}$  is the correlation between the first independent and the dependent  
4 variable,  $r_{x_2y}$  is the correlation between the second independent variable and the dependent  
5 variable and  $r_{x_1x_2}$  is the correlation between the two independent variables.

6 Hence the standard regression coefficients, which are the relative importance of (in our case)  
7 winter precipitation and summer temperature for annual balance only depends on the  
8 correlations among winter precipitation, summer temperature and annual balance

9 The difference between linear models and the climate index is that winter precipitation and  
10 summer temperature are individually weighted when using linear models, whereas the two  
11 independent variables are equally weighted when employing the climate index. Hence, the  
12 relative importances of summer temperature and winter precipitation are allowed to be  
13 different using linear models, whereas they are artificially kept similar using climate index  
14 models. Linear models were compared to models based on climate indices using F-tests.

15 In contrast to p-values and confidence bounds, Bayesian credible intervals are simple to  
16 interpret. We used the simplest possible Bayesian model, namely setting a uniform prior for  
17 the two standard regression coefficients for winter precipitation and summer temperature.  
18 This results in posterior distributions for the parameter estimates that are proportional to the  
19 maximum likelihood estimates of the parameter values. Bayesian credible intervals are simple  
20 to interpret and indicate the parameter space within which a parameter is found with a certain  
21 probability. In this study, we interpreted the relative importance of summer temperature and  
22 winter precipitation as different, when the median of the posterior distribution of one  
23 parameter was outside the 2.5 and 97.5 percentiles of the posterior distribution of the other  
24 parameter.

25

### 26 **2.2.3 Additive models**

27 In contrast to linear models, where coefficients link independent and dependent variables, this  
28 linking is achieved by a smoothing curve in additive models

$$29 \quad y = a + f_1(x_1) + f_2(x_2) \quad (7)$$

1 (Zuur et al., 2009; Fig. 2). We used cubic regressions splines with three knots as smoothing  
2 terms. The number of knots was kept low to ensure monotony of the smoothing terms. The  
3 additive models were compared to linear models and climate index models by F- tests.

4 With the three statistical models proposed, we assume that errors in mass balance  
5 measurements are random and that climate data are error free. If the errors in mass balance  
6 measurements contain a systematic component, the estimates of relative importance of  
7 summer temperature and winter precipitation for annual balance are biased. If annual balances  
8 are systematically overestimated, the relative importance of summer temperature for annual  
9 balance is systematically underestimated.

#### 11 **2.2.4 Cross-Validation and analysis in running windows**

12 All the models were tested by calculating leave-one-out cross-validation (jack-knifing, e.g.  
13 Efron and Gong, 1983) and h-block cross-validation (Burman et al., 1994) where h-samples  
14 are left out on either side of the sample to be predicted. In this study we set h to 2. H-block  
15 cross-validation is a powerful method to test effects of temporal autocorrelation in time-  
16 series. However, preliminary autocorrelation calculations revealed no significant ( $p < 0.05$ )  
17 AR(1) autocorrelation coefficients. We calculated cross-validated mean absolute deviations  
18 and coefficients of determination.

19 After calculating models for the entire observation period, we wanted to assess if the relative  
20 importance of summer temperature and winter precipitation changed through time and if these  
21 changes were consistent among the glaciers. For this purpose, we ran models in 25-year  
22 moving windows. The significance of changes in variance explained was again tested with F-  
23 Tests. According to these tests, additive models were never superior to linear models.

#### 25 **2.2.5 Comparison to climate modes**

26 Preliminary analysis in running windows showed changes of relative importance of summer  
27 temperature and winter precipitation for annual balances that were consistent for all glaciers  
28 in southern Norway. We therefore assessed if these results were influenced by two large scale  
29 patterns of oceanic and atmospheric variability over the north Atlantic realm The North  
30 Atlantic Oscillation (NAO), an atmospheric pattern with an approximately decadal cyclicity

1 (Hurrell et al., 2001; Wanner et al., 2001) and the Atlantic Multidecadal Oscillation (AMO), a  
2 pattern in sea-surface temperature that is linked to changes in thermohaline ocean circulation  
3 with a cyclicity of 65 -70 years (Schlesinger and Ramankutty, 1994; Trenberth and Shea,  
4 2006). The NAO mainly influences the strength and tracks of the westerlies and thereby the  
5 amount of winter precipitation in north-western Europe.

6 Nesje et al. (2000) and Marzeion and Nesje (2012) found strong and significant ( $p < 0.05$ )  
7 correlations between NAO-index and annual mass balances of glaciers in southern Norway,  
8 with correlations decreasing with increasing distance to the coast. For northern Norway,  
9 Marzeion and Nesje found not significant or significantly negative ( $p < 0.05$ ) correlations  
10 between NAO-index and annual mass balances. In this study, we adopt a different approach to  
11 assess the influence of the NAO on annual mass balances. We wanted to assess if the relative  
12 importance of summer temperature and winter precipitation were depending on the NAO.  
13 Most of the glacier mass balance series investigated were shorter than 50 years. We therefore  
14 investigated the effects of changes in NAO by dividing the time series into two subsets with  
15 NAO-indices above and below the median of the NAO-index for the period in which mass  
16 balance measurements were available. We then estimated the relative importance of summer  
17 temperature and winter precipitation for the annual mass balance for these two subsets. We  
18 also wanted to assess if there were differences between the correlations between the NAO-  
19 index and winter mass balances and annual balances for years with above and below median  
20 NAO-index. We also used the ratio of the standard deviation of the winter balance to the  
21 standard deviation of the annual balance ( $sBw/sBa$ ) and the ratio of the standard deviation of  
22 the winter balance to the standard deviation of the annual balance ( $sBs/sBa$ ) (e.g. Andreassen  
23 et al. (2005)) to see if these ratios were different for mass balance data of years with above  
24 and below median NAO-index.

25 Considering the period 1946 – 2010, the average monthly November through April  
26 precipitation in Bergen was 230 mm for the years with above median NAO-index and 170  
27 mm in the years with below median NAO-index, which is significantly lower ( $p < 0.05$ ).

28 The longest mass balance series started in 1946. The AMO was generally positive from ca.  
29 1930 to 1962 and from 1997 to the present, whereas it was negative between 1963 and 1996.  
30 In the negative subset of the AMO, the correlation between the NAO-index and extended  
31 winter precipitation in Bergen was  $r = 0.82$  ( $p < 0.05$ ), whereas it was  $r = 0.56$  ( $p < 0.05$ ) for  
32 the years with predominantly positive AMO-index. The average November through April

1 precipitation in Bergen was not differing between the two subsets (200mm/month). The  
2 average May through September temperature from Bergen-Florida for the positive AMO  
3 subset was 14.4°C, whereas it was 12.6°C in the negative AMO subset. Average T MJJAS for  
4 the period 1949 – 1962 was 13.8°C, which is as well significantly ( $p < 0.05$ ) higher than the  
5 average temperature in the negative AMO subset. As summer temperatures in Bergen were  
6 significantly ( $p < 0.05$ ) higher in the positive AMO subset, we wanted to test if this altered the  
7 relative importance of summer temperature and winter precipitation for annual balances. This  
8 analysis was only carried out for the two long data series starting in 1946 and 1949. The data  
9 series were divided into two subsets of years of predominantly positive (1946/1949 – 1962,  
10 1997 – 2010) and negative (1963 – 1996) AMO. We also estimated the ratios  $sBw/sBa$  and  
11  $sBs/sBa$  (e.g. Andreassen et al. 2005) with AMO+ and AMO-.

12 All calculations were done in R (R Core Team 2014) and its add-on packages lmodel2  
13 (Legendre, 2014), and mgcv (Wood, 2014).

14

### 15 **3 Results**

#### 16 **3.1 Model performance**

17 The employed statistical models explained large proportions of the variance of annual mass  
18 balances (Table 1). For the maritime glaciers, the models explained more than 70% of the  
19 variance. The variance explained for continental glaciers varied between 50% and 70%. Table  
20 1 shows input variables, model types, variance explained by the most parsimonious models  
21 and standard regression coefficients of linear models (i.e. the relative importance of summer  
22 temperature and winter precipitation) and their Bayesian credible intervals. Cross-validated  $r^2$   
23 using leave-one-out cross-validation and h-block cross-validation were comparable to  
24 apparent  $r^2$ . The only exception was Ålfotbreen, where an additive model was most  
25 parsimonious. Cross-validated  $r^2$  was reduced by 0.1, i.e. the variance explained was reduced  
26 by 10% and linear models had higher  $r^2$  under cross-validation. Cross-validated mean  
27 absolute deviations were also lowest for the models chosen, except for Ålfotbreen where  
28 again linear models yielded lowest mean absolute deviations.

### 3.2 Relative importance of summer temperature and winter precipitation for the annual mass balance

For Storbreen, Engabreen and Storglaciären, the statistical models using climate indices as input variables were most parsimonious. These are the only glaciers where standard regression coefficients of linear models were not different. Hence, linear models were also assigning about similar weights to summer temperature and winter precipitation for these three glaciers. For the maritime glaciers Rembesdalsskåka and Nigardsbreen, linear models indicated a higher relative importance of winter precipitation than of summer temperature, whereas for the continental glaciers Hellstugubreen and Gråsubreen, the relative importance of summer temperature was higher than the relative importance of winter precipitation. For the maritime Ålfotbreen, an additive model was explaining significantly ( $p < 0.05$ ) more of the total variance than a linear model. The smooth terms of summer temperature and winter precipitation are shown in Figure 2. The slope of the smooth for temperature was flatter than the slope of a linear model for below average temperatures and steeper than the slope of a linear model for above average temperatures. Hence the expected sensitivity of the annual mass balance for a change of  $1^{\circ}\text{C}$  increased with increasing temperatures. In contrast, the slope of the smooth for precipitation was steeper than the slope of a linear model for below average precipitation values and was flatter than the slope of a linear model for above average precipitation levels. The expected sensitivity of the annual mass balance for a change in precipitation decreased with increasing precipitation.

### 3.3 Temporal changes in the relative importance of summer temperature and winter precipitation

Temporal changes of relative importance of summer temperature and winter precipitation are shown in Fig. 3b-i. The relative importance of winter precipitation, as indicated by standard regression coefficients of winter precipitation in 25-year running windows, was lowest at the end of the observation period. The relative importance of summer temperature, as indicated by standard regression coefficients of summer temperature in 25-year running windows, increased towards the end of the observation period (Fig. 3b-i).

Winter precipitation was more important than summer temperature for the annual balance of continental glaciers in southern Norway (STO, HEL, and GR) for the 25-year windows

1 centred between 1977 and 1985. For STO, the period of higher relative importance of winter  
2 precipitation than relative importance of summer temperature was extended up to the 25-year  
3 window centred around 1990 (Fig 3e). For the maritime glaciers in southern Norway, the  
4 Bayesian credible intervals of the standard regression coefficients (relative importances) were  
5 not overlapping for 25-year windows centred before 1990, but were overlapping for the last  
6 five running windows.

7 Storbreen indicated about equal importance of winter precipitation and summer temperature  
8 for 25-year windows ending prior to 1990 (Fig. 3e). The relative importance of summer  
9 temperature was higher than the relative importance of winter precipitation for 25-year  
10 windows ending in the 1980s for Storglaciären (Fig. 3i).

### 11 **3.4 Influences of NAO and AMO on annual balances**

12 The mass balance models for years with above and below median NAO-index, respectively,  
13 were different in terms of variance explained and in terms of relative importance assigned to  
14 summer temperature and winter precipitation. They also differed from models covering the  
15 entire measurement period.

16 For years with above median NAO, models for Ålfotbreen, Rembesdalsskåka, Nigardsbreen  
17 and Storbreen explained as much of the variance of the mass balance as models for the entire  
18 data series, whereas for Hellstugubreen and Gråsubreen, the variance explained was reduced  
19 compared to the models for the entire period. Interestingly, for Ålfotbreen standard regression  
20 coefficients for winter precipitation and summer temperature were not different. For the phase  
21 with below median NAO-index, models for Ålfotbreen, Rembesdalsskåka and Nigardsbreen  
22 explained less of the variance than in the entire period and standard regression coefficients for  
23 precipitation and temperature were not different, whereas models for Gråsubreen and  
24 Hellstugubreen explained more of the variance than in the entire period, and together with  
25 Storbreen displayed a higher importance of summer temperature than winter precipitation.  
26 The two glaciers with long data series had an average mass loss of 0.54 m water equivalents  
27 per year (m w.e./yr.) when the NAO-index was low, but an average gain of 0.03 m w.e./yr. for  
28 Storglaciären and an average loss of 0.08 m w.e./yr. for Storbreen with high NAO-index.

29 For all glaciers, except for ALF, the ratio  $sBs/sBa$  was lower in years with above median  
30 NAO-index than for the entire data series and the ratio  $sBw/sBa$  was higher than for the entire  
31 data series for REM, STO, HEL, GR and STORGL (Fig 4). For years with below median

1 NAO-index, the ratio  $sBs/sBa$  was higher than in the entire data series and  $sBw/sBa$  was  
2 lower than in the entire data series except for ALF and ENG (Fig. 4).

3 Correlations between NAO-index and winter and annual balance were different for the  
4 subsets of years with above and below median NAO-index (Fig. 5). For glaciers in southern  
5 Norway, the correlation between NAO-index and winter and annual balance was higher than  
6 for the entire time series for years with above median NAO-index and was lower than for the  
7 entire series for years with below median NAO-index.

8 Changes in relative importances of winter precipitation and summer temperature were also  
9 found for the AMO+ and AMO- phases. The mass balance models for positive and negative  
10 AMO were differing for Storbreen in southern Norway (Table 1), whereas they remained  
11 unchanged for Storglaciären in northern Sweden. For Storbreen, the influence of winter  
12 precipitation was significantly higher than the influence of summer temperature with negative  
13 AMO-index, whereas the situation was opposite with positive AMO-index (Table 1). For both  
14 glaciers, the average annual mass balance was different in the two phases defined by positive  
15 and negative AMO indices: Storbreen lost an average of 0.5 m w.e./yr. and Storglaciären 0.48  
16 m w.e./yr. when the AMO-index was positive, whereas the loss was reduced to averages of  
17 0.15 m w.e./yr. and 0.02 m w.e./yr. for Storbreen and Storglaciären, respectively, when the  
18 AMO-index was negative. The AMO also affected the standard deviation ratios. For  
19 Storbreen, the ratios  $sBs/sBa$  and  $sBw/sBa$  were equal when the AMO was in its negative  
20 phase (Fig 4). During the positive phase of the AMO,  $sBs/sBa$  was higher than  $sBw/sBa$ .

21

22

## 23 4 Discussion

### 24 4.1 Model Performance

25 We used simple statistical models that are only taking into account summer temperature and  
26 winter precipitation to model annual mass balances. Even though these models are simplistic,  
27 they explain large proportions of the variance of annual balances, and are therefore  
28 appropriate to estimate relative importance of summer temperature and winter precipitation  
29 for annual balances. Climate sensitivities of Engabreen (Schuler et al. 2005),  
30 Rembesdalsskåka (Giesen and Oerlemans, 2011) and Storbreen (Andreassen and Oerlemans,



1 2009) show that summer balances are largely unaffected by changes in precipitation, which  
2 suggest minor importance of summer precipitation for summer balance. Still other important  
3 components such as the direct effect of radiation are not entirely accounted for when only  
4 using summer temperature to model ablation.

5 The model performance is increased for coastal maritime glaciers. This might have several  
6 reasons: i) precipitation is highly variable in space and therefore precipitation from Bergen is  
7 possibly more appropriate for coastal glaciers than for continental glaciers. Still, using  
8 precipitation from meteorological stations closer to the continental glaciers did not improve  
9 the model performance for continental glaciers. ii) Processes not represented in our model are  
10 more important in summer (radiation) than in winter (wind redistribution of snow).

## 12 **4.2 Relative importance of summer temperature and winter precipitation**

13 Our results showed, as also demonstrated in other studies (Andreassen and Oerlemans, 2009;  
14 Giesen and Oerlemans, 2010; Laumann and Nesje, 2009a, 2009b, 2014; Oerlemans, 1992),  
15 that the annual glacier mass balance on near coastal, maritime glaciers was mainly controlled  
16 by winter precipitation and that the annual mass balance on the inland, continental glaciers  
17 was mainly controlled by summer temperature (Andreassen et al., 2005; Nesje et al., 1995).  
18 Hence standard regression coefficients of linear models are proved to be good estimators of  
19 the relative importance of temperature and precipitation for annual balances. The relative  
20 importance as determined by standard regression coefficients display similar patterns as the  
21 standard deviation ratios presented by Andreassen et al. (2005) and also shown in Fig 4. The  
22 exceptions are NIG, and STO. For NIG, standard regression coefficients indicate higher  
23 relative importance of winter precipitation compared to summer temperature, but standard  
24 deviation ratios are similar. Standard regression coefficients suggest equal relative importance  
25 of summer temperature and winter precipitation for STO, whereas the standard deviation ratio  
26  $sBs/sBa$  is higher than  $sBw/sBa$ . For both NIG and STO, climate sensitivities estimated by de  
27 Woul and Hock (2005) and Rasmussen and Conway (2005) using degree day models differ:  
28 de Woul and Hock (2005) estimate the precipitation increase needed to level a temperature  
29 increase of 1 K to be 30% and 50% for NIG and STO, respectively, whereas Rasmussen and  
30 Conway found lower values of 25% and 28%. Engelhardt et al. (2013) also modelled mass  
31 balances of NIG and STO using degree day models. Modelled annual balances showed a

1 strong positive bias for NIG and a strong negative bias for STO. Hence assessing the relative  
2 importance of winter precipitation and summer temperature on annual balances of NIG and  
3 STO seems difficult.

#### 4 **4.3 Temporal changes of relative importance of summer temperature and** 5 **winter precipitation and their relation to large scale oceanic and** 6 **atmospheric patterns**

7 As shown in this study, the relative importance of summer temperature and winter  
8 precipitation for annual balances is not constant in time. Temporal changes in relative  
9 importance of summer temperature and winter precipitation are consistent for entire southern  
10 Norway (Fig. 3), suggesting common large scale forcing of the relative importance of summer  
11 temperature and winter precipitation.

12 Maritime glaciers had a consistently positive mass balance between 1988 and 1996 and  
13 continental glaciers were no longer losing mass (Fig. 3, Nesje et al., 2000; Andreassen et al.  
14 2005; Nesje and Matthews, 2012). Looking at the 25-year windows centred between 1978 and  
15 1984, we found that winter precipitation was more important than summer temperature for all  
16 glaciers including the continental glaciers in southern Norway, although the differences were  
17 not significant for the continental Gråsubreen. For the three continental glaciers in southern  
18 Norway, this phase was characterised by a marked decrease in relative importance of summer  
19 temperature and a marked increase in relative importance of winter precipitation.

20 In this phase, the AMO-index was consistently negative and the NAO-indexes were  
21 consistently positive between 1988 and 1996 (Fig. 3). In tendency, negative AMO indices  
22 were associated with reduced summer temperatures over Europe and positive NAO-indexes  
23 were associated with increased zonal flow in winter, entailing more winter precipitation in  
24 Northern Europe. Hence, the large-scale oceanic and atmospheric patterns were favourable for  
25 glacier growth.

26 As another example, in the 2000s all glaciers except Engabreen and Nigardsbreen generally  
27 experienced negative mass balances and mass balances of Engabreen and Nigardsbreen were  
28 at equilibrium. In this period, the importance of summer temperature for the annual mass  
29 balance was increased (Fig. 3), even though these 25-year windows still contained the years  
30 1988 – 1996 with their transient mass surplus. The increasing relative importance of summer  
31 temperature and decreasing relative importance of winter precipitation for the annual balance

1 at the end of the measurement period is consistent with more negative summer balances and  
2 less positive winter balances found for glaciers in southern Norway (e.g. Engelhardt et al.  
3 2013). The AMO-index changed sign in the late 1990s and summer temperatures were in  
4 general higher than between 1985 and 1995.

5 For glaciers in the European Alps, Huss et al. (2010) found pronounced mass loss during  
6 phases of positive AMO-index and mass gain in phases of negative AMO-index, which is  
7 similar to finds in this study. The phases of increased glacier melt are, however, not  
8 simultaneous in the Swiss Alps and in Scandinavia. In the Swiss Alps, a pronounced mass  
9 loss lasting to the present started in the late 1980s, whereas continental glaciers in  
10 Scandinavia lost mass between the start of the measurements and 1987 and all glaciers in  
11 Scandinavia lost mass after about 1998. This difference is most probably caused by the fact  
12 that changes in melt rates are most influential for mass balances in the Alps (Huss et al.,  
13 2010), whereas a decade with predominantly positive NAO-indexes began in the late 1980s  
14 (1988/89 winter) associated with increased relative importance of winter precipitation for  
15 Scandinavian glaciers (Fig. 3). For a more detailed comparison of the influences of the NAO  
16 on mass balances in the Alps and in Scandinavia, the reader is referred to Six et al. (2001) and  
17 Marzeion and Nesje (2012).

#### 19 **4.4 Direct influence of NAO and AMO on annual mass balances**

20 Clear differences are found between the subsets with above median and below median NAO-  
21 index. In winters with high NAO-index, stronger westerly flow and increased precipitation is  
22 expected (e.g. Wanner et al. 2001). The mass balance models of the maritime glaciers  
23 explained more of the total variance with high NAO-index and the relative importance of  
24 winter precipitation for the total mass balance was increased. This was according to  
25 expectations, as increased winter precipitation is expected to increase the importance of the  
26 winter precipitation for mass balance models.

27 For all glaciers, the correlation between NAO-index and winter and annual mass balance was  
28 higher for years with above median NAO-index (Fig. 5). Additionally, the coefficient of  
29 determination between winter balance and NAO-index was decreased for the subset of years  
30 with below median NAO-index (Fig. 5). This means that the reduction in coefficient of  
31 determination between NAO-index and annual balance was not only caused by an increased

1 importance of the summer balance for the annual balance, but also by a loss of accordance  
2 between NAO-index and winter balance. This loss in accordance is only partly caused by  
3 lower accordance among precipitation in Bergen and winter balances, but mainly by a  
4 consistently decreased correlation between the NAO-index and precipitation in Bergen.  
5 Consequently the NAO-index is only a good predictor for winter balances of glaciers in  
6 southern Norway in years with above median NAO-index. This is reiterating a find by Six et  
7 al. (2001), who do not recommend to model glacier mass balances solely based on the NAO-  
8 index. Unstable relations between the NAO-index and glacier length changes in Scandinavia  
9 as well as in the Alps were also found by Imhof et al. (2011).

10 For the two glaciers with long mass balance time-series, the influence of the NAO seemed  
11 equal to the influence of the AMO, as the difference between the average mass balances in the  
12 two NAO levels considered was about equal to the difference in the two AMO states. The  
13 AMO states only include consecutive years, whereas individual years were assigned to the  
14 NAO-index. The phase between ca. 1987 and 1995 with major mass gain for maritime  
15 glaciers and neutral mass balances for continental glaciers was characterised by negative  
16 AMO-index and predominantly positive NAO-index, that were both favourable for glaciers.

17 The relation between AMO and NAO seems rather complex and depends on the time scale  
18 considered (Li et al., 2013; Peings and Magnusdottir, 2014). On short time scales, the  
19 atmospheric NAO pattern influences the sea surface temperature, whereas on longer time  
20 scales, the sea-surface temperature AMO pattern drives the atmospheric NAO. Hence Li et al.  
21 (2013) find the NAO to lead the AMO by 16 years and state that the NAO is an excellent  
22 predictor for AMO and thereby Northern Hemisphere temperature, whereas Peings and  
23 Magnusdottir (2014) find ‘that the multidecadal fluctuations of the wintertime North Atlantic  
24 Oscillation (NAO) are tied to the AMO, with an opposite signed relationship between the  
25 polarities of the AMO and the NAO. Our statistical analyses suggest that the AMO signal  
26 precedes the NAO by 10–15 years ...’

27 The association of negative AMO and positive NAO seems to be typical (Peings and  
28 Magnusdottir 2014), whereas positive AMO favours negative NAO and blocking situations.  
29 For the time period 1965 – 1998, with negative AMO, only 10 years have a negative NAO-  
30 index, whereas for the considerably shorter phase 1999 – 2010 already 6 years had a negative  
31 NAO-index. Hence, the two modes favouring glacier mass gain and mass loss, respectively,  
32 tended to occur simultaneously. However, the influence of AMO and NAO should not be

1 overestimated, as similar weather patterns still result in different amounts of precipitation and  
2 in different levels of temperature (Jacobeit et al., 2003; Kuettel et al., 2011). Kuettel et al.  
3 (2011), for instance, attribute 60% of the changes of winter precipitation over southern  
4 Norway between the periods 1900-1949 and 1950-1999 to changes within weather patterns  
5 and only 40% to changes in frequencies of weather patterns.

6

## 7 **5 Conclusions**

8 We used simple statistical models to assess the relative importance of summer temperature  
9 and winter precipitation for annual balances of eight glaciers in Scandinavia. The relative  
10 importances found using statistical models were comparable to estimates of relative  
11 importance obtained using different methods. Most importantly, the relative importance of  
12 summer temperature and winter precipitation for annual balances varied through time. Winter  
13 precipitation was most important when the Atlantic Multidecadal Oscillation Index was  
14 negative and the North Atlantic Oscillation Index was positive. Towards present, the relative  
15 importance of winter precipitation decreased for all glaciers while the relative importance of  
16 summer temperature was increasing. The influence of NAO and AMO on the relative  
17 importance of summer temperature and winter precipitation for annual balance was confirmed  
18 considering subsets of different NAO and AMO levels, with increasing relative importance of  
19 winter precipitation in years with NAO+ and AMO- and increased relative importance of  
20 summer temperature in years with AMO+ and NAO-.

21

## 22 **Acknowledgements**

23 We would like to thank two reviewers and Pascal Hänggi for comments on an earlier version  
24 of this manuscript.

25

## 1 **References**

- 2 Andreassen, L.M., Elvehøy, H., Johannesson, T., Oerlemans J., Beldring S. and van den  
3 Broeke M.: Modelling the climate sensitivity of Storbreen and Engabreen, Norway.  
4 Norwegian Water Resources and Energy Directorate. Report no. 3, NVE, Oslo, 2006.
- 5 Andreassen, L. M., Elvehoy, H., Kjollmoen, B., Engeset, R. V. and Haakensen, N.: Glacier  
6 mass-balance and length variation in Norway, *Ann. Glaciol.*, 42, 317–325, 2005.
- 7 Andreassen, L. M. and Oerlemans, J.: Modelling Long-Term Summer and Winter Balances  
8 and the Climate Sensitivity of Storbreen, Norway, *Geogr. Ann. Ser. -Phys. Geogr.*, 91A, 233–  
9 251, 2009.
- 10 Andreassen, L.M. and Winsvold, S.H.: Inventory of Norwegian Glaciers. Norwegian Water  
11 Resources and Energy Directorate. Rapport 38-2012, NVE, Oslo, 2012.
- 12 Bakke, J., Dahl, S. O., Paasche, O., Lovlie, R. and Nesje, A.: Glacier fluctuations,  
13 equilibrium-line altitudes and palaeoclimate in Lyngen, northern Norway, during the  
14 Lateglacial and Holocene, *Holocene*, 15(4), 518–540, doi:10.1191/0959683605hl815rp, 2005.
- 15 Burman, P., Chow, E. and Nolan, D.: A Cross-Validatory Method for Dependent Data,  
16 *Biometrika*, 81, 351–358, doi:10.1093/biomet/81.2.351, 1994.
- 17 Efron, B. and Gong, G.: A Leisurely Look at the Bootstrap, the Jackknife, and Cross-  
18 Validation, *Am. Stat.*, 37(1), 36–48, doi:10.2307/2685844, 1983.
- 19 Engelhardt, M., Schuler, T. V. and Andreassen, L. M.: Glacier mass balance of Norway 1961-  
20 2010 calculated by a temperature-index model, *Ann. Glaciol.*, 54(63), 32–40,  
21 doi:10.3189/2013AoG63A245, 2013.
- 22 Giesen, R. H. and Oerlemans, J.: Response of the ice cap Hardangerjokulen in southern  
23 Norway to the 20th and 21st century climates, *Cryosphere*, 4, 191–213, doi:10.5194/tc-4-191-  
24 2010, 2010.
- 25 Hock, R., Radic, V. and De Woul, M.: Climate sensitivity of Storglaciaren, Sweden: an  
26 intercomparison of mass-balance models using ERA-40 re-analysis and regional climate  
27 model data, *Ann. Glaciol.*, 46, , 2007.

- 1 Hurrell, J. W., Kushnir, Y. and Visbeck, M.: The North Atlantic Oscillation, *Science*,  
2 291(5504), 603–605, doi:10.1126/science.1058761, 2001.
- 3 Huss, M., Hock, R., Bauder, A. and Funk, M.: 100-year mass changes in the Swiss Alps  
4 linked to the Atlantic Multidecadal Oscillation, *Geophys. Res. Lett.*, 37, L10501,  
5 doi:10.1029/2010GL042616, 2010.
- 6 Imhof, P., Nesje, A. and Nussbaumer, S. U.: Climate and glacier fluctuations at  
7 Jostedalsbreen and Folgefonna, southwestern Norway and in the western Alps from the  
8 “Little Ice Age” until the present: The influence of the North Atlantic Oscillation, *Holocene*,  
9 22, 235–247, 2012.
- 10 IPCC: Climate Change 2007: The Physical Science Basis Working Group I Contribution to  
11 the Fourth Assessment Report of the IPCC, Camb. Univ. Press [online] Available from:  
12 [http://www.cambridge.org/no/academic/subjects/earth-and-environmental-](http://www.cambridge.org/no/academic/subjects/earth-and-environmental-science/climatology-and-climate-change/climate-change-2007-physical-science-basis-working-group-i-contribution-fourth-assessment-report-ipcc)  
13 [science/climatology-and-climate-change/climate-change-2007-physical-science-basis-](http://www.cambridge.org/no/academic/subjects/earth-and-environmental-science/climatology-and-climate-change/climate-change-2007-physical-science-basis-working-group-i-contribution-fourth-assessment-report-ipcc)  
14 [working-group-i-contribution-fourth-assessment-report-ipcc](http://www.cambridge.org/no/academic/subjects/earth-and-environmental-science/climatology-and-climate-change/climate-change-2007-physical-science-basis-working-group-i-contribution-fourth-assessment-report-ipcc) (Accessed 13 December 2014),  
15 2007.
- 16 IPCC: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I  
17 to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, Camb.  
18 Univ. Press [online] Available from: [http://www.cambridge.org/no/academic/subjects/earth-](http://www.cambridge.org/no/academic/subjects/earth-and-environmental-science/climatology-and-climate-change/climate-change-2013-physical-science-basis-working-group-i-contribution-fifth-assessment-report-intergovernmental-panel-climate-change?format=PB)  
19 [and-environmental-science/climatology-and-climate-change/climate-change-2013-physical-](http://www.cambridge.org/no/academic/subjects/earth-and-environmental-science/climatology-and-climate-change/climate-change-2013-physical-science-basis-working-group-i-contribution-fifth-assessment-report-intergovernmental-panel-climate-change?format=PB)  
20 [science-basis-working-group-i-contribution-fifth-assessment-report-intergovernmental-panel-](http://www.cambridge.org/no/academic/subjects/earth-and-environmental-science/climatology-and-climate-change/climate-change-2013-physical-science-basis-working-group-i-contribution-fifth-assessment-report-intergovernmental-panel-climate-change?format=PB)  
21 [climate-change?format=PB](http://www.cambridge.org/no/academic/subjects/earth-and-environmental-science/climatology-and-climate-change/climate-change-2013-physical-science-basis-working-group-i-contribution-fifth-assessment-report-intergovernmental-panel-climate-change?format=PB) (Accessed 13 December 2014), 2013.
- 22 Jacobeit, J., Wanner, H., Luterbacher, J., Beck, C., Philipp, A. and Sturm, K.: Atmospheric  
23 circulation variability in the North-Atlantic-European area since the mid-seventeenth century,  
24 *Clim. Dyn.*, 20, 341–352, 2003.
- 25 Jones, P. D., Jonsson, T. and Wheeler, D.: Extension to the North Atlantic Oscillation using  
26 early instrumental pressure observations from Gibraltar and south-west Iceland, *Int. J.*  
27 *Climatol.*, 17, 1433–1450, 1997.
- 28 Kjøllmoen, B. (ed.): Glaciological investigations in Norway 2010. Norwegian Water  
29 Resources and Energy Directorate. Rapport, 3.NVE, Oslo, 2011.

- 1 Kuettel, M., Luterbacher, J. and Wanner, H.: Multidecadal changes in winter circulation-  
2 climate relationship in Europe: frequency variations, within-type modifications, and long-  
3 term trends, *Clim. Dyn.*, 36(5-6), 957–972, , 2011.
- 4 Laumann, T. and Nesje, A.: A simple method of simulating the future frontal position of  
5 Briksdalsbreen, western Norway, *Holocene*, 19, 221–228, 2009a.
- 6 Laumann, T. and Nesje, A.: The impact of climate change on future frontal variations of  
7 Briksdalsbreen, western Norway, *J. Glaciol.*, 55, 789–796, 2009b.
- 8 Laumann, T. and Nesje, A.: Sporteggbreen, western Norway, in the past, present and future:  
9 Simulations with a two-dimensional dynamical glacier model, *Holocene*, 24, 842–852, 2014.
- 10 Legendre, P.: lmodel2: Model II Regression. [online] Available from: [http://cran.r-](http://cran.r-project.org/web/packages/lmodel2/index.html)  
11 [project.org/web/packages/lmodel2/index.html](http://cran.r-project.org/web/packages/lmodel2/index.html) (Accessed 13 December 2014), 2014.
- 12 Legendre, P. and Legendre, L.: *Numerical Ecology*, 3rd ed., Elsevier, Amsterdam., 2012.
- 13 Li, J., Sun, C. and Jin, F.-F.: NAO implicated as a predictor of Northern Hemisphere mean  
14 temperature multidecadal variability, *Geophys. Res. Lett.*, 40, 5497–5502,  
15 doi:10.1002/2013GL057877, 2013.
- 16 Marzeion, B. and Nesje, A.: Spatial patterns of North Atlantic Oscillation influence on mass  
17 balance variability of European glaciers, *Cryosphere*, 6(3), 661–673, doi:10.5194/tc-6-661-  
18 2012, 2012.
- 19 Nesje, A.: Briksdalsbreen in western Norway: AD 1900-2004 frontal fluctuations as a  
20 combined effect of variations in winter precipitation and summer temperature, *Holocene*, 15,  
21 1245–1252, 2005.
- 22 Nesje, A.: Latest Pleistocene and Holocene alpine glacier fluctuations in Scandinavia, *Quat.*  
23 *Sci. Rev.*, 28(21-22), 2119–2136, doi:10.1016/j.quascirev.2008.12.016, 2009.
- 24 Nesje, A., Bakke, J., Dahl, S. O., Lie, O. and Matthews, J. A.: Norwegian mountain glaciers  
25 in the past, present and future, *Glob. Planet. Change*, 60, 10–27, 2008.



- 1 Nesje, A., Johannessen, T. and Birks, H.: Briksdalsbreen, Western Norway - Climatic Effects  
2 on the Terminal Response of a Temperate Glacier Between AD 1901 and 1994, *Holocene*, 5,  
3 343–347, 1995.
- 4 Oerlemans, J.: Climate Sensitivity of Glaciers in Southern Norway - Application of an  
5 Energy-Balance Model to Nigardsbreen, Hellstugubreen and Alftobreen, *J. Glaciol.*, 38, 223–  
6 232, 1992.
- 7 Oerlemans, J.: A flowline model for Nigardsbreen, Norway: projection of future glacier  
8 length based on dynamic calibration with the historic record, *Ann. Glaciol.*, 24, 382–9, 1997.
- 9 Peings, Y. and Magnúsdóttir, G.: Forcing of the wintertime atmospheric circulation by the  
10 multidecadal fluctuations of the North Atlantic ocean, *Environ. Res. Lett.*, 9, 034018,  
11 doi:10.1088/1748-9326/9/3/034018, 2014.
- 12 Rasmussen, L. A., Andreassen, L. M. and Conway, H.: Reconstruction of mass balance of  
13 glaciers in southern Norway back to 1948, *Ann. Glaciol.*, 46, 255–60, 2007.
- 14 Rasmussen, L. A. and Conway, H.: Influence of upper-air conditions on glaciers in  
15 Scandinavia, in *Annals of Glaciology*, *Ann. Glaciol.*, 42, 402–408, 2005.
- 16 Schlesinger, M. and Ramankutty, N.: An Oscillation in the Global Climate System of Period  
17 65-70 Years, *Nature*, 367(6465), 723–726, doi:10.1038/367723a0, 1994.
- 18 Schuler, T. V., Hock, R., Jackson, M., Elvehøy, H., Braun, M., Brown, I. and Hagen, J.-O.:  
19 Distributed mass-balance and climate sensitivity modelling of Engabreen, Norway, *Ann.*  
20 *Glaciol.*, 42, 395–401, 2005.
- 21 Six, D., Reynaud, L. and Letreguilly, A.: Alpine and Scandinavian glaciers mass balances,  
22 their relations with the North Atlantic Oscillation., *Comptes Rendus Acad. Sci. Ser. II Fasc. -*  
23 *Sci.*, 333(11), 693–698, doi:10.1016/S1251-8050(01)01697-4, 2001.
- 24 Trenberth, K. E. and Shea, D. J.: Atlantic hurricanes and natural variability in 2005, *Geophys.*  
25 *Res. Lett.*, 33, L12704, doi:10.1029/2006GL026894, 2006.
- 26 Wanner, H., Bronnimann, S., Casty, C., Gyalistras, D., Luterbacher, J., Schmutz, C.,  
27 Stephenson, D. B. and Xoplaki, E.: North Atlantic Oscillation - Concepts and studies, *Surv.*  
28 *Geophys.*, 22, 321–382, doi:10.1023/A:1014217317898, 2001.

1 Wood, S.: mgcv: Mixed GAM Computation Vehicle with GCV/AIC/REML smoothness  
2 estimation. [online] Available from: <http://cran.r-project.org/web/packages/mgcv/index.html>  
3 (Accessed 13 December 2014), 2014.

4 De Woul, M. and Hock, R.: Static mass-balance sensitivity of Arctic glaciers and ice caps  
5 using a degree-day approach, *Ann. Glaciol.*, 42, 217–224, 2005.

6 Zuur, A. F., Ieno, E. N., Walker, N. J., Saveliev, A. A. and Smith, G. M.: *Mixed Effects*  
7 *Models and Extensions in Ecology with R*, Springer, Berlin., 2009.

8

9

1 Table 1. Table of most parsimonious statistical models. Input variables used and model types  
2 are indicated along with apparent and cross-validated variance explained. Cross-validated  
3 mean absolute deviations and relative importance of summer temperature (LM Coef T) and  
4 winter precipitation (LM Coef T) are indicated along with uncertainties of estimates of relative  
5 importances. Relative importance of summer temperature and winter precipitation and  
6 apparent variance explained are also indicated for subsets only including years with above  
7 (NAO+) and below (NAO-) median NAO-index, years with negative AMO-index (AMO-)  
8 and for STO and STORGL years with positive AMO-index (AMO+). ALF (Ålfotbreen),  
9 REM (Rembesdalsskåka), NIG (Nigardsbreen), STORBR (Storbreen), HEL (Hellstugubreen),  
10 GR (Gråsubreen), ENG (Engabreen), STORGL (Storglaciären), Am : Additive Model, LM:  
11 Linear Model, CI: Climate Index, NAO: North Atlantic Oscillation, AMO: Atlantic  
12 Multidecadal Oscillation.

13


14 Only available in separate file.

15

16

17


1



2


3 Figure 1. Map of glaciers and annual means of temperature and precipitation. Glaciers:  
4 Ålfotbreen (ALF), Rembesdals-skåka (REM), Nigardsbreen (NIG), Storbreen (STO),  
5 Hellstugubreen (HEL), Gråsubreen (GR), Engabreen (ENG) and Storglaciären (STORGL).  
6 1961-90 normal mean annual temperature (b) and mean annual precipitation (c)  
7 (<http://met.no/Klima/Klimastatistikk>).

8



1

2 **Figure 2. Additive model for Ålfotbreen. a) Smooth term (S (T MJJAS); black) and linear**  
 3 **model (red) for summer temperature (T MJJAS). b) Smooth term (S (P NDJFM); black) and**  
 4 **linear model (red) for winter precipitation (P NDJFM). Dotted lines indicate confidence**  
 5 **bounds.**




1  
 2 **Figure 3. a) Cumulative mass balances of Ålfotbreen (ALF), Rembesdalskåka (REM),**  
 3 **Nigardsbreen (NIG), Storbreen (STO), Hellstugubreen (HEL), Gråsubreen (GR), Engabreen**  
 4 **(ENG) and Storglaciären (STORGL). Data: nve.no/bre (Norwegian glaciers) and**  
 5 **bolin.su.se/data/tarfala (Storglaciären, northern Sweden).**

1 b) – i) Relative importance (standard regression coefficients) of winter precipitation and  
2 summer temperature in 25-year moving windows. Blue (red) lines median estimates standard  
3 regression coefficients (relative importance) of winter precipitation (summer temperature).  
4 Light blue and pink shadings indicate 2.5% and 97.5% quantiles of Bayesian credible  
5 intervals of standard regression coefficients (relative importance). Results are presented as  
6 25-year centred windows. j) Atlantic Multidecadal Oscillation Index  
7 (<http://www.esrl.noaa.gov/psd/data/timeseries/AMO/>, 30-year loess-smoothed). k) North  
8 Atlantic Oscillation Index (Jones et al. 1997, updated).

9

10


11



1

2 Figure 4. Standard deviation ratios. Ratios between standard deviations of winter balances  
 3 (sBw) and annual balances (sBa, sBw/sBa) and summer balances (sBs) and annual balances  
 4 (sBs/sBw). Standard deviation ratios are shown for the entire measurement period (central  
 5 symbols) and for periods of above (left symbols) and below (right symbols) median NAO-  
 6 index, respectively. sBw: standard deviation of winter mass balance, sBs: standard deviation  
 7 of summer mass balance; sBa: standard deviation of annual mass balance.





1

2 **Figure 5. Coefficients of determination ( $r^2$ ) among mass balances and North Atlantic**  
 3 **Oscillation (NAO) Index (Jones et al., 1997, updated). Coefficients of determination between**  
 4 **winter mass balances and annual mass balances and winter NAO-index. Coefficients of**  
 5 **determinations are shown for the entire measurement period (central symbols) and for periods**  
 6 **of above (left symbols) and below (right symbols) median NAO-index, respectively. Bw:**  
 7 **winter mass balance, Ba: annual mass balance; NAO: NAO-index.**