

Analysis of the mass balance time series of glaciers in the Italian Alps

L. Carturan^{*1,6}, C. Baroni^{2,3,6}, M. Brunetti⁴, A. Carton^{5,6}, G. Dalla Fontana^{1,6}, M. C. Salvatore^{2,6}, T. Zanoner^{5,6,7},
G. Zuecco¹

¹ Dipartimento Territorio e Sistemi Agro-Forestali, University of Padova, Viale dell'Università 16, 35020, Legnaro, Padova, Italy

² Dipartimento di Scienze della Terra, University of Pisa, Via S. Maria 53, 56126, Pisa, Italy

³ IGG-CNR, Via Giuseppe Moruzzi 1, 56124, Pisa, Italy

⁴ ISAC-CNR, Via Gobetti 101, 40129, Bologna, Italy

⁵ Dipartimento di Geoscienze, University of Padova, Via Gradenigo 6, 35131, Padova, Italy

⁶ Comitato Glaciologico Italiano, Corso Massimo d'Azeglio 42, 10125, Torino, Italy

⁷ CNR-IRPI, Strada delle Cacce 73, 10135, Torino, Italy

Corresponding author: luca.carturan@unipd.it

Abstract

This work presents an analysis of the mass balance series of nine Italian glaciers, which were selected based on the length, continuity and reliability of observations. All glaciers experienced mass loss in the observation period, which is variable for the different glaciers and ranges between 10 and 47 years. The longest series display increasing mass loss rates, which were mainly due to increased ablation during longer and warmer ablation seasons. The mean annual mass balance (B_a) in the decade from 2004 to 2013 ranged from -1788 mm to -763 mm w.e. y^{-1} . Low-altitude glaciers with low elevation ranges are more out of balance than the higher, larger and steeper glaciers, which maintain residual accumulation areas in their upper reaches. The response of glaciers is mainly controlled by the combination of Oct-May precipitations and Jun-Sep temperatures, but rapid geometric adjustments and atmospheric changes lead to modifications in their response to climatic variations. In particular, a decreasing correlation of B_a with the Jun-Sep temperatures and an increasing correlation with Oct-May precipitations are observed for some glaciers. In addition, the Oct-May temperatures tend to become significantly correlated with B_a , possibly indicating a decrease in the fraction of solid precipitation, and/or increased ablation, during the accumulation season. Because most of the monitored glaciers have no more accumulation area, their observations series are at risk due to their impending extinction, thus requiring a soon replacement.

35

36 **1 Introduction**

37 The mass balance of glaciers is a key variable for monitoring strategies of the Earth climate system because
38 it is the direct and undelayed response of glaciers to atmospheric conditions. Other reactions of glaciers to
39 climatic changes, such as the fluctuations of the front, are more easy and immediate to measure but
40 represent indirect, delayed and filtered signals (WGMS, 2008; Zemp et al., 2005).

41 The direct glaciological method (Østrem and Brugman, 1991) is the standardized method in worldwide
42 glacier monitoring strategies. This method consists of in-situ measurements of the surface accumulation
43 and ablation, taken at single points and then extrapolated and integrated to yield the glacier-wide surface
44 mass balance (Kaser et al., 2003; Cogley et al., 2011). The World Glacier Monitoring Service (WGMS)
45 collects and publishes mass balance data of glaciers obtained by the glaciological method as part of global
46 climate-related observation systems (Zemp et al., 2009; WGMS, 2012 and 2013, and earlier issues).

47 The European Alps are one of the regions of the world with the highest density of glaciers that are subject
48 to mass balance observations. Twenty five glaciers have ongoing and continuous mass balance series with
49 at least 10 years of observations, and 11 of them are longer than 30 years
50 (<http://www.wgms.ch/metadatabrowser.html>, last access: 27 September 2015). In the Italian Alps, nine
51 glaciers have ongoing and continuous mass balance series longer than 10 years, and only one glacier (the
52 Careser Glacier) has a series longer than 30 years.

53 The mass balance series of the glaciers in the Italian Alps have not yet been reviewed and analysed jointly.
54 The Italian glaciers may have a peculiar behaviour compared to the glaciers from other regions of the
55 European Alps, because of the differences in glacier characteristics, climatic features and trends of
56 meteorological variables (Brunetti et al., 2006 and 2009; Auer et al., 2007). Differences may occur in the
57 response of the glaciers in different sub-regions of the Italian Alps or with different characteristics, which
58 have not been recognized. It is also interesting to highlight possible feedbacks in the response of Italian
59 glaciers to atmospheric changes.

60 Therefore, this work aims to i) analyse and compare the direct mass balance series of the glaciers in the
61 Italian Alps, ii) understand the behaviour of the measured glaciers in relation to the observed climatic
62 trends, and iii) highlight possible future requirements for the mass balance monitoring strategy in the
63 Italian Alps.

64

65 **2 Available mass balance series**

66 In this work, we analyse the glaciers with at least 10 years of continuous and ongoing mass balance
67 measurements, which were obtained using the direct glaciological methods and published in peer reviewed
68 journals or in the WGMS publications (CGI, 1914–1977 and 1978–2011; Baroni et al., 2012, 2013, 2014;
69 WGMS, 2012 and 2013, and earlier issues). "Continuous" indicates the series with data gaps <10%,
70 and "ongoing" indicates that the mass balance observations have been performed in the last two years (i.e.,
71 the 2012 and 2013 hydrological years). These criteria ensure the comparability of the series, a sufficient
72 length in the temporal analyses and reliability of the measurements and calculations.

73 Nine monitored glaciers fulfil these characteristics in the Italian Alps and are clustered in three geographic
74 areas (Fig. 1). The two monitored glaciers in the Gran Paradiso Group (Western Alps), i.e., Grand Etrèt
75 (since 2002) and Ciardoney (since 1992), are rather small (area < 1 km²) and have low mean elevations and
76 low elevation ranges (Table 1). Snowfall is the prevailing feeding source, but windborne snow and
77 avalanching also contribute to snow accumulation.

78 The longest series of mass balance measurements in the Italian Alps has been collected on the Careser
79 Glacier, in the Ortles-Cevedale (Eastern Alps, Fig. 1) since 1967. Currently, this glacier is undergoing rapid
80 shrinking and fragmentation in smaller units. It is characterized by a flat surface, prevailing southern
81 exposure, quite low mean elevation and feeding by snowfall. Its area decreased from 5 km² in 1967 to 1.6
82 km² in 2012 (Carturan et al., 2013a). In the 1980s, observations started in two other glaciers of the Ortles-
83 Cevedale: Fontana Bianca and Sforzellina. These two small mountain glaciers (area <0.5 km²) have different
84 characteristics: Fontana Bianca is rather steep with negligible debris cover and mainly fed by snowfall,
85 whereas Sforzellina is flatter, debris-covered in its lower part and fed by avalanches in its upper part. In the
86 2000s, mass balance observations began in the Lunga Glacier and in the southern branch of La Mare Glacier,
87 which are larger valley glaciers (1.9 and 2.2 km², respectively) that reach higher elevations (3378 and 3518
88 m, respectively) and mainly fed by snowfall.

89 In Val Ridanna (Breonie Occidentali Group, Eastern Alps, Fig. 1), the measurements began in 1996 in the
90 Pendente Glacier and were extended to the Malavalle Glacier in 2002. The first glacier is a 0.9 km² wide
91 mountain glacier, characterized by a flat surface, low mean elevation, southern exposure and significant
92 accumulation from windborne snow. The second glacier is a much larger (6.9 km²) valley glacier with higher
93 mean and maximum elevation and mainly fed by snowfall.

94

95 **3 Methods**

96

97 **3.1 Mass balance measurements and calculations**

98 Point measurements of the annual mass balance in the ablation area consist of repeated readings of
99 ablation stakes, which are made of aluminium, wood or plastic and drilled into the ice/firn using hand drills
100 or steam drills. In the accumulation area, the depth of the snow at the end of the ablation season is
101 measured using hand probes, and its density is determined in snowpits. Snow depth soundings and density
102 measurements in the snowpits or by hand coring devices are also used for winter mass balance
103 measurements, which are performed on all glaciers except Sforzellina. The summer mass balance is derived
104 by subtracting the winter mass balance from the annual mass balance. The density of measuring points
105 varies among different glaciers in relation to their extent, accessibility and complexity of the mass balance
106 distribution (Fig. 2). The ablation stake density ranges from 4 points km⁻² (Malavalle Glacier) to 45 points
107 km⁻² (Sforzellina Glacier). The density of snow depth soundings for the winter mass balance determination
108 ranges from 15 points km⁻² (Malavalle Glacier) to 142 points km⁻² (Fontana Bianca Glacier).

109 Point measurements are interpolated and extrapolated to the entire area of the glaciers using different
110 procedures. In the Grand Etrèt and Ciardoney glaciers, each ablation stake is assumed representative of a
111 specific part of the glacier, where the mass balance distribution is assumed homogeneous. Then, a
112 weighted mean is calculated, using the area of the homogeneous parts into which the glacier is subdivided

113 as weights (<http://www.pngp.it/>, last access: 27 September 2015; <http://www.nimbus.it/>, last access: 27
114 September 2015). In Malavalle and Pendente glaciers, the area is divided into “sub-catchments”, and for
115 each sub-catchment, a linear regression of point balances vs. altitude is calculated and used for the
116 spatialization (<http://www.provinz.bz.it/wetter/glacierreport.asp>, last access: 27 September 2015). In
117 Careser Glacier, the even distribution of the mass balance and good coverage of measurement points
118 enable the use of automatic interpolation algorithms (Spatial Analyst Tools) in the ESRI-ArcGIS software.
119 Manual drawing of balance isolines is used for the remaining Sforzellina, Fontana Bianca, La Mare and
120 Lunga glaciers (Catasta and Smiraglia, 1993; Cannone et al., 2008;
121 <http://www.provinz.bz.it/wetter/glacierreport.asp>; Carturan, 2016). The measurements are performed
122 close to the time of maximum and minimum mass balance during the year, when the glacier and
123 atmospheric conditions are favourable for field surveys. The “floating-date” time measurement system is
124 used for all glaciers (Cogley et al., 2011).

125 Typical random errors reported in the literature for glacier-wide mass balance estimates obtained with
126 these methods are of about ± 200 mm w.e. y^{-1} (Lliboutry, 1974; Braithwaite and Olesen, 1989; Cogley and
127 Adams, 1998; Cogley, 2009). The accuracy indicated by the investigators carrying out mass balance
128 measurements in the nine Italian glaciers range between ± 0.05 and 0.30 m w.e. y^{-1} (WGMS, 2015; Carturan,
129 2016). Assessments based on the comparison between the direct and the geodetic mass balance have been
130 published for the Careser, La Mare and Lunga glaciers, indicating a good match of the two methods and
131 revealing that a calibration of the direct mass balance results is not required (Carturan et al., 2013a; Galos
132 et al., 2015; Carturan, 2016).

133

134 **3.2 Meteorological series**

135 The climatic variables used in this work consist of synthetic records of the monthly mean temperature and
136 total monthly precipitation, which are obtained for the centre of the three main geographic areas described
137 in Sect. 2, using the procedure reported in Brunetti et al. (2012). Starting from sparse meteorological data
138 recorded at meteorological stations, the synthetic meteorological series are generated using the anomaly
139 method (New et al., 2000; Mitchell and Jones, 2005). This method is based on the assumption that the
140 spatio-temporal structure of the signal of a meteorological variable over a given area can be described by
141 the superimposition of two fields: the climatological normals over a given reference period (i.e., the
142 climatologies) and the departures from them (i.e., the anomalies). The climatologies are linked to the
143 geographic features of the territory and characterized by remarkable spatial gradients; the departures are
144 linked to the climate variability and change, and they are generally characterized by higher spatial
145 coherence.

146 Under this assumption, the climatologies and anomalies can be reconstructed in completely independent
147 manners and based on different data sets. For climatologies, the priority is the high spatial resolution, and a
148 short time span (few decades) is sufficient. A lower spatial resolution is sufficient for the anomalies, but
149 more importance is given to the data quality and availability of long records. Thus, all series that were used
150 for the anomaly component were subjected to homogenization.

151 The interpolation methods are different for the two components. The climatologies, which are
152 characterized by high spatial gradients, were reconstructed using the procedure in Brunetti et al. (2014),
153 exploiting the relationship between the meteorological variable and the physical characteristics of the
154 terrain. The anomalies, which are characterized by higher spatial coherence, were reconstructed using

155 weighted averages as described in Brunetti et al. (2006). The weights are horizontal and vertical distance
156 weighting functions, with the addition of an angular weight that accounts for the anisotropy in the
157 distribution of stations around the sites. Finally, the two fields were superimposed to obtain the temporal
158 series in absolute values for each site.

159

160 **3.3 Analyses of the mass balance and meteorological series**

161 The time series of annual mass balance (B_a), winter mass balance (B_w), summer mass balance (B_s) and
162 Accumulation Area Ratio (AAR, i.e., the ratio of the area of the accumulation zone to the area of the glacier)
163 were analysed and compared to highlight the possible trends, break points, common behaviour and
164 peculiarities of single glaciers and/or single years. To highlight the systematic differences among the
165 glaciers, the mean values of B_a , B_w , B_s and AAR were calculated in the common period of observation from
166 2004 to 2013. The decadal means of B_a for the Italian glaciers were compared to the decadal means for a
167 sample of nine representative glaciers of the European Alps (Zemp et al., 2005). The correlations among the
168 B_a series of different glaciers and among B_a of single glaciers with the respective series of B_w and B_s were
169 subsequently computed, to identify possible groups of glaciers with similar behaviours and to understand
170 the relative importance of the seasonal components of mass balance.

171 Linear trends and moving averages were calculated for the time series of air temperature and precipitation
172 to highlight the climatic drivers of the observed glacier changes. In particular, we focused on the
173 precipitation of the accumulation season, from October to May (Oct-May), and on the air temperature of
174 the ablation season, from June to September (Jun-Sep) (Pelto, 2008; Carturan et al., 2013b), computing
175 their correlation with B_a and performing multiple linear regression analyses. For the four glaciers with the
176 longest mass balance series (Careser, Fontana Bianca, Sforzellina and Ciardoney), we performed a moving
177 correlation analysis of B_a vs. the seasonal and annual temperature and precipitation, to recognize possible
178 changes and/or trends in their response and sensitivity to climatic fluctuations, e.g., ascribable to
179 geometric adjustments.

180

181 **4 Results and discussion**

182 **4.1 Analysis of mass balance series**

183 The longest available series for the glaciers in the Italian Alps clearly show a trend towards more negative
184 B_a in the observation period (Fig. 3), and one or two change points, which were identified using the
185 'Changepoint' R package (Killick and Eckley, 2014). In particular, the series of Careser Glacier shows three
186 phases: i) the period from 1967 to 1980 with near equilibrium conditions (mean $B_a = -132$ mm w.e. γ^{-1} , STD
187 = 540 mm w.e.); ii) the period from 1981 to 2002 with imbalanced conditions (mean $B_a = -1192$ mm w.e. γ^{-1} ,
188 STD = 517 mm w.e.); and iii) the period after 2002 with stronger imbalance (mean $B_a = -1926$ mm w.e. γ^{-1} ,
189 STD = 725 mm w.e.). A sudden transition in 1980-81 is also clearly visible in the series of AAR, which
190 documents the nearly complete disappearance of the accumulation area of this glacier, which is likely
191 responsible for the step change in its mass balance series during that period.

192 The transition of 2002-03 is also observable for the Fontana Bianca, Sforzellina and Pendente glaciers,
193 whose measurements started in the 1980s and 1990s. Their mean B_a values changed from -599, -868 and -
194 703 mm w.e. γ^{-1} , before 2002, to -1257, -1471 and -1308 mm w.e. γ^{-1} after 2002, respectively. This

195 transition is less obvious for the Ciardoney Glacier, which experienced a notably negative mass balance
196 already in 1998 and 1999. The B_w and B_s series have some gaps but suggest that the increased mass loss
197 rates were mainly ascribable to increased ablation (and associated positive feedbacks) instead of decreased
198 snow accumulation. These results are consistent with previous works, which indicate that the mass changes
199 of the glaciers in the Alps, at the annual and decadal scale, are mainly driven by the summer balance (e.g.,
200 Schoner et al., 2000; Vincent et al., 2004; Zemp et al., 2008; Huss et al., 2015).

201 The AAR series show that the accumulation area almost vanished from all glaciers in the 2000s except the
202 years 2001, 2010 and 2013, when several glaciers were close to balanced-budget conditions mainly as a
203 result of the increased B_w . In these years, the highest increase in AAR occurred in the Fontana Bianca
204 Glacier, which is steep and exposed to the east. On the contrary, the AAR did not significantly increase in
205 the neighbouring Careser Glacier, which is flatter and mainly exposed to the south (Table 1). This behaviour
206 is uncommon for flat glaciers because they should be more sensitive to variations of the Equilibrium Line
207 Altitude (ELA) than the steeper glaciers (Benn and Evans, 2010), and reveals that the Careser Glacier is
208 almost completely below the current ELA, also in the years of ELA minima. La Mare and Malavalle glaciers,
209 which are larger and cover a wider elevation range (Table 1), show more persistent accumulation areas,
210 although their size is too small to ensure balanced-budget conditions.

211 The B_a and B_s values of different glaciers tend to diverge in years with largely negative mass balance and
212 converge in years closer to equilibrium (1993, 2001, 2010 and 2013, Fig. 3). Reinforcing processes and
213 feedbacks likely amplify the differences among the glaciers in imbalanced years, particularly the decrease in
214 the glacier-average albedo caused by the early disappearance of snow from low-lying, flat and less
215 topographic-shielded glaciers, and by the accumulation of dust and debris on the surface. B_w also shows the
216 alternation of years with small/large variability among the glaciers, but this behaviour cannot be clearly
217 related to the magnitude of the snow accumulation, as observed in the two high-accumulation years 2009
218 (high variability) and 2013 (low variability). In this case, the spatial variability of the precipitation during the
219 accumulation season, which is larger than the spatial variability of air temperature in the ablation season,
220 determines the interannual variability of B_w for single glaciers, which is further controlled by snow
221 redistribution processes. Snow redistribution appears more effective for the Pendente, Grand Etrèt and
222 Ciardoney glaciers, leading to over-accumulation in snow-rich winters (e.g., in 2009) and larger interannual
223 variability of B_w . Correlation coefficients calculated between B_w and October-May precipitations range
224 between 0.73 and 0.78 are significant at the 0.05 level only for Careser, La Mare, Lunga, Fontana Bianca
225 and Malavalle, while they are not statistically significant for Pendente, Grand Etrèt and Ciardoney, in line
226 with the hypothesised higher importance of snow redistribution processes in these three glaciers.

227 In the period from 2004 to 2013, significantly higher B_w is observed for Pendente and Grand Etrèt,
228 compared with the other glaciers in the same geographic area (Table 2), explaining the persistence of these
229 two ice bodies at such low altitude (Table 1). In the same period, the Careser Glacier had the lowest
230 average B_a and AAR, whereas the Malavalle and La Mare glaciers had the highest average B_a , B_s and AAR,
231 retaining accumulation areas in their upper parts. However, the mean AARs were remarkably low for all
232 analysed glaciers, and far from balanced-budget conditions ($AAR_0 = 0.55 - 0.58$, Dyurgerov et al., 2009;
233 Mernild et al., 2013). Overall, low-altitude and flat glaciers with low elevation ranges are more out of
234 balance than the steeper glaciers at higher altitude with higher elevation ranges, as acknowledged in
235 various other studies (e.g., Furbish and Andrews, 1984; Benn and Evans, 2010; Carturan et al., 2013b;
236 Fischer et al., 2015).

237 At the regional scale, the spatial representativeness of five Italian mass balance glaciers can be assessed on
238 the basis of the geodetic mass balance calculations performed by Carturan et al., (2013b). In the period
239 from the 1980s to the 2000s, the average geodetic mass balance rate of the 112 glaciers in the Ortles-
240 Cevedale Group has been $-0.69 \text{ m w.e. } \gamma^{-1}$. If we consider the average geodetic mass balance in the same
241 period as an index of the spatial representativeness for single glaciers, we obtain in decreasing order: i) La
242 Mare with $-0.64 \text{ m w.e. } \gamma^{-1}$, ii) Sforzellina with $-0.86 \text{ m w.e. } \gamma^{-1}$, iii) Fontana Bianca with $-0.90 \text{ m w.e. } \gamma^{-1}$, iv)
243 Lunga with $-1.00 \text{ m w.e. } \gamma^{-1}$, and v) Careser with $-1.43 \text{ m w.e. } \gamma^{-1}$. Geodetic calculations only exist for few
244 areas in the Italian Alps (e.g., Galos et al., 2015) and do not include the other four mass balance glaciers
245 analysed in this study. Therefore it was not possible to evaluate their spatial representativeness at the
246 regional scale. Similarly, quantitative assessments of the representativeness of all the nine glaciers at the
247 scale of the entire Italian Alps will require further investigations, integrating in-situ measurements,
248 remotely sensed observations and numerical modelling (WGMS, 2015).

249 The response of Italian glaciers to the climatic conditions of the last decades is similar to that of nine
250 representative glaciers of the entire European Alps (Zemp et al., 2005; Fig. 4), although single glaciers
251 display different mass loss rates (Table 2). The Italian glaciers display $\sim 200\text{-}250 \text{ mm w.e. } \gamma^{-1}$ more negative
252 B_a until 2002 and $\sim 200 \text{ mm w.e. } \gamma^{-1}$ less negative B_a since then. Therefore, it can be assessed that the mean
253 B_a values for the Italian and “European” glaciers are fairly similar. Comparable results were obtained by
254 Huss et al., (2015), who compared the decadal mean B_a of glaciers from France, Switzerland, Austria and
255 Italy. These comparisons may be affected by the loss of spatial representativeness of some glaciers (e.g.
256 Careser in the Italian Alps and Sarennes in the French Alps) and by the different subsets of Italian glaciers
257 which are useable in the four different sub-periods. In the last decade, the inclusion of La Mare and
258 Malavalle glaciers in the Italian subset and the concurrent sharp decrease of B_a for the Sarennes, St. Sorlin
259 and Gries glaciers explain the different behaviours of the two groups of glaciers. However, the smaller
260 Italian glaciers (average area = 1.79 km^2) may have a shorter response time to climatic changes, adjusting
261 their geometry faster than the larger glaciers (average area = 3.63 km^2) which are representative of the
262 European Alps (Hoelzle et al., 2003; Abermann et al., 2009). The rapid shrinking and fragmentation of
263 Careser Glacier is a good example: in the last decade, its area has halved, and it completely lost the parts
264 subject to higher ablation (Carturan et al., 2013a). Changes in the general atmospheric circulation and
265 spatial distribution of precipitation could also have played a role and will be discussed in Sect. 4.2.

266 There is a generally high correlation among the B_a values of the analysed glaciers (Table 3). The series of
267 Careser, Fontana Bianca and La Mare glaciers show a highly significant correlation with most other glaciers,
268 even if they have different characteristics or are far away. On the contrary, the Lunga Glacier shows a lower
269 correlation and lower statistical significance with the glaciers of the same mountain group. However, it has
270 the shortest series, and most importantly, it does not include the highly negative B_a of 2003, which certainly
271 increases the correlation among other glaciers. There are notably high correlations in the Ortles Cevedale
272 between Careser and La Mare and between Fontana Bianca and La Mare glaciers. A similarly high
273 correlation is observed between Pendente and Malavalle glaciers in Val Ridanna, whereas there is a much
274 lower correlation between the two glaciers of the Gran Paradiso Group, which suggests that differences in
275 local topo-climatic factors can be decisive on such small ice bodies (e.g., Kuhn, 1995; DeBeer and Sharp,
276 2009; Carturan et al., 2013c; Scotti et al., 2014; Colucci and Guglielmin, 2015).

277 For most glaciers, B_a is more correlated to B_s than to B_w (Table 4), which further confirms the importance of
278 summer ablation. The relevance of the snow redistribution and over-accumulation on the Pendente and
279 Grand Etrèt glaciers is indicated by the higher correlation of their B_a with B_w . On La Mare Glacier, the two
280 seasonal components have similar correlations with B_a . However, these results are influenced by the length

281 of the observation period and the presence/absence of extreme years with high accumulation (e.g., 2001)
282 or high ablation (e.g., 2003) in the observation series of individual glaciers. For the analysed glaciers, no
283 significant correlation was found between B_s and B_w .

284

285 4.2 Climatic controls

286 In the period from 1961 to 2013, there are highly significant warming trends for the Jun-Sep air
287 temperature (Fig. 5a, b and c); they are highest in the Gran Paradiso Group ($0.40^{\circ}\text{C decade}^{-1}$) and lowest in
288 Val Ridanna ($0.35^{\circ}\text{C decade}^{-1}$). The three phases in the longer B_a and B_s series of glaciers (Fig. 3) can be
289 recognized as periods with stationary Jun-Sep temperature, separated by switches in the early 1980s and
290 after the peak of 2003. The warming trends are lower in the accumulation season and range from 0.25 to
291 $0.27^{\circ}\text{C decade}^{-1}$ (Fig. 5d, e and f), but thermal inversions at the valley weather stations could have partially
292 masked the warming at the altitude of the glaciers in this season. The transition towards higher Oct-May
293 temperature occurred in the late 1980s, after a minimum in the first half of the same decade. A distinct
294 warm peak in Oct-May temperature occurred in 2007.

295 The precipitation does not show any significant trend in the accumulation season (Fig. 6d, e and f). The
296 moving averages display oscillations of 10-20% above and below the 1961-1990 mean, which lasted
297 approximately 10-15 years and were higher in the Gran Paradiso Group than in the Ortles-Cevedale and Val
298 Ridanna. Periods with below-average precipitation are recognized in the 1960s, first half of 1970s, and
299 1990s, whereas periods with above-average precipitation occurred in the second half of 1970s and the first
300 half of 1980s. The last 10-15 years were characterized by precipitation close to the mean, with important
301 maxima in 2001, 2009 and 2013, and minima in 2007 and 2012. Similarly to the findings from Durand et al.,
302 (2009a and b) and Eckert et al., (2011) for the French and western Swiss Alps, a change point in winter
303 precipitation of Ortles-Cevedale and Val Ridanna series were identified in 1977, corresponding to an
304 increase of about 10-12%. Linear trends of summer precipitation are positive but not statistically significant.
305 The interannual variability of the Jun-Sep precipitation is remarkably higher in the Gran Paradiso Group (Fig.
306 6a, b and c).

307 Large scale circulation patterns, such as the North Atlantic Oscillation (NAO) and the Northern Hemisphere
308 blocking frequency, are connected with the temporal and spatial variability of winter precipitation in the
309 Alps (Quadrelli et al., 2001). Several studies highlighted a contrasting behaviour of precipitation anomalies
310 in the Oct-May period between the northern and southern Alps, i.e. opposite correlation with indexed large
311 scale circulation patterns (e.g., Quadrelli et al., 2001; Schmidli et al., 2002; Brunetti et al., 2006) and
312 opposite long-term trends in the seasonal precipitation totals (e.g., Brunetti et al., 2006 and 2009; Auer et
313 al., 2007). This characteristic, and the tendency towards a decreasing NAO index in the last two decades
314 (http://www.cpc.ncep.noaa.gov/products/precip/CWlink/pna/JFM_season_ao_index.shtml, last access: 4
315 October 2015), leading to increased winter precipitation in the southern side of the Alps, may provide an
316 additional explanation for the different behaviour of “European” and “Italian” glaciers shown in Fig. 4.
317 Opposite effects of the NAO on the winter precipitation and glacier mass balance in the northern and
318 southern parts of the Eastern Alps were also reported, for example, by Marzeion and Nesje (2012).

319 A correlation analysis of B_w , B_s and B_a versus the seasonal and annual mean NAO index was performed,
320 using the mass balance data of glaciers from Italy and from other nations of the European Alps (Table 5).
321 The results show prevailing negative correlations between the B_w of Italian glaciers and the NAO in the
322 accumulation season, whereas positive correlations prevail in other nations, with the exception of Gries

323 Glacier which however is close to the Italian border. These results are in agreement with the mentioned
324 literature and with our hypothesis of opposite effects of recent NAO trends in the winter precipitation of
325 the northern and southern sides of the Alps. In line with the findings of Reichert et al., (2001), Six et al.,
326 (2001), and Thibert et al., (2013), a negative correlation was calculated between B_s/B_a and the NAO in the
327 accumulation season. If a causal relationship can be hypothesised for the Italian glaciers, related to the
328 albedo feedback from wet/dry winters (with low/high NAO, respectively), the same cannot be stated for
329 glaciers in other countries, due to the prevalent positive correlation of their B_w with the winter NAO. The
330 examination of meteorological series confirms that increased ablation and the related feedbacks are the
331 main causes of the increased imbalance of the analysed Italian glaciers, as observed in Sect. 4.1. This result
332 is further corroborated by the higher correlation of B_a with the Jun-Sep temperature than with the Oct-May
333 precipitation, at least for the glaciers with longer observation series (Careser, Fontana Bianca, Sforzellina
334 and Ciardoney, Table 6). B_a of glaciers with shorter observation series is not significantly correlated with the
335 Jun-Sep temperature; instead, two of them (La Mare and Lunga) show a correlation with the Oct-May
336 precipitation. Combining the Oct-May precipitation and Jun-Sep temperature in a multiple linear regression
337 model leads to highly significant coefficients for both variables, even when the single seasonal components
338 are not correlated with B_a (e.g., for the Pendente and La Mare glaciers). Approximately two-thirds of the B_a
339 variance can be explained by the multiple linear regression. The poorest results were obtained for the two
340 glaciers in Val Ridanna (Pendente and Malavalle) and the Grand Etrèt Glacier. As the first two glaciers are
341 close to the main Alpine divide, they likely benefit from the high orographic uplift that locally enhances
342 precipitation (Schwarb, 2000), but which cannot be accounted for by the multiple regression model due to
343 the lack of weather stations in that area. In addition, the multiple regression model does not account for
344 accumulation by windborne snow on the Pendente and Grand Etrèt glaciers.

345 The Careser, Fontana Bianca and Pendente glaciers display significant negative correlations between their
346 B_a and the Oct-May temperature. For the Careser Glacier, there is also a negative correlation between B_w
347 and the Oct-May temperature. Normally, in this period, most precipitation falls as snow, and the glaciers
348 have negligible ablation and low temperature sensitivity (Oerlemans and Reichert, 2000). However,
349 increasing temperature starts to lead to significant ablation in this period and to reduce the fraction of solid
350 precipitation as clearly detectable in the ablation season (Carturan et al., 2013b). An emblematic example is
351 the warm accumulation season of 2006-07, when the liquid precipitation reached 3000-3100 m a.s.l. (24
352 October 2006), and ice ablation exceeded 50 cm at 3000 m a.s.l. on the Careser and La Mare glaciers.

353 The correlation between B_a of Careser Glacier and the Oct-May temperature starts to become significant in
354 the late 1980s, as shown in the moving correlation analyses (30-year time window in Fig. 7). In the first 20
355 years, the correlation was absent or not statistically significant. These results are consistent with the
356 discussed effects of increasing temperature on the ablation and partitioning between liquid and solid
357 precipitation (Beniston et al., 2003). Reducing the window size from 30 to 15 years leads to a noisier signal
358 and in this case the correlation between B_a and Oct-May temperature does not reach the 95% significance
359 thresholds. However, it is interesting to remark the reversal of the correlation sign from positive in the first
360 years to negative in the last years.

361 The four glaciers in Fig. 7 (Careser, Fontana Bianca, Sforzellina and Ciardoney) share a common trend
362 towards i) a non-significant moving correlation between B_a and the Jun-Sep temperature and ii) a
363 significant moving correlation between B_a and Oct-May precipitation. This behaviour is probably related to
364 the snow-rich accumulation seasons of 2001, 2009 and 2013, and to the fact that the ablation season is
365 already so warm that i) summer snow falls mostly above the highest reaches of the glaciers, which reduces

366 the interannual variability of summer melt, and ii) conditions close to balanced-budget only occur after
367 snow-rich accumulation seasons.

368 Rapid geometric changes may also lead to a non-linear response of B_a to atmospheric changes, at least for
369 some glaciers. For example, the multiple regression residuals of the Careser Glacier, which were mostly
370 positive in the 1980s, 1990s and 2000s, became predominantly negative after 2008 (Fig. 8). This change
371 may suggest that the rapid modifications occurred in the latest years could have induced a negative
372 feedback, reducing the mass loss rate of the glacier, whose current surface and shape are strongly different
373 from the recent past (inset in Fig. 8). Because the multiple regression model does not use the Oct-May
374 temperature as an explanatory variable, it cannot account for the effects of the warm accumulation season
375 of 2006-07, which led to a very low B_w , early disappearance of winter snow and positive albedo feedback.
376 Therefore, the year 2007 results in highly positive regression residuals.

377

378 **4.3 Future requirements**

379 A common characteristic for all glaciers analysed is their very low mean AAR in the last decade (Table 2).
380 Accumulation areas were almost inexistent in most glaciers, indicating that they will soon disappear, even
381 without additional warming. Some glaciers are displaying morphological changes that indicate their
382 impending extinction, such as rapid disintegration (e.g., Careser Glacier, Fig. 8) and surface lowering in the
383 upper accumulation area (e.g., Fontana Bianca Glacier). The AARs of approximately 0.25 indicate that
384 accumulation areas still exist in the larger and higher-reaching La Mare and Malavalle glaciers. However,
385 given that balanced-budget conditions require AAR close to 0.55, large mass loss and areal reduction are
386 also expected for these two glaciers to reach equilibrium with the climatic conditions of the last ten years.

387 The forthcoming vanishing of the monitored glaciers put the continuation of their mass balance
388 observations at risk. Recently-started monitoring programs in larger and higher-reaching glaciers, such as
389 Malavalle and La Mare, will ensure continued observations in Val Ridanna and the Ortles-Cevedale. In line
390 with the recommendations from the WGMS (Zemp et al., 2009), similar observation programs should start
391 in other large and high-reaching glaciers of the Italian Alps, e.g., in the Gran Paradiso group (to substitute
392 Ciardoney and Grand Etrèt) and in other mountain groups. Both the initiation of observations over new
393 glaciers and the replacement of vanishing glaciers will require an assessment of the spatial
394 representativeness of single glaciers through the comparison of the current mass loss rates over wide
395 geographic areas (Haeberli et al., 2013). This assessments can be obtained using modern techniques such
396 as the multi-temporal differencing of digital elevation models, which enable the comparison of mass loss
397 rates in the last years/decades, by means of the geodetic method, over entire regions or mountain ranges
398 (e.g., Paul and Haeberli, 2008; Abermann et al., 2011; Carturan et al., 2013b; Berthier et al., 2014; Fischer et
399 al., 2015). The geodetic mass balance should also help to control the glacier-wide B_a series measured with
400 the direct glaciological method, and to construct a constant-geometry mass balance record (Elsberg et al.,
401 2001) to be connected to climatic drivers.

402

403 **5 Conclusions**

404 In this work, we have analysed the time series of the glaciers with mass balance observations in the Italian
405 Alps. Based on the results of the analyses, the following conclusions can be drawn:

- 406 ▪ All examined glaciers are experiencing imbalanced conditions, and the longer series show sustained
407 negative trends of B_a .
- 408 ▪ The observed behaviour was mainly caused by increased ablation, led by warmer temperature and
409 related feedbacks. The total precipitation does not show any significant trend, but the fraction of
410 solid precipitation decreased as a consequence of the warmer temperature.
- 411 ▪ The B_a of the analysed glaciers is mainly correlated to B_s , except for two glaciers where windborne
412 snow enhances the importance of B_w . For most glaciers, approximately two thirds of the B_a variance
413 can be explained by multiple linear regression, using the Oct-May precipitation and Jun-Sep
414 temperature as independent variables.
- 415 ▪ The monitored Italian glaciers have comparable mass loss rates to a sample of representative
416 glaciers of the entire European Alps. However, the moving correlation analyses and time series of
417 residuals from multiple linear regressions suggest that the smaller (and thinner) Italian glaciers may
418 be reacting faster to atmospheric changes.
- 419 ▪ Most monitored glaciers have no more accumulation area and are at risk of extinction, even
420 without additional warming. Therefore, they will soon require a replacement with larger and higher
421 glaciers that retain accumulation areas.
- 422 ▪ Regional assessments of the mass loss rates using the geodetic method are required to identify
423 possible replacing glaciers, evaluate their spatial representativeness and enable the transitions
424 from replaced to replacing glaciers, as suggested by Haeberli et al. (2013).

425

426 **Author contribution**

427 M. Brunetti processed the meteorological data and prepared the synthetic meteorological series used in
428 this work. Thomas Zanoner compiled the database of the mass balance data and geometric characteristics
429 of the glaciers. L. Carturan and Giulia Zuecco performed the temporal and statistical analyses of the mass
430 balance series. L. Carturan prepared the manuscript with contributions from all co-authors.

431

432 **Acknowledgments**

433 This study was funded by the Italian MIUR Project (PRIN 2010-11): "Response of morphoclimatic system
434 dynamics to global changes and related geomorphological hazards" (local and national coordinators G.
435 Dalla Fontana and C. Baroni). The authors acknowledge the regional environmental agencies of Piemonte,
436 Val d'Aosta, Lombardia, Autonomous Provinces of Trento and Bolzano, which provided the meteorological
437 and topographic data used in this study. Many thanks to Stephan Galos (University of Innsbruck) for sharing
438 information on glaciers in Ötztal and South Tyrol. We would also like to thank all investigators and
439 contributing institutions for the collection and free exchange of the glacier mass balance series.

440

441 **References**

442 Abermann, J., Lambrecht, A., Fischer, A., and Kuhn, M.: Quantifying changes and trends in glacier area and
443 volume in the Austrian Ötztal Alps (1969-1997-2006), *The Cryosphere*, 3, 205-215, doi:10.5194/tc-3-205-
444 2009, 2009.

445 Abermann, J., Kuhn, M., and Fischer, A.: Climatic controls of glacier distribution and glacier changes in
446 Austria. *Ann. Glaciol.*, 52(59), 83-90, 2011.

447 Auer, I., Böhm, R., Jurkovic, A., Lipa, W., Orlik, A., Potzmann, R., Schöner, W., Ungersböck, M., Matulla, C.,
448 Briffa, K., Jones, P., Efthymiadis, D., Brunetti, M., Nanni, T., Maugeri, M., Mercalli, L., Mestre, O., Moisselin,
449 J.-M., Begert, M., Müller-Westermeier, G., Kveton, V., Bochnicek, O., Stastny, P., Lapin, M., Szalai, S.,
450 Szentimrey, T., Cegnar, T., Dolinar, M., Gajic-Capka, M., Zaninovic, K., Majstorovic, Z., and Nieplova, E.:
451 HISTALP - Historical Instrumental climatological Surface Time series of the greater ALPine Region, *Int. J.*
452 *Climatol.*, 27, 17-46, doi: 10.1002/joc.1377, 2007.

453 Baroni, C., Bondesan, A., and Mortara, G.: Report of the Glaciological Survey of 2011 - Relazioni della
454 campagna glaciologica 2011, *Geogr. Fis. Din. Quat.*, 35(2), 211-279, doi:10.4461/GFDQ.2012.35.19, 2012.

455 Baroni, C., Bondesan, A., and Mortara, G.: Report of the Glaciological Survey of 2012 - Relazioni della
456 campagna glaciologica 2012, *Geogr. Fis. Din. Quat.*, 36(2), 303-374, doi: 10.4461/GFDQ.2013.36.24, 2013.

457 Baroni, C., Bondesan, A., and Mortara, G.: Report of the Glaciological Survey 2013. Relazioni della
458 Campagna Glaciologica 2013, *Geogr. Fis. Din. Quat.*, 37, 163-227, doi: 10.4461/GFDQ.2014.37.16, 2014.

459 Beniston, M., Keller, F., Koffi, B., and Goyette, S.: Estimates of snow accumulation and volume in the Swiss
460 Alps under changing climatic conditions, *Theor. Appl. Climatol.*, 76, 125–140, doi: 10.1007/s00704-003-
461 0016-5, 2003.

462 Benn, D. I. and Evans, D. J. A. (Eds.): *Glaciers and Glaciation*, Hodder Education, London, UK, 802 pp., 2010.

463 Berthier, E., Vincent, C., Magnússon, E., Gunnlaugsson, Á. Þ., Pitte, P., Le Meur, E., Masiokas, M., Ruiz, L.,
464 Pálsson, F., Belart, J. M. C., and Wagnon, P.: Glacier topography and elevation changes derived from
465 Pléiades sub-meter stereo images, *The Cryosphere*, 8, 2275-2291, doi:10.5194/tc-8-2275-2014, 2014.

466 Braithwaite, R. J., and Olesen, O. B.: Detection of climate signal by inter-stake correlations of annual
467 ablation data Qamanârssûp Sermia, West Greenland, *J. Glaciol.*, 35(120), 253-259, 1989.

468 Brunetti, M., Maugeri, M., Nanni, T., Auer, I., Böhm, R., and Schöner, W.: Precipitation variability and
469 changes in the Greater Alpine region over the 1800–2003 period, *J. Geophys. Res.*, 111, D11107, doi:
470 10.1029/2005/D06674, 2006.

471 Brunetti, M., Lentini, G., Maugeri, M., Nanni, T., Auer, I., Böhm, R., and Schöner, W.: Climate variability and
472 change in the Greater Alpine Region over the last two centuries based on multi-variable analysis, *Int. J.*
473 *Climatol.*, 29(15), 2197-2225, doi: 10.1002/joc.1857, 2009.

474 Brunetti, M., Lentini, G., Maugeri, M., Nanni, T., Simolo, C., and Spinoni, J.: Projecting North Eastern Italy
475 temperature and precipitation secular records onto a high resolution grid, *Phys. Chem. Earth*, 40-41, 9-22,
476 doi: 10.1016/j.pce.2009.12.005, 2012.

477 Brunetti, M., Maugeri, M., Nanni, T., Simolo, C., and Spinoni, J.: High-resolution temperature climatology
478 for Italy: interpolation method intercomparison. *Int. J. Climatol.*, 34, 1278-1296, doi: 10.1002/joc.3764,
479 2014.

480 Cannone, N., Diolaiuti, G., Guglielmin, M., and Smiraglia, C.: Accelerating climate change impacts on alpine
481 glacier forefield ecosystems in the European Alps, *Ecol. Appl.*, 18(3), 637-648, doi: 10.1890/07-1188.1, 2008.

482 Carturan, L., Baroni, C., Becker, M., Bellin, A., Cainelli, O., Carton, A., Casarotto, C., Dalla Fontana, G., Godio,
483 A., Martinelli, T., Salvatore, M. C., and Seppi, R.: Decay of a long-term monitored glacier: Careser Glacier
484 (Ortles-Cevedale, European Alps), *The Cryosphere*, 7, 1819-1838, doi:10.5194/tc-7-1819-2013, 2013a.

485 Carturan, L., Filippi, R., Seppi, R., Gabrielli, P., Notarnicola, C., Bertoldi, L., Paul, F., Rastner, P., Cazorzi, F.,
486 Dinale, R., and Dalla Fontana, G.: Area and volume loss of the glaciers in the Ortles-Cevedale group (Eastern
487 Italian Alps): controls and imbalance of the remaining glaciers, *The Cryosphere*, 7, 1339-1359,
488 doi:10.5194/tc-7-1339-2013, 2013b.

489 Carturan, L., Baldassi, G. A., Bondesan, A., Calligaro, S., Carton, A., Cazorzi, F., Dalla Fontana, G., Francese, R.,
490 Guarnieri, A., Milan, N., Moro, D., and Tarolli, P.: Current behavior and dynamics of the lowermost Italian
491 glacier (Montasio Occidentale, Julian Alps), *Geogr. Ann. A*, 95(1), 79-96, doi: 10.1111/geoa.12002, 2013c.

492 Carturan, L.: New measurement series for the future monitoring of glacier mass balance in the European
493 Alps: methods and results on La Mare Glacier (Ortles-Cevedale, Italy), *J. Glaciol.*, in review, 2016.

494 Casartelli, G., Kappenberger, G., and Smiraglia, C.: Accumulo e ablazione sui ghiacciai delle alpi lombarde e
495 svizzere: Risultati di alcuni recenti bilanci di massa, *Rivista Geografica Italiana*, 103(1), 1-30, 1996.

496 Catasta, G., and Smiraglia, C.: The mass balance of a cirque glacier in the Italian Alps (Ghiacciaio della
497 Sforzellina, Ortles-Cevedale Group), *J. Glaciol.*, 39(131), 87-90, 1993.

498 CGI (Comitato Glaciologico Italiano): Reports of the glaciological surveys, *Bollettino del Comitato*
499 *Glaciologico Italiano, Series I and II*, 1–25, 1914–1977.

500 CGI (Comitato Glaciologico Italiano): Reports of the glaciological surveys, *Geogr. Fis. Din. Quat.*, 1–35, 1978–
501 2011.

502 Cogley, J. G.: Geodetic and direct mass-balance measurements: comparison and joint analysis. *Ann. Glaciol.*,
503 50(50), 96-100, 2009.

504 Cogley, J. G., and Adams, W. P.: Mass balance of glaciers other than the ice sheets, *J. Glaciol.*, 44(147), 315-
505 325, 1998.

506 Cogley, J. G., Hock, R., Rasmussen, L. A., Arendt, A. A., Bauder, A., Braithwaite, R. J., Jansson, P., Kaser, G.,
507 Moller, M., Nicholson, L., and Zemp, M.: Glossary of Glacier Mass Balance and Related Terms, IHP-VII
508 Technical Documents in Hydrology No. 86, IACS Contribution No. 2, UNESCO-IHP, Paris, 2011.

509 Colucci, R. R., and Guglielmin, M.: Precipitation–temperature changes and evolution of a small glacier in the
510 southeastern European Alps during the last 90 years, *Int. J. Climatol.*, 35(10), 2783–2797,
511 doi: 10.1002/joc.4172, 2015.

512 Debeer, C. M., and Sharp, M. J.: Topographic influences on recent changes of very small glaciers in the
513 Monashee Mountains, British Columbia, Canada, *J. Glaciol.*, 55(192), 691–700, doi:
514 10.3189/002214309789470851, 2009.

515 Durand, Y., Laternser, M., Giraud, G., Etchevers, P., Lesaffre, B., and Mérindol, L.: Reanalysis of 44 years of
516 climate in the French Alps (1958–2002): methodology, model validation, climatology, and trends for air
517 temperature and precipitation, *J. Appl. Meteorol. Climatol.*, 48(3), 429–449, 2009a.

518 Durand, Y., Giraud, G., Laternser, M., Etchevers, P., Mérindol, L., and Lesaffre, B.: Reanalysis of 47 years of
519 climate in the French Alps (1958–2005): climatology and trends for snow cover, *J. Appl. Meteorol. Climatol.*,
520 48(12), 2487–2512, 2009b.

521 Dyurgerov, M., Meier, M. F., and Bahr, D. B.: A new index of glacier area change: a tool for glacier
522 monitoring, *J. Glaciol.*, 55(192), 710–716, 2009.

523 Eckert, N., Baya, H., Thibert, E., and Vincent, C.: Extracting the temporal signal from a winter and summer
524 mass-balance series: application to a six-decade record at Glacier de Sarennes, French Alps, *J. Glaciol.*,
525 57(201), 134–150, 2011.

526 Elsberg, D. H., Harrison, W. D., Echelmeyer, K. A., and Krimmel, R. M.: Quantifying the effect of climate and
527 surface change on glacier mass balance, *J. Glaciol.*, 47, 649–658, 2001.

528 Fischer, M., Huss, M., and Hoelzle, M.: Surface elevation and mass changes of all Swiss glaciers 1980–2010,
529 *The Cryosphere*, 9, 525–540, doi:10.5194/tc-9-525-2015, 2015.

530 Furbish, D. J., and Andrews, J. T.: The use of hypsometry to indicate long-term stability and response of
531 valley glaciers to changes in mass transfer, *J. Glaciol.*, 30, 199–211, 1984.

532 Galos, S. P., Klug, C., Prinz, R., Rieg, L., Dinale, R., Sailer, R., and Kaser, G.: Recent glacier changes and
533 related contribution potential to river discharge in the Vinschgau / Val Venosta, Italian Alps, *Geogr. Fis. Din.*
534 *Quat.*, 38, 143–154, doi: 10.4461/GFDQ.2015.38.13, 2015.

535 Haeberli, W., Huggel, C., Paul, F., and Zemp, M.: Glacial responses to climate change, in: *Treatise on*
536 *Geomorphology*, Academic Press, 13, San Diego, California, 152–175, 2013.

537 Hoelzle, M., Haeberli, W., Dischl, M., and Peschke, W.: Secular glacier mass balances derived from
538 cumulative glacier length changes, *Global Planet. Change*, 36, 295–306, doi: 10.1016/S0921-
539 8181(02)00223-0, 2003.

540 Huss, M., Dhulst, L., and Bauder, A.: New long-term mass balance series for the Swiss Alps, *J. Glaciol.*,
541 61(227), 551–562, doi: 10.3189/2015JoG15j015, 2015.

542 Kaser, G., Fountain, A., and Jansson, P.: A manual for monitoring the mass balance of mountain glaciers,
543 IHP-VI Technical Documents in Hydrology No. 59, UNESCO-IHP, Paris, 2003.

544 Killick, R., Eckley, I. A.: Changepoint: An R package for changepoint analysis, *J. Stat. Softw.*, 58(3), 1–19. DOI:
545 10.18637/jss.v058.i03, 2014.

546 Kuhn, M.: The mass balance of very small glaciers, *Zeitschrift für Gletscherkunde und Glazialgeologie*, 31(1–
547 2), 171–179, 1995.

- 548 Lliboutry, L.: Multivariate statistical analysis of glacier annual balances, *J. Glaciol.*, 13, 371-392, 1974.
- 549 Marzeion, B. and Nesje, A.: Spatial patterns of North Atlantic Oscillation influence on mass balance
550 variability of European glaciers, *The Cryosphere*, 6, 661-673, doi:10.5194/tc-6-661-2012, 2012.
- 551 Mernild, S. H., Lipscomb, W. H., Bahr, D. B., Radić, V., and Zemp, M.: Global glacier changes: a revised
552 assessment of committed mass losses and sampling uncertainties, *The Cryosphere*, 7, 1565-1577,
553 doi:10.5194/tc-7-1565-2013, 2013.
- 554 Mitchell, T. D., and Jones, P. D.: An improved method of constructing a database of monthly climate
555 observations and associated high-resolution grids, *Int. J. Climatol.* 25, 693–712, doi: 10.1002/joc.1181, 2005.
- 556 New, M., Hulme, M., and Jones, P. D.: Representing twentieth century space-time variability. Part 2:
557 development of 1901–96 monthly grids of surface climate, *J. Climate*, 13, 2217–2238, 2000.
- 558 Oerlemans, J., and Reichert, B. K.: Relating glacier mass balance to meteorological data by using a seasonal
559 sensitivity characteristic, *J. Glaciol.*, 46(152), 1–6, doi: 10.3189/172756500781833269, 2000.
- 560 Østrem, G., and Brugman, M.: Mass balance measurement techniques. A manual for field and office work,
561 National Hydrology Research Institute (NHRI), Science Report, (4), Saskatoon, Canada, 224 pp., 1991.
- 562 Paul, F., and Haeberli, W.: Spatial variability of glacier elevation changes in the Swiss Alps obtained from
563 two digital elevation models, *Geophys. Res. Lett.*, 35, L21502, doi:10.1029/2008GL034718, 2008.
- 564 Pelto, M. S.: Glacier annual balance measurement, forecasting and climate correlations, North Cascades,
565 Washington 1984–2006, *The Cryosphere*, 2, 13-21, doi:10.5194/tc-2-13-2008, 2008
- 566 Quadrelli, R., Lazzeri, M., Cacciamani, C., and Tibaldi, S.: Observed winter Alpine precipitation variability and
567 links with large-scale circulation patterns, *Clim. Res.*, 17(3), 275-284, doi:10.3354/cr017275, 2001.
- 568 Reichert, B. K., Bengtsson, L., and Oerlemans, J.: Midlatitude forcing mechanisms for glacier 25 mass
569 balance investigated using general circulation models, *J. Climate*, 14, 3767–3784, 2001.
- 570 Salvatore, M. C., Zanoner, T., Baroni, C., Carton, A., Banchieri, F. A., Viani, C., Giardino, M., and Perotti, L.:
571 The state of Italian glaciers: a snapshot of the 2006-2007 hydrological period, *Geogr. Fis. Dinam. Quat.*, 38,
572 175-198, 2015.
- 573 Schmidli, J., Schmutz, C., Frei, C., Wanner, H., and Schär, C.: Mesoscale precipitation variability in the region
574 of the European Alps during the 20th century, *Int. J. Climatol.*, 22(9), 1049-1074, doi: 10.1002/joc.769, 2002.
- 575 Schöner, W., Auer, I., and Böhm, R.: Climate variability and glacier reaction in the Austrian eastern Alps,
576 *Ann. Glaciol.*, 31(1), 31-38, 2000.
- 577 Schwarb, M.: The alpine precipitation climate, Ph.D. thesis, ETH Zürich, Switzerland, 131 pp., 2000.
- 578 Scotti, R., Brardinoni, F., and Crosta, G. B.: Post-LIA glacier changes along a latitudinal transect in the
579 Central Italian Alps, *The Cryosphere*, 8, 2235-2252, doi:10.5194/tc-8-2235-2014, 2014.
- 580 Six, D., Reynaud, L., and Letréguilly, A.: Alpine and Scandinavian glaciers mass balances, their relations with
581 the North Atlantic Oscillation, *Comptes Rendus de l'Académie des Sciences, Series IIA, Earth Planet. Sc.*, 333,
582 693–698, 2001.

583 Thibert, E., Eckert, N., and Vincent, C.: Climatic drivers of seasonal glacier mass balances: an analysis of 6
584 decades at Glacier de Sarennes (French Alps), *The Cryosphere*, 7(1), 47-66, 2013.

585 Vincent, C., Kappenberger, G., Valla, F., Bauder, A., Funk, M., and Le Meur, E.: Ice ablation as evidence of
586 climate change in the Alps over the 20th century, *J. Geophys. Res.*, 109(D10), D10104, doi:
587 10.1029/2003JD003857, 2004.

588 WGMS: Global glacier changes: facts and figures, edited by: Zemp, M., Roer, I., Kaab, A., Hoelzle, M., Paul,
589 F., and Haerberli, W., UNEP, World Glacier Monitoring Service, Zürich, Switzerland, 2008.

590 WGMS: Fluctuations of Glaciers 2005-2010, Volume X, edited by: Zemp, M., Frey, H., Gärtner-Roer, I.,
591 Nussbaumer, S. U., Hoelzle, M., Paul, F., and W. Haerberli: ICSU(WDS)/IUGG(IACS)/UNEP/UNESCO/WMO,
592 World Glacier Monitoring Service, Zurich, Switzerland, 336 pp., publication based on database version:
593 doi:10.5904/wgms-fog-2012-11, 2012.

594 WGMS: Glacier Mass Balance Bulletin No. 12 (2010-2011), edited by: Zemp, M., Nussbaumer, S. U., Naegeli,
595 K., Gärtner-Roer, I., Paul, F., Hoelzle, M., and Haerberli, W.: ICSU(WDS)/IUGG(IACS)/UNEP/UNESCO/WMO,
596 World Glacier Monitoring Service, Zurich, Switzerland, 106 pp., publication based on database version: doi:
597 10.5904/wgms-fog-2013-11, 2013.

598 WGMS: Fluctuations of Glaciers Database. World Glacier Monitoring Service, Zurich, Switzerland.
599 DOI:10.5904/wgms-fog-2015-11, 2015. Online access: <http://dx.doi.org/10.5904/wgms-fog-2015-11>

600 Zemp, M., Frauenfelder, R., Haerberli, W., and Hoelzle, M.: Worldwide glacier mass balance measurements:
601 general trends and first results of the extraordinary year 2003 in Central Europe, *Data of Glaciological*
602 *Studies [Materialy glyatsiologicheskikh issledovaniy]*, 99, Moscow, Russia, 3–12, 2005.

603 Zemp, M., Paul, F., Hoelzle, M., and Haerberli, W.: Glacier fluctuations in the European Alps 1850-2000: an
604 overview and spatio-temporal analysis of available data, in *The darkening peaks: Glacial retreat in scientific*
605 *and social context*, edited by: Orlove, B., Wiegandt, E., and Luckman, B. H., University of California Press,
606 Berkeley, 152-167, 2008.

607 Zemp, M., Hoelzle, M., and Haerberli, W.: Six decades of glacier mass-balance observations: a review of the
608 worldwide monitoring network, *Ann. Glaciol.*, 50(50), 101-111, doi: 10.3189/172756409787769591, 2009.

609

610

611

612

Tables

613 Table 1 – Physical characteristics of the Italian glaciers with the mass balance series analysed in this study
 614 (year 2006, NextData - DATAGRALP project: [http://www.nextdataproject.it/?q=en/content/special-project-](http://www.nextdataproject.it/?q=en/content/special-project-datagrulp)
 615 [datagrulp](http://www.nextdataproject.it/?q=en/content/special-project-datagrulp), last access: 27 September 2015; Salvatore et al., 2015).

Glacier	Geographic area	Area (km ²)	Minimum elevation (m a.s.l.)	Maximum elevation (m a.s.l.)	Median elevation (m a.s.l.)	Prevailing aspect	Average slope (°)	First survey year
Grand Etrèt	Gran Paradiso	0.47	2667	3190	2894	N	23	2002
Ciardoney	Gran Paradiso	0.59	2855	3170	3039	E-NE	18	1992
Fontana Bianca	Ortles-Cevedale	0.48	2889	3342	3166	E	23	1984
Sforzellina	Ortles-Cevedale	0.29	2790	3046	2868	NW	16	1987
Lunga	Ortles-Cevedale	1.86	2678	3378	3128	NE	19	2004
Careser	Ortles-Cevedale	2.39	2868	3279	3069	S	11	1967
La Mare (southern branch)	Ortles-Cevedale	2.16	2652	3518	3215	NE	21	2003
Pendente	Val Ridanna	0.95	2621	3064	2781	S	15	1996
Malavalle	Val Ridanna	6.92	2512	3441	2971	SE	14	2002

616

617

618

619 Table 2 – Mean values (and STD in brackets) of B_w , B_s , B_a and AAR for nine Italian glaciers in the period from
 620 2004 to 2013 (Car = Careser, FB = Fontana Bianca, Pen = Pendente, Cia = Ciardoney, Sfo = Sforzellina, GE =
 621 Grand Etrèt, Lun = Lunga, Mar = La Mare, Mal = Malavalle). Values expressed in mm w.e. except AAR that is
 622 in percent.

	Car	Mar	FB	Sfo	Lun	Pen	Mal	Cia	GE
B_w (9 years)	927 (330)	989 (301)	1085 (338)	\	991 (222)	1537 (425)	1194 (256)	1052 (421)	1472 (578)
B_s (9 years)	-2740 (368)	-1758 (303)	-2183 (457)	\	-2151 (368)	-2857 (525)	-2087 (386)	-2510 (378)	-2396 (321)
B_a (10 years)	-1788 (590)	-763 (395)	-1088 (642)	-1399 (505)	-1195 (466)	-1231 (692)	-825 (484)	-1419 (646)	-946 (648)
AAR (10 years)	1 (3)	25 (14)	11 (22)	\	12 (16)	4 (8)	23 (17)	3 (5)	\

623

624

625

626 Table 3 – Correlation matrix of B_a for nine Italian glaciers. * and ** indicate Spearman correlation significant
627 at the 0.05 and 0.01 level, respectively.

	Car	FB	Pen	Cia	Sfo	GE	Lun	Mar	Mal
Car	1.00								
FB	0.82**	1.00							
Pen	0.85**	0.76**	1.00						
Cia	0.87**	0.86**	0.55*	1.00					
Sfo	0.82**	0.75**	0.65**	0.81**	1.00				
GE	0.74**	0.77**	0.66*	0.62*	0.69*	1.00			
Lun	0.70*	0.73*	0.60	0.48	0.49	0.77**	1.00		
Mar	0.90**	0.96**	0.70*	0.90**	0.82**	0.80**	0.71*	1.00	
Mal	0.87**	0.84**	0.97**	0.61*	0.65*	0.68*	0.61	0.80**	1.00

628

629

630

631 Table 4 – Correlation coefficients of B_a vs. B_w and B_s . * and ** indicate Spearman correlation significant at
632 the 0.05 and 0.01 level, respectively.

	No of years	B_w	B_s
Car	40	0.46**	0.94**
FB	22	0.24	0.84**
Pen	12	0.84**	0.67*
Cia	22	0.51*	0.76**
GE	12	0.84**	0.66*
Lun	10	0.64*	0.69*
Mar	10	0.66*	0.64*
Mal	9	0.48	0.85**

633

634

635

636 Table 5 – Correlation coefficients of B_w , B_s and B_a vs. seasonal and annual NAO. Five-year triangular moving
 637 averages have been applied to the time series before correlation analyses. *, ** and *** indicate Spearman
 638 correlation significant at the 0.10, 0.05 and 0.01 level, respectively.

639

Nation	Glacier	Winter balance			Summer balance			Annual balance		
		DJF NAO	Oct-May NAO	Annual NAO	DJF NAO	Oct-May NAO	Annual NAO	DJF NAO	Oct-May NAO	Annual NAO
I	Car	-0.51**	-0.34	-0.16	-0.23	-0.05	0.34	-0.30*	-0.13	0.23
I	FB	-0.11	0.18	0.00	0.22	0.10	0.25	0.07	0.35	0.41*
I	Pen	-0.70	-0.90*	-0.70	-1.00**	-0.90*	-1.00**	0.12	0.15	0.23
I	Cia	0.17	0.34	0.19	-0.03	0.10	0.27	0.13	0.27	0.36
I	GE	-0.81***	-0.74**	-0.79**	-0.88***	-0.76**	-0.91***	-0.86***	-0.76**	-0.83***
I	Lun	-0.94**	-0.94**	-0.94**	-0.94**	-0.77	-0.94**	-0.89**	-0.83*	-0.89**
I	Mar	-1.00***	-0.89**	-1.00***	0.14	0.03	0.14	-0.79**	-0.82**	-0.89***
I	Mal	-0.90*	-0.70	-0.90*	-0.50	-0.80	-0.50	-0.81***	-0.74**	-0.45
I	Sfo							0.59***	0.59***	0.61***
F	St. Sorlin							-0.44***	-0.38***	-0.04
F	Sarennes	0.36***	0.43***	0.49***	-0.53***	-0.50***	-0.19	-0.41***	-0.37***	-0.05
CH	Silvretta	0.28*	0.13	-0.05	-0.57***	-0.45***	-0.18	-0.38***	-0.28**	0.02
CH	Gries	-0.39***	-0.22	-0.18	-0.58***	-0.53***	-0.25*	-0.49***	-0.39***	-0.03
A	Sonnblick							-0.48***	-0.45***	-0.07
A	Vernagt	0.27*	0.51***	0.49***	-0.42***	-0.38**	-0.09	-0.36**	-0.26*	0.07
A	Kesselwand							-0.42***	-0.38***	-0.07
A	Hintereis							-0.55***	-0.44***	-0.10

640

641

642

643 Table 6 - Spearman correlation coefficients and multiple regression results of B_a vs. seasonal mean
 644 temperature and precipitation. *, ** and *** indicate 0.05, 0.01 and 0.001 significance levels.

Air temperature – Correlation coefficients									
	Car	FB	Sfo	Cia	Pen	GE	Mal	Mar	Lun
No of years	47	27	27	22	18	12	12	11	10
Jun-Sep	-0.77***	-0.49**	-0.52**	-0.64**	-0.40	0.01	-0.24	-0.16	0.49
Oct-May	-0.37**	-0.42*	-0.10	-0.18	-0.49*	-0.18	-0.57	-0.28	-0.33
Year	-0.68***	-0.49**	-0.30	-0.49*	-0.69**	-0.31	-0.71**	-0.27	-0.14

Precipitation – Correlation coefficients									
	Car	FB	Sfo	Cia	Pen	GE	Mal	Mar	Lun
No of years	47	27	27	22	18	12	12	11	10
Jun-Sep	-0.15	-0.02	0.09	0.20	0.00	0.02	-0.04	0.26	-0.05
Oct-May	0.28	0.40*	0.32	0.47*	0.37	0.57	0.43	0.67*	0.71*
Year	0.11	0.36	0.39*	0.53*	0.34	0.64*	0.34	0.83**	0.53

Multiple linear regression - Coefficients									
	Car	FB	Sfo	Cia	Pen	GE	Mal	Mar	Lun
No of years	47	27	27	22	18	12	12	11	10
Jun-Sep temperature	-776.453 (***)	-663.487 (***)	-575.225 (***)	-796.739 (***)	-496.521 (**)	-63.899	-355.106	-668.941 (**)	115.265
Oct-May precipitation	2.186 (***)	3.342 (***)	2.915 (***)	3.315 (***)	2.380 (*)	2.897 (**)	3.051 (*)	4.122 (**)	2.666 (*)
Intercept	-3265.013 (***)	-3311.632 (***)	-3176.797 (***)	1753.826	1011.719	-2707.559	19.212	- 3380.905 (***)	- 2619.872 (**)
% of explained variance	75.6	68.7	72.1	73.7	51.5	56.8	56.0	78.4	64.5


645

646

647

Figures

648




649

650 Figure 1. Geographic setting of the glaciers with mass balance measurements analysed in this work
651 (Microsoft® BingTM Maps).


652

Val Ridanna





- Mass balance measurement points
- Contours (height interval = 50 m)

Ortles-Cevedale







Gran Paradiso



653


654

Figure 2. Surface topography and measurement network of the nine glaciers analysed in this study.



656 Figure 3. Time series of B_w , B_s , B_a and AAR for the nine Italian glaciers analysed.

657




658

659 Figure 4. Cumulative mass balance for the nine Italian glaciers and for a set of nine other glaciers
660 representative of the European Alps. Dotted and dashed lines indicate the average B_a for the two groups of
661 glaciers in the periods from 1967 to 1983, 1984 to 1993, 1994 to 2003 and 2004 to 2013.

662


663



664

665 Figure 5. Left column: mean ablation season (Jun-Sep) air temperature anomalies in (a) Gran Paradiso, (b)
 666 Ortles-Cevedale, and (c) Val Ridanna. Right column: mean accumulation season (Oct-May) air temperature
 667 anomalies in (d) Gran Paradiso, (e) Ortles-Cevedale, and (f) Val Ridanna. Reference period: 1961-1990. All
 668 linear trends are significant at the 0.001 level.


669



670

671 Figure 6. Left column: ablation season (Jun-Sep) total precipitation anomalies in (a) Gran Paradiso, (b)
 672 Ortles-Cevedale, and (c) Val Ridanna. Right column: accumulation season (Oct-May) total precipitation
 673 anomalies in (d) Gran Paradiso, (e) Ortles-Cevedale, and (f) Val Ridanna. Reference period: 1961-1990. None
 674 of the linear trends is significant at the 0.05 level.

675




676

677 Figure 7. Bootstrapped moving correlation coefficient between annual mass balance and seasonal values of
 678 air temperature and precipitation. Shaded straight lines indicate significance at 95% level.

679

680



681

682 Figure 8. Plot of residuals of the multiple linear regression of B_a vs. Oct-May precipitation and Jun-Sep
683 temperature on the Careser Glacier. Multiple regression coefficients are reported in Table 6. The inset
684 shows the extent of the glacier in three different years.

685

686

687

688

689

690

691

692

693

694

695

696

697