

1 **COVER LETTER**

2
3 We would like to thank the Editor and both Reviewers for their comments to the revised version of our
4 manuscript. In response to their suggestions, we did the following changes to the manuscript:

- 5 1) two figures have been added, showing the lengthening of the ablation season and the time series
6 of the DJF NAO index
7 2) a description of the analyses concerning the NAO index has been added in Section 3.3.
8 3) section 4.1 was reorganized in unnumbered subparts to improve clarity
9 4) a point has been added in the Conclusions, concerning the results of the NAO index analysis
10 5) we implemented the suggested changes and answered to the specific comments made by the
11 reviewers, as detailed in the following of this document. The authors responses are reported in
12 italic right below the reviewer comments. Line and page numbers are referred to the revised paper
13 (first review)

14
15 **Reviewer 1**

16 The authors have revised their paper based on the recommendation of two reviewers. Both reviews were
17 generally positive but requested some changes regarding the presentation of the results, as well as
18 additional analysis related to the spatial representativeness of the measurements and their interpretation.
19 In my opinion, the authors have adequately addressed all comments and have added the some new and
20 interesting analysis. The paper thus can be accepted subject to some minor corrections:

21 - Line 470: I am not sure if The Cryosphere accepts referencing papers that are not yet accepted (Carturan,
22 2016).

23 *According to the guidelines for authors, works in review should also be included in the reference list.*

24 - Line 472: What is “a good match”? Based on which method the authors did decide that no calibration of
25 the series is required. Some more details would be helpful here.

26 *We meant that the discrepancy between the geodetic and the glaciological method is lower than the lowest
27 detectable bias following Zemp and others, (2013). Sentence and reference added in the text.*

28 - Line 589: Here, the authors just state the results of their analysis. I would like to see at least one sentence
29 that interprets these numbers and tells the reader what they actually mean.

30 *Ok, sentence added.*

31 - Line 675-677: The sentence is not very clear. It would benefit from reformulation.

32 *The sentence was reformulated accordingly.*

33
34 **Reviewer 2**

35 This paper is under the second review process and I understand that the main changes recommended by
36 the 2 reviewers have been introduced almost accordingly. In this new version, the connection of the mass
37 balance terms of investigated glacier with NAO anomalies is quantitatively analyzed. Correlation matrixes
38 have been supplemented. I think the paper has gain in strength but nevertheless still needs some
39 improvements in lying on more substantial analyses. Here follow some points that could be developed in
40 that way:

41 Line 362: It is stated in the abstract that ablations season have lengthened but I have missed later in the
42 core of the paper where this is mentioned and how it is demonstrated. Can the authors provide an ablation
43 duration time plot for few glaciers showing this trend? Or a time plot of the number of days with positive
44 temperatures at the glacier elevation along the period of record of the mass balance series?

45 *A new figure (Figure 6) was added, displaying the number of days per year with maximum air temperature*
46 *exceeding 0°C at 3000 m a.s.l., calculated from the series of the Careser diga weather station (2600 m a.s.l.,*
47 *Ortles-Cevedale Group). A sentence was added in the text (Section 4.2) to comment the results.*

48 Line 478 and section 3.2 therein: As one of the covariates now analyzed with temperatures and
49 precipitations time series, NAO anomaly series worth a specific paragraph here, and I suggest particularly to
50 detail which monthly standardized series has been used, the choice of the cumulated anomalies (DJF,
51 December to February; etc...) and the retained data process for smoothing.

52 Figure 5 could be completed with a time plot of the raw-annual (DJF) NAO anomalies, the smoothed signal
53 and trends to display the well-known break points in 1970 and 1990.

54 *A description of the analyses concerning the NAO index has been added in Section 3.3. A new figure*
55 *depicting the time series of the winter NAO was added (Figure 8).*

56 Line 526 and section 4.1. I wonder how this section could be somewhat reorganized and divided in
57 (unnumbered) subparts. I get a little bit lost by the succeeding results of Bw, Ba, Bs and AAR series...

58 *This section was reorganized in unnumbered subparts to improve clarity.*

59 Lines 634-642. I would strongly encourage the authors to derive at sensitivity of summer balance to
60 temperature from the correlations of Bs to temperature, and to analyze the result for instance with respect
61 to "median elevation" of the glacier provided in Table 1.

62 *We calculated the correlation between Bs and temperature. Unfortunately, it is statistically significant only*
63 *for 3 out of 8 glaciers, i.e. only for the ones with the longer Bs series. The suggested analysis and*
64 *intercomparison would probably require longer Bs series for obtaining meaningful results.*

65 Line 650. Remarkable is the finding of the change point for winter precipitations in 1977 in Ortles-Cevedale
66 and Val Ridanna ranges. Until present this has been identified to have a rather regional significance limited
67 to the western Alps.

68 *Ok, emphasized in the text.*

69 Line 667. Indicate here or in section 3.2/3.3 the way NAO monthly anomalies are smoothed (5
70 years=current +/- 2 years?) and the way the authors chose this low-pass filter. Noteworthy to mention in
71 the text is that smoothing is proceed to highlight mainly possible convergent low-frequency patterns which
72 are not detectable at the annual scale.

73 *Ok, added in the text (Section 3.3).*

74 Some comments/questions arise from the new analyses:

75 - Most of the studied Eastern Italian glaciers have their Bw anti-correlated with DJF NAO anomalies. This is
76 in conflict with the result from Marzeion and Nesje (2012) who reported positive connections.

77 *The mentioned authors report positive correlation between Bw and DJF NAO anomalies along the northern
78 boundary of the Eastern Alps, whereas there is anti-correlation in the central and southern parts.*

79 - Some glaciers have stronger connection with the winter NAO through their summer balance (Pendente) or
80 with their winter balance (La Mare, Careser), some glaciers neither on summer or winter balance
81 components (Ciardoney). For Pendente, the links are balanced and cancel in the annual balance. The figure
82 then is quite composite. When significant, the link is always negative with the winter balance. However,
83 this link is found (surprisingly) positive for Vernagtferner and Sarenne.

84 *This is not surprising because previous works cited in our paper report contrasting behaviour of
85 precipitation anomalies in the Oct-May period between the northern and southern Alps, i.e. opposite
86 correlation with indexed large scale circulation patterns (e.g., Quadrelli et al., 2001; Schmidli et al., 2002,
87 Brunetti et al., 2006; Marzeion and Nesje, 2012).*

88 - How local precipitations are connected to the larger NAO signal? This may help to make clear the
89 preceding question.

90 *The winter precipitation anomalies of the three geographic areas where the Italian mass balance glaciers
91 are located (Section 2) are also anti-correlated with the winter NAO (Spearman correlation significant at the
92 0.01 level). Sentence added in the text (Section 4.2).*

93 Line 752. Conclusion

94 Add between points 3-4 a conclusion on the NAO analysis; Important is that some of the mass balance
95 series are connected to the synoptic signal held by NAO index through both summer and winter
96 components, sometimes with a strong link. But sometimes the link is weak and there is not a unique figure
97 and an understandable spatial structure.

98 *Ok, point added in the Conclusions.*

99

100

101

102

103

104

105

Analysis of the mass balance time series of glaciers in the Italian Alps

106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139

L. Carturan^{*1,6}, C. Baroni^{2,3,6}, M. Brunetti⁴, A. Carton^{5,6}, G. Dalla Fontana^{1,6}, M. C. Salvatore^{2,6}, T. Zanoner^{5,6,7}, G. Zuecco¹

¹ Dipartimento Territorio e Sistemi Agro-Forestali, University of Padova, Viale dell'Università 16, 35020, Legnaro, Padova, Italy

² Dipartimento di Scienze della Terra, University of Pisa, Via S. Maria 53, 56126, Pisa, Italy

³ IGG-CNR, Via Giuseppe Moruzzi 1, 56124, Pisa, Italy

⁴ ISAC-CNR, Via Gobetti 101, 40129, Bologna, Italy

⁵ Dipartimento di Geoscienze, University of Padova, Via Gradenigo 6, 35131, Padova, Italy

⁶ Comitato Glaciologico Italiano, Corso Massimo d'Azeglio 42, 10125, Torino, Italy

⁷ CNR-IRPI, Strada delle Cacce 73, 10135, Torino, Italy

Corresponding author: luca.carturan@unipd.it

Abstract

This work presents an analysis of the mass balance series of nine Italian glaciers, which were selected based on the length, continuity and reliability of observations. All glaciers experienced mass loss in the observation period, which is variable for the different glaciers and ranges between 10 and 47 years. The longest series display increasing mass loss rates, which were mainly due to increased ablation during longer and warmer ablation seasons. The mean annual mass balance (B_a) in the decade from 2004 to 2013 ranged from -1788 mm to -763 mm w.e. y^{-1} . Low-altitude glaciers with low elevation ranges are more out of balance than the higher, larger and steeper glaciers, which maintain residual accumulation areas in their upper reaches. The response of glaciers is mainly controlled by the combination of Oct-May precipitations and Jun-Sep temperatures, but rapid geometric adjustments and atmospheric changes lead to modifications in their response to climatic variations. In particular, a decreasing correlation of B_a with the Jun-Sep temperatures and an increasing correlation with Oct-May precipitations are observed for some glaciers. In addition, the Oct-May temperatures tend to become significantly correlated with B_a , possibly indicating a decrease in the fraction of solid precipitation, and/or increased ablation, during the accumulation season. Because most of the monitored glaciers have no more accumulation area, their observations series are at risk due to their impending extinction, thus requiring a soon replacement.

140

141 **1 Introduction**

142 The mass balance of glaciers is a key variable for monitoring strategies of the Earth climate system because
143 it is the direct and undelayed response of glaciers to atmospheric conditions. Other reactions of glaciers to
144 climatic changes, such as the fluctuations of the front, are more easy and immediate to measure but
145 represent indirect, delayed and filtered signals (WGMS, 2008; Zemp et al., 2005).

146 The direct glaciological method (Østrem and Brugman, 1991) is the standardized method in worldwide
147 glacier monitoring strategies. This method consists of in-situ measurements of the surface accumulation
148 and ablation, taken at single points and then extrapolated and integrated to yield the glacier-wide surface
149 mass balance (Kaser et al., 2003; Cogley et al., 2011). The World Glacier Monitoring Service (WGMS)
150 collects and publishes mass balance data of glaciers obtained by the glaciological method as part of global
151 climate-related observation systems (Zemp et al., 2009; WGMS, 2012 and 2013, and earlier issues).

152 The European Alps are one of the regions of the world with the highest density of glaciers that are subject
153 to mass balance observations. Twenty five glaciers have ongoing and continuous mass balance series with
154 at least 10 years of observations, and 11 of them are longer than 30 years
155 (<http://www.wgms.ch/metadatabrowser.html>, last access: 27 September 2015). In the Italian Alps, nine
156 glaciers have ongoing and continuous mass balance series longer than 10 years, and only one glacier (the
157 Careser Glacier) has a series longer than 30 years.

158 The mass balance series of the glaciers in the Italian Alps have not yet been reviewed and analysed jointly.
159 The Italian glaciers may have a peculiar behaviour compared to the glaciers from other regions of the
160 European Alps, because of the differences in glacier characteristics, climatic features and trends of
161 meteorological variables (Brunetti et al., 2006 and 2009; Auer et al., 2007). Differences may occur in the
162 response of the glaciers in different sub-regions of the Italian Alps or with different characteristics, which
163 have not been recognized. It is also interesting to highlight possible feedbacks in the response of Italian
164 glaciers to atmospheric changes.

165 Therefore, this work aims to i) analyse and compare the direct mass balance series of the glaciers in the
166 Italian Alps, ii) understand the behaviour of the measured glaciers in relation to the observed climatic
167 trends, and iii) highlight possible future requirements for the mass balance monitoring strategy in the
168 Italian Alps.

169

170 **2 Available mass balance series**

171 In this work, we analyse the glaciers with at least 10 years of continuous and ongoing mass balance
172 measurements, which were obtained using the direct glaciological methods and published in peer reviewed
173 journals or in the WGMS publications (CGI, 1914–1977 and 1978–2011; Baroni et al., 2012, 2013, 2014;
174 WGMS, 2012 and 2013, and earlier issues). "Continuous" indicates the series with data gaps <10%,
175 and "ongoing" indicates that the mass balance observations have been performed in the last two years (i.e.,
176 the 2012 and 2013 hydrological years). These criteria ensure the comparability of the series, a sufficient
177 length in the temporal analyses and reliability of the measurements and calculations.

178 Nine monitored glaciers fulfil these characteristics in the Italian Alps and are clustered in three geographic
179 areas (Fig. 1). The two monitored glaciers in the Gran Paradiso Group (Western Alps), i.e., Grand Etrèt
180 (since 2002) and Ciardoney (since 1992), are rather small (area < 1 km²) and have low mean elevations and
181 low elevation ranges (Table 1). Snowfall is the prevailing feeding source, but windborne snow and
182 avalanching also contribute to snow accumulation.

183 The longest series of mass balance measurements in the Italian Alps has been collected on the Careser
184 Glacier, in the Ortles-Cevedale (Eastern Alps, Fig. 1) since 1967. Currently, this glacier is undergoing rapid
185 shrinking and fragmentation in smaller units. It is characterized by a flat surface, prevailing southern
186 exposure, quite low mean elevation and feeding by snowfall. Its area decreased from 5 km² in 1967 to 1.6
187 km² in 2012 (Carturan et al., 2013a). In the 1980s, observations started in two other glaciers of the Ortles-
188 Cevedale: Fontana Bianca and Sforzellina. These two small mountain glaciers (area <0.5 km²) have different
189 characteristics: Fontana Bianca is rather steep with negligible debris cover and mainly fed by snowfall,
190 whereas Sforzellina is flatter, debris-covered in its lower part and fed by avalanches in its upper part. In the
191 2000s, mass balance observations began in the Lunga Glacier and in the southern branch of La Mare Glacier,
192 which are larger valley glaciers (1.9 and 2.2 km², respectively) that reach higher elevations (3378 and 3518
193 m, respectively) and mainly fed by snowfall.

194 In Val Ridanna (Breonie Occidentali Group, Eastern Alps, Fig. 1), the measurements began in 1996 in the
195 Pendente Glacier and were extended to the Malavalle Glacier in 2002. The first glacier is a 0.9 km² wide
196 mountain glacier, characterized by a flat surface, low mean elevation, southern exposure and significant
197 accumulation from windborne snow. The second glacier is a much larger (6.9 km²) valley glacier with higher
198 mean and maximum elevation and mainly fed by snowfall.

199

200 **3 Methods**

201

202 **3.1 Mass balance measurements and calculations**

203 Point measurements of the annual mass balance in the ablation area consist of repeated readings of
204 ablation stakes, which are made of aluminium, wood or plastic and drilled into the ice/firn using hand drills
205 or steam drills. In the accumulation area, the depth of the snow at the end of the ablation season is
206 measured using hand probes, and its density is determined in snowpits. Snow depth soundings and density
207 measurements in the snowpits or by hand coring devices are also used for winter mass balance
208 measurements, which are performed on all glaciers except Sforzellina. The summer mass balance is derived
209 by subtracting the winter mass balance from the annual mass balance. The density of measuring points
210 varies among different glaciers in relation to their extent, accessibility and complexity of the mass balance
211 distribution (Fig. 2). The ablation stake density ranges from 4 points km⁻² (Malavalle Glacier) to 45 points
212 km⁻² (Sforzellina Glacier). The density of snow depth soundings for the winter mass balance determination
213 ranges from 15 points km⁻² (Malavalle Glacier) to 142 points km⁻² (Fontana Bianca Glacier).

214 Point measurements are interpolated and extrapolated to the entire area of the glaciers using different
215 procedures. In the Grand Etrèt and Ciardoney glaciers, each ablation stake is assumed representative of a
216 specific part of the glacier, where the mass balance distribution is assumed homogeneous. Then, a
217 weighted mean is calculated, using the area of the homogeneous parts into which the glacier is subdivided

218 as weights (<http://www.pngp.it/>, last access: 27 September 2015; <http://www.nimbus.it/>, last access: 27
219 September 2015). In Malavalle and Pendente glaciers, the area is divided into “sub-catchments”, and for
220 each sub-catchment, a linear regression of point balances vs. altitude is calculated and used for the
221 spatialization (<http://www.provinz.bz.it/wetter/glacierreport.asp>, last access: 27 September 2015). In
222 Careser Glacier, the even distribution of the mass balance and good coverage of measurement points
223 enable the use of automatic interpolation algorithms (Spatial Analyst Tools) in the ESRI-ArcGIS software.
224 Manual drawing of balance isolines is used for the remaining Sforzellina, Fontana Bianca, La Mare and
225 Lunga glaciers (Catasta and Smiraglia, 1993; Cannone et al., 2008;
226 <http://www.provinz.bz.it/wetter/glacierreport.asp>; Carturan, 2016). The measurements are performed
227 close to the time of maximum and minimum mass balance during the year, when the glacier and
228 atmospheric conditions are favourable for field surveys. The “floating-date” time measurement system is
229 used for all glaciers (Cogley et al., 2011).

230 Typical random errors reported in the literature for glacier-wide mass balance estimates obtained with
231 these methods are of about ± 200 mm w.e. γ^{-1} (Lliboutry, 1974; Braithwaite and Olesen, 1989; Cogley and
232 Adams, 1998; Cogley, 2009). The accuracy indicated by the investigators carrying out mass balance
233 measurements in the nine Italian glaciers range between ± 0.05 and 0.30 m w.e. γ^{-1} (WGMS, 2015; Carturan,
234 2016). Assessments based on the comparison between the direct and the geodetic mass balance have been
235 published for the Careser, La Mare and Lunga glaciers, indicating that the discrepancy between the two
236 methods is lower than the lowest detectable bias (following Zemp and others, 2013), a good match of the
237 two methods and revealing that a calibration of the direct mass balance results is not required (Carturan et
238 al., 2013a; Galos et al., 2015; Carturan, 2016).

239

240 **3.2 Meteorological series**

241 The climatic variables used in this work consist of synthetic records of the monthly mean temperature and
242 total monthly precipitation, which are obtained for the centre of the three main geographic areas described
243 in Sect. 2, using the procedure reported in Brunetti et al. (2012). Starting from sparse meteorological data
244 recorded at meteorological stations, the synthetic meteorological series are generated using the anomaly
245 method (New et al., 2000; Mitchell and Jones, 2005). This method is based on the assumption that the
246 spatio-temporal structure of the signal of a meteorological variable over a given area can be described by
247 the superimposition of two fields: the climatological normals over a given reference period (i.e., the
248 climatologies) and the departures from them (i.e., the anomalies). The climatologies are linked to the
249 geographic features of the territory and characterized by remarkable spatial gradients; the departures are
250 linked to the climate variability and change, and they are generally characterized by higher spatial
251 coherence.

252 Under this assumption, the climatologies and anomalies can be reconstructed in completely independent
253 manners and based on different data sets. For climatologies, the priority is the high spatial resolution, and a
254 short time span (few decades) is sufficient. A lower spatial resolution is sufficient for the anomalies, but
255 more importance is given to the data quality and availability of long records. Thus, all series that were used
256 for the anomaly component were subjected to homogenization.

257 The interpolation methods are different for the two components. The climatologies, which are
258 characterized by high spatial gradients, were reconstructed using the procedure in Brunetti et al. (2014),
259 exploiting the relationship between the meteorological variable and the physical characteristics of the

260 terrain. The anomalies, which are characterized by higher spatial coherence, were reconstructed using
261 weighted averages as described in Brunetti et al. (2006). The weights are horizontal and vertical distance
262 weighting functions, with the addition of an angular weight that accounts for the anisotropy in the
263 distribution of stations around the sites. Finally, the two fields were superimposed to obtain the temporal
264 series in absolute values for each site.

265

266 **3.3 Analyses of the mass balance and meteorological series**

267 The time series of annual mass balance (B_a), winter mass balance (B_w), summer mass balance (B_s) and
268 Accumulation Area Ratio (AAR, i.e., the ratio of the area of the accumulation zone to the area of the glacier)
269 were analysed and compared to highlight the possible trends, break points, common behaviour and
270 peculiarities of single glaciers and/or single years. To highlight the systematic differences among the
271 glaciers, the mean values of B_a , B_w , B_s and AAR were calculated in the common period of observation from
272 2004 to 2013. The decadal means of B_a for the Italian glaciers were compared to the decadal means for a
273 sample of nine representative glaciers of the European Alps (Zemp et al., 2005). The correlations among the
274 B_a series of different glaciers and among B_a of single glaciers with the respective series of B_w and B_s were
275 subsequently computed, to identify possible groups of glaciers with similar behaviours and to understand
276 the relative importance of the seasonal components of mass balance.

277 Linear trends and moving averages were calculated for the time series of air temperature and precipitation
278 to highlight the climatic drivers of the observed glacier changes. In particular, we focused on the
279 precipitation of the accumulation season, from October to May (Oct-May), and on the air temperature of
280 the ablation season, from June to September (Jun-Sep) (Pelto, 2008; Carturan et al., 2013b), computing
281 their correlation with B_a and performing multiple linear regression analyses. For the four glaciers with the
282 longest mass balance series (Careser, Fontana Bianca, Sforzellina and Ciardoney), we performed a moving
283 correlation analysis of B_a vs. the seasonal and annual temperature and precipitation, to recognize possible
284 changes and/or trends in their response and sensitivity to climatic fluctuations, e.g., ascribable to
285 geometric adjustments. A correlation analysis of B_w , B_s and B_a versus the seasonal (December to February
286 and October to May) and annual mean North Atlantic Oscillation (NAO) index was performed, using the
287 mass balance data of glaciers from Italy and from other nations of the European Alps. Five-year (i.e.,
288 current +/- 2 years) triangular moving averages have been applied to the time series before correlation
289 analyses, to highlight possible convergent low-frequency patterns which are not detectable at the annual
290 scale.

291

292 **4 Results and discussion**

293 **4.1 Analysis of mass balance series**

294 **Annual balance**

295 The longest available series for the glaciers in the Italian Alps clearly show a trend towards more negative
296 B_a in the observation period (Fig. 3), and one or two change points, which were identified using the
297 'Changepoint' R package (Killick and Eckley, 2014). In particular, the series of Careser Glacier shows three
298 phases: i) the period from 1967 to 1980 with near equilibrium conditions (mean $B_a = -132$ mm w.e. y^{-1} , STD
299 = 540 mm w.e.); ii) the period from 1981 to 2002 with imbalanced conditions (mean $B_a = -1192$ mm w.e. y^{-1} ,

300 STD = 517 mm w.e.); and iii) the period after 2002 with stronger imbalance (mean $B_a = -1926$ mm w.e. y^{-1} ,
301 STD = 725 mm w.e.). ~~A sudden transition in 1980-81 is also clearly visible in the series of AAR, which~~
302 ~~documents the nearly complete disappearance of the accumulation area of this glacier, which is likely~~
303 ~~responsible for the step change in its mass balance series during that period.~~

304 The transition of 2002-03 is also observable for the Fontana Bianca, Sforzellina and Pendente glaciers,
305 whose measurements started in the 1980s and 1990s. Their mean B_a values changed from -599, -868 and -
306 703 mm w.e. y^{-1} , before 2002, to -1257, -1471 and -1308 mm w.e. y^{-1} after 2002, respectively. This
307 transition is less obvious for the Ciardoney Glacier, which experienced a notably negative mass balance
308 already in 1998 and 1999.

309 Seasonal balance and AAR

310 The B_w and B_s series have some gaps but suggest that the increased mass loss rates were mainly ascribable
311 to increased ablation (and associated positive feedbacks) instead of decreased snow accumulation. These
312 results are consistent with previous works, which indicate that the mass changes of the glaciers in the Alps,
313 at the annual and decadal scale, are mainly driven by the summer balance (e.g., Schonert et al., 2000;
314 Vincent et al., 2004; Zemp et al., 2008; Huss et al., 2015).

315 The AAR series show that the accumulation area almost vanished from all glaciers in the 2000s except the
316 years 2001, 2010 and 2013, when several glaciers were close to balanced-budget conditions mainly as a
317 result of the increased B_w . In these years, the highest increase in AAR occurred in the Fontana Bianca
318 Glacier, which is steep and exposed to the east. On the contrary, the AAR did not significantly increase in
319 the neighbouring Careser Glacier, which is flatter and mainly exposed to the south (Table 1). This behaviour
320 is uncommon for flat glaciers because they should be more sensitive to variations of the Equilibrium Line
321 Altitude (ELA) than the steeper glaciers (Benn and Evans, 2010), and reveals that the Careser Glacier is
322 almost completely below the current ELA, also in the years of ELA minima. La Mare and Malavalle glaciers,
323 which are larger and cover a wider elevation range (Table 1), show more persistent accumulation areas,
324 although their size is too small to ensure balanced-budget conditions.

325 The B_a and B_s values of different glaciers tend to diverge in years with largely negative mass balance and
326 converge in years closer to equilibrium (1993, 2001, 2010 and 2013, Fig. 3). Reinforcing processes and
327 feedbacks likely amplify the differences among the glaciers in imbalanced years, particularly the decrease in
328 the glacier-average albedo caused by the early disappearance of snow from low-lying, flat and less
329 topographic-shielded glaciers, and by the accumulation of dust and debris on the surface. B_w also shows the
330 alternation of years with small/large variability among the glaciers, but this behaviour cannot be clearly
331 related to the magnitude of the snow accumulation, as observed in the two high-accumulation years 2009
332 (high variability) and 2013 (low variability). In this case, the spatial variability of the precipitation during the
333 accumulation season, which is larger than the spatial variability of air temperature in the ablation season,
334 determines the interannual variability of B_w for single glaciers, which is further controlled by snow
335 redistribution processes. Snow redistribution appears more effective for the Pendente, Grand Etrèt and
336 Ciardoney glaciers, leading to over-accumulation in snow-rich winters (e.g., in 2009) and larger interannual
337 variability of B_w . Correlation coefficients calculated between B_w and October-May precipitations range
338 between 0.73 and 0.78 are significant at the 0.05 level only for Careser, La Mare, Lunga, Fontana Bianca
339 and Malavalle, while they are not statistically significant for Pendente, Grand Etrèt and Ciardoney, in line
340 with the hypothesised higher importance of snow redistribution processes in these three glaciers.

341

342 Comparisons in the common period from 2004 to 2013 and spatial representativeness

343 In the period from 2004 to 2013, significantly higher B_w is observed for Pendente and Grand Etrèt,
344 compared with the other glaciers in the same geographic area (Table 2), explaining the persistence of these
345 two ice bodies at such low altitude (Table 1). In the same period, the Careser Glacier had the lowest
346 average B_a and AAR, whereas the Malavalle and La Mare glaciers had the highest average B_a , B_s and AAR,
347 retaining accumulation areas in their upper parts. However, the mean AARs were remarkably low for all
348 analysed glaciers, and far from balanced-budget conditions ($AAR_0 = 0.55 - 0.58$, Dyurgerov et al., 2009;
349 Mernild et al., 2013). Overall, low-altitude and flat glaciers with low elevation ranges are more out of
350 balance than the steeper glaciers at higher altitude with higher elevation ranges, as acknowledged in
351 various other studies (e.g., Furbish and Andrews, 1984; Benn and Evans, 2010; Carturan et al., 2013b;
352 Fischer et al., 2015).

353 At the regional scale, the spatial representativeness of five Italian mass balance glaciers can be assessed on
354 the basis of the geodetic mass balance calculations performed by Carturan et al., (2013b). In the period
355 from the 1980s to the 2000s, the average geodetic mass balance rate of the 112 glaciers in the Ortles-
356 Cevedale Group has been -0.69 m w.e. y^{-1} . If we consider the average geodetic mass balance in the same
357 period as an index of the spatial representativeness for single glaciers, we obtain in decreasing order: i) La
358 Mare with -0.64 m w.e. y^{-1} , ii) Sforzellina with -0.86 m w.e. y^{-1} , iii) Fontana Bianca with -0.90 m w.e. y^{-1} , iv)
359 Lunga with -1.00 m w.e. y^{-1} , and v) Careser with -1.43 m w.e. y^{-1} . These results confirm that a proper
360 assessment of the spatial representativeness is required when inferring regional-scale mass balance
361 estimates using single glaciers. Geodetic calculations only exist for few areas in the Italian Alps (e.g., Galos
362 et al., 2015) and do not include the other four mass balance glaciers analysed in this study. Therefore it was
363 not possible to evaluate their spatial representativeness at the regional scale. Similarly, quantitative
364 assessments of the representativeness of all the nine glaciers at the scale of the entire Italian Alps will
365 require further investigations, integrating in-situ measurements, remotely sensed observations and
366 numerical modelling (WGMS, 2015).

367 Comparison with other glaciers in the European Alps

368 The response of Italian glaciers to the climatic conditions of the last decades is similar to that of nine
369 representative glaciers of the entire European Alps (Zemp et al., 2005; Fig. 4), although single glaciers
370 display different mass loss rates (Table 2). The Italian glaciers display ~ 200 - 250 mm w.e. y^{-1} more negative
371 B_a until 2002 and ~ 200 mm w.e. y^{-1} less negative B_a since then. Therefore, it can be assessed that the mean
372 B_a values for the Italian and “European” glaciers are fairly similar. Comparable results were obtained by
373 Huss et al., (2015), who compared the decadal mean B_a of glaciers from France, Switzerland, Austria and
374 Italy. These comparisons may be affected by the loss of spatial representativeness of some glaciers (e.g.
375 Careser in the Italian Alps and Sarennes in the French Alps) and by the different subsets of Italian glaciers
376 which are useable in the four different sub-periods. In the last decade, the inclusion of La Mare and
377 Malavalle glaciers in the Italian subset and the concurrent sharp decrease of B_a for the Sarennes, St. Sorlin
378 and Gries glaciers explain the different behaviours of the two groups of glaciers. However, the smaller
379 Italian glaciers (average area = 1.79 km²) may have a shorter response time to climatic changes, adjusting
380 their geometry faster than the larger glaciers (average area = 3.63 km²) which are representative of the
381 European Alps (Hoelzle et al., 2003; Abermann et al., 2009). The rapid shrinking and fragmentation of
382 Careser Glacier is a good example: in the last decade, its area has halved, and it completely lost the parts
383 subject to higher ablation (Carturan et al., 2013a). Changes in the general atmospheric circulation and
384 spatial distribution of precipitation could also have played a role and will be discussed in Sect. 4.2.

385 Correlation analyses

386 There is a generally high correlation among the B_a values of the analysed glaciers (Table 3). The series of
387 Careser, Fontana Bianca and La Mare glaciers show a highly significant correlation with most other glaciers,
388 even if they have different characteristics or are far away. On the contrary, the Lunga Glacier shows a lower
389 correlation and lower statistical significance with the glaciers of the same mountain group. However, it has
390 the shortest series, and most importantly, it does not include the highly negative B_a of 2003, which certainly
391 increases the correlation among other glaciers. There are notably high correlations in the Ortles Cevedale
392 between Careser and La Mare and between Fontana Bianca and La Mare glaciers. A similarly high
393 correlation is observed between Pendente and Malavalle glaciers in Val Ridanna, whereas there is a much
394 lower correlation between the two glaciers of the Gran Paradiso Group, which suggests that differences in
395 local topo-climatic factors can be decisive on such small ice bodies (e.g., Kuhn, 1995; DeBeer and Sharp,
396 2009; Carturan et al., 2013c; Scotti et al., 2014; Colucci and Guglielmin, 2015).

397 For most glaciers, B_a is more correlated to B_s than to B_w (Table 4), which further confirms the importance of
398 summer ablation. The relevance of the snow redistribution and over-accumulation on the Pendente and
399 Grand Etrèt glaciers is indicated by the higher correlation of their B_a with B_w . On La Mare Glacier, the two
400 seasonal components have similar correlations with B_a . However, these results are influenced by the length
401 of the observation period and the presence/absence of extreme years with high accumulation (e.g., 2001)
402 or high ablation (e.g., 2003) in the observation series of individual glaciers. For the analysed glaciers, no
403 significant correlation was found between B_s and B_w .

404

405 **4.2 Climatic controls**

406 In the period from 1961 to 2013, there are highly significant warming trends for the Jun-Sep air
407 temperature (Fig. 5a, b and c); they are highest in the Gran Paradiso Group ($0.40^\circ\text{C decade}^{-1}$) and lowest in
408 Val Ridanna ($0.35^\circ\text{C decade}^{-1}$). The three phases in the longer B_a and B_s series of glaciers (Fig. 3) can be
409 recognized as periods with stationary Jun-Sep temperature, separated by switches in the early 1980s and
410 after the peak of 2003. The warming trends are lower in the accumulation season and range from 0.25 to
411 $0.27^\circ\text{C decade}^{-1}$ (Fig. 5d, e and f), but thermal inversions at the valley weather stations could have partially
412 masked the warming at the altitude of the glaciers in this season. The transition towards higher Oct-May
413 temperature occurred in the late 1980s, after a minimum in the first half of the same decade. A distinct
414 warm peak in Oct-May temperature occurred in 2007. The warming trend led to increased duration of the
415 ablation period. The number of days per year with maximum temperature exceeding 0°C , extrapolated at
416 3000 m a.s.l. from the series of the Careser diga weather station (2600 m a.s.l. in the Ortles-Cevedale Group)
417 with a lapse rate of $0.65^\circ\text{C}/100\text{ m}$, increased from 160-170 in the 1960s-1970s to about 190 in the late
418 1990s and 2000s (Fig. 6).

419 The precipitation does not show any significant trend in the accumulation season (Fig. ~~6d~~7d, e and f). The
420 moving averages display oscillations of 10-20% above and below the 1961-1990 mean, which lasted
421 approximately 10-15 years and were higher in the Gran Paradiso Group than in the Ortles-Cevedale and Val
422 Ridanna. Periods with below-average precipitation are recognized in the 1960s, first half of 1970s, and
423 1990s, whereas periods with above-average precipitation occurred in the second half of 1970s and the first
424 half of 1980s. The last 10-15 years were characterized by precipitation close to the mean, with important
425 maxima in 2001, 2009 and 2013, and minima in 2007 and 2012. Similarly to the findings from Durand et al.,
426 (2009a and b) and Eckert et al., (2011) for the French and western Swiss Alps, a-change points in winter

427 precipitation of Ortles-Cevedale and Val Ridanna series were identified in 1977, corresponding to an
428 increase of about 10-12%. ~~This finding is remarkable because, until present, this change point has been~~
429 ~~identified to have a rather regional significance limited to the western Alps.~~ Linear trends of summer
430 precipitation are positive but not statistically significant. The interannual variability of the Jun-Sep
431 precipitation is remarkably higher in the Gran Paradiso Group (Fig. ~~6a7a~~, b and c).

432 Large scale circulation patterns, such as the North Atlantic Oscillation (NAO) and the Northern Hemisphere
433 blocking frequency, are connected with the temporal and spatial variability of winter precipitation in the
434 Alps (Quadrelli et al., 2001). Several studies highlighted ~~a~~-contrasting behaviour of precipitation anomalies
435 in the Oct-May period between the northern and southern Alps, i.e. opposite correlation with indexed large
436 scale circulation patterns (e.g., Quadrelli et al., 2001; Schmidli et al., 2002, Brunetti et al., 2006) and
437 opposite long-term trends in the seasonal precipitation totals (e.g., Brunetti et al., 2006 and 2009; Auer et
438 al., 2007). This characteristic, and the tendency towards a decreasing NAO index in the last two decades
439 (<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/pna/new.nao.shtml>~~http://www.cpc.ncep.noaa.g~~
440 ~~ov/products/precip/CWlink/pna/JFM_season_nao_index.shtml~~, last access: ~~4–11 October–February~~
441 ~~20152016~~; Fig. 8), leading to increased winter precipitation in the southern side of the Alps, may provide an
442 additional explanation for the different behaviour of “European” and “Italian” glaciers shown in Fig. 4.
443 Opposite effects of the NAO on the winter precipitation and glacier mass balance in the northern and
444 southern parts of the Eastern Alps were also reported, for example, by Marzeion and Nesje (2012).

445 ~~Our A-correlation analysis of B_w , B_s and B_a versus the seasonal and annual mean NAO index confirm that~~
446 ~~was performed, using the mass balance data of glaciers from Italy and from other nations of the European~~
447 ~~Alps (Table 5). The results show pp~~prevailing negative correlations ~~exists~~ between the B_w of Italian glaciers
448 and the NAO in the accumulation season, whereas positive correlations prevail in other nations (Table 5),
449 with the exception of Gries Glacier which however is close to the Italian border. ~~The winter precipitation~~
450 ~~anomalies of the three geographic areas where the Italian mass balance glaciers are located (Section 2) are~~
451 ~~also anti-correlated with the winter NAO (Spearman correlation significant at the 0.01 level).~~ ~~These results~~
452 ~~are in agreement with the mentioned literature and with our hypothesis of opposite effects of recent NAO~~
453 ~~trends in the winter precipitation of the northern and southern sides of the Alps.~~ In line with the findings of
454 Reichert et al., (2001), Six et al., (2001), and Thibert et al., (2013), a negative correlation was calculated
455 between B_s/B_a and the NAO in the accumulation season. ~~If a causal relationship can be hypothesised ff~~
456 ~~for the Italian glaciers~~ ~~the, related to the~~ albedo feedback from wet/dry winters (with low/high NAO,
457 respectively) ~~can at least partly explain this behaviour.~~ ~~the same cannot be stated ff~~ for glaciers in other
458 countries ~~however, given the, due to the~~ prevailing ~~lent~~ positive correlation of their B_w with the winter NAO,
459 ~~the link between B_s/B_a and the winter NAO is not so obvious and deserves additional analyses.~~

460 The examination of meteorological series confirms that increased ablation and the related feedbacks are
461 the main causes of the increased imbalance of the analysed Italian glaciers, as observed in Sect. 4.1. This
462 result is further corroborated by the higher correlation of B_a with the Jun-Sep temperature than with the
463 Oct-May precipitation, at least for the glaciers with longer observation series (Careser, Fontana Bianca,
464 Sforzellina and Ciardoney, Table 6). B_a of glaciers with shorter observation series is not significantly
465 correlated with the Jun-Sep temperature; instead, two of them (La Mare and Lunga) show a correlation
466 with the Oct-May precipitation. Combining the Oct-May precipitation and Jun-Sep temperature in a
467 multiple linear regression model leads to highly significant coefficients for both variables, even when the
468 single seasonal components are not correlated with B_a (e.g., for the Pendente and La Mare glaciers).
469 Approximately two-thirds of the B_a variance can be explained by the multiple linear regression. The
470 poorest results were obtained for the two glaciers in Val Ridanna (Pendente and Malavalle) and the Grand

471 Etrèt Glacier. As the first two glaciers are close to the main Alpine divide, they likely benefit from the high
472 orographic uplift that locally enhances precipitation (Schwarb, 2000), but which cannot be accounted for by
473 the multiple regression model due to the lack of weather stations in that area. In addition, the multiple
474 regression model does not account for accumulation by windborne snow on the Pendente and Grand Etrèt
475 glaciers.

476 The Careser, Fontana Bianca and Pendente glaciers display significant negative correlations between their
477 B_a and the Oct-May temperature. For the Careser Glacier, there is also a negative correlation between B_w
478 and the Oct-May temperature. Normally, in this period, most precipitation falls as snow, and the glaciers
479 have negligible ablation and low temperature sensitivity (Oerlemans and Reichert, 2000). However,
480 increasing temperature starts to lead to significant ablation in this period and to reduce the fraction of solid
481 precipitation as clearly detectable in the ablation season (Carturan et al., 2013b). An emblematic example is
482 the warm accumulation season of 2006-07, when the liquid precipitation reached 3000-3100 m a.s.l. (24
483 October 2006), and ice ablation exceeded 50 cm at 3000 m a.s.l. on the Careser and La Mare glaciers.

484 The correlation between B_a of Careser Glacier and the Oct-May temperature starts to become significant in
485 the late 1980s, as shown in the moving correlation analyses (30-year time window in Fig. 79). In the first 20
486 years, the correlation was absent or not statistically significant. These results are consistent with the
487 discussed effects of increasing temperature on the ablation and partitioning between liquid and solid
488 precipitation (Beniston et al., 2003). Reducing the window size from 30 to 15 years leads to a noisier signal
489 and in this case the correlation between B_a and Oct-May temperature does not reach the 95% significance
490 thresholds. However, it is interesting to remark the reversal of the correlation sign from positive in the first
491 years to negative in the last years.

492 The four glaciers in Fig. 7-9 (Careser, Fontana Bianca, Sforzellina and Ciardoney) share a common trend
493 towards i) a non-significant moving correlation between B_a and the Jun-Sep temperature and ii) a
494 significant moving correlation between B_a and Oct-May precipitation. This behaviour is probably related to
495 the snow-rich accumulation seasons of 2001, 2009 and 2013, and to the fact that the ablation season is
496 already so warm that i) summer snow falls mostly above the highest reaches of the glaciers, which reduces
497 the interannual variability of summer melt, and ii) conditions close to balanced-budget only occur after
498 snow-rich accumulation seasons.

499 Rapid geometric changes may also lead to a non-linear response of B_a to atmospheric changes, at least for
500 some glaciers. For example, the multiple regression residuals of the Careser Glacier, which were mostly
501 positive in the 1980s, 1990s and 2000s, became predominantly negative after 2008 (Fig. 810). This change
502 may suggest that the rapid modifications occurred in the latest years could have induced a negative
503 feedback, reducing the mass loss rate of the glacier, whose current surface and shape are strongly different
504 from the recent past (inset in Fig. 810). Because the multiple regression model does not use the Oct-May
505 temperature as an explanatory variable, it cannot account for the effects of the warm accumulation season
506 of 2006-07, which led to a very low B_w , early disappearance of winter snow and positive albedo feedback.
507 Therefore, the year 2007 results in highly positive regression residuals.

508

509 4.3 Future requirements

510 A common characteristic for all glaciers analysed is their very low mean AAR in the last decade (Table 2).
511 Accumulation areas were almost inexistent in most glaciers, indicating that they will soon disappear, even

512 without additional warming. Some glaciers are displaying morphological changes that indicate their
513 impending extinction, such as rapid disintegration (e.g., Careser Glacier, Fig. 810) and surface lowering in
514 the upper accumulation area (e.g., Fontana Bianca Glacier). The AARs of approximately 0.25 indicate that
515 accumulation areas still exist in the larger and higher-reaching La Mare and Malavalle glaciers. However,
516 given that balanced-budget conditions require AAR close to 0.55, large mass loss and areal reduction are
517 also expected for these two glaciers to reach equilibrium with the climatic conditions of the last ten years.

518 The forthcoming vanishing of the monitored glaciers put the continuation of their mass balance
519 observations at risk. Recently-started monitoring programs in larger and higher-reaching glaciers, such as
520 Malavalle and La Mare, will ensure continued observations in Val Ridanna and the Ortles-Cevedale. In line
521 with the recommendations from the WGMS (Zemp et al., 2009), similar observation programs should start
522 in other large and high-reaching glaciers of the Italian Alps, e.g., in the Gran Paradiso group (to substitute
523 Ciardoney and Grand Etrèt) and in other mountain groups. Both the initiation of observations over new
524 glaciers and the replacement of vanishing glaciers will require an assessment of the spatial
525 representativeness of single glaciers through the comparison of the current mass loss rates over wide
526 geographic areas (Haeberli et al., 2013). This assessments can be obtained using modern techniques such
527 as the multi-temporal differencing of digital elevation models, which enable the comparison of mass loss
528 rates in the last years/decades, by means of the geodetic method, over entire regions or mountain ranges
529 (e.g., Paul and Haeberli, 2008; Abermann et al., 2011; Carturan et al., 2013b; Berthier et al., 2014; Fischer et
530 al., 2015). The geodetic mass balance should also help to control the glacier-wide B_a series measured with
531 the direct glaciological method, and to construct a constant-geometry mass balance record (Elsberg et al.,
532 2001) to be connected to climatic drivers.

533

534 5 Conclusions

535 In this work, we have analysed the time series of the glaciers with mass balance observations in the Italian
536 Alps. Based on the results of the analyses, the following conclusions can be drawn:

- 537 ▪ All examined glaciers are experiencing imbalanced conditions, and the longer series show sustained
538 negative trends of B_a .
- 539 ▪ The observed behaviour was mainly caused by increased ablation, led by warmer temperature and
540 related feedbacks. The total precipitation does not show any significant trend, but the fraction of
541 solid precipitation decreased as a consequence of the warmer temperature.
- 542 ▪ The B_a of the analysed glaciers is mainly correlated to B_s , except for two glaciers where windborne
543 snow enhances the importance of B_w . For most glaciers, approximately two thirds of the B_a variance
544 can be explained by multiple linear regression, using the Oct-May precipitation and Jun-Sep
545 temperature as independent variables.
- 546 ▪ The monitored Italian glaciers have comparable mass loss rates to a sample of representative
547 glaciers of the entire European Alps. However, the moving correlation analyses and time series of
548 residuals from multiple linear regressions suggest that the smaller (and thinner) Italian glaciers may
549 be reacting faster to atmospheric changes.
- 550 ▪ Large scale circulation patterns, such as the NAO, have opposite effects in the northern and
551 southern sides of the European Alps. Most of the Italian mass balance series are anti-correlated to

552 [the synoptic signal held by the NAO index, through both winter and summer components,](#)
553 [sometimes with a strong link. However, in some cases the link is weak or absent and there is not a](#)
554 [clear spatial structure.](#)

555 ▪ Most monitored glaciers have no more accumulation area and are at risk of extinction, even
556 without additional warming. Therefore, they will soon require a replacement with larger and higher
557 glaciers that retain accumulation areas.

558 ▪ Regional assessments of the mass loss rates using the geodetic method are required to identify
559 possible replacing glaciers, evaluate their spatial representativeness and enable the transitions
560 from replaced to replacing glaciers, as suggested by Haerberli et al. (2013).

561

562 **Author contribution**

563 M. Brunetti processed the meteorological data and prepared the synthetic meteorological series used in
564 this work. Thomas Zanoner compiled the database of the mass balance data and geometric characteristics
565 of the glaciers. L. Carturan and Giulia Zuecco performed the temporal and statistical analyses of the mass
566 balance series. L. Carturan prepared the manuscript with contributions from all co-authors.

567

568 **Acknowledgments**

569 This study was funded by the Italian MIUR Project (PRIN 2010-11): "Response of morphoclimatic system
570 dynamics to global changes and related geomorphological hazards" (local and national coordinators G.
571 Dalla Fontana and C. Baroni). The authors acknowledge the regional environmental agencies of Piemonte,
572 Val d'Aosta, Lombardia, Autonomous Provinces of Trento and Bolzano, which provided the meteorological
573 and topographic data used in this study. Many thanks to Stephan Galos (University of Innsbruck) for sharing
574 information on glaciers in Ötztal and South Tyrol. We would also like to thank all investigators and
575 contributing institutions for the collection and free exchange of the glacier mass balance series. [Etienne](#)
576 [Bertier and two anonymous referees provided useful suggestions in the review process.](#)

577

578 **References**

579 Abermann, J., Lambrecht, A., Fischer, A., and Kuhn, M.: Quantifying changes and trends in glacier area and
580 volume in the Austrian Ötztal Alps (1969-1997-2006), *The Cryosphere*, 3, 205-215, doi:10.5194/tc-3-205-
581 2009, 2009.

582 Abermann, J., Kuhn, M., and Fischer, A.: Climatic controls of glacier distribution and glacier changes in
583 Austria. *Ann. Glaciol.*, 52(59), 83-90, 2011.

584 Auer, I., Böhm, R., Jurkovic, A., Lipa, W., Orlik, A., Potzmann, R., Schöner, W., Ungersböck, M., Matulla, C.,
585 Briffa, K., Jones, P., Efthymiadis, D., Brunetti, M., Nanni, T., Maugeri, M., Mercalli, L., Mestre, O., Moisselin,
586 J.-M., Begert, M., Müller-Westermeier, G., Kveton, V., Bochnicek, O., Stastny, P., Lapin, M., Szalai, S.,
587 Szentimrey, T., Cegnar, T., Dolinar, M., Gajic-Capka, M., Zaninovic, K., Majstorovic, Z., and Nieplova, E.:

- 588 HISTALP - Historical Instrumental climatological Surface Time series of the greater ALPine Region, *Int. J.*
589 *Climatol.*, 27, 17-46, doi: 10.1002/joc.1377, 2007.
- 590 Baroni, C., Bondesan, A., and Mortara, G.: Report of the Glaciological Survey of 2011 - Relazioni della
591 campagna glaciologica 2011, *Geogr. Fis. Din. Quat.*, 35(2), 211-279, doi:10.4461/GFDQ.2012.35.19, 2012.
- 592 Baroni, C., Bondesan, A., and Mortara, G.: Report of the Glaciological Survey of 2012 - Relazioni della
593 campagna glaciologica 2012, *Geogr. Fis. Din. Quat.*, 36(2), 303-374, doi: 10.4461/GFDQ.2013.36.24, 2013.
- 594 Baroni, C., Bondesan, A., and Mortara, G.: Report of the Glaciological Survey 2013. Relazioni della
595 Campagna Glaciologica 2013, *Geogr. Fis. Din. Quat.*, 37, 163-227, doi: 10.4461/GFDQ.2014.37.16, 2014.
- 596 Beniston, M., Keller, F., Koffi, B., and Goyette, S.: Estimates of snow accumulation and volume in the Swiss
597 Alps under changing climatic conditions, *Theor. Appl. Climatol.*, 76, 125–140, doi: 10.1007/s00704-003-
598 0016-5, 2003.
- 599 Benn, D. I. and Evans, D. J. A. (Eds.): *Glaciers and Glaciation*, Hodder Education, London, UK, 802 pp., 2010.
- 600 Berthier, E., Vincent, C., Magnússon, E., Gunnlaugsson, Á. Þ., Pitte, P., Le Meur, E., Masiokas, M., Ruiz, L.,
601 Pálsson, F., Belart, J. M. C., and Wagnon, P.: Glacier topography and elevation changes derived from
602 Pléiades sub-meter stereo images, *The Cryosphere*, 8, 2275-2291, doi:10.5194/tc-8-2275-2014, 2014.
- 603 Braithwaite, R. J., and Olesen, O. B.: Detection of climate signal by inter-stake correlations of annual
604 ablation data Qamanârssûp Sermia, West Greenland, *J. Glaciol.*, 35(120), 253-259, 1989.
- 605 Brunetti, M., Maugeri, M., Nanni, T., Auer, I., Böhm, R., and Schöner, W.: Precipitation variability and
606 changes in the Greater Alpine region over the 1800–2003 period, *J. Geophys. Res.*, 111, D11107, doi:
607 10.1029/2005/D06674, 2006.
- 608 Brunetti, M., Lentini, G., Maugeri, M., Nanni, T., Auer, I., Böhm, R., and Schöner, W.: Climate variability and
609 change in the Greater Alpine Region over the last two centuries based on multi-variable analysis, *Int. J.*
610 *Climatol.*, 29(15), 2197-2225, doi: 10.1002/joc.1857, 2009.
- 611 Brunetti, M., Lentini, G., Maugeri, M., Nanni, T., Simolo, C., and Spinoni, J.: Projecting North Eastern Italy
612 temperature and precipitation secular records onto a high resolution grid, *Phys. Chem. Earth*, 40-41, 9-22,
613 doi: 10.1016/j.pce.2009.12.005, 2012.
- 614 Brunetti, M., Maugeri, M., Nanni, T., Simolo, C., and Spinoni, J.: High-resolution temperature climatology
615 for Italy: interpolation method intercomparison. *Int. J. Climatol.*, 34, 1278-1296, doi: 10.1002/joc.3764,
616 2014.
- 617 Cannone, N., Diolaiuti, G., Guglielmin, M., and Smiraglia, C.: Accelerating climate change impacts on alpine
618 glacier forefield ecosystems in the European Alps, *Ecol. Appl.*, 18(3), 637-648, doi: 10.1890/07-1188.1, 2008.
- 619 Carturan, L., Baroni, C., Becker, M., Bellin, A., Cainelli, O., Carton, A., Casarotto, C., Dalla Fontana, G., Godio,
620 A., Martinelli, T., Salvatore, M. C., and Seppi, R.: Decay of a long-term monitored glacier: Careser Glacier
621 (Ortles-Cevedale, European Alps), *The Cryosphere*, 7, 1819-1838, doi:10.5194/tc-7-1819-2013, 2013a.
- 622 Carturan, L., Filippi, R., Seppi, R., Gabrielli, P., Notarnicola, C., Bertoldi, L., Paul, F., Rastner, P., Cazorzi, F.,
623 Dinale, R., and Dalla Fontana, G.: Area and volume loss of the glaciers in the Ortles-Cevedale group (Eastern

- 624 Italian Alps): controls and imbalance of the remaining glaciers, *The Cryosphere*, 7, 1339-1359,
625 doi:10.5194/tc-7-1339-2013, 2013b.
- 626 Carturan, L., Baldassi, G. A., Bondesan, A., Calligaro, S., Carton, A., Cazorzi, F., Dalla Fontana, G., Francese, R.,
627 Guarnieri, A., Milan, N., Moro, D., and Tarolli, P.: Current behavior and dynamics of the lowermost Italian
628 glacier (Montasio Occidentale, Julian Alps), *Geogr. Ann. A*, 95(1), 79-96, doi: 10.1111/geoa.12002, 2013c.
- 629 Carturan, L.: [Replacing monitored glaciers undergoing extinction: a new measurement series on La Mare](#)
630 [Glacier \(Ortles-Cevedale, Italy\)](#)~~New measurement series for the future monitoring of glacier mass balance~~
631 [in the European Alps: methods and results on La Mare Glacier \(Ortles-Cevedale, Italy\)](#), *J. Glaciol.*, in review,
632 2016.
- 633 Casartelli, G., Kappenberger, G., and Smiraglia, C.: Accumulo e ablazione sui ghiacciai delle alpi lombarde e
634 svizzere: Risultati di alcuni recenti bilanci di massa, *Rivista Geografica Italiana*, 103(1), 1-30, 1996.
- 635 Catasta, G., and Smiraglia, C.: The mass balance of a cirque glacier in the Italian Alps (Ghiacciaio della
636 Sforzellina, Ortles-Cevedale Group), *J. Glaciol.*, 39(131), 87-90, 1993.
- 637 CGI (Comitato Glaciologico Italiano): Reports of the glaciological surveys, *Bollettino del Comitato*
638 *Glaciologico Italiano*, Series I and II, 1–25, 1914–1977.
- 639 CGI (Comitato Glaciologico Italiano): Reports of the glaciological surveys, *Geogr. Fis. Din. Quat.*, 1–35, 1978–
640 2011.
- 641 Cogley, J. G.: Geodetic and direct mass-balance measurements: comparison and joint analysis. *Ann. Glaciol.*,
642 50(50), 96-100, 2009.
- 643 Cogley, J. G., and Adams, W. P.: Mass balance of glaciers other than the ice sheets, *J. Glaciol.*, 44(147), 315-
644 325, 1998.
- 645 Cogley, J. G., Hock, R., Rasmussen, L. A., Arendt, A. A., Bauder, A., Braithwaite, R. J., Jansson, P., Kaser, G.,
646 Moller, M., Nicholson, L., and Zemp, M.: Glossary of Glacier Mass Balance and Related Terms, IHP-VII
647 Technical Documents in Hydrology No. 86, IACS Contribution No. 2, UNESCO-IHP, Paris, 2011.
- 648 Colucci, R. R., and Guglielmin, M.: Precipitation–temperature changes and evolution of a small glacier in the
649 southeastern European Alps during the last 90 years, *Int. J. Climatol.*, 35(10), 2783–2797,
650 doi: 10.1002/joc.4172, 2015.
- 651 Debeer, C. M., and Sharp, M. J.: Topographic influences on recent changes of very small glaciers in the
652 Monashee Mountains, British Columbia, Canada, *J. Glaciol.*, 55(192), 691–700, doi:
653 10.3189/002214309789470851, 2009.
- 654 Durand, Y., Laternser, M., Giraud, G., Etchevers, P., Lesaffre, B., and Mérindol, L.: Reanalysis of 44 years of
655 climate in the French Alps (1958–2002): methodology, model validation, climatology, and trends for air
656 temperature and precipitation, *J. Appl. Meteorol. Climatol.*, 48(3), 429–449, 2009a.
- 657 Durand, Y., Giraud, G., Laternser, M., Etchevers, P., Mérindol, L., and Lesaffre, B.: Reanalysis of 47 years of
658 climate in the French Alps (1958–2005): climatology and trends for snow cover, *J. Appl. Meteorol. Climatol.*,
659 48(12), 2487–2512, 2009b.

- 660 Dyurgerov, M., Meier, M. F., and Bahr, D. B.: A new index of glacier area change: a tool for glacier
661 monitoring, *J. Glaciol.*, 55(192), 710-716, 2009.
- 662 Eckert, N., Baya, H., Thibert, E., and Vincent, C.: Extracting the temporal signal from a winter and summer
663 mass-balance series: application to a six-decade record at Glacier de Sarennes, French Alps, *J. Glaciol.*,
664 57(201), 134-150, 2011.
- 665 Elsberg, D. H., Harrison, W. D., Echelmeyer, K. A., and Krimmel, R. M.: Quantifying the effect of climate and
666 surface change on glacier mass balance, *J. Glaciol.*, 47, 649–658, 2001.
- 667 Fischer, M., Huss, M., and Hoelzle, M.: Surface elevation and mass changes of all Swiss glaciers 1980–2010,
668 *The Cryosphere*, 9, 525-540, doi:10.5194/tc-9-525-2015, 2015.
- 669 Furbish, D. J., and Andrews, J. T.: The use of hypsometry to indicate long-term stability and response of
670 valley glaciers to changes in mass transfer, *J. Glaciol.*, 30, 199–211, 1984.
- 671 Galos, S. P., Klug, C., Prinz, R., Rieg, L., Dinale, R., Sailer, R., and Kaser, G.: Recent glacier changes and
672 related contribution potential to river discharge in the Vinschgau / Val Venosta, Italian Alps, *Geogr. Fis. Din.*
673 *Quat.*, 38, 143-154, doi: 10.4461/GFDQ.2015.38.13, 2015.
- 674 Haeberli, W., Huggel, C., Paul, F., and Zemp, M.: Glacial responses to climate change, in: *Treatise on*
675 *Geomorphology*, Academic Press, 13, San Diego, California, 152-175, 2013.
- 676 Hoelzle, M., Haeberli, W., Dischl, M., and Peschke, W.: Secular glacier mass balances derived from
677 cumulative glacier length changes, *Global Planet. Change*, 36, 295–306, doi: 10.1016/S0921-
678 8181(02)00223-0, 2003.
- 679 Huss, M., Dhulst, L., and Bauder, A.: New long-term mass balance series for the Swiss Alps, *J. Glaciol.*,
680 61(227), 551-562, doi: 10.3189/2015JoG15j015, 2015.
- 681 Kaser, G., Fountain, A., and Jansson, P.: A manual for monitoring the mass balance of mountain glaciers,
682 IHP-VI Technical Documents in Hydrology No. 59, UNESCO-IHP, Paris, 2003.
- 683 Killick, R., Eckley, I. A.: Changepoint: An R package for changepoint analysis, *J. Stat. Softw.*, 58(3), 1-19. DOI:
684 10.18637/jss.v058.i03, 2014.
- 685 Kuhn, M.: The mass balance of very small glaciers, *Zeitschrift für Gletscherkunde und Glazialgeologie*, 31(1–
686 2), 171–179, 1995.
- 687 Lliboutry, L.: Multivariate statistical analysis of glacier annual balances, *J. Glaciol.*, 13, 371-392, 1974.
- 688 Marzeion, B. and Nesje, A.: Spatial patterns of North Atlantic Oscillation influence on mass balance
689 variability of European glaciers, *The Cryosphere*, 6, 661-673, doi:10.5194/tc-6-661-2012, 2012.
- 690 Mernild, S. H., Lipscomb, W. H., Bahr, D. B., Radić, V., and Zemp, M.: Global glacier changes: a revised
691 assessment of committed mass losses and sampling uncertainties, *The Cryosphere*, 7, 1565-1577,
692 doi:10.5194/tc-7-1565-2013, 2013.
- 693 Mitchell, T. D., and Jones, P. D.: An improved method of constructing a database of monthly climate
694 observations and associated high-resolution grids, *Int. J. Climatol.* 25, 693–712, doi: 10.1002/joc.1181, 2005.

- 695 New, M., Hulme, M., and Jones, P. D.: Representing twentieth century space-time variability. Part 2:
696 development of 1901–96 monthly grids of surface climate, *J. Climate*, 13, 2217–2238, 2000.
- 697 Oerlemans, J., and Reichert, B. K.: Relating glacier mass balance to meteorological data by using a seasonal
698 sensitivity characteristic, *J. Glaciol.*, 46(152), 1–6, doi: 10.3189/172756500781833269, 2000.
- 699 Østrem, G., and Brugman, M.: Mass balance measurement techniques. A manual for field and office work,
700 National Hydrology Research Institute (NHRI), Science Report, (4), Saskatoon, Canada, 224 pp., 1991.
- 701 Paul, F., and Haeberli, W.: Spatial variability of glacier elevation changes in the Swiss Alps obtained from
702 two digital elevation models, *Geophys. Res. Lett.*, 35, L21502, doi:10.1029/2008GL034718, 2008.
- 703 Pelto, M. S.: Glacier annual balance measurement, forecasting and climate correlations, North Cascades,
704 Washington 1984–2006, *The Cryosphere*, 2, 13-21, doi:10.5194/tc-2-13-2008, 2008
- 705 Quadrelli, R., Lazzeri, M., Cacciamani, C., and Tibaldi, S.: Observed winter Alpine precipitation variability and
706 links with large-scale circulation patterns, *Clim. Res.*, 17(3), 275-284, doi:10.3354/cr017275, 2001.
- 707 Reichert, B. K., Bengtsson, L., and Oerlemans, J.: Midlatitude forcing mechanisms for glacier 25 mass
708 balance investigated using general circulation models, *J. Climate*, 14, 3767–3784, 2001.
- 709 Salvatore, M. C., Zanoner, T., Baroni, C., Carton, A., Banchieri, F. A., Viani, C., Giardino, M., and Perotti, L.:
710 The state of Italian glaciers: a snapshot of the 2006-2007 hydrological period, *Geogr. Fis. Dinam. Quat.*, 38,
711 175-198, 2015.
- 712 Schmidli, J., Schmutz, C., Frei, C., Wanner, H., and Schär, C.: Mesoscale precipitation variability in the region
713 of the European Alps during the 20th century, *Int. J. Climatol.*, 22(9), 1049-1074, doi: 10.1002/joc.769, 2002.
- 714 Schöner, W., Auer, I., and Böhm, R.: Climate variability and glacier reaction in the Austrian eastern Alps,
715 *Ann. Glaciol.*, 31(1), 31-38, 2000.
- 716 Schwarb, M.: The alpine precipitation climate, Ph.D. thesis, ETH Zürich, Switzerland, 131 pp., 2000.
- 717 Scotti, R., Brardinoni, F., and Crosta, G. B.: Post-LIA glacier changes along a latitudinal transect in the
718 Central Italian Alps, *The Cryosphere*, 8, 2235-2252, doi:10.5194/tc-8-2235-2014, 2014.
- 719 Six, D., Reynaud, L., and Letréguilly, A.: Alpine and Scandinavian glaciers mass balances, their relations with
720 the North Atlantic Oscillation, *Comptes Rendus de l'Académie des Sciences, Series IIA, Earth Planet. Sc.*, 333,
721 693–698, 2001.
- 722 Thibert, E., Eckert, N., and Vincent, C.: Climatic drivers of seasonal glacier mass balances: an analysis of 6
723 decades at Glacier de Sarennes (French Alps), *The Cryosphere*, 7(1), 47-66, 2013.
- 724 Vincent, C., Kappenberger, G., Valla, F., Bauder, A., Funk, M., and Le Meur, E.: Ice ablation as evidence of
725 climate change in the Alps over the 20th century, *J. Geophys. Res.*, 109(D10), D10104, doi:
726 10.1029/2003JD003857, 2004.
- 727 WGMS: Global glacier changes: facts and figures, edited by: Zemp, M., Roer, I., Kaab, A., Hoelzle, M., Paul,
728 F., and Haerberli, W., UNEP, World Glacier Monitoring Service, Zürich, Switzerland, 2008.

729 WGMS: Fluctuations of Glaciers 2005-2010, Volume X, edited by: Zemp, M., Frey, H., Gärtner-Roer, I.,
730 Nussbaumer, S. U., Hoelzle, M., Paul, F., and W. Haeberli: ICSU(WDS)/IUGG(IACS)/UNEP/UNESCO/WMO,
731 World Glacier Monitoring Service, Zurich, Switzerland, 336 pp., publication based on database version:
732 doi:10.5904/wgms-fog-2012-11, 2012.

733 WGMS: Glacier Mass Balance Bulletin No. 12 (2010-2011), edited by: Zemp, M., Nussbaumer, S. U., Naegeli,
734 K., Gärtner-Roer, I., Paul, F., Hoelzle, M., and Haeberli, W.: ICSU(WDS)/IUGG(IACS)/UNEP/UNESCO/WMO,
735 World Glacier Monitoring Service, Zurich, Switzerland, 106 pp., publication based on database version: doi:
736 10.5904/wgms-fog-2013-11, 2013.

737 WGMS: Fluctuations of Glaciers Database. World Glacier Monitoring Service, Zurich, Switzerland.
738 DOI:10.5904/wgms-fog-2015-11, 2015. Online access: <http://dx.doi.org/10.5904/wgms-fog-2015-11>

739 Zemp, M., Frauenfelder, R., Haeberli, W., and Hoelzle, M.: Worldwide glacier mass balance measurements:
740 general trends and first results of the extraordinary year 2003 in Central Europe, Data of Glaciological
741 Studies [Materialy glyatsiologicheskikh issledovaniy], 99, Moscow, Russia, 3–12, 2005.

742 Zemp, M., Paul, F., Hoelzle, M., and Haeberli, W.: Glacier fluctuations in the European Alps 1850-2000: an
743 overview and spatio-temporal analysis of available data, in The darkening peaks: Glacial retreat in scientific
744 and social context, edited by: Orlove, B., Wiegandt, E., and Luckman, B. H., University of California Press,
745 Berkeley, 152-167, 2008.

746 Zemp, M., Hoelzle, M., and Haeberli, W.: Six decades of glacier mass-balance observations: a review of the
747 worldwide monitoring network, *Ann. Glaciol.*, 50(50), 101-111, doi: 10.3189/172756409787769591, 2009.

748 [Zemp, M., Thibert, E., Huss, M., Stumm, D., Rolstad Denby, C., Nuth, C., Nussbaumer, S. U., Moholdt, G.,](#)
749 [Mercer, A., Mayer, C., Joerg, P. C., Jansson, P., Hynek, B., Fischer, A., Escher-Vetter, H., Elvehøy, H., and](#)
750 [Andreassen, L. M.: Reanalysing glacier mass balance measurement series. *The Cryosphere*, 7, 1227-1245,](#)
751 [doi:10.5194/tc-7-1227-2013, 2013.](#)

752

753

754

755

Tables

756 Table 1 – Physical characteristics of the Italian glaciers with the mass balance series analysed in this study
 757 (year 2006, NextData - DATAGRALP project: [http://www.nextdatapoint.it/?q=en/content/special-project-](http://www.nextdatapoint.it/?q=en/content/special-project-datagrulp)
 758 [datagrulp](http://www.nextdatapoint.it/?q=en/content/special-project-datagrulp), last access: 27 September 2015; Salvatore et al., 2015).

Glacier	Geographic area	Area (km ²)	Minimum elevation (m a.s.l.)	Maximum elevation (m a.s.l.)	Median elevation (m a.s.l.)	Prevailing aspect	Average slope (°)	First survey year
Grand Etrèt	Gran Paradiso	0.47	2667	3190	2894	N	23	2002
Ciardoney	Gran Paradiso	0.59	2855	3170	3039	E-NE	18	1992
Fontana Bianca	Ortles-Cevedale	0.48	2889	3342	3166	E	23	1984
Sforzellina	Ortles-Cevedale	0.29	2790	3046	2868	NW	16	1987
Lunga	Ortles-Cevedale	1.86	2678	3378	3128	NE	19	2004
Careser	Ortles-Cevedale	2.39	2868	3279	3069	S	11	1967
La Mare (southern branch)	Ortles-Cevedale	2.16	2652	3518	3215	NE	21	2003
Pendente	Val Ridanna	0.95	2621	3064	2781	S	15	1996
Malavalle	Val Ridanna	6.92	2512	3441	2971	SE	14	2002

759

760

761

762 Table 2 – Mean values (and STD in brackets) of B_w , B_s , B_a and AAR for nine Italian glaciers in the period from
 763 2004 to 2013 (Car = Careser, FB = Fontana Bianca, Pen = Pendente, Cia = Ciardoney, Sfo = Sforzellina, GE =
 764 Grand Etrèt, Lun = Lunga, Mar = La Mare, Mal = Malavalle). Values expressed in mm w.e. except AAR that is
 765 in percent.

	Car	Mar	FB	Sfo	Lun	Pen	Mal	Cia	GE
B_w (9 years)	927 (330)	989 (301)	1085 (338)	\	991 (222)	1537 (425)	1194 (256)	1052 (421)	1472 (578)
B_s (9 years)	-2740 (368)	-1758 (303)	-2183 (457)	\	-2151 (368)	-2857 (525)	-2087 (386)	-2510 (378)	-2396 (321)
B_a (10 years)	-1788 (590)	-763 (395)	-1088 (642)	-1399 (505)	-1195 (466)	-1231 (692)	-825 (484)	-1419 (646)	-946 (648)
AAR (10 years)	1 (3)	25 (14)	11 (22)	\	12 (16)	4 (8)	23 (17)	3 (5)	\

766

767

768

769 Table 3 – Correlation matrix of B_a for nine Italian glaciers. * and ** indicate Spearman correlation significant
 770 at the 0.05 and 0.01 level, respectively.

	Car	FB	Pen	Cia	Sfo	GE	Lun	Mar	Mal
Car	1.00								
FB	0.82**	1.00							
Pen	0.85**	0.76**	1.00						
Cia	0.87**	0.86**	0.55*	1.00					
Sfo	0.82**	0.75**	0.65**	0.81**	1.00				
GE	0.74**	0.77**	0.66*	0.62*	0.69*	1.00			
Lun	0.70*	0.73*	0.60	0.48	0.49	0.77**	1.00		
Mar	0.90**	0.96**	0.70*	0.90**	0.82**	0.80**	0.71*	1.00	
Mal	0.87**	0.84**	0.97**	0.61*	0.65*	0.68*	0.61	0.80**	1.00

771

772

773

774 Table 4 – Correlation coefficients of B_a vs. B_w and B_s . * and ** indicate Spearman correlation significant at
775 the 0.05 and 0.01 level, respectively.

	No of years	B_w	B_s
Car	40	0.46**	0.94**
FB	22	0.24	0.84**
Pen	12	0.84**	0.67*
Cia	22	0.51*	0.76**
GE	12	0.84**	0.66*
Lun	10	0.64*	0.69*
Mar	10	0.66*	0.64*
Mal	9	0.48	0.85**

776

777

778

779 Table 5 – Correlation coefficients of B_w , B_s and B_a vs. seasonal and annual NAO. Five-year triangular moving
 780 averages have been applied to the time series before correlation analyses. *, ** and *** indicate Spearman
 781 correlation significant at the 0.10, 0.05 and 0.01 level, respectively.

782

Nation	Glacier	Winter balance			Summer balance			Annual balance		
		DJF NAO	Oct-May NAO	Annual NAO	DJF NAO	Oct-May NAO	Annual NAO	DJF NAO	Oct-May NAO	Annual NAO
I	Car	-0.51**	-0.34	-0.16	-0.23	-0.05	0.34	-0.30*	-0.13	0.23
I	FB	-0.11	0.18	0.00	0.22	0.10	0.25	0.07	0.35	0.41*
I	Pen	-0.70	-0.90*	-0.70	-1.00**	-0.90*	-1.00**	0.12	0.15	0.23
I	Cia	0.17	0.34	0.19	-0.03	0.10	0.27	0.13	0.27	0.36
I	GE	-0.81***	-0.74**	-0.79**	-0.88***	-0.76**	-0.91***	-0.86***	-0.76**	-0.83***
I	Lun	-0.94**	-0.94**	-0.94**	-0.94**	-0.77	-0.94**	-0.89**	-0.83*	-0.89**
I	Mar	-1.00***	-0.89**	-1.00***	0.14	0.03	0.14	-0.79**	-0.82**	-0.89***
I	Mal	-0.90*	-0.70	-0.90*	-0.50	-0.80	-0.50	-0.81***	-0.74**	-0.45
I	Sfo							0.59***	0.59***	0.61***
F	St. Sorlin							-0.44***	-0.38***	-0.04
F	Sarennes	0.36***	0.43***	0.49***	-0.53***	-0.50***	-0.19	-0.41***	-0.37***	-0.05
CH	Silvretta	0.28*	0.13	-0.05	-0.57***	-0.45***	-0.18	-0.38***	-0.28**	0.02
CH	Gries	-0.39***	-0.22	-0.18	-0.58***	-0.53***	-0.25*	-0.49***	-0.39***	-0.03
A	Sonnblick							-0.48***	-0.45***	-0.07
A	Vernagt	0.27*	0.51***	0.49***	-0.42***	-0.38**	-0.09	-0.36**	-0.26*	0.07
A	Kesselwand							-0.42***	-0.38***	-0.07
A	Hintereis							-0.55***	-0.44***	-0.10

783

784

785

786 Table 6 - Spearman correlation coefficients and multiple regression results of B_a vs. seasonal mean
787 temperature and precipitation. *, ** and *** indicate 0.05, 0.01 and 0.001 significance levels.

Air temperature – Correlation coefficients									
	Car	FB	Sfo	Cia	Pen	GE	Mal	Mar	Lun
No of years	47	27	27	22	18	12	12	11	10
Jun-Sep	-0.77***	-0.49**	-0.52**	-0.64**	-0.40	0.01	-0.24	-0.16	0.49
Oct-May	-0.37**	-0.42*	-0.10	-0.18	-0.49*	-0.18	-0.57	-0.28	-0.33
Year	-0.68***	-0.49**	-0.30	-0.49*	-0.69**	-0.31	-0.71**	-0.27	-0.14

Precipitation – Correlation coefficients									
	Car	FB	Sfo	Cia	Pen	GE	Mal	Mar	Lun
No of years	47	27	27	22	18	12	12	11	10
Jun-Sep	-0.15	-0.02	0.09	0.20	0.00	0.02	-0.04	0.26	-0.05
Oct-May	0.28	0.40*	0.32	0.47*	0.37	0.57	0.43	0.67*	0.71*
Year	0.11	0.36	0.39*	0.53*	0.34	0.64*	0.34	0.83**	0.53

Multiple linear regression - Coefficients									
	Car	FB	Sfo	Cia	Pen	GE	Mal	Mar	Lun
No of years	47	27	27	22	18	12	12	11	10
Jun-Sep temperature	-776.453 (***)	-663.487 (***)	-575.225 (***)	-796.739 (***)	-496.521 (**)	-63.899	-355.106	-668.941 (**)	115.265
Oct-May precipitation	2.186 (***)	3.342 (***)	2.915 (***)	3.315 (***)	2.380 (*)	2.897 (**)	3.051 (*)	4.122 (**)	2.666 (*)
Intercept	-3265.013 (***)	-3311.632 (***)	-3176.797 (***)	1753.826	1011.719	-2707.559	19.212	- 3380.90 5 (***)	- 2619.872 (**)
% of explained variance	75.6	68.7	72.1	73.7	51.5	56.8	56.0	78.4	64.5


788

789

790

Figures

791





792

793 Figure 1. Geographic setting of the glaciers with mass balance measurements analysed in this work
794 (Microsoft® BingTM Maps).




795

Val Ridanna






- Mass balance measurement points
- Contours (height interval = 50 m)

Ortles-Cevedale







Gran Paradiso



796


797

Figure 2. Surface topography and measurement network of the nine glaciers analysed in this study.



799 Figure 3. Time series of B_w , B_s , B_a and AAR for the nine Italian glaciers analysed.

800




801

802 Figure 4. Cumulative mass balance for the nine Italian glaciers and for a set of nine other glaciers
803 representative of the European Alps. Dotted and dashed lines indicate the average B_a for the two groups of
804 glaciers in the periods from 1967 to 1983, 1984 to 1993, 1994 to 2003 and 2004 to 2013.

805


806



807

808 Figure 5. Left column: mean ablation season (Jun-Sep) air temperature anomalies in (a) Gran Paradiso, (b)
 809 Ortles-Cevedale, and (c) Val Ridanna. Right column: mean accumulation season (Oct-May) air temperature
 810 anomalies in (d) Gran Paradiso, (e) Ortles-Cevedale, and (f) Val Ridanna. Reference period: 1961-1990. All
 811 linear trends are significant at the 0.001 level.

812









813

814 Figure 6. Days per year with maximum air temperature exceeding 0°C at 3000 m a.s.l., calculated from the
 815 series of the Careser diga weather station (2600 m a.s.l., Ortles-Cevedale Group).

816

817

818




819

820 | Figure 67. Left column: ablation season (Jun-Sep) total precipitation anomalies in (a) Gran Paradiso, (b)
 821 | Ortles-Cevedale, and (c) Val Ridanna. Right column: accumulation season (Oct-May) total precipitation
 822 | anomalies in (d) Gran Paradiso, (e) Ortles-Cevedale, and (f) Val Ridanna. Reference period: 1961-1990. None
 823 | of the linear trends is significant at the 0.05 level.

824

825

826



827

828






829

830

831

Figure 8. Winter NAO index from 1961 to 2013

(<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/pna/new.ao.shtml>, last access: 11 February 2016).




832

833 | Figure 79. Bootstrapped moving correlation coefficient between annual mass balance and seasonal values
 834 of air temperature and precipitation. Shaded straight lines indicate significance at 95% level.

835

836



837

838 | Figure 810. Plot of residuals of the multiple linear regression of B_a vs. Oct-May precipitation and Jun-Sep
839 temperature on the Careser Glacier. Multiple regression coefficients are reported in Table 6. The inset
840 shows the extent of the glacier in three different years.

841

842

843

844

845

846

847

848

849

850

851

852

853