

Growth of a young pingo in the Canadian Arctic observed by RADARSAT-2 interferometric satellite radar

Sergey V. Samsonov¹, Trevor C. Lantz², Steven V. Kokelj³, and Yu Zhang¹

¹Canada Centre for Mapping and Earth Observation, Natural Resources Canada, 560 Rochester Street, Ottawa, ON K1A0E4, Canada.

²School of Environmental Studies, University of Victoria, Victoria, BC, Canada.

³Northwest Territories Geoscience Office, Government of the Northwest Territories, Yellowknife, NWT, Canada.

Correspondence to: Sergey V. Samsonov (sergey.samsonov@canada.ca)

Abstract. Advancements in radar technology are increasing our ability to detect earth surface deformation in permafrost environments. In this paper we use satellite Differential Interferometric Synthetic Aperture Radar (DInSAR) to describe the growth of a previously unreported pingo in the Tuktoyaktuk Coastlands. High-resolution RADARSAT-2 imagery (2011-2014) analyzed with the
5 Multidimensional Small Baseline Subset (MSBAS) DInSAR revealed a maximum 2.7 cm/year of domed uplift located in a drained lake basin. Satellite measurements suggest that this feature is one of the largest diameter pingos in the region that is presently growing. Observed changes in elevation were modeled as a 348×290 m uniformly loaded elliptical plate with clamped edge. Analysis of historical aerial photographs showed that ground uplift at this location initiated sometime be-
10 tween 1935 and 1951 following drainage of the residual pond. Uplift is largely due to the growth of intrusive ice, because the 9% expansion of pore water associated with permafrost aggradation into saturated sands is not sufficient to explain the observed short- and long-term deformation rates. The modeled thickness of permafrost using the Northern Ecosystem Soil Temperature (NEST) was consistent with the maximum height of this feature and the 1972-2014 elevation changes estimated
15 from aerial photographs, suggesting that permafrost aggradation is resulting in the freezing a sub-pingo water lens. Seasonal variations in the uplift rate seen in the DInSAR data closely match the modeled seasonal pattern in the deepening rate of freezing front. This study demonstrates that interferometric satellite radar can successfully contribute to understanding the dynamics of terrain uplift in response to permafrost aggradation and ground ice development in remote polar environments.
20 However, the present-day growth rate is smaller than predicted by the modeling and no clear growth is observed at other smaller pingos in contrast with field studies performed mainly before the 1990s. Further investigation of this apparent discrepancy complemented by field investigations may help define limitations of our processing methodology and DInSAR data.

1 Introduction

25 Pingos are conical ice-cored hills that grow in the permafrost environment (Mackay, 1998 and references within). There are over 11000 pingos in the world. Of these about 1350 are concentrated on the Tuktoyaktuk Coastal Plain in the western Canadian Arctic, 500 are located elsewhere in Canada, 1500 in Alaska, 6000 or more in Russia, and at least several hundred in the other parts of the world, such as Fennoscandia, Greenland, Spitsbergen, Mongolia, and Tibetan Plateau of China (Mackay, 30 1998; Wu et al., 2005; Grosse and Jones, 2011; Yoshikawa et al., 2013). Numerous near circular collapsed features interpreted as pingo remnants, most being of Holocene age, have been observed in former permafrost areas, such as Ireland, the United Kingdom, countries of Eastern Europe and China (Jin et al., 2007). Pingo-like features have also been observed on Mars using the High Resolution Imaging Science Experiment (HiRISE) camera on board the Mars Reconnaissance Orbiter 35 (Dundas et al., 2008; Burr et al., 2009; Soare et al., 2014). Understanding the genesis, growth and collapse of pingos provides insight into the landscape forming processes associated with permafrost aggradation and ground ice development on Earth, Mars and possibly other planets. Seminal work by Dr J.R. Mackay on genesis and growth of pingos has provided tremendous insight into the landscape forming processes associated with permafrost aggradation and ground ice development (Mackay, 40 1977, 1979, 1987, 1990, 1992, 1997, 1998). This research has provided context for our work, which explores the potential of SAR technology and processing techniques to describe surface deformation that would arise from processes associated with pingo growth.

The great majority (i.e. $\sim 98\%$) of hydrostatic or closed system pingos develop in drained lake basins in association with permafrost aggradation in unfrozen saturated sandy lake sediments (Gurney, 1998; Mackay, 1998; Jones et al., 2011). Rapid lake drainage, caused either by coastal erosion or thermal erosion of ice wedges, exposes the lake bottom to subfreezing air temperatures (Fig 1, Mackay, 1992). This initiates the process of permafrost aggradation into the unfrozen lake-bottom sediments. Downward aggradation of permafrost progresses at a rate inversely proportional to the square root of time (Freitag and McFadden, 1997). The water-ice phase transition produces 9% volume expansion, i.e. $V_i = \frac{\rho_w}{\rho_i} V_w = \frac{999.97}{916.70} V_w \approx 1.09 V_w$, where ρ_w , ρ_i , V_w , and V_i are water and ice density and volume respectively. During downward freezing of saturated sands, given free drainage, there will be little or no heave because of pore water expulsion and flow of water in the direction of lower pressure (Mackay, 1997). In a drained lake, expelled pore water typically flows to a residual pond, which is the usual site of pingo growth (Mackay, 1998). A closed system is formed when 55 the residual pond freezes to the bottom. In this case, increasing pressure derived from pore water expelled during permafrost aggradation into the surrounding lake bottom can heave the relatively small area of thin permafrost beneath the residual pond to initiate pingo growth. If the addition of water to this location exceeds the rate of downward freezing, then a pressurized, sub-pingo water lens forms and heaves the pingo upward. Freezing of the pressurized water forms intrusive ice. If the

60 water pressure drops due to escape of water through a hydraulic fracture or a change in the ground
water supply, segregated ice may form (Mackay, 1998).

Thick permafrost, sandy sediments, abundant lakes underlain by talik (i.e. unfrozen zones) and
relatively frequent lake drainage contribute to the high concentration of pingos in the Tuktoyaktuk
Coastlands (Mackay, 1998). The maximum thickness of lake ice in this region is about 2 m, so
65 lakes with greater depths are underlain by a talik (unfrozen sediments) (Mackay, 1998). The mean
basal radius of most pingos is about 50 m while the maximum radius is typically less than 300 m
(Mackay, 1979). Their heights vary between a few meters and approximately 55 m, with a mean
of 5 m (Grosse and Jones, 2011; Jones et al., 2011). Pingo basal area is typically established in the
first few years of growth, determined primarily by characteristics of the residual pond where growth
70 has occurred. Subsequently a pingo grows higher with a minimal increase in its basal area (Mackay,
1998). The long-term monitoring of numerous pingos by Mackay (1977); ? has shown that growth
rates can vary significantly through time in response to changes in: 1) rates of ground water flow to
the sub-pingo water lens, 2) release of pressurized water along hydraulic fractures and 3) variation
in rates of downward freezing (Mackay, 1977, 1998). Pingo growth may continue as long as ground
75 water continues to be supplied to the area beneath a pingo. It may take more than a thousand years
for a talik to refreeze, so a pingo in the Tukotyaktuk coastlands may continue to grow for hundreds
to thousands of years.

Previous studies of pingo growth and subsidence have been primarily based on annual or semi-
annual field measurements of heights at benchmark locations and have required monumental field
80 effort (Mackay, 1977, 1998). The subtle temporal changes of discrete topographic features such as
pingos could not be easily detected by the early coarse resolution Synthetic Aperture Radars, such
as ERS-1/2, ENVISAT or ALOS, that were capable of mapping large-scale ground deformation in
permafrost areas (Liu et al., 2010; Chen et al., 2012, 2013). However, availability of high resolution
SAR data from the newer RADARSAT-2 satellite and advanced processing methodology allowed
85 us to identify a very large-diameter pingo that has not been previously described in the Tuktoyaktuk
Coastlands and for the first time, monitor pingo growth rate with high temporal and spatial resolution
over a period of four years.

2 Data and processing methodology

Historical aerial photographs, high resolution Differential Interferometric Synthetic Aperture Radar
90 (DInSAR) data, and field measurements were used to characterize the observed feature, located in
the Tuktoyaktuk Coastlands, Northwest Territories, Canada.

2.1 Historical aerial photographs

To constrain the timing of the lake drainage event and to estimate long-term changes in elevation at this site, the following historical aerial photographs from the Natural Resources Canada Air Photo
95 Library were analyzed: 1935 (A5023: 88R), 1951 (A12918: 092-093), 1972 (A22961: 029-030) and
2004 (A31864: 072-073). Summit Evolution Photogrammetric Workstation was used to create soft-
copy stereophoto models from 1972 and 2004 aerial photographs. The scale of these images allowed
us to resolve differences in height of ~ 0.5 m. Absolute orientation was completed using a 2004 or-
thophoto from the Mackenzie Valley Airphoto Project supported by the Northwest Territories Centre
100 for Geomatics, Government of the Northwest Territories, and a Digital Elevation Model (DEM) pro-
vided by the National Topographic Service, Government of Canada. To estimate the elevation at 300
(1972) and 118 (2004) locations inside the former lake basin we used Summit Evolution's stereoplot-
ter feature and DTM Collection tool to interface with ArcGIS and record coordinates and elevation
values. This point data was interpolated to a continuous DEM for each time period using the Natural
105 Neighbor tool in ArcGIS.

2.2 Differential Interferometric Synthetic Aperture Radar

To map the growth of this feature sixteen ascending and twenty descending Ultra-Fine 3 (U3) images
with a spatial resolution of 1.6×2.8 m and coverage of 20×20 km (Table 1, Fig 3) were acquired
by the RADARSAT-2 satellite between June 2011 and September 2014 over the Tuktoyaktuk Coast-
110 lands. DInSAR processing was performed with GAMMA software (Wegmuller and Werner, 1997).
The 3×3 spatial averaging (i.e. multilooking) in range and azimuth directions. A single master for
each set was selected and remaining images were re-sampled into the master geometry. The to-
pographic phase was removed using the 20 m resolution DEM (GeoGratis Web Services, Natural
Resources Canada) with an approximate reported elevation accuracy of 5 m. Differential interfero-
115 grams were filtered using the adaptive filtering with filtering function based on local fringe spectrum
(Goldstein and Werner, 1998) and unwrapped using the minimum cost flow algorithm (Costantini,
1998). The residual orbital ramp was observed in many interferograms and baseline refinement was
performed to correct it. Minor interpolation of each interferogram was performed in order to increase
the coverage reduced by decorrelation. This process is described in greater detail in Samsonov et al.
120 (2011a). From forty ascending and sixty four descending highly coherent interferograms the ascend-
ing and descending line-of-sight linear deformation rates were computed with the Small Baseline
Subset (SBAS) algorithm (Samsonov et al., 2011b). Then, ascending and descending interferograms
were resampled to a common lat/long grid with GMT scripts (<http://gmt.soest.hawaii.edu/>) of a re-
duced spatial extent that covers only the study site with the uniform spatial sampling of 10 m, and

125 analyzed with the Multidimensional Small Baseline Subset (MSBAS) algorithm (Samsonov and
d'Oreye, 2012; Samsonov et al., 2014):

$$\begin{pmatrix} \hat{A} \\ \lambda L \end{pmatrix} \begin{pmatrix} V_E \\ V_U \end{pmatrix} = \begin{pmatrix} \hat{\Phi} \\ 0 \end{pmatrix} \quad (1)$$

where matrix \hat{A} consist of time intervals between consecutive SAR acquisitions, V_E and V_U represents
unknown horizontal east-west and vertical velocities that are to be determined, $\hat{\Phi}$ represents
130 observed DInSAR data, λ is a regularization parameter and L is a first-difference operator. Thus, MS-
BAS software computed vertical and horizontal deformation rates between consecutive acquisitions
for each pixel by simultaneously processing ascending and descending DInSAR data. Deformation
time series were then reconstructed from the computed deformation rates and the mean rate was
estimated by fitting a straight line to the time series. Only vertical deformation time series and the
135 mean vertical deformation rate were used in further analysis.

2.3 Field measurements

During a field visit in August, 2014 this site was surveyed using a Nikon Nivo 5.M Total Station
whose position was determined using resection from two benchmarks. The position of one of these
benchmarks was determined using a Trimble R4 GNSS system. Post processing the data from this
140 static survey using Natural Resources Canada's Precise Point Positioning yielded a benchmark with
an orthometric height of 4.388 ± 0.001 m. This point was used to correct the elevations obtained from
the Total Station survey.

3 Results: Geodetic and field observations

The DEM, RADARSAT-2 satellite data coverage, locations of known pingos from the Canada Base
145 Map (GeoGratis Web Services, Natural Resources Canada) and suspected pingos visually identified
in high resolution optical imagery are shown in Fig 2.

The observed elevated feature is located in the middle of a large drained lake basin (Fig 4). The
earliest available aerial photograph shows that in 1935 the study site was covered by a small pond
(Fig 4). By 1951, the pond was no longer visible and an area comparable with present conditions had
150 been uplifted. This photographic evidence suggests that the growth of this feature began sometime
during 1935-1951, however the lake had drained at an unknown prior time. The DEMs computed
from 1972, 2004, and 2014 data (Fig 5) show that elevation of this feature gradually increased from
about 8 m in 1972 to 12 m in 2014.

The ascending and descending line-of-sight linear deformation rates (Fig 6) were calculated with
155 the 2σ measurement precision of 1.1 and 0.7 cm/year respectively. In spite of the presence of many
known pingos in the area (black and brown circles in Figs 2 and 6), deformation pattern resembling

growth of pingo was only observed at the study site outlined in red. This signal is particularly clear in the descending map, which was computed from a large set of sixty four interferograms. Precision of the ascending map is lower due to the smaller number of available images, but it still fully confirms deformation signals observed in the descending map. We could not detect uplift at other known pinguos because of their size comparable to the spatial sampling of deformation products (10 m) and low resolution DEM used to remove the topographic phase.

Vertical deformation time series from 2011 to 2014 calculated with MSBAS-DInSAR for the study site show maximum cumulative uplift of 9 cm over the 3.2 year time span (Fig 7). Over this period, the maximum annual linear rate of 2.7 cm/year was observed in the center of the feature (Fig 8) and a mean uplift rate of the entire feature was 1.5 cm/year. In order to better capture seasonal variability in the uplift rate linear and harmonic functions were simultaneously fitted to the cumulative deformation time series:

$$d(t) = A \sin\left(\frac{2\pi}{T}t + \psi\right) + Bt + C \quad (2)$$

where $d(t)$ is the cumulative deformation (uplift) measured in cm as a function of time t measured in years, T is the period equal to 1 year, A is the magnitude of the seasonal component equal to 0.4 cm, ψ is the phase at $t=0$ equal to 2.6 radian, B is the linear rate equal to 2.6 cm and C is the intercept equal to -5190.6 cm. The uplift rate was estimated by differentiation of the fitted function. The modeled seasonal uplift rate reaches maximum values in July and minimum values in February. Based on these observations this near-circular feature, approximately 500 m in diameter, was interpreted as a young pingo deforming the ground surface as a result of heave generated by the maintenance of a pressurized sub-pingo water lens and growth of intrusive and/or segregated ground ice.

4 Deformation rate modeling

In order to estimate shape and location of pingo and annual pressure change inside the sub-pingo water lens the vertical deformation rate was modeled by representing the pingo-like feature as an elastic uniformly loaded elliptical plate with a clamped edge (Timoshenko and Woinowsky-Krieger, 1959):

$$\Delta w = \Delta w_0 \left(1 - \frac{X^2}{a^2} - \frac{Y^2}{b^2}\right)^2$$

$$\Delta w_0 = \frac{\Delta q}{D} \frac{1}{\left(\frac{24}{a^4} + \frac{24}{b^4} + \frac{16}{a^2b^2}\right)}$$

$$D = \frac{Eh^3}{12(1-\nu^2)} \quad (3)$$

$$X = x \cos \alpha + y \sin \alpha$$

$$Y = -x \sin \alpha + y \cos \alpha$$

where Δw is the observed annual deformation rate, x, y and X, Y are the spatial coordinates in geodetic and rotated coordinate systems, a and b are the semi-major and semi-minor axis, α is the tilt angle, Δq is the annual rate of change in the effective driving upward pressure of sub-permafrost pore water, E is the Young's module, h is the total thickness of the overburden and pingo ice, and ν is the Poisson's ratio. Suitability of an elastic uniformly loaded circular plate model for describing the surface deformation associated with pingo growth and schematics was discussed in detail by Mackay (1987), and its schematic diagram can be found in Fig 3. Optimization based on simplex algorithm (Nelder and Mead, 1965) allowed estimation of the model parameters and their precision.

The best-fit model shown in Fig 8 corresponds to an elliptical plate centered at $69.3728^\circ N, 133.1688^\circ W$ with semi-major and semi-minor axes of 348 ± 2 m and 290 ± 2 m, tilted 27 ± 3 degrees clockwise, with magnitude of $\frac{\Delta q}{D} = 1.46 \pm 0.02 \times 10^{-10} \text{ m}^{-2}$. Assuming the total thickness of the overburden and pingo ice h ranges from 11-22 m (from NEST modeling, see below), Young's modulus E ranges 0.8-3.5 GPa (Schmeltz et al., 2002; Yang et al., 2015) and Poisson's ratio ν equals 0.4, the pressure increase Δq ranges 15-540 Pa/year, which is sufficient to uplift the ice column by 0.2-6.0 cm/year. The observed average uplift rate of 1.5 cm/year is within this range. The quantity $\frac{\Delta q}{D}$ was measured with the precision of $\sim 1\%$.

205 5 Permafrost aggradation modeling

The growth of a pingo is due to downward freezing of a pressurized sub-pingo water lens (Mackay, 1979). To approximate long-term change in the thickness of ice and the growth rates of pingos, Mackay (e.g. 1979, 1998) applied a simplified Stefan solution (Ingersoll et al., 1954) with a one-year time step. In this study, we used the Northern Ecosystem Soil Temperature (NEST) model (Zhang et al., 2003) to investigate the long-term patterns and rates of permafrost aggradation, total permafrost thickness, and the associated ground uplift. NEST is a one-dimensional process-based transient permafrost model that considers the effects of climate, vegetation, snow, and soil conditions on ground thermal dynamics based on energy and water transfer through the soil-vegetation-atmosphere system (Zhang et al., 2003). Ground temperature is calculated by solving the one-dimensional heat conduction equation. The upper boundary condition (the ground surface or snow surface when snow is present) is determined based on the surface energy balance, and the lower boundary condition (at a depth of 120 m) is defined based on the geothermal heat flux (0.09 W/m^2). The thickness of the snowpack is determined based on snow density and the amount of snow on the ground (water equivalent), calculated as the cumulative difference between snowfall and snowmelt. The profile of snow density is simulated considering compaction and destructive metamorphism. Soil water dynamics are simulated considering water input (rainfall and snowmelt), output (evaporation and transpiration), and distribution among soil layers. Soil thawing and freezing, and associated changes in fractions of ice and liquid water, are determined based on energy conservation. The NEST model

has been tested and applied at different scales to predict thermal conditions for areas across Canada
225 (Zhang et al., 2005, 2006, 2008, 2012, 2014). A more detailed description of the model can be found
in Zhang et al. (2003).

In this study NEST is used to model two scenarios. The first scenario (pingo scenario) models
permafrost aggradation involving growth of near pure ice. This scenario provides an analogue for
the growth of pingo ice over a pressurized sub-pingo water lens, which is assumed to be continuously
230 supplied with expelled pore water from permafrost aggradation in the surrounding sediments (Fig.
1) (Mackay, 1977). In this scenario we assumed that ground materials consist of a saturated porous
material with 99% porosity to represent a growing ice core over a sub-pingo water lens. As the model
was developed for terrestrial conditions, we used saturated porous material to represent sub-pingo
water lens. The freezing of this water results in the growth of pingo ice. Given these assumptions
235 and a reasonable approximation for the date of pingo inception, the modeled thickness of permafrost
growth as near pure ice should approximate maximum height of the feature.

The second scenario is a hypothesis for the aggradation of permafrost into saturated sand without
the development of pingo ice (non-pingo scenario). In this case ground materials consist of saturated
sand with 33% porosity. In both scenarios we assumed that the residual pond drained and permafrost
240 aggradation began in 1940. However, since we do not know the exact date when the residual pond
drained, we additionally ran the model for the two scenarios using start dates of 1930 and 1950. We
also do not know when the larger basin drained within which the pingo has developed.

The model input data of daily air temperature and precipitation were compiled from observations
at Tuktoyaktuk weather station (about 10 km away from the study site and observations began in
245 1948) with gaps filled based on observations at Inuvik (about 115 km away from the study area). Be-
fore 1948, the climate data were filled using the data from 1960-1967. Vapor pressure was estimated
using daily minimum air temperature and daily solar radiation was estimated based on latitude, day
of year and diurnal temperature range (Zhang et al., 2012). For the pingo scenario, the model was
initialized using a surface temperature typical for a drained lake bottom of 4° C (Mackay, 1987). For
250 the non-pingo scenario, the model was initialized using the climate data in 1960, a typical climate
year of the area.

5.1 Modeling results

In the pingo scenario, the modeled thickness of permafrost was 10.7 (9.9-11.2) m by the end of 2014,
which is very similar to the maximum relief of the observed feature (12 m) (Fig 2). The modeled
255 growth of ice thickness from 1972-2014 was 3.8 (3.6-4.3) m, which is comparable to the 4 m of
vertical heave that occurred over that time period. The modeled growth rate of ice thickness during
2011-2014 was 7.4 (7.1-8.0) cm/year, which is larger than the satellite observed maximum uplift rate
(2.7 cm/year).

In the non-pingo scenario, the modeled freezing front was at the depth of 22.6 m by the end of 260 2014. The 9% volume expansion due to freezing of soil pore water to a modeled depth of 22.6 m produced 0.67 (0.61 - 0.71) m of uplift by the end of 2014. This is much smaller than the actual height of the observed feature (12 m) (Fig 2). Similarly, uplift during 1972-2014 was only 0.24 (0.22-0.27) m, which is much smaller than the 4 m of vertical heave observed during that period. The maximum uplift rate during 2011-2014 was only 0.46 (0.43-0.50) cm/year, which is much smaller than the 265 observed maximum uplift rate of 2.7 cm/year (Fig 8).

The time series of the modeled permafrost aggradation rates (i.e. deepening rates of the freezing front), air temperature, and DInSAR observed uplift rates for the 2011-2014 period are shown in Fig 9. After seven decades of permafrost aggradation the pingo (ice) and non-pingo (saturated sand) modeling scenarios showed slower deepening rate of the freezing front in February and April in 270 comparison with a more rapid rate of deepening in July and October (Fig 9), respectively. Due to a lag in the downward propagation of the winter cooling wave, the freezing front descends faster in summer than winter. The 10-year difference in the model initiation time resulted in less than a 10% difference in the depths of the freezing front and less than a month of phase shift, which are considered insignificant.

275 The satellite observed seasonal uplift rate reaches peak values in late July (red dashed line in Fig 9), which coincides with the annual maximum rate of downward permafrost growth as predicted by the NEST model for the pingo scenario. Active-layer thaw in summer typically yields surface settlement in areas of permafrost terrain. These DInSAR results suggest that the seasonal pattern of satellite observed land uplift is likely driven by increased water pressure beneath the pingo and subsequent freezing of the water lens. In order to produce the observed rate of uplift during the 280 2011-2014 monitoring period, and a height increase of 4 m during 1972-2014, permafrost aggradation must have been accompanied by the pore water expulsion from the surrounding areas and the growth of intrusive and/or segregated ice beneath the pingo feature. It is likely that pressure conditions beneath the growing pingo feature are not always conducive to growth of intrusive ice, so 285 that downward aggradation of permafrost beneath the pingo is yielding both lens ice and pore ice (Mackay, 1979). Since the basin within which the pingo is growing may have drained long before 1940, the rate of permafrost aggradation we model in sands does not likely represent the current rate of permafrost aggradation into the surrounding drained lake basin.

6 Discussion and Conclusions

290 The relief and size of this feature, in conjunction with observed rates of deformation suggest that uplift has occurred at least in part due to presence of a sub-pingo water lens and growth of intrusive ice. For this to occur, pore water must have been expelled by permafrost aggrading into the surrounding drained lake basin by aggrading permafrost to a growing pingo. This process would in-

crease pore water pressure and maintain a sub-pingo water lens to enable growth of intrusive ice and
295 heave of the ground surface, consistent with the common growth mechanism of hydrostatic pingos
in Tuktoyaktuk Coastlands (Mackay, 1998).

Based on DInSAR observations, the pingo volume increase during 2011-2014 is $4672 \text{ m}^3/\text{year}$. If
this change in volume was caused in association with the formation of intrusive ice by freezing of
the sub-pingo water lens, the surrounding area has to be larger than $1.2 \times 10^6 \text{ m}^2$, or $\sim 56\%$ of the
300 whole lake basin, assuming that only $60\% \times 9\%$ of the pore water is expelled (Mackay, 1987). This
lake basin is large enough to supply such amount of water for the growth of the pingo.

Although significant attention has been given to climate warming and permafrost degradation in
the arctic regions, landscape forming processes associated with permafrost aggradation and ground
ice development are still very important. In spite of a large number (i.e. > 11000) of pingos on
305 Earth their growth has never been detected by satellite radar interferometry. Previous studies of
pingo dynamics were largely based on annual or semi-annual field observations performed over a
long period of time (Mackay, 1998). Since the size of most pingos is comparable to the spatial
resolution of early SAR sensors their dynamics could not have been easily detected by such sensors.
Seasonal land cover changes (i.e. vegetation growth, snow cover) further complicated interferometric
310 analysis. The availability of high-resolution satellite radar data and novel processing techniques,
coupled with field validation provide new opportunities to study the dynamics of pingos and other
permafrost processes associated with ground ice development and degradation at high temporal and
spatial resolutions. We demonstrated that DInSAR data allows discrimination of ground movements
characteristic of processes involved in pingo development.

315 Historical aerial photographs, field measurements and modeling suggest that the satellite detected
feature is an actively growing pingo that developed over a residual pond in a large drained lake
basin. Uplift of the ground at this location initiated sometime between 1935 and 1951 following lake
drainage and is caused at least in part by the growth of intrusive ice, because the 9% expansion of
pore water during permafrost aggradation in saturated sands with no free drainage is not sufficient
320 to explain the observed short- and long-term deformation rates. The estimated rates and patterns
of uplift determined by DInSAR are within the range of conditions determined by Mackay (1998)
during the long-term study of numerous growing pingos in the study region. Thermal modeling
of permafrost growth involving the freezing of a water lens indicated that modeled thickness of
ice growth from 1940 to 2014 is comparable with the maximum height of this feature. Similarly,
325 the modeled thickness of ice growth from 1972 to 2014 is comparable with changes in elevation
of the feature estimated from aerial photographs. Pattern of variation in the uplift rate seen in the
DInSAR data also matches the modeled seasonal change in the deepening rate of freezing front.
Seasonal changes in the rate of pingo ice growth have also been observed in alternating clear and
bubble bands in pingo ice (Mackay, 1990). The model of a pingo as an elastic uniformly loaded
330 elliptical plate with clamped edge provided a good fit to the observed feature. The location, and

semi-major and semi-minor radii of the modeled feature are 69.3728°N , 133.1688°W , 348 ± 2 m and 290 ± 2 m. Our observations indicate that this pingo is one of the largest (diameter) actively growing pingos in this region. DInSAR could also be used for studying other known pingos, pingo remnants and pingo-like features observed on Mars. Future monitoring of this and other ground-ice related landforms by interferometric satellite radar, which can yield high resolution spatial and temporal data can compliment field-based investigations and contribute to a better understanding of permafrost related processes, including the impact of climate change.

This analysis of ground deformation in the Tuktoyaktuk Coastlands (Fig 6) only shows active deformation, resembling pingo growth, at one single site. However, earlier field measurements performed by Mackay (1998) showed that many of the smaller pingos in the Tuktoyaktuk Coastlands are actively growing. The majority of RADARSAT-2 interferograms used in this study were contaminated by various sources of noise, including decorrelation caused by seasonal changes and snow cover, atmospheric disturbances and residual orbital ramps, which resulted in reduced precision. However, a few interferograms acquired in ideal conditions (i.e. small spatial baseline, frozen ground, no snow) were of an exceptionally high quality. In these interferograms no active deformation processes of similar spatial extent and magnitude were observed except for the pingo studied here. The discrepancy between long-term field measurements and these results is likely caused by the inability of our processing methodology to resolve deformation at smaller spatial scales. It is recommended to use high-resolution DEM for removing the topographic phase for improving DInSAR spatial resolution.

Acknowledgements. We would like to acknowledge field assistance provided by Abra Martin, Ciara Sharpe, Robin Felix, Doug Panaktalok and Pierre Berube. Assistance with mapping was provided by Rebecca Segal. Funding support for this project was provided by the NWT CIMP program, the Natural Science and Engineering Research Council, and the Canada Foundation for Innovation, and Polar Knowledge Canada S&T Program (project 1516-186). Post-processing was performed and figures were plotted with R, GMT and GNUPLOT software. RADARSAT-2 Data and Products ©MacDONALD, DETTWILER AND ASSOCIATES LTD. (2011-2014) - All Rights Reserved. RADARSAT is an official mark of the Canadian Space Agency. We thank C.R. Burn and N. Short for their valuable comments on an earlier version of this manuscript.

References

- 360 Burr, D., Tanaka, K., and Yoshikawa, K.: Pingos on Earth and Mars, *Planetary and Space Science*, 57, 541–555, doi:10.1016/j.pss.2008.11.003, 2009.
- Chen, F., Lin, H., Li, Z., Chen, Q., and Zhou, J.: Interaction between permafrost and infrastructure along the Qinghai-Tibet Railway detected via jointly analysis of C- and L-band small baseline SAR interferometry., *Remote Sensing of Environment*, 123, 532–540, doi:10.1016/j.rse.2012.04.020, 2012.
- 365 Chen, F., Lin, H., Zhou, W., Hong, T., and Wang, G.: Surface deformation detected by ALOS PALSAR small baseline SAR interferometry over permafrost environment of Beiluhe section, Tibet Plateau, China, *Remote Sensing of Environment*, 138, 10–18, 2013.
- Costantini, M.: A novel phase unwrapping method based on network programming, *IEEE Transactions on Geoscience and Remote Sensing*, 36, 813–821, 1998.
- 370 Dundas, C., Mellon, M., McEwen, A., Lefort, A., Keszthelyi, L., and Thomas, N.: HiRISE observations of fractured mounds: Possible Martian pingos, *Geophysical Research Letters*, 35, doi:10.1029/2007GL031798, 2008.
- Freitag, D. and McFadden, T.: *Introduction to Cold Regions Engineering*, ASCE Publications, 1997.
- Goldstein, R. and Werner, C.: Radar interferogram filtering for geophysical applications, *Geophysical Research Letters*, 25, 4035–4038, 1998.
- 375 Grosse, G. and Jones, B.: Spatial distribution of pingos in northern Asia, *The Cryosphere*, 5, 13–33, doi:10.5194/tc-5-13-2011, 2011.
- Gurney, S.: Aspects of the genesis and geomorphology of pingos: perennial permafrost mounds, *Progress in Physical Geography*, 22, 307–324, 1998.
- 380 Ingersoll, L., Zobel, O., and Ingersoll, A.: *Heat conduction: with engineering, geological and other applications*,., Madison University of Wisconsin Press, Madison, 325p, 1954.
- Jin, H., Chang, X., and Wang, S.: Evolution of permafrost on the Qinghai-Xizang (Tibet) Plateau since the end of the late Pleistocene, *Journal of Geophysical Research*, 112, doi:10.1029/2006JF000521, 2007.
- Jones, B., Grosse, G., Hinkel, K., Arp, C., Walker, S., Beck, R., and Galloway, J.: Assessment of pingo distribution and morphometry using an IfSAR derived digital surface model, western Arctic Coastal Plain, Northern Alaska, *Geomorphology*, 138, 1–14, doi:10.1016/j.geomorph.2011.08.007, 2011.
- 385 Liu, L., Zhang, T., and Wahr, J.: InSAR measurements of surface deformation over permafrost on the North Slope of Alaska., *Journal of Geophysical Research - Earth Surface*, 115, doi:dx.doi.org/10.1029/2009JF001547, 2010.
- 390 Mackay, J.: Pulsating pingos, Tuktoyaktuk Peninsula, N.W.T., *Canadian Journal of Earth Sciences*, 14, 209–222, doi:10.1139/e77-023, 1977.
- Mackay, J.: Pingos of the Tuktoyaktuk Peninsula Area, Northwest Territories, *Geographie physique et Quaternaire*, 33, 3–61, 1979.
- Mackay, J.: Some mechanical aspects of pingo growth and failure, western Arctic coast, Canada, *Canadian Journal of Earth Sciences*, 24, 1108–1119, 1987.
- 395 Mackay, J.: Seasonal growth banks in pingo ice., *Canadian Journal of Earth Science*, 27, 1115–1125, 1990.

- Mackay, J.: Arctic ecosystems in semi-arid regions: Implications for resource management., chap. Lake stability in an ice-rich permafrost environment: examples from the western Arctic coast., pp. 1–25, Environment Canada, Saskatoon, Sask., NHRI Symposium Series 7, 1992.
- 400 Mackay, J.: A full-scale field experiment (1978-1995) on the growth of permafrost by means of lake drainage, western Arctic coast: a discussion of the method and some results, *Canadian Journal of Earth Sciences*, 34, 17–33, doi:10.1139/e17-002, 1997.
- Mackay, J.: Pingo growth and collapse, Tuktoyaktuk Peninsula Area, Western Arctic Coast, Canada: A long-term field study, *Geographie physique et Quaternaire*, 3, 271–323, 1998.
- 405 Nelder, J. and Mead, R.: A simplex method for function minimization, *Computational Journal*, 7, 308–313, 1965.
- Samsonov, S. and d’Oreye, N.: Multidimensional time series analysis of ground deformation from multiple InSAR data sets applied to Virunga Volcanic Province, *Geophysical Journal International*, 191, 1095–1108, doi:10.1111/j.1365-246X.2012.05669.x, 2012.
- 410 Samsonov, S., Beavan, J., Gonzalez, P., Tiampo, K., and J., F.: Ground deformation in the Taupo Volcanic Zone, New Zealand observed by ALOS PALSAR interferometry, *Geophysical Journal International*, 187, 147–160, 2011a.
- Samsonov, S., van der Kooij, M., and Tiampo, K.: A simultaneous inversion for deformation rates and topographic errors of DInSAR data utilizing linear least square inversion technique, *Computers and Geosciences*, 415 37, 1083–1091, 2011b.
- Samsonov, S., d’Oreye, N., González, P., Tiampo, K., Ertolahti, L., and Clague, J.: Rapidly accelerating subsidence in the Greater Vancouver region from two decades of ERS-ENVISAT-RADARSAT-2 DInSAR measurements, *Remote Sensing of Environment*, 143, 180 – 191, doi:10.1016/j.rse.2013.12.017, 2014.
- Schmeltz, M., Rignot, E., and MacAyea, D.: Tidal flexure along ice-sheet margins: comparison of InSAR with 420 an elastic-plate model, *Annals of Glaciology*, 34, 202–208, 2002.
- Soare, R., Conway, S., Dohm, J., and El-Maarry, M.: Possible open-system (hydraulic) pingos in and around the Argyre impact region of Mars, *Earth and Planetary Science Letters*, 398, 2014.
- Timoshenko, S. and Woinowsky-Krieger, S.: *Theory of Plates and Shells*, (Engineering Societies Monographs), McGraw-Hill Book Company, Inc., 1959.
- 425 Wegmuller, U. and Werner, C.: GAMMA SAR processor and interferometry software, in: *The 3rd ERS symposium on space at the service of our environment*, Florence, Italy, 1997.
- Wu, Z., Barosh, P., Hu, D., Wu, Z., Peisheng, Y., Qisheng, L., and Chunjing, Z.: Migrating pingos in the permafrost region of the Tibetan Plateau, China and their hazard along the Golmud-Lhasa railway, *Engineering Geology*, 79, 267–287, doi:10.1016/j.enggeo.2005.02.003, 2005.
- 430 Yang, Z., Still, B., and Ge, X.: Mechanical properties of seasonally frozen and permafrost soils at high strain rate, *Cold Regions Science and Technology*, 113, 12–19, doi:10.1016/j.coldregions.2015.02.008, 2015.
- Yoshikawa, K., Sharkhuu, N., and Sharkhuu, A.: Groundwater hydrology and stable isotope analysis of an open-system pingo in Northwestern Mongolia, *Permafrost and Periglacial Processes*, 24, 175–183, 2013.
- Zhang, Y., Chen, W., and Cihlar, J.: A process-based model for quantifying the impact of climate change on 435 permafrost thermal regimes, *Journal of Geophysical Research (Atmospheres)*, 108, 4695, 2003.

- Zhang, Y., Chen, W., Smith, S., Riseborough, D., and Cihlar, J.: Soil temperature in Canada during the twentieth century: complex responses to atmospheric climate change., *Journal of Geophysical Research*, 110, doi:10.1029/2004JD004910, 2005.
- 440 Zhang, Y., Chen, W., and Riseborough, D.: Temporal and spatial changes of permafrost in Canada since the end of the Little Ice Age, *Journal of Geophysical Research*, 111, doi:10.1029/2006JD007284, 2006.
- Zhang, Y., Chen, W., and Riseborough, D.: Disequilibrium response of permafrost thaw to climate warming in Canada over 1850-2100, *Geophysical Research Letters*, 35, doi:10.1029/2007GL032117, 2008.
- 445 Zhang, Y., Li, J., Wang, X., Chen, W., Sladen, W., Dyke, L., Dredge, L., Poitevin, J., McLennan, D., Stewart, H., Kowalchuk, S., Wu, W., Kershaw, G. P., and Brook, R.: Modelling and mapping permafrost at high spatial resolution in Wapusk National Park, Hudson Bay Lowlands, *Canadian Journal of Earth Sciences*, 49, 925–937, doi:10.1139/E2012-031, 2012.
- Zhang, Y., Olthof, I., Fraser, R., and Wolfe, S.: A new approach to mapping permafrost and change incorporating uncertainties in ground conditions and climate projections, *The Cryosphere*, 8, 2177–2194, doi:10.5194/tc-8-2177-2014, 2014.

Table 1. RADARSAT-2 Ultra-Fine 3 (U3) Synthetic Aperture Radar images used in this study, θ is azimuth and ϕ is incidence angle, N is number of images and M is number of interferograms computed for each data set. Time series span 19 June 2011 - 7 September 2014 interval when both ascending and descending data are available.

DInSAR set	Orbit	Coverage	θ°	ϕ°	N	M
RADARSAT-2, U3	asc	20110612-20140924	344	32	16	61
RADARSAT-2, U3	dsc	20110619-20140907	-165	32	17	62
Total:		20110619-20140907			31	123




Figure 1. Growth of close system pingo (modified from Mackay (1998)): (a) lake underlain by basin of unfrozen saturated sands; (b) lake after rapid drainage, permafrost aggradation, pore water expulsion, hydrostatic pressure increase beneath residual pond with thin layer of permafrost; (c) growing pingo underlain by sub-pingo water lens.




Figure 2. Study site located in Northwest Territories, Canada is outlined in red. Background is 90 m resolution Digital Elevation Model (GeoGratis Web Services, Natural Resources Canada). Ascending and descending RADARSAT-2 Ultra-fine 3 (U3) frames are outlined in black. Black circles show locations of pingos identified on Canada Base Map (GeoGratis Web Services, Natural Resources Canada). Brown circles show locations of suspected pingos visually identified in high resolution optical imagery. Names of known pingos are shown in black. Extent in top-left corner shows location of site in North America.




Figure 3. Perpendicular baseline and time span of ascending (top) and descending (bottom) interferograms used in this study.




Figure 4. (top) Historical aerial photographs of study site (National Air Photo Library, Natural Resources Canada) from 1935, 1950, 1972, and 2004 showing that the study site was occupied by a pond in 1935. For reference length of a is 560 m and length of b is 770 m. (bottom) Photograph of study site taken in August, 2014 from helicopter (viewing from east).




Figure 5. Elevation (>5 m) in 1972 and 2004 from stereo-photo analysis and in 2014 from Total Station surveys. Background is 14 August, 2014 radar backscatter intensity.




Figure 6. Line-of-sight linear deformation rate during 2011-2014 period computed from ascending (top) and descending (bottom) RADARSAT-2 U3 imagery. Study site is outlined in red. Black circles show locations of known pingos from Canada Base Map (GeoGratis Web Services, Natural Resources Canada). Brown circles show locations of suspected pingos visually identified in high resolution optical imagery. Decorrelated areas are shown in white. *R* is reference region. Extents in top-left corner show deformation rate histograms with red arrows marking 95% confidence interval.




Figure 7. Selected images of vertical ground deformation time series during 19 June, 2011 - 7 September 2014 computed from ascending and descending RADARSAT-2 data using Multidimensional Small Baseline Subset (MSBAS) technique. Decorrelated areas are shown in white. *R* is reference region. Gray image shows 14 August, 2014 radar backscatter intensity.




Figure 8. (top) Observed data, model and residual of vertical linear deformation rate. Decorrelated areas are shown in white. Best fit model corresponds to ellipse located at $69.3728^{\circ}\text{N}, 133.1688^{\circ}\text{W}$ with semi-major and semi-minor radii of 348 ± 2 m and 290 ± 2 m, tilted 27 ± 3 degrees clockwise, with magnitude of $\frac{\Delta a}{D} = 1.46 \pm 0.02 \times 10^{-10} \text{ m}^{-2}$. (bottom) Latitudinal and longitudinal profiles across uplifting area. Observed data is plotted in black, modeled data is plotted in red.




Figure 9. (top) Observed and modeled (according to equation 2) uplift time series for area experiencing maximum uplift and (bottom) modeled deepening rate of freezing front for pingo (blue line) and non-pingo (brown line) scenarios assuming lake drained in 1940, modeled (according to equation 2) uplift rate (red dashed line), and mean daily air temperature observed at Tuktoyaktuk weather station located 10 km north-east from study site (gray line). Acceleration of uplift is observed in July when deepening rate of freezing front is largest and air temperature reaches annual maximum.