

Answer to reviewer's comments

We would like to thank the three reviewers for their time and constructive comments. We understand and sympathize with the request for more details from all reviewers, but are unable to satisfy all such requests because of the nature of a TC Brief Communication. Having chosen this format, we are restricted to three figures and a maximum of four journal pages as well as only 20 references. Below we address all comments and suggestions (reviewer's comments in black, our reply in blue) and we uploaded a revised version of the manuscript. Within the available space, we have tried to accommodate the request for more details.

Reviewer 1: Ted Scambos

The authors have identified a newly-propagating rift in the southern Larsen C using satellite images. The rift, formerly stable with the rift tip lying just inside a suture zone of marine ice, has grown significantly in 2013-2014 as revealed by Landsat images. The authors describe some scenarios for future growth and eventual calving of a major iceberg from the Larsen C, which they show would adjust the strain field significantly.

The paper requires a table of the Landsat images used. There is a significant change in the imagery between Landsat 7 and Landsat 8, but this is not mentioned or discussed. Improved radiometry could lead to a more sensitive detection of the rift tip, and falsely appear as a rapid growth of the rift. How did they insure that the interpretations were consistent across sensors? Overall, the authors were understandably eager to get this out in front of the community; to do so, they cut short a lot of descriptive details, a lot of analysis of the actual observations, a lot of details of image processing, tables of images, data points, etc. and a lot of other analyses that are straightforward but time-consuming to do.

These are understandable comments but the reasons behind the brief description of the methods is more to do with the format of the paper than any desire for speed. In short we included only what was necessary. We do not believe that the improved radiometric resolution of Landsat 8 over Landsat 7 will have made any real difference in sensitivity of rift tip detection. This will be much more sensitive to spatial resolution (which is identical) and illumination (which is not controlled). We also find no rapid growth of the tip between our use of Landsat 7 and Landsat 8 data. There may be a lot of further analyses that are straightforward but time consuming, but we don't believe that these would alter the results, which are quite clear and unambiguous. We have modified the text in places to clarify some of the points above. We also included a table of data points as supplementary material.

Data that might have been brought to bear: MODIS thermal images (identify date ranges of rapid progression of the fracture, and investigate what might have triggered them); TerraSAR images (same goal). IceBridge data (radar for ice thickness and detailed surface topo). Repeat ice velocity mapping. More analysis of past calvings and rift progressions. Oceanographic data (speed of sub-shelf water flow, from moorings within the ice shelf?) It is likely that the author team is working on some of these things during the review process.

Great suggestions for further work. We are content that the paper is complete as it stands, and lack the space for further work in this manuscript.

I should probably note a potential deal-breaker for the paper's hypothesis up front: Figure 1 and Figure 2 show a fracture in the same new zone, downstream of the current one, which has

been there for some time (see http://nsidc.org/data/iceshelves_images/ and <http://nsidc.org/MMS/moa/moamap.html>). If a rift has occurred in the same area in the past and not continued to propagate further and cause a calving, then the point of this study is a bit moot. Note the many structural similarities between the downstream 'stalled' rift and the new rift. Looking at current maps of flow speed, the downstream rift was in the new rift's position approximately 60 years ago. A check of earlier Landsat images (LIMA, Landsat TM, Landsat MSS) would at least indicate if the downstream rift has been there for decades.

It is correct that there is a surface feature in the Trail flow unit further downstream, but the essential difference is that the propagation of it is stalled on both ends. Landsat imagery indicates that it is not an open rift as the one discussed in this manuscript, but rather the surface expression of a basal crevasse, which can form on the ice shelf (Luckman et al, 2012; McGrath et al., 2012).

The open rift that we describe in this paper reaches from close to the calving front at Gipps ice rise through the suture zone, which is a deviation from the previous pattern of rift propagation in this area. Thus we think that the here described rift and the feature you describe cannot be compared.

P863 L15 – I think 'medium' is subjective, perhaps just say 'at 250 m pixel scale' (note also that pixels are not equal to resolution, picky point but often forgotten).

Corrected

P863-864 L26-L1 – 'minimized by careful control of brightness and contrast' - a less subjective approach would be to high-pass filter the data and then match histograms of the image area of the shelf with a reference image. I admit, it's not likely to change the measurements much, but it sounds better than 'we turned the knobs'.

We have amended the text to clarify the way in which image contrast was handled, and agree that a procedure based on high-pass filtering is highly unlikely to change the measurements.

P864 – it reads as though the selection of the two calving scenarios was also somewhat subjective. More of a discussion here of past calvings of the Larsen C could strengthen the choice. Icebridge data or surface radar would help justify the image-based picks. For Scenario 2, using the ice front as a guide seems very ad hoc. Since you have a numerical model, is there not some indication of where a fracture might propagate?

We chose rift scenarios carefully to minimise subjectivity, but there will always be an element of choice. A model-based projection requires a completely different type of model to the one discussed here.

We chose the first scenario based on the features within the satellite image which is more or less the shortest connection via the weaknesses to the front. The ice bridge data available in this region does not resolve these features, so we cannot say for sure that they are basal crevasses. However, ground based radar surveys in different parts of the ice shelf across similar features proved them to be connected to basal crevasses (Luckman et al, 2012; McGrath et al., 2012).

A previous calving in 1986 followed a similar feature, which can be seen from the shape of the calving front in 1988 as shown in figure 1. The second scenario assumes a more extreme event, which did not happen before in that way. However, there are also rifts visible reaching from the kink in the calving front towards the interior of the ice shelf, and the last large

calving which occurred in 2008 was delineated by a similar straight line towards the front at the northern end.

P865 L10-11, please give a quantitative speed to go with ‘modestly’. What is the basis for saying ‘previously appeared to resist transverse fractures’? If you mean that the rifttips terminated in the suture zone, this might not be how you want to say it, because ‘previously’ implies that something changed to permit rift propagation. L12, I think a look at more available data might constrain the timing of a rapid jump more accurately.

Again, what are the quantitative speeds, or lower limits, for all of the time intervals that rift propagation was mapped?

We would prefer not to give rift propagation rates, as we only have discrete measurements in time and therefore only know the mean rate of rift propagation. The instantaneous rates may be much faster than this. The rift has been growing slowly while in the suture zone, and we assume this is due to the warmer and thus softer ice, which can better accommodate stress by ductile behavior (see Kulesa et al. 2014). Furthermore we assume that the fracturing through the Trail inlet has happened more or less instantly, like a calving event would. Thus it is difficult to put rates to the propagation and we changed the wording in the manuscript to better reflect this.

P865 Numerical model. Did you also look at the present-day stresses with the rift in place (but not broken away)? If this is truly a fracture through the ice sheet, it will significantly alter the stress field on either side – this experiment should be run. This brings up a more general question about rift propagation – as a rift moves across a shelf, the ‘lever arm’ of ice that has been set apart by the fracture grows. Wind and ocean stress should concentrate at the propagating tip, accumulating from a wide area (the area of the future berg). I note also that you don’t cite or discuss Joughin and MacAyeal in regards to rift propagation (2005 GRL). This would be a good one to look at, and would point directly to the importance of repeat velocity mapping near the riftzone.

We think that you are right that velocity mapping of the area would be a sensible thing to do and is a good way to monitor the current development of the rift and we will consider doing so in the future.

The numerical model we are using here is a diagnostic model for ice shelf dynamics and we used it to determine the stability of the calving front according to the criterion we defined earlier in Kulesa et al.(2014). We did not plan to model the propagation of the rift, although we agree that this may be done with a different model. We are aware of the Joughin and MacAyeal paper but decided not to discuss it here as our focus is not on modeling rift propagation.

P866 I think it is clear that there is much more that could be said about the results of the numerical model. Was there a flow speed change with the new geometries? Does the shear stress present at Bawden Ice Rise change? Is there support for the guess at the calving front for Scenario I or II? What front shape (of several plausible ones) tends to maximize the amount of low, or high, stress-flow ice at the front?

We have added velocity maps from the model as a figure to this reply (figure R1). The change in flow speed is not spectacular and there is still a large impact of Bawden ice rise on the

stress field, which can be seen in the new figure 3 in the revised manuscript, which now also shows the contour line of zero second principal stress.

P866 L23 – ‘development of the rift width – a bit awkward, just say ‘spreading rate’? The following sentence is interesting, and related to a thought I had looking at rifts in Figure 1 – the presence of marine ice in a suture can reduce fracture penetration in that zone, but stress leading to fracture might be propagated on the far side of the suture, causing a ‘leap-frog’ fracture. This seems to be the case downstream of the new rift.

We thought that spreading rate could be misread as propagation, so we would prefer to keep the phrase as it stands.

You are raising an interesting point here, however, even if this is what happened further downstream (the feature you saw as a deal-breaker), there is still the difference that in case of the described rift the suture zone did fail, and that we describe an open rift and not the surface expression of a basal crevasse

Figure 1 – How was the background image produced? It appears to be a shaded relief of the Peninsula (from Alison’s excellent DEM) above the grounding line, and a high-pass filtered MODIS image below that. That is not described.

Yes, we did add the reference for Alison’s data set and altered the figure caption, apologies for the oversight.

Figure 2 – I believe the point labeled Dec 2012 should be Dec 2013? I think it would be important to map this progression better and show the January 2014 point and the August 2014 point on the image. An additional figure showing the rift expansion at the 2011 rift tip point would be interesting as well.

No it’s really 2012, but the Dec. 2013 position is would appear almost identical on the figure. We chose the points to highlight based entirely on significant jumps in the sequence. Adding further points would simply over-write the ones on the map. Due to space limitation we could not add another figure to the manuscript.

Reviewer 2: Catherine Walker

Using MODIS and LandSat imagery, Jansen and co-authors monitor and report upon the propagation of a rift in the Larsen C Ice Shelf that grew rapidly during 2014 after having been mostly stagnant/stable before that year with its rift tip lying in/near a suture zone likely filled with marine ice. The authors then discuss two possible directions for future propagation, likely calving scenarios, and the associated adjustment of the strain field at the ice front, enabling discussion of future stability. This is a neat and important observation that will be interesting to monitor going forward. The paper highlights the very recent development on Larsen C ice shelf and thus I can understand why it’s been submitted so quickly after the observation! However, while the observation and possible calving scenarios are interesting and noteworthy, the rigorousness with which the event(s) are placed into context, analyzed, and detailed is somewhat lacking. With a few fixes, I think this manuscript will be ready for primetime. In order for significant conclusions to be drawn from this work at this stage, though, it is my feeling that some of this analysis needs to be undertaken in a more rigorous manner. Mostly, I think at this point the authors just need to add more detail to further our understanding of rift propagation and ice shelf dynamics near the front.

Thanks for the supportive comments. We do not detect a specific suggestion within this paragraph to which we can respond. However, we repeat that a lot of the reason for lack of detail lies in the nature of a TC Brief Communication which is limited in length and number of figures. We do not agree that the analysis needs more rigour.

It would be helpful, perhaps coming from my specific point of view, to have a table or list of the observed propagation rates for the rift in recent years so that it's obvious that the rate of propagation increased dramatically. While it's clear from the author's finding that the rift had a large increase in rifting rate, it would be interesting to be able to see the change in rate as it crossed the Joerg Peninsula suture zone and approached the Trail Inlet flow unit. Otherwise all the reader knows about the large change in rift propagation rate is that it covered approx. 20 km in 8 months (or 2.5 km yr⁻¹ between August 2014 and January 2015).

The rift tip is only detectable at sufficient precision in high spatial resolution optical data. Satellite radar suffers from insufficient contrast and MODIS data is too coarse. Thus Landsat is the only source of data available, we have used every available image, and all points are plotted in Figure 2 (graph, not map). Between available images, we cannot know the instantaneous rate of rift propagation because our sampling is not regular nor frequent. Nevertheless, the line joining our observations serves to illustrate the mean rate of rift propagation with clearly does vary, and can be related generally to specific regions of the shelf (see discussion). Nevertheless, this is a reasonable comment, so we have amended both the text and the caption to clarify. We do not think it appropriate to calculate propagation rates, but included the data table as a supplementary file.

How fast is this when compared to other years (as far back as the imagery allows)? What is the background rifting rate? Did it grow rapidly at any other time since its initiation or is this the first time in its history that it has exhibited a large jump in size? Related to this point, how do other rifts behave that are nearby this rift? Have any of those rifts, which appear to be similar in structure, exhibited this large jump behavior in their past before becoming more stagnant? Is this observation reminiscent of any other rift propagation events on other ice shelves?

The rift tip we investigate in this paper is traveling along a crevasse feature, which has not moved further into the interior of the ice shelf since it has been documented on remote sensing data, which reaches back to the KATE 200 mosaic from 1975 presented in Skvarka et al., (1994). In Glasser et al., (2009), it is also clearly visible that the tip is at the margin of the suture zone.

A more thorough investigation of previous propagation rates of this rift, or a comparison with neighbouring rifts or rifts on other ice shelves would certainly be interesting, but would much more appropriately be the subject of another paper, and would certainly not fit within this limited length paper. The paper here focusses on one particular rift, its recent development, and its potential impact on stability.

P. 863, L. 3: Expand on this? What is the usual pattern? Are there no other instances of rifts passing through suture zones?

The usual pattern is that they stop at the suture zone. No other rift has been observed penetrating the suture zone. Of course the calving event in 1986 must have gone through it, but that was about 50 kilometers further downstream when it happened.

P. 863, L. 15-22: More description is necessary with regard to the use of the imagery to monitor the rift. How was the MODIS imagery used? Specifically, how was the near-real-time data used to monitor the general propagation and likely future path of the rift? Was the additional length of the rift wide enough to be visible in MODIS imagery? While the authors state that Landsat data at high spatial resolution was used to assess the rift length in all images unobscured by clouds between Nov. 2010 and January 2015, how many images exactly were used? What was the temporal resolution of the cloud-free/useable images? Did smaller-timescale changes occur? Perhaps more helpful here than a full-on description (it might make for slightly wordy paragraph), perhaps just a table of the Landsat/MODIS images used would be fine. Additionally, again perhaps from my own point of view, but it would be helpful to understand how the changes between Landsat 7 and 8 might have affected the measurements since both were used in the study.

We agree that a table of Landsat images would be a good idea and have included it as supplementary material. For the comparability of the sensors please see above in the reply to reviewer 1. As stated in the text, the MODIS data was used only to monitor the general propagation of the rift and predict where it might go. The spatial resolution is not sufficient to add to the rift propagation series (Figure 2). We have clarified this point in the text. We provided the link to the NRT archive simply because this was our source of data and stating it would allow others to repeat the analysis. As we have stated in previous responses, only 15 cloud free images were available and we used all of them. The text has been clarified on this point. The table shows that the transition between Landsat 7 and 8 did not affect our measurements.

How was the starting point of Nov. 2010 selected? Since the rift was first observed prior to 2010, why was it deemed not relevant to track its propagation back to its earliest observation? This probably won't change the overall outcome of this particular paper, but in general if you want to discuss changes in the behavior of a rift, why not start at the beginning? The reasoning behind starting in 2010 (rather than its earliest observation) isn't clear.

The paper focusses on the recent rift propagation and its impact. We chose November 2010 to show three years of data before the recent more rapid propagation. The text has been amended to clarify this point.

P. 863, L. 26- P. 864, L. 1: How were the brightness and contrast controlled? Were there limits set for detectability? What imagery software was used? This may have been mentioned elsewhere, but it should be included here in this section regarding satellite observations.

We have amended the text a little to clarify this point for reviewer 1. However, we don't think this issue is particularly important. Using any image processing software it is possible to optimize contrast at the rift tip. We have no reason to believe that this issue would have impact on the results beyond the stated estimate error of a few tens of meters.

P. 864, L. 5-6: How do you differentiate between surface features and basal crevasses? Was radar used to determine the orientation of the basal crevasses?

Features like these were mapped in other regions of the ice shelf: Surface and basal crevasses are distinguishable due to their spatial scales, please see Luckman et al. (2012) or McGrath et al. (2012) and the answers to reviewer 1. We assume that the surface features are the surface

expression (troughs) of basal crevasses. The rift has been so far propagating along such a feature which originated from Kenyon Peninsula.

P. 864, L. 8-11: Could you expand on the determination of the two Scenarios? Are there any other scenarios? Why are these the most likely? Here it would be relevant to discuss, perhaps, previous calving patterns of Larsen C. These scenarios are described as if they are test cases, but perhaps with more explanation of how they were determined, it would be more clear as to why these were chosen as the two likely scenarios.

As stated in the reply to the reviewer 1 above, both scenarios are based on visible zones of weakness (most likely basal crevasses) and previous calving events followed a similar patterns, only further downstream.

P. 865, L. 10-22: Because this is a paper reporting the recent development of a rift, I would expect a longer results section describing the rift activity and history, placing the current development into context. This paragraph highlights the point that I wanted to make above - it would greatly improve the manuscript as an observation paper to describe the actual observations in detail, rather than just giving a few quantitative values (e.g., 40 km yr⁻¹ widening rate or 20 km in 8 months. . . what are these when placed in context?). Specifically in Line 11, what does 'modestly' mean? What observation led you to the conclusion that the rift 'previously appeared to resist transverse fractures'? This isn't clear. In Line 12, what quantitatively constitutes "dramatically"? Overall this section and the entire manuscript needs to be more focused on quantitative values -while the qualitative observation of its growth and the numerical modeling results are an interesting snapshot, it would be relevant to present these observations and modeling results in the context of quantitative observations.

We agree that the history of the rift is not described in great detail, as we are focusing on the current processes. We do not agree that more numbers would improve the paper. The rates are based on sporadically available imagery, and more numbers would just give a false sense of accuracy. The focus of this paper is the obvious change in the pattern, that the rift penetrated the suture zone, in contrast to all observations made before, and the stability of the calving front after possible calving scenarios.

P. 866: It would be great to enhance the discussion of the numerical model results as it appears to yield very interesting results! With such a model, many different scenarios could be investigated and discussed. Especially, though, it would be great to know more about the set up for the scenarios and the outcomes. Can/does the model show a change in velocity after calving in Scenario I/II/both? Is there any scenario where the velocity of the flow did not change? How does the stress around the rift itself appear? Does it change? Thinking about it from a non-modeling perspective, I would expect the rift to alter the stress field. Does it have any effect on the outcome in Scenario I or II?

We added a summary figure of the calculated flow velocities for both the reference run and both scenarios to this discussion (figure R1), but unfortunately there's no room for this in the manuscript.

Of course the rift is altering the stress field, but this is not relevant for the results with the new calving fronts, where the former rift position delineates the new calving front. We did not run a simulation with the propagated rift, but did simulations with soft fillings in the rift prior to its propagation through the suture zone to investigate the influence on the stress field; you can find these results in Kulesa et al. (2014) in the supplementary material.

It was not our aim to model rift propagation, we wanted to focus on the calving front stability after the calving scenarios.

P. 866, L. 22-24: Here it would be relevant to have a table or list of the rate per year. This would highlight the observed change in rate nicely, in addition to exhibit what is meant by “grown intermittently”. I think a table of values for both rift length and rift width would be a great addition.

We agree that a table of the imagery used and data points would be useful and upload it as supplementary material. We do not agree that presenting rift propagation rates would be appropriate for reasons stated above.

P. 866-867: Perhaps it is outside the scope of this manuscript (likely so), but the subject has me wondering, do the authors have a feeling for why this sudden increase in rift propagation rate occurred or why the rift suddenly jumped across a suture zone? This isn't covered much in the paper as it stands now, and it would be interesting to possibly understand a little bit more about possible reasons for this sudden change, if any.

Here we can only speculate, and this is why we did not include this in the paper, which is focused on the consequences of a possible calving. It might be that the influence of the marine ice layer is weakening due to basal melting (Kulesa et al., 2014; Holland et al., 2015).

Figure 2: Though it might make the figure unwieldy, it would be neat to see the appearance of this region in Nov. 2010 in contrast to this image, so the reader could observe changes not only in the length/shape of the rift in question but also its neighbors. Also, would it be possible to label propagation rates on the plot? Additionally, I think the Dec 2012 label should be Dec 2013?

We considered all of these suggestions carefully. A comparison with the Nov. 2010 image reveals nothing more than that the rift propagated through the suture zone. Changes in neighbouring rifts are not significant over the same time period. We therefore do not think that this would be a useful addition to Fig. 2. For reasons already given, we do not think it appropriate to present propagation rates because these are only mean values between the available observations, and are not necessarily indicative of true propagation rates, especially where images are infrequent. Yes, the label is correct. We labelled only those points that showed significant advance of the rift tip.

Short Comment: Maurice Pelto

Jansen et al (2015) provide a compelling observation of rift extension on Larsen C Ice Shelf that could have important implications. This is an important finding and will prompt further investigation of this feature. Most of the comments below are simply a request for more detail that would help us learn more from this interesting dynamic change in the ice shelf that could have large consequences. This includes possibly referencing other ice shelves that experienced ice losses that could have had similar changes in flow stress fields (flow angles) besides Larsen B. Providing a brief example of model validation is essential. It would be worth noting briefly whether or not there are any velocity output differences between models or with present observations. The model does not have to be reviewed in detail as that is in previous papers.

We include here (figure R1) a comparison of the modeled flow velocities of the reference run and the velocity data of Rignot et al. (2011), which are easier to compare than the point results in Haug et al., (2010) you suggested. The model does not quite capture the strong gradients at Gipps ice rise, as we only have a soft ice filling in the rift and not an open rift. In the central part it agrees better. As we are restricted to three figures in the manuscript we did not include this figure in the paper.

862-21: A different specific example of a rift tip ending at a confluence flow unit would be useful.

The tips of basal crevasses are aligning at the suture zones across the entire Larsen C. The open rifts in this southern area of Larsen C all stopped opening before they reached the margin of the suture zone but there are some examples in the north of the ice shelf of open Rifts stalled by suture zones which are containing marine ice (McGrath et al., 2012).

863-3: Given previous satellite imagery, has a rift not propagated across the suture zone before? This needs to be stipulated along with the interval that imagery was observed. Does not need to be detailed, and can be done at 865-11.

We refer to Glasser et al. (2009), who show a nice overview of structures on the Larsen C. A calving event did occur in 1986, where the suture zone was cut through, but this was 50 km downstream, 25 km downstream of today's calving front.

864-2: Has the width changed near the actual width tip as it has propagated, for example 500 m from the tip how wide is it now compared to at the initiation of the expansion, can be reported later.

As the rift is very thin close to its tip it is not possible to resolve these changes on satellite imagery, so we decided to measure rift width changes at a position further down the rift. The rift tip is detected where its width is just visible, which is inevitably less than one pixel in size.

864-9: What are the existing weaknesses?

The existing weaknesses are most likely basal crevasses in the central front of the ice shelf. We refer to Luckman et al. (2012) to explain our assumption that such features are most likely basal crevasses. Please see also the answers to the reviewer comments above.

864-25: This model has in other studies been validated with comparison of simulated and observed velocities. Details of the model do not have to be reviewed here; however, some means of validation needs to be offered. Haug et al (2010) Figure 3 provides a velocity field for validation.

Please see comment above and figure R1.

865-11: If not addressed earlier refer to the period of observations in satellite images that the rift had not crossed the suture zone.

Since the calving in 1986 the suture zone has not been crossed, we added this to the text.

865-19: Is this November 2010? What is the current rift width at this point and what does that imply? Over what length has the rift width reached a value of twice the ice surface elevation or some critical width versus thickness?

Yes, November 2010, we added this to the manuscript. The current rift tip is given in the supplementary table. As stated earlier, the rift width at the detected tip is necessarily on the order of a pixel in size, so this kind of criteria cannot be investigated.

865-21: How does the actual velocity change as it crosses into the new flow unit? Haug et al (2010) Figure 3 provides a velocity for this region to address this.

We do not see how this is relevant to the paper.

866-7: What were and are the angles? The difference needs to be better illustrated and quantified in Figure 3 it is hard to accurately identify the difference field.

The stress-flow angles may be read approximately from the figure using the colour scale. The important observation is that 'stable' values are a deeper shade of red than 'un-stable' values. We believe that this is sufficiently explained and illustrated and that the implication in the figure is clear. But to make things a little clearer, we added indicators for the direction of the first principal stress to the figure.

866-17: What angles are very low? Is there any thickness or velocity output from the models that would provide further insight to future changes?

There are certainly no thickness changes, as the model is diagnostic, so there is no evolution of geometry. The velocity changes are summarized in the figure 1R below, but in our view they are not essential for the stability of the calving front.

867-11: Is Larsen B the best analog since surface melt played such a key role there? Do either George VI (Figure 2 and 5; Holt et al (2013), Wordie (Figure 4.3; Cook and Vaughan, 2010) or Wilkins (Figure 5, Braun et al (2009)) provide a good example in terms of rift development, ice rise impact or changing flow angle versus calving front? If not no need to cite.

We think that surface melt alone cannot explain the retreat of the calving front of Larsen B. The region between former Larsen A and B is subject to strong surface melt as well and is still there. Thus the geometrical setting and the ice dynamics have to play a role as well.

It would certainly interesting to extend the analysis to other ice shelves, but we haven't done so at the moment and it would be beyond the scope of this Brief Communication manuscript.

References

- Cook, A. J., T. I. Murray, A. Luckman, D. G. Vaughan, and N. E. Barrand. 2012. Antarctic Peninsula 100 m Digital Elevation Model Derived from ASTER GDEM. Boulder, Colorado USA: National Snow and Ice Data Center. <http://dx.doi.org/10.7265/N58K7711>.
- Glasser, N. F., Kulesa, B., Luckman, A., Jansen, D., King, E. C., Sammonds, P. R., Scambos, T. A., and Jezek, K. C., 2009. Surface structure and stability of the Larsen C Ice Shelf, Antarctic Peninsula. *Journal of Glaciology* 55 (191), 400–410. URL <http://dx.doi.org/10.3189/002214309788816597>
- Holland, P. R., Brisbourne, A., Corr, H. F. J., McGrath, D., Purdon, K., Paden, J., Fricker, H. A., Paolo, F. S., and Fleming, A. H., Jan. 2015. Atmospheric and oceanic forcing of Larsen C Ice Shelf thinning. *The Cryosphere Discussions* 9 (1), 251-299.
- Kulesa, B., Jansen, D., Luckman, A. J., King, E. C., and Sammonds, P. R., 2014. Marine ice regulates the future stability of a large Antarctic ice shelf. *Nature communications*, doi: 10.1038/ncomms4707
- Luckman, A. J., Jansen, D., Kulesa, B., King, E. C., Sammonds, P.R., and Benn, D. I., 2012. Basal crevasses in Larsen C Ice Shelf and implication for their global abundance. *TheCryosphere*, 6, 113-123, doi: 10.5194/tc-6-113-2012.
- McGrath, D., Steffen, K., Holland, P.R., Scambos, T. S., Rajaram, H., Abdalati, W., Rignot, E., 2014. The structure and effect of suture zones in the Larsen C Ice Shelf, Antarctica. *Journal of Geophysical Research: Earth Surface*, 119, 588-602, doi: 10.1002/2013JF002935.
- Rignot, E., J. Mouginot, and B. Scheuchl., 2011. MEaSuREsInSAR-Based Antarctica Ice Velocity Map. Version 1. Boulder, Colorado, USA: National Snow and Ice Data Center.
- Skvarca, P., 1994. Changes and surface features of the Larsen Ice Shelf, Antarctica, derived from Landsat and Kosmos mosaics. *Annals of Glaciology*, 20, 6-12.




Figure R1: Comparison of the velocity fields. (a) Simulated velocity field for the reference simulation with the 2015 calving front. (b) Observed velocities (Rignot et al., 2011). (c) simulated velocities for Scenario I. (d) Simulated velocities for Scenario II. Background image is MODIS Aqua, Dec. 3rd 2014. The DEM of the Antarctic Peninsula: Cook et al. (2012).