

A daily, 1-km resolution dataset of downscaled Greenland ice sheet surface mass balance (1958-2015)

Brice Noël¹, Willem Jan van de Berg¹, Horst Machguth^{2,3,4}, Stef Lhermitte⁵,
Ian Howat⁶, Xavier Fettweis⁷, and Michiel R. van den Broeke¹

¹Institute for Marine and Atmospheric research Utrecht, Utrecht University, Utrecht, Netherlands.

²Department of Geography, University of Zurich, Zurich Switzerland.

³Department of Geosciences, University of Fribourg, Fribourg, Switzerland.

⁴Geological Survey of Denmark and Greenland GEUS, Copenhagen Denmark.

⁵KU Leuven, Department of Earth & Environmental Sciences, Leuven, Belgium.

⁶Byrd Polar Research Center and School of Earth Sciences, Ohio State University, Columbus, USA.

⁷Department of Geography, University of Liège, Liège, Belgium.

Correspondence to: Brice Noël (B.P.Y.Noel@uu.nl)

Abstract.

This study presents a dataset of daily, 1-km resolution Greenland ice sheet (GrIS) surface mass balance (SMB) covering the period 1958-2015. Applying corrections for elevation, bare ice albedo and accumulation bias, the high-resolution product is statistically downscaled
5 from the native daily output of the polar regional climate model RACMO2.3 at 11-km. The dataset includes all individual SMB components projected to a down-sampled version of the Greenland Ice Mapping Project (GIMP) digital elevation model and ice mask. The 1-km mask better resolves narrow ablation zones, valley glaciers, fjords and disconnected ice caps. Relative to the 11-km product, the more detailed representation of isolated glaciated areas
10 leads to increased precipitation over the southeastern GrIS. In addition, the downscaled product shows a significant increase in runoff owing to better resolved low-lying marginal glaciated regions. The combined corrections for elevation and bare ice albedo markedly improve model agreement with a newly compiled dataset of ablation measurements.

1 Introduction

During the last two decades, the Greenland ice sheet (GrIS) experienced significant mass loss as a result of increased meltwater runoff and sustained high solid ice discharge from marine-terminating outlet glaciers (Van den Broeke et al., 2009; Rignot et al., 2008, 2011; Sasgen et al., 2012; Shepherd et al., 2012; Enderlin et al., 2014). To fill spatial and temporal gaps in the scarce in-situ observations, regional climate models (RCMs) are often used to produce maps of the GrIS surface mass balance (SMB; Van Angelen et al. (2013); Burgess et al. (2010); Ettema et al. (2010a,b); Fettweis (2007); Fettweis et al. (2005, 2011); Noël et al. (2015); Lucas-Picher et al. (2012)). RCMs explicitly calculate the individual SMB components (Lenaerts et al., 2012), i.e. precipitation, runoff and sublimation, over the entire ice sheet (Fig. 1) at high spatial and temporal resolution and over extended periods. However, the current spatial resolution of RCMs, typically 5 to 20 km, remains too coarse to accurately resolve glaciated areas in topographically complex regions such as small isolated ice caps and marginal outlet glaciers flowing into narrow fjords. In these regions, the relatively coarse elevation and land ice masks used in RCMs might result in runoff underestimation (Franco et al., 2012; Noël et al., 2015), hampering realistic regional SMB estimates. Performing higher-resolution simulations to address these issues would require a substantial computational effort and is thus restricted to case studies of small regions and relatively short time periods.

As an alternative, statistical downscaling can be applied to RCM output. Previously, this method has been applied to the GrIS using global reanalysis and climate data (Hanna et al., 2005, 2008, 2011). Machguth et al. (2013) downscaled near-surface temperature and precipitation from 3 different RCMs (11-25 km spatial resolution) to force a glacier mass balance model on a 250 m grid derived from the Greenland Ice Mapping Project (GIMP) digital elevation model (DEM) (Howat et al., 2014), accurately resolving local glaciers and ice caps of Greenland. Vertical gradients of climate parameters were iteratively calibrated to enable the mass balance model to generate a realistic melt distribution for the period 1980-2010, but the very high resolution restricted the analysis to a few regions. Franco et al. (2012) statistically downscaled GrIS SMB by interpolating each component of the Modèle Atmosphérique Régional (MAR) from the original 25 km grid to a 15 km resolution. This method used local daily vertical gradients, except for precipitation, to correct for elevation differences between MAR and a down-sampled version of the 5 km DEM from Bamber et al. (2001). The elevation correction significantly reduced SMB biases. However, a resolution of 15 km remains insufficient to resolve the rugged topography at the ice sheet margins; to address this issue, near-km resolution is necessary.

Here, we present a new dataset of daily, 1-km resolution GrIS SMB components (precipitation, melt, runoff, refreezing, sublimation and snowdrift erosion) covering the period

1958-2015. The SMB product is statistically downscaled from data of the polar regional climate model RACMO2.3 at 11-km (Fig. 1), using an elevation dependent technique based on the elevation and ice mask from the GIMP DEM (Howat et al., 2014), down-sampled to 1-km. The following section briefly describes RACMO2.3, the GIMP DEM, observational datasets and MODIS bare ice albedo product used to evaluate and correct the downscaled dataset. The downscaling algorithm is explained in Section 3. Downscaled SMB is evaluated using ablation and accumulation measurements in Section 4. Section 5 discusses the downscaling results for four different regions and for the entire ice sheet. The added value, limitations and uncertainties of the downscaling method are argued in Section 6, followed by conclusions in Section 7.

2 Model and data

2.1 The regional climate model RACMO2

A detailed description of the Regional Atmospheric Climate Model (RACMO2) is presented in Van Meijgaard et al. (2008). RACMO2 incorporates the atmospheric dynamics and physics modules from the High Resolution Limited Area Model (HIRLAM) and the European Centre for Medium-range Weather Forecasts Integrated Forecast System (ECMWF-IFS, Undèn et al. (2002)). The polar version of RACMO2 is developed by the Institute for Marine and Atmospheric Research (IMAU), Utrecht University, and is especially adapted for use over ice sheets and other glaciated regions. Polar RACMO2 is interactively coupled to a multi-layer snow module, accounting for firn densification, meltwater percolation, re-freezing and runoff (Ettema et al., 2010a); an albedo scheme with prognostic snow grain size (Kuipers Munneke et al., 2011) and a drifting snow module, simulating snow erosion and the drifting snow contribution to sublimation (Lenaerts et al., 2012). Recently, RACMO2.1 has been updated to RACMO2.3 as discussed in Van Wessem et al. (2014) and Noël et al. (2015). Model evaluation against SMB measurements, collected in the accumulation and ablation zones of the GrIS, showed generally improved agreement (Noël et al., 2015). The native 11-km climate run is forced at the lateral boundaries by ERA-40 (1958-1978, Uppala et al. (2005)) and ERA-Interim (1979-2015, Stark et al. (2007); Dee et al. (2011)) reanalyses and uses the 5 km DEM and ice mask from Bamber et al. (2001).

2.2 GIMP DEM

To downscale RACMO2.3 output, we use the ice mask and topography from the GIMP DEM, described in Howat et al. (2014), and currently considered to be one of the most complete ice masks for Greenland (Rastner et al., 2012). A 1-km ice mask and DEM are obtained by averaging the original 90 m GIMP grid-cells in each 1-km pixel covering Greenland. A

1-km resolution is deemed an acceptable trade-off between improved resolution, i.e. a 121 fold improvement compared to the 11-km grid, and manageable data handling given the daily time resolution, time span (1958-2015) and the number of SMB components. As an example, Figure 2a shows the topography and ice mask from RACMO2.3 at 11-km in central east Greenland (blue box 1 in Fig. 1) and Figure 2b the GIMP DEM at 1-km. The latter much better resolves small scale landforms such as narrow fjords and calving glacier tongues. Integrated over the contiguous GrIS, the ice-covered area of $1.69 \cdot 10^6 \text{ km}^2$ for the 1-km grid represents a 0.5% decrease relative to the 11-km mask. For our SMB calculations, we only consider grounded ice, i.e. we discarded floating ice pixels using a 1-km version of the 90 m grounded ice mask used in Enderlin and Howat (2013).

2.3 Ablation and accumulation measurements

To evaluate the daily downscaled SMB product, we use 1155 SMB measurements collected in the GrIS ablation (1073) and accumulation (182) zones. The ablation dataset (Machguth et al., 2016) was compiled as part of the Programme for Monitoring of the Greenland Ice Sheet (PROMICE) (Van As et al., 2011) and includes stake and AWS measurements retrieved from 213 sites (yellow dots in Fig. 1). Accumulation observations were derived from 182 sites including snow pits and firn cores (Bales et al., 2001, 2009) as well as airborne radar measurements (Overly et al., 2015) (white dots in Fig. 1). We exclusively selected data having a temporal overlap with RACMO2.3 simulations (1958-2015). We rejected observations from sites with a $> 100 \text{ m}$ height bias relative to the representative elevation of the 1-km GIMP topography.

To compare modelled and downscaled SMB with observations, different selection approaches were applied in the ablation and accumulation zones, as described in Noël et al. (2015). In the accumulation zone, we select the closest grid-cell on the 11-km and 1-km grids to represent modelled and downscaled SMB, respectively. In the ablation zone, an altitude correction is applied by selecting the grid-cell with the smallest elevation bias among the closest pixel and its eight adjacent neighbours.

2.4 MODIS bare ice albedo

A 1-km version of the 500 m MODerate-resolution Imaging Spectroradiometer (MODIS) 16-day Albedo product (MCD43A3) is used to retrieve estimates of bare ice albedo in the GrIS ablation zone. Bare ice albedo is estimated as the average of the 5% lowest surface albedo measurements for the period 2000-2015. A similar ice albedo product is used in RACMO2.3 based on MODIS observations between 2001 and 2010 (Noël et al., 2015). In RACMO2.3, bare ice albedo ranges from 0.3, i.e. dark bare ice exposed in the low ablation zone, to 0.55 under persistent snow cover in the GrIS accumulation zone. Bare ice albedo

120 of glaciated pixels with no valid MODIS estimate are set to 0.47.

3 Methods

The daily, 1-km SMB product consists of statistically downscaled output from a previously conducted RACMO2.3 simulation at 11-km, covering the period 1958-2015. RACMO2.3 settings and lateral forcing are described in Noël et al. (2015). The downscaling algorithm
125 corrects the interpolated SMB components using their local regression to elevation. Figure 3 shows the spatial correlation of individual SMB components with elevation on the 11-km RACMO2.3 grid. The spatial correlation is calculated for each grid-box using 8 adjacent ice-covered pixels.

The elevation correction is exclusively applied to the SMB components which show a significant and spatially homogeneous correlation with elevation, i.e. melt, runoff and sublimation
130 (Fig. 3). These SMB components decrease with decreasing air temperature, represented by a negative correlation with elevation (Fig. 3b, d and e). Although precipitation negatively correlates with elevation over most of the ice sheet, the correlation remains small and highly heterogeneous at the margins (Fig. 3a). Snowdrift erosion exhibits a noisy correlation pattern. Therefore, daily precipitation and snowdrift erosion are bi-linearly interpolated to the
135 1-km ice mask without elevation corrections. Refreezing exhibits a marked bimodal correlation pattern (not shown), gradually increasing with height in the ablation zone, where pore space is more abundant, and decreasing towards the ice sheet interior due to limited meltwater supply. For this reason, and in order to have a consistent liquid water balance,
140 daily refreezing is calculated as a residual:

$$RF = RA + ME - RU \quad (1)$$

where RF is the residual refreezing, RA is rainfall, ME is surface melt, and RU is melt-
145 water runoff.

Daily SMB values are obtained by summing the individually downscaled components:

$$SMB = P_{\text{tot}} - RU - SU - ER \quad (2)$$

150

where P_{tot} is total precipitation (liquid and solid), RU is meltwater runoff, SU is total sublimation (from surface and drifting snow) and ER is drifting snow erosion.

3.1 Elevation dependent downscaling

The downscaling algorithm interpolates daily SMB components to the 1-km topography and ice mask in three successive steps (Fig. 4a).

First, the local dependence on elevation is calculated on the original RACMO2.3 11-km grid. Regression parameters are computed on a daily basis and are, therefore, only valid for that specific day. A local regression slope, b_{11km} (mmWE per m, Fig. 4a), is calculated for each ice-covered RACMO2.3 grid-point using the maximum amount of points available, i.e. we use a total of six to nine ice-covered grid cells, being the current one and minimum five to maximum eight adjacent pixels. This minimum number is chosen after testing the downscaling sensitivity to the number of regression cells used, as discussed in Section 3.2. An approximation of the SMB components at mean sea level, a_{11km} (mmWE, Fig. 4a), is then obtained using b_{11km} and the current pixel. The regression is applied to the current grid cell to prevent local estimates of a_{11km} to significantly differ from the original RACMO2.3 value. Local regression parameters for melt and runoff are only computed for pixels experiencing ablation. Moreover, erroneous positive regression slopes, i.e. increasing melt rates with altitude, are discarded until the following stage.

Next, valid estimates of b_{11km} and a_{11km} are extrapolated iteratively on the 11-km grid to fully cover the more extensive 1-km ice mask. To that end, daily regression parameters are extrapolated outwards of the 11-km ice mask by averaging b_{11km} from at least 3 ice-covered pixels from the eight cells surrounding the current one.

Finally, the extrapolated fields of b_{11km} and a_{11km} are bi-linearly interpolated to the 1-km ice mask, providing estimates of b_{1km} and a_{1km} . The downscaled SMB components ($X_{v0.2}$), i.e. runoff, melt and sublimation, are then computed as a linear function of the high-resolution topography as:

$$X_{v0.2} = a_{1km} + b_{1km} \times \text{elevation}_{1km} \quad (3)$$

The downscaled dataset that is based on the above elevation dependent technique is hereafter referred to as version v0.2.

3.2 Sensitivity experiment

Figure 5 shows the difference between 11-km and downscaled GrIS integrated daily runoff in summer 2011. Each line represents a different minimum number of grid-cells, ranging from 3 to 9, used to estimate the local regression of runoff with elevation (b_{11km} ; Fig. 4a). The results are moderately sensitive to the number of regression points used except for the 9 cells setting (current pixel and its 8 neighbours). The latter systematically underestimates

runoff at the beginning and the end of the melt season as it discards all low-lying glaciated pixels at the edge of the GrIS, which experience early melt and largest values of runoff. The standard deviation between the different settings (~ 0.2 Gt/day) is significantly smaller than the difference between 11-km and 1-km runoff (~ 0.6 Gt/day). The more regression points used, the smoother the runoff to elevation gradient field becomes, lowering the downscaled runoff and bringing it closer to the 11-km model output. Conversely, a small number of regression points can lead to spuriously large local gradients. To prevent the downscaling algorithm from substantially converging to, or diverging away from, 11-km RACMO2.3 output, we adopted a setting of minimum 6 regression points, which is closest to the average value of the different experiments (± 0.1 Gt/day).

3.3 Melt and runoff adjustments

RACMO2.3 uses a prescribed bare ice albedo field, typically ranging from 0.30 in the low ablation zone to 0.55 under persistent snow cover. It is based on the 5% lowest MODIS values of surface albedo averaged for the period 2001-2010 (Noël et al., 2015). A comparison with a similar 1-km MODIS product averaged for 2000-2015, ranging from 0.15 to 0.55, shows a systematic overestimation of ice albedo at 11-km, especially for low-lying marginal glacier tongues as shown in e.g. Fig. 12i. This causes melt energy to be underestimated during the melt season. To correct for this, downscaled melt and runoff are adjusted by estimating the missing amount of ice melt (ME_{add}) resulting from underestimated absorption of downward shortwave radiation (SW_d). In addition, as RACMO2.3 calculates radiative fluxes on a horizontal plane, the direct fraction of SW_d is corrected for the slope and orientation of each 1-km glaciated grid-cell, as described in Weiser et al. (2016). For simplicity, we assume SW_d to be equally partitioned between diffuse and direct radiation, and that the sun is exactly in the South at noon. This assumption is purely pragmatic; on the basis of data availability, it could be further refined in future versions of the downscaling procedure. Figs. 6b and 8 show that ablation underestimation in v0.2 is restricted to the low ablation zone ($SMB < -4$ mWE), where bare ice is exposed for long episodes in summer. Therefore, the following corrections are only applied to the ablation zone on days of melting bare ice when both surface runoff and melt are non-zero in the downscaled product v0.2:

$$ME_{\text{add}} = \Delta\alpha \times 0.5 \left(\frac{SW_{d \text{ 1-km}}}{L_f} + \xi \frac{SW_{d \text{ 1-km}}}{L_f} \right) \quad (4)$$

where ME_{add} (mmWE per day) is the additional amount of ice melt calculated at 1-km; $\Delta\alpha$ (-) is the difference between the averaged bare ice albedo retrieved from the set of regression cells used to downscale runoff at 11-km and the MODIS albedo product at 1-km; $SW_{d \text{ 1-km}}$ is the modelled daily cumulated downward shortwave radiation bi-linearly inter-

polated to 1-km; L_f is the latent heat of fusion ($3.337 \cdot 10^5$ J/kg) and ξ (-) is the correction
 225 factor for a tilted plane (Fig. 4b), applied to the direct component of downward shortwave
 radiation:

$$\xi = \frac{\cos(\zeta^*)}{\cos(\zeta)}$$

$$\begin{aligned} 230 \quad \zeta^* &= \sin(\zeta)\cos(a)\cos(\sigma)\cos(\Theta) + \sin(\zeta)\sin(\sigma)\sin(\Theta) \\ &+ \cos(\zeta)\cos(\sigma) \end{aligned}$$

$$\zeta = \arccos\left(\sin(\phi)\sin(\delta) + \cos(H)\cos(\phi)\cos(\delta)\right) \quad (5)$$

235

where ζ^* is the solar angle of incidence for a tilted plane, ζ is the solar zenith angle, a is
 the azimuth of the tilted plane, σ is the local surface slope, Θ is the orientation, ϕ is the
 latitude, δ is the solar declination and H is the hour angle set to 0 at noon (Fig. 4b). All
 angles are expressed in radians.

240

The bare ice albedo bias correction aims at minimizing the misfit between downscaled SMB
 v0.2 and in-situ measurements (Fig. 6b) by estimating the missing runoff in the low ab-
 lation zone. Additional runoff RU_{add} is calculated by applying a daily specific fraction Γ
 (-) to ME_{add} , estimating the melt contribution to surface runoff. Γ is defined as the ratio
 245 between daily downscaled runoff and melt in v0.2 estimated using elevation dependence only:

$$\text{RU}_{\text{add}} = \Gamma \times \text{ME}_{\text{add}} \quad (6)$$

Assuming that the residual misfit between reconstructed and observed SMB (ΔSMB , Fig. 6b)
 250 for the different ablation sites can be ascribed to underestimated runoff in the low ablation
 zone of the GrIS, RU_{add} is then scaled by a factor f_{scale} (-), obtained by computing a
 least-square fit minimising the difference between ΔSMB and RU_{add} using all ablation
 measurements:

$$255 \quad \Delta\text{SMB} = f_{\text{scale}} \times \text{RU}_{\text{add}}$$

$$f_{\text{scale}} = \frac{\sum \Delta\text{SMB} \times \text{RU}_{\text{add}}}{\sum (\text{RU}_{\text{add}})^2} \quad (7)$$

The least square fit yields a value of $f_{\text{scale}} = 1.176$ for the GrIS. This means that RU_{add} , i.e.
 260 accounting for elevation and bare ice albedo corrections, has yet to be increased by $\sim 18\%$

to optimise the agreement between downscaled and in-situ SMB (Fig. 6c). The fact that $f_{\text{scale}} > 1$ strongly suggests that additional processes might play a role in enhancing surface ablation, e.g. underestimation of modelled sensible heat flux from warm air advection along the GrIS periphery (Noël et al., 2015; Fausto et al., 2016) and uncertainties in cloud representation (Van Tricht et al., 2016). However, as the statistical downscaling approach is not designed to correct for these physical processes, we adopted the empirical approach presented above. The adjusted amount of runoff ($\text{RU}_{\text{v1.0}}$) is obtained by adding the missing runoff to the downscaled runoff ($\text{RU}_{\text{v0.2}}$).

$$\text{RU}_{\text{v1.0}} = \text{RU}_{\text{v0.2}} + f_{\text{scale}} \times \text{RU}_{\text{add}} \quad (8)$$

The corrected melt ($\text{ME}_{\text{v1.0}}$) is obtained in a similar fashion and refreezing ($\text{RF}_{\text{v1.0}}$) is estimated as a residual between adjusted melt, runoff and rainfall:

$$\text{ME}_{\text{v1.0}} = \text{ME}_{\text{v0.2}} + \text{ME}_{\text{add}} \quad (9)$$

$$\text{RF}_{\text{v1.0}} = \text{RA} + \text{ME}_{\text{v1.0}} - \text{RU}_{\text{v1.0}} \quad (10)$$

The downscaled SMB dataset resulting from the combined elevation correction and runoff adjustment is referred to as version v1.0 in the following sections.

4 Evaluation of daily downscaled SMB

Figure 6 evaluates the original RACMO2.3 SMB at 11-km (a), the 1-km raw downscaled SMB version v0.2 (b) and the 1-km corrected downscaled SMB version v1.0 (c) (mWE per year) with 1073 observations from 213 ablation sites (yellow dots in Fig. 1). The observational period was matched with the modelled and downscaled SMB using the exact number of days. Each blue star corresponds to the cumulative SMB for a duration ranging from 10 days to a full hydrological year. The downscaled SMB v0.2 agrees better with observations compared to the RACMO2.3 output at 11-km (Figs. 6a and b): we find a significant decrease of the RMSE (190 mmWE or -16%) and a smaller bias (100 mmWE or -21%). The deviation from unity of the regression slope decreases from 0.28 to 0.21 (-25%), and the variance explained increases from 47% to 61%. When applying the bare ice albedo and local orientation corrections, we find further significant improvements relative to version v0.2 (Fig. 6c), with now 78% of the variance explained and a significant decrease in RMSE (270 mmWE or -27%) and bias (310 mmWE or -84%). Red stars represent data from PROMICE station QAS_L (61.03°N, 46.85°W, 310 m.a.s.l; yellow dot in Fig. 11a) situated in an extremely

narrow ablation zone (~ 10 km) at the southwestern tip of Greenland. Here, modelled ablation gradients at 11-km are strongly underestimated in RACMO2.3 and are only marginally better resolved at 1-km. At this site, the additional corrections are especially important to obtain agreement with observations.

Figure 7 compares annual mean observed and downscaled SMB (v1.0) along 8 different SMB transects. There is good agreement for most transects, except for Helheim glacier (66.41N, -38.34W). The downscaled product fails at reproducing the quasi-constant ablation rate (~ -1 mWE) characterizing the Helheim transect. The reason for this low SMB gradient is not clear at present; it may be due to uncertainties in individual observation covering relatively short periods, i.e. 1 or 2 months, which are only limited to the melt season (July-August). Another possible explanation is that Helheim glacier experiences pronounced winter accumulation at low elevations, potentially caused by drifting snow transport, limiting summer ablation. In addition, Nioghalvjerds-fjorden and Storstrømmen transects (Figs. 7a-b) also show significant remaining biases between in-situ and downscaled SMB at elevations lower than 200 m. We hypothesize that these SMB measurements are located on floating glacier tongues with melt ponds, resulting in very low satellite albedo, while stake measurements are performed between ponds on brighter surfaces. As a result, the bare ice albedo correction could be overestimated.

In the accumulation zone, a small improvement is also found compared to v0.2 (Fig. 8), but accumulation remains underestimated. The SMB bias and RMSE are reduced by 0.7 (-2%) and 1.8 mmWE (-3%) whereas the regression slope and variance explained remain unchanged. In the accumulation zone, SMB is mostly driven by precipitation which is bi-linearly interpolated to 1-km without elevation correction. In addition, changes in sublimation are small due to the relatively homogeneous topography of the ice sheet interior, limiting SMB changes through downscaling. Despite significant improvements in the cloud scheme of RACMO2.3 (Noël et al., 2015), clouds become saturated and start to produce precipitation at too low elevations, resulting in overestimated precipitation at the margins, e.g. southeast Greenland, while the ice sheet interior experiences too dry conditions. This precipitation bias is currently being investigated, and we aim to resolve it in the upcoming version RACMO2.4. To overcome the systematic negative SMB bias of RACMO2.3 in the GrIS accumulation zone (-37.5 mmWE/yr, Fig. 8), the daily total precipitation v0.2 is adjusted to correct for underestimation in the ice sheet accumulation zone ($SMB > 0$ mmWE/yr):

$$PR_{v1.0} = PR_{v0.2} + \frac{PR_{v0.2}}{PR_{v0.2}^2} \times \sigma_{SMB} \quad (11)$$

where $PR_{v1.0}$ is the daily adjusted total precipitation v1.0, $PR_{v0.2}$ is the daily bi-linearly

interpolated total precipitation v0.2, $PR_{v0.2}^a$ is the annual cumulative bi-linearly interpolated total precipitation v0.2 and σ_{SMB} is the accumulation zone SMB bias in the downscaled product v1.0.

The final $SMB_{v1.0}$ product is reconstructed as:

$$SMB_{v1.0} = PR_{v1.0} - RU_{v1.0} - SU - ER \quad (12)$$

5 High-resolution SMB patterns: case studies

Table 1 lists annual mean modelled and downscaled SMB components (Gt per year) integrated over four different regions (blue boxes in Fig. 1) as well as over the entire GrIS. These regions were selected for their specific climates, rough topography and narrow glaciated features which were not well resolved at 11-km. Figures 9, 10, 11 and 12 show the ice sheet mask for the selected regions at 11-km (red cells) and 1-km (orange cells) as well as peripheral glaciers and ice caps at 1-km (blue cells), the elevation bias between the 1-km and 11-km DEMs, and the bare ice albedo bias between the 1-km MODIS product and RACMO2.3 at 11-km; the latter figures moreover show the main SMB components at both resolutions for the two downscaled products (v0.2 and v1.0). In the following sections, we discuss the impact of downscaling on regional SMB. Here, SMB components are exclusively integrated over the contiguous GrIS; the SMB of detached ice caps will be discussed in a forthcoming paper.

5.1 Central east Greenland

Central east Greenland (blue box 1 in Fig. 1) is characterised by a large body of interconnected valley glaciers, mostly terminating in narrow glacial fjords. Figure 9a, e and i underline the inability of the 11-km mask to properly represent many glaciated areas, local topography or bare ice albedo. In the 1-km mask, the ice covered area increases ($\sim 2\%$) while the elevation bias can locally exceed 500 m over glacial valleys and small scale promontories (Tab. 1 and Fig. 9e); the average elevation bias is 80 m. These differences affect SMB in two ways. First, precipitation increases by 2.6 Gt/yr (12%) in v0.2 (Tab. 1 and Fig. 9b and f), exclusively caused by the expansion of glaciated area (no elevation correction is applied). Another 1.6 Gt/yr (6%) of precipitation is added in v1.0 (Fig. 9j) to compensate for the systematic negative SMB bias in the GrIS accumulation zone, as discussed in Section 4. For both downscaling versions, changes in runoff mirror the elevation change between the two resolutions (Fig. 9e), highlighting the high sensitivity of runoff to elevation. In version v0.2, integrated runoff increases by 7.7 Gt/yr (48%) (Fig. 9c and g). Furthermore, Fig. 9i reveals

a systematic overestimation of bare ice albedo at 11-km. Correcting for this further increases runoff over the glaciers tongues (Fig. 9k), accounting for ~ 13 Gt/yr (55%) of additional runoff with respect to v0.2 (Tab. 1). Negligible changes in sublimation and drifting snow are found (Tab. 1). As a consequence, integrated SMB on the 1-km mask decreases by 5.3 Gt/yr in version v0.2 (Fig. 9d and h) and by 16.6 Gt/yr in version v1.0 (Fig. 9l). This analysis for central east Greenland demonstrates the importance of accurately reproducing small scale topography and ice albedo to realistically capture local SMB variations.

5.2 Central west Greenland

The 11-km resolution DEM provides a reasonable representation of the wide, gently sloping western ablation zone of the GrIS, where most glaciers are land-terminating. The northern part of the selected area includes several marine-terminating glaciers which are better represented at 1-km (Fig. 10d and h).

Owing to negligible difference in glaciated area, precipitation remains almost unchanged for the two resolutions and versions (~ 15 Gt/yr). In both downscaled versions, enhanced runoff is mostly obtained over narrow, low-lying glaciers tongues and detached ice caps (Fig. 10c, g and k) where most of the elevation and ice albedo biases are found (Fig. 10e, i). On the ice sheet, the elevation correction increased runoff by about 1 Gt/yr (5%) (Fig. 10g) while an additional ~ 2 Gt/yr (10%) (Fig. 10k) can be ascribed to the ice albedo correction (Tab. 1).

5.3 South Greenland

Southeast Greenland (blue box 3 in Fig. 1) is a rugged region (Fig. 10e), characterized by multiple topographically-forced precipitation maxima (Fig. 11b and f) and narrow marginal ablation zones (Fig. 11c, g and k). Similar to central east Greenland, the larger glaciated area (+6.5%, Fig. 11a) at 1-km enhances integrated precipitation by ~ 6 Gt/yr (+7%) in v0.2 and 8.4 Gt/yr (+9%) in v1.0. Increased runoff (2.2 Gt/yr or 5% in version v0.2) at the southern margins can be ascribed to additional melt production over the better resolved narrow ablation zones (Fig. 11g and k) combined with a moderate mean elevation difference (~ 17 m) between both resolutions. In v0.2, the ice mask expansion explains most of the integrated SMB changes, leading to an overall mass gain of 3.3 Gt/yr.

Fig. 6b reveals considerable ablation underestimation in southern Greenland, expressed as a systematic SMB bias of 2 to 4 mWE relative to measurements collected at PROMICE station QAS_L (red dots in Fig. 6a). The main reason for this underestimation is that SMB at this location is characterized by a rare combination of high snowfall and strong summer melt.

The remaining ablation underestimation in v0.2 can be partly ascribed to an overestimated

bare ice albedo (0.47) prescribed in RACMO2.3 (Noël et al., 2015); observed albedo at QAS_L frequently falls to 0.2 during the melt season (Fausto et al., 2016). As a result, the additional bare ice albedo correction significantly improves runoff at station QAS_L (Fig. 6c). Integrated over region 3, runoff increases by another ~ 13 Gt/yr (29%) relative to v0.2 (Fig. 11k). The increased marginal mass loss leads to the expansion of the southern ablation zone towards higher elevations (Fig. 11k and l), in line with local observations (Fig. 6c).

5.4 North Greenland

In north Greenland (blue box 4 in Fig. 1), the climate is dry, and most glaciers are marine-terminating. The ice sheet surface is relatively smooth and homogeneous. The wide ablation zone is reasonably well captured at 11-km, leading to a modest deviation in elevation (~ 43 m) (Fig. 12e). However, the ice-covered area decreases by $\sim 11\%$ between both resolutions as the 11-km grid contained erroneous floating glacier tongues (Fig. 12a). The ice area reduction at 1-km affects precipitation (-0.8 Gt/yr or -12%) (Fig. 12b and f) and runoff (-3.1 Gt/yr or -35%) (Fig. 12c and g), resulting in a small SMB increase (2.3 Gt/yr) in version v0.2 (Fig. 12d and h). Large bare ice albedo discrepancies can be found on five major glaciers (Fig. 12i) where runoff increases substantially (~ 2 Gt/yr or 34%) in version v1.0, further decreasing the integrated SMB by 1.0 Gt/yr compared to v0.2 (Fig. 12h and l).

5.5 Greenland ice sheet

Although similar in area, the 1-km ice sheet mask better resolves peripheral glaciers at the GrIS margins than RACMO2.3 at 11-km. GrIS integrated precipitation increases by 16.6 Gt/yr ($+2\%$) in v0.2, most of which can be ascribed to ice area expansion in the east (2.6 Gt/yr) and south of Greenland (5.8 Gt/yr), where precipitation is large. An additional 56.2 Gt/yr ($+8\%$) is obtained in v1.0 when correcting for the accumulation zone SMB bias. The smooth topography of the ice sheet interior results in a small elevation difference of 4 m between both resolutions. Significant elevation biases are mostly restricted to peripheral glaciers and narrow ablation zones at the GrIS margins. As a result, runoff increases by 13.6 Gt/yr ($+5\%$) in version v0.2. Accounting for the bare ice albedo bias in RACMO2.3 further increases runoff by 69.3 Gt/yr in version v1.0, leading to a much improved agreement with ablation measurements. Of our selected areas, central east and south Greenland contribute 25% and 18% to the total runoff increase in the downscaled product v1.0 owing to the many low-lying glaciers tongues that can only be resolved at 1-km. Due to their smoother topography, north and centre west Greenland contribute much less to the runoff change ($\sim 3\%$ and 1% , respectively). Integrated over the contiguous ice

sheet, SMB is not significantly affected by the elevation dependence for which enhanced precipitation (16.6 Gt/yr) yearly balances the moderate increase in runoff (13.6 Gt/yr). In contrast, the bare ice albedo and precipitation corrections substantially increase marginal runoff (82.9 Gt/yr) and accumulation (72.8 Gt/yr), resulting in a decrease of SMB of -11.1 Gt/yr (-3%) relative to the 11-km product.

6 Added value, limitations and uncertainties

The downscaled SMB v1.0 is the first dataset to provide daily SMB estimates for all outlet glaciers of the GrIS at a 1-km resolution and for 58 years (1958-2015). Relative to the original RACMO2.3 output, this dataset improves local SMB values (Fig. 7) and produces more realistic SMB patterns over rugged glaciated areas along the GrIS margins (Figs. 9-12). Figs. 6 and 8 show that SMB v1.0 is an overall improvement on the original RACMO2.3. To further investigate this, Fig. 13 shows the annual mean SMB RMSE (model vs. observations) of the 11-km SMB field in RACMO2.3 (red), the downscaled product v0.2 (green) and v1.0 (blue) as a function of observed SMB, binned in 0.5 m w.e. intervals. In the ablation zone (SMB < 0), the SMB RMSE is reduced by 29-65% in v1.0 relative to the 11-km product, owing to the elevation correction in v0.2 (9-23%) and the additional albedo correction (20-42%). In the accumulation zone, the elevation dependence (9%) and the precipitation adjustment (19%) also contribute to reduce the SMB RMSE by 28% in v1.0. The largest RMSE reduction occurs in the lower GrIS ablation zone, where improvements in topography and bare ice albedo in v1.0 are greatest.

Although significantly improved, the downscaled SMB v1.0 is likely to be locally underestimated for four reasons: a) the bare ice albedo correction is evenly applied to both snow covered and bare ice regions experiencing surface melt and runoff, as no relevant proxy, reflecting day-to-day snow coverage, could be derived from RACMO2.3. However, this issue should have a limited effect on the magnitude of downscaled melt and runoff since the albedo correction is most efficient in summer, when the snow cover of low-lying glaciers has likely melted; b) the MODIS ice albedo product at 1-km becomes less accurate at high latitudes, likely suffering from bare soil contamination resulting from mixed reflectance signals recorded in both the tundra and ice covered regions. Note that floating glacier tongues also show too low surface albedo, e.g. Petermann glacier (yellow dot in Fig. 12a)), resulting from mixed signals from adjacent dark melt pond and brighter dry ice. The resulting albedo underestimation over low-lying floating tongues below 200 m leads to overestimated ablation (~ 0.2 mWE/yr; Figs. 7a and b); c) the average 1-km MODIS ice albedo product for 2000-2015 used in the melt correction remains constant in time and might underestimate the bare ice albedo prior to 2000 as the period 2000-2015 encompasses multiple record high

melt years; d) the degradation of MODIS Terra sensors (Polashenski et al., 2015). These limitations underline the high sensitivity of the downscaled product to the input fields used to initialize the downscaling procedure, i.e. RCM version used, the resulting modelled SMB components, bare ice albedo records, ablation measurements, topography and ice mask. The downscaled SMB v1.0 presents an estimated uncertainty of ~ 6 Gt/yr in the GrIS ablation zone, which was estimated by integrating the SMB bias in v1.0 (30 mmWE, Fig. 6c) over the ablation zone of the contiguous ice sheet (~ 202.000 km²).

We anticipate that the new, 1-km Greenland SMB product is especially useful for studies that address the mass balance of Greenland outlet glaciers that are too steep and/or narrow to be properly resolved at the typical horizontal resolution of regional climate models (~ 5 -15 km). Future downscaled products can have even higher resolution (100m) and will be based on further improved RCM output fields of precipitation and melt.

7 Conclusions

The relatively coarse spatial resolution currently used in RCMs remains insufficient to properly resolve small scale variations in elevation and ice cover at the ice sheet margins, significantly affecting the calculation of melt and runoff. In the present study, we statistically downscale individual SMB components from RACMO2.3 at 11-km to a 1-km ice mask and topography derived from the GIMP DEM, using a daily specific elevation dependence. Moreover, runoff and melt are corrected for biases in bare ice albedo in RACMO2.3. Precipitation and snowdrift erosion are bi-linearly interpolated without applying an elevation correction. Total precipitation is also adjusted to compensate for the dry accumulation bias of RACMO2.3 in the ice sheet interior. Downscaled daily SMB is then retrieved for the period 1958-2015 by summing daily downscaled precipitation, runoff, sublimation and drifting snow erosion. An evaluation of the downscaled SMB product against observations, collected both in the ablation and accumulation zones of the GrIS, shows improved agreement. In the ablation zone, the variance explained by the downscaled product v1.0 increased by 31% relative to the original RACMO2.3 11-km output, mainly through better resolved narrow outlet glaciers at the GrIS margins.

Integrated over the GrIS, precipitation increased by 16.6 Gt/yr due to the larger glaciated area in south and east Greenland at 1-km; an additional correction of 56.2 Gt/yr must account for the accumulation bias in the ice sheet interior in RACMO2.3. Likewise, a 13.6 Gt/yr increase in runoff is attributed to elevation corrections on the 1-km topography and another 69.3 Gt/yr extra runoff can be ascribed to underestimated bare ice albedo over narrow outlet glaciers at the GrIS margins. A small area in central east Greenland alone, characterized by multiple narrow glacier tongues poorly resolved at 11-km, accounts for \sim

25% of the total additional runoff.

Acknowledgements. B. Noël, W. J. van de Berg, and M. R. van den Broeke acknowledge support from the Polar Programme of the Netherlands Organization for Scientific Research (NWO/ALW) and the Netherlands Earth System Science Centre (NESSC). I. Howat and the GIMP project are supported by the U.S. National Aeronautics and Space Administration (NASA). H. Machguth acknowledges support from the Programme for Monitoring of the Greenland Ice Sheet (PROMICE), funded by the Danish Energy Agency's (DANCEA) program.

References

- 515 R. C. Bales, J. R. McConnell, E. Mosley-Thompson, and B. Csatho. Accumulation over the Greenland ice sheet from historical and recent records. *Journal of Geophysical Research*, 106(D24): 33813 – 33825, 2001. doi:10.1029/2001JD900153.
- R. C. Bales, Q. Guo, D. Shen, J. R. McConnell, G. Du, J. F. Burkhart, V. B. Spikes, E. Hanna, and J. Cappelen. Annual accumulation for Greenland updated using ice core data developed during
520 2000–2006 and analysis of daily coastal meteorological data. *Journal of Geophysical Research*, 114(D6):D06116, 2009. doi:10.1029/2008JD011208.
- J. L. Bamber, S. Ekholm, and W. B. Krabill. A new, high-resolution digital elevation model of Greenland fully validated with airborne laser altimeter data. *Journal of Geophysical Research*, 106:6733 – 6745, 2001. doi:10.1029/2000JB900365.
- 525 E. W. Burgess, R. R. Forster, J. E. Box, E. Mosley-Thompson, D. H. Bromwich, R. C. Bales, and L. C. Smith. A spatially calibrated model of annual accumulation rate on the Greenland Ice Sheet (1958–2007). *Journal of Geophysical Research*, 115:F02004, 2010. doi:10.1029/2009JF001293.
- D. P. Dee, S. M. Uppala, A. J. Simmons, P. Berrisford, P. Poli, S. Kobayashi, U. Andrae, M. A. Balmaseda, G. Balsamo, P. Bauer, P. Bechtold, A. C. M. Beljaars, L. van de Berg, J. Bidlot, N. Bormann, C. Delsol, R. Dragani, M. Fuentes, A. J. Geer, L. Haimberger, S. B. Healy,
530 H. Hersbach, E. V. Hólm, L. Isaksen, P. Kållberg, M. Köhler, M. Matricardi, A. P. McNally, B. M. Monge-Sanz, J.-J. Morcrette, B.-K. Park, C. Peubey, P. de Rosnay, C. Tavolato, J.-N. Thépaut, and F. Vitart. The ERA-Interim reanalysis: configuration and performance of the data assimilation system. *Quarterly Journal of the Royal Meteorological Society*, 137:553 – 597, 2011. doi:10.1002/qj.828.
- 535 E. M. Enderlin and I. M. Howat. Submarine melt rate estimates for floating termini of Greenland outlet glaciers (2000–2010). *Journal of Glaciology*, 59(213):67 – 75(9), 2013. doi:http://dx.doi.org/10.3189/2013JoG12J049.
- E. M. Enderlin, I. M. Howat, S. Jeong, M.-J. Noh, J. H. van Angelen, and M. R. van den Broeke. An improved mass budget for the Greenland ice sheet. *Geophysical Research Letters*, 43(3):866 – 872, 2014. doi:10.1002/2013GL059010.
- 540 J. Ettema, M. R. van den Broeke, E. van Meijgaard, and W. J. van de Berg. Climate of the Greenland ice sheet using a high-resolution climate model - Part2: Near-surface climate and energy balance. *The Cryosphere*, 4:529 – 544, 2010a. doi:10.5194/tc-4-529-2010.
- 545 J. Ettema, M. R. van den Broeke, E. van Meijgaard, W. J. van de Berg, J. E. Box, and K. Steffen. Climate of the Greenland ice sheet using a high-resolution climate model – Part 1: Evaluation. *The Cryosphere*, 4:511 – 527, 2010b. doi:10.5194/tc-4-511-2010.
- R. S. Fausto, D. van As, J. E. Box, W. Colgan, P. L. Langen, and R. H. Mottram. The implication of nonradiative energy fluxes dominating Greenland ice sheet exceptional ablation area surface
550 melt in 2012. *Geophysical Research Letters*, 43:1944 – 8007, 2016. doi:10.1002/2016GL067720.
- X. Fettweis. Reconstruction of the 1979–2006 Greenland ice sheet surface mass balance using the regional climate model MAR. *The Cryosphere*, 1:21 – 40, 2007. doi:10.5194/tc-1-21-2007.
- X. Fettweis, H. Gallée, F. Lefebvre, and J.-P. van Ypersele. Greenland surface mass balance simulated

- by a regional climate model and comparison with satellite-derived data in 1990-1991. *Climate Dynamics*, 24:623 – 640, 2005. doi:10.1007/s00382-005-0010-y.
- 555 X. Fettweis, M. Tedesco, M. van den Broeke, and J. Ettema. Melting trends over the Greenland ice sheet (1958-2009) from spaceborne microwave data and regional climate models. *The Cryosphere*, 5:359 – 375, 2011. doi:10.5194/tc-5-359-2011.
- B. Franco, X. Fettweis, C. Lang, and M. Erpicum. Impact of spatial resolution on the modelling
560 of the Greenland ice sheet surface mass balance between 1990–2010, using the regional climate model MAR. *The Cryosphere*, 6:695 – 711, 2012. doi:10.5194/tc-6-695-2012.
- E. Hanna, P. Huybrechts, I. Janssens, J. Cappelen, K. Steffen, and A. Stephens. Runoff and mass balance of the Greenland ice sheet: 1958-2003. *Journal of Geophysical Research*, 110:D13108, 2005. doi:10.1029/2004JD005641.
- 565 E. Hanna, P. Huybrechts, K. Steffen, J. Cappelen, R. Huff, C. Shuman, T. Irvine-Fynn, S. Wise, and M. Griffiths. Increased Runoff from Melt from the Greenland Ice Sheet: A Response to Global Warming. *Journal of Climate*, 21:331 – 341, 2008. doi:10.1175/2007JCLI1964.1.
- E. Hanna, P. Huybrechts, J. Cappelen, K. Steffen, R. C. Bales, E. Burgess, J. R. McConnell, J. P. Steffensen, M. Van den Broeke, L. Wake, G. Bigg, M. Griffiths, and D. Savas. Greenland Ice Sheet
570 surface mass balance 1870 to 2010 based on Twentieth Century Reanalysis, and links with global climate forcing. *Journal of Geophysical Research*, 116:D24121, 2011. doi:10.1029/2011JD016387.
- I. M. Howat, A. Negrete, and B. E. Smith. The Greenland Ice Mapping Project (GIMP) land classification and surface elevation data sets. *The Cryosphere*, 8:1509 – 1518, 2014. doi:10.5194/tc-8-1509-2014.
- 575 P. Kuipers Munneke, M. R. van den Broeke, J. T. M. Lenaerts, M. G. Flanner, A. S. Gardner, and W. J. van de Berg. A new albedo parameterization for use in climate models over the Antarctic ice sheet. *Journal of Geophysical Research*, 116:D05114, 2011. doi:10.1029/2010JD015113.
- J. T. M. Lenaerts, M. R. van den Broeke, J. H. Angelen, E. van Meijgaard, and S. J. Déry. Drifting snow climate of the Greenland ice sheet: a study with a regional climate model. *The Cryosphere*,
580 6:891 – 899, 2012. doi:10.5194/tc-6-891-2012.
- P. Lucas-Picher, M. Wulff-Nielsen, J. H. Christensen, Gudfinna Adalgeirsdóttir, and Ruth M. and S. B. Simonsen. Very high resolution regional climate model simulations over Greenland: Identifying added value. *Journal of Geophysical Research*, 117:D02108, 2012. doi:10.1029/2011JD016267.
- H. Machguth, P. Rastner, T. Bolch, N. Mölg, L. Sandberg Sørensen, G. Adalgeirsdottir, J. H. van
585 Angelen, M. R. van den Broeke, and X. Fettweis. The future sea-level rise contribution of Greenland’s glaciers and ice caps. *Environmental Research Letters*, 8(2):025005, 2013. doi:10.1088/1748-9326/8/2/025005.
- H. Machguth, H. Thomsen, A. Weidick, A. P. Ahlstrøm, J. Abermann, M. L. Andersen, S. Andersen, A. A. Bjørk, J. E. Box, R. J. Braithwaite, C. E. Bøggild, M. Citterio, P. Clement, W. Colgan, R. S. Fausto, K. G. S. Gubler, B. Hasholt, B. Hynek, N. Knudsen, S. Larsen, S. Mernild,
590 J. Oerlemans, H. Oerter, O. Olesen, C. Smeets, K. Steffen, M. Stober, S. Sugiyama, D. van As, M. van den Broeke, and R. S. van de Wal. Greenland surface mass balance observations from the ice sheet ablation area and local glaciers. *Journal of Glaciology*, pages 1 – 27, 2016.

doi:10.1017/jog.2016.75.

- 595 B. Noël, W. J. van de Berg, E. van Meijgaard, P. Kuipers Munneke, R. S. W. van deWal, and M.R. van den Broeke. Evaluation of the updated regional climate model RACMO2.3: summer snowfall impact on the Greenland Ice Sheet. *The Cryosphere*, 9:1831 – 1844, 2015. doi:10.5194/tc-9-1831-2015.
- T. B. Overly, R. L. Hawley, V. Helm, E. M. Morris, and R. N. Chaudhary. Greenland annual
600 accumulation along the EGIG line, 1959–2004, from ASIRAS airborne radar and detailed neutron-probedensity measurements. *The Cryosphere Discussion*, 9:6791 – 6828, 2015. doi:10.5194/tcd-9-6791-2015, 2015.
- C. M. Polashenski, J. E. Dibb, M. G. Flanner, J. Y. Chen, Z. R. Courville, A. M. Lai, J. J. Schauer, M. M. Shafer, and M. Bergin. Neither dust nor black carbon causing apparent albedo decline in
605 Greenland’s dry snow zone: Implications for MODIS C5 surface reflectance. *Geophysical Research Letters*, 42(21):9319 – 9327, 2015. doi:10.1002/2015GL065912.
- P. Rastner, T. Bolch, N. Mölg, H. Machguth, R. Le Bris, and F. Paul. The first complete inventory of the local glaciers and ice caps on Greenland. *The Cryosphere*, 6:1483 – 1495, 2012. doi:10.5194/tc-6-1483-2012.
- 610 E. Rignot, J. E. Box, E. Burgess, and E. Hanna. Mass balance of the Greenland ice sheet from 1958 to 2007. *The Cryosphere*, 35:L20502, 2008. doi:10.1029/2008GL035417.
- E. Rignot, I. Velicogna, M. R. van den Broeke, A. Monaghan, and J. Lenaerts. Acceleration of the contribution of the Greenland and Antarctic ice. *Geophysical Research Letters*, 38:L05503/1 – L05503/5, 2011. doi:10.1029/2011GL046583.
- 615 I. Sasgen, M. R. van den Broeke, J. L. Bamber, E. Rignot, L. Sandberg Sørensenf, B. Wouters, Z. Martinec, I. Velicogna, and S. B. Simonsen. Timing and origin of recent regional ice-mass loss in Greenland. *Earth and Planetary Science Letters*, 333 - 334:293 – 303, 2012. doi:10.1016/j.epsl.2012.03.033.
- A. Shepherd, E. R. Ivins, G. A. V. R. Barletta, M. J. Bentley, S. Bettadpur, K. H. Briggs, D. H.
620 Bromwich, R. Forsberg, N. Galin, M. Horwath, S. Jacobs, I. Joughin, M. A. King, J. T. M. Lenaerts, J. Li, S. R. M. Ligtenberg, A. Luckman, S. B. Luthcke, M. McMillan, R. Meister, G. Milne, J. Mouginot, A. Muir, J. P. Nicolas, J. Paden, A. J. Payne, H. Pritchard, E. Rignot, H. Rott, L. Sandberg Sørensen, T. A. Scambos, B. Scheuchl, E. J. O. Schrama, B. Smith, A. V. Sundal, J. H. van Angelen, W. J. van de Berg, M. R. van den Broeke, D. G.
625 Vaughan, I. Velicogna, J. Wahr, P. L. Whitehouse, D. J. Wingham, D. Yi, D. Young, and H. J. Zwally. A Reconciled Estimate of Ice-Sheet Mass Balance. *Science*, 338(6111):1183 – 1189, 2012. doi:10.1126/science.1228102.
- J. D. Stark, Exeter Met Office, C. J. Donlon, M. J. Martin, and M. E. McCulloch. OSTIA: An operational, high resolution, real time, global sea surface temperature analysis system. *OCEANS 2007 - Europe*, pages 1 – 4, 2007. doi:10.1109/OCEANSE.2007.4302251. Conference Publications.
- 630 P. Undèn, L. Rontu, H. Järvinen, P. Lynch, J. Calvo, G. Cats, J. Cuxart, K. Eerola, C. Fortelius, J. A. Garcia-Moya, C. Jones, G. Lenderlink, A. McDonald, R. Mcgrath, B. Navascues, N. W. Nielsen, V. Degaard, E. Rodriguez, M. Rummukainen, K. Sattler, B. H. Sass, H. Savijarvi, B. W.

Schreur, R. Sigg, and H. The. HIRLAM-5. *Scientific Documentation*, 2002. Technical Report.

- 635 S. M. Uppala, P. W. K  llberg, A. J. Simmons, U. Andrae, V. Da Costa Bechtold, M. Fiorino,
J. K. Gibson, J. Haseler, A. Hernandez, G. A. Kelly, X. Li, K. Onogi, S. Saarinen, N. Sokka,
R. P. Allan, E. Andersson, K. Arpe, M. A. Balmaseda, A. C. M. Beljaars, L. Van De Berg,
J. Bidlot, N. Bormann, S. Caires, F. Chevallier, A. Dethof, M. Dragosavac, M. Fisher, M. Fuentes,
S. Hagemann, E. H  lm, B. J. Hoskins, L. Isaksen, P. A. E. M. Janssen, R. Jenne, A. P. McNally,
640 J-F. Mahfouf, J-J. Morcrette, N. A. Rayner, R. W. Saunders, P. Simon, A. Ster, K. E. Trenberth,
A. Untch, D. Vasiljevic, P. Viterbo, and J. Woollen. The ERA-40 re-analysis. *Quarterly Journal
of the Royal Meteorological Society*, 131:2961 – 3012, 2005.
- J. H. Van Angelen, M. R. van den Broeke, B. Wouters, and J. T. M. Lenaerts. Contemporary (1969-
2012) evolution of the climate and surface mass balance of the Greenland ice sheet. *Surveys in*
645 *Geophysics*, 2013. doi:10.1007/s10712-013-9261-z.
- D. Van As, R. S. Fausto, A. P. Ahlstr  m, S. B. Andersen, M. L. Andersen, M. Citterio, K. Edelvang,
P. Gravesen, H. Machguth, F. M. Nick, S. Nielsen, and A. Weidick. Temperature and ablation
records from the Programme for Monitoring of the Greenland Ice Sheet (PROMICE). *Geological*
Survey of Denmark and Greenland Bulletin, 23:73 – 76, 2011. URL [www.geus.dk/publications/](http://www.geus.dk/publications/bull)
650 [bull](http://www.geus.dk/publications/bull).
- M. R. Van den Broeke, P. Smeets, and J. Ettema. Surface layer climate and turbulent exchange in
the ablation zone of the west Greenland ice sheet. *International Journal of Climatology*, 29:2309
– 2323, 2009. doi:10.1002/joc.1815.
- E. Van Meijgaard, L. H. van Uft, W. J. van de Berg, F. C. Bosveld, B. van den Hurk, G. Lenderink,
655 and A. P. Siebesma. *Technical Report 302: The KNMI regional atmospheric climate model
RACMO version 2.1*. Royal Netherlands Meteorological Institute, De Bilt, 2008.
- K. Van Tricht, S. Lhermitte, J. T. M. Lenaerts, I. V. Gorodetskaya, T. S. L’Ecuyer, B. No  l, M. R.
van den Broeke, D. D. Turner, and N. P. M. van Lipzig. Clouds enhance Greenland ice sheet
meltwater runoff. *Nature communications*, 7(10266), 2016. doi:10.1038/ncomms10266.
- 660 J. M. Van Wessem, C. H. Reijmer, J. T. M. Lenaerts, W. J. van de Berg, M. R. van den Broeke, and
E. van Meijgaard. Updated cloud physics improve the modelled near-surface climate of Antarctica
of a regional atmospheric climate model. *The Cryosphere*, 8:125 – 135, 2014. doi:10.5194/tc-8-
125-2014.
- U. Weiser, M. Olefs, W. Sch  ner, G. Weyss, and B. Hynek. Correction of broadband snow albedo
665 measurements affected by unknown slope and sensor tilts. *The Cryosphere*, 10:775 – 790, 2016.
doi:10.5194/tc-10-775-2016.




Fig. 1. Annual mean SMB modelled by RACMO2.3 at 11-km over the GrIS and surrounding ice caps for the period 1958-2015. This figure also depicts the location of 213 ablation measuring sites (yellow dots) and 182 accumulation sites (white dots) used for downscaled SMB evaluation as well as the four GrIS marginal regions (blue boxes), discussed in Section 5. Letters refer to the different transects shown in Fig.7.




Fig. 2. Elevation and ice mask (yellow) as prescribed in RACMO2.3 at 11-km (left) and derived from the GIMP DEM down-sampled to 1-km (right) over central east Greenland (blue box 1 in Fig. 1).

1958-2015	Regions	Centre east			Centre west			South			North			GrIS		
Resolution	Unit	11km	1km	Δ	11km	1km	Δ	11km	1km	Δ	11km	1km	Δ	11km	1km	Δ
SMB $v_{0.2}$	<i>Gt/yr</i>	5.0	-0.3	-5.3	-4.6	-5.4	-0.8	44.3	47.6	3.3	-2.6	-0.3	2.3	349.3	351.3	2.0
Runoff $v_{0.2}$	<i>Gt/yr</i>	16.1	23.8	7.7	18.3	19.2	0.9	42.4	44.6	2.2	8.9	5.8	-3.1	284.1	297.7	13.6
Precip $v_{0.2}$	<i>Gt/yr</i>	22.6	25.2	2.6	15.0	15.2	0.2	91.4	97.2	5.8	6.9	6.1	-0.8	675.4	692.0	16.6
SMB $v_{1.0}$	<i>Gt/yr</i>	5.0	-11.6	-16.6	-4.6	-6.7	-2.1	44.3	37.3	-7.0	-2.6	-1.3	1.3	349.3	338.2	-11.1
Runoff $v_{1.0}$	<i>Gt/yr</i>	16.1	36.7	20.6	18.3	21.1	2.8	42.4	57.5	15.1	8.9	7.7	-1.2	284.1	367.0	82.9
Precip $v_{1.0}$	<i>Gt/yr</i>	22.6	26.8	4.2	15.0	15.8	0.8	91.4	99.8	8.4	6.9	7.0	0.1	675.4	748.2	72.8
Sublimation	<i>Gt/yr</i>	2.1	2.1	0.0	1.6	1.6	0.0	4.4	4.7	0.3	0.8	0.7	-0.1	41.3	41.9	0.6
Snow drift	<i>Gt/yr</i>	-0.5	-0.4	0.1	-0.3	-0.2	0.1	0.2	0.3	0.1	-0.1	-0.1	0.0	0.7	1.1	0.4
Ice area	10^4 km^2	5.9	6.0	0.1	2.7	2.7	-0.02	7.7	8.2	0.5	3.5	3.1	-0.4	170.3	169.4	-0.9

Table 1. Table listing (top) the annual mean integrated SMB components (Gt/year) covering the period 1958-2015 over four different regions, centre east ($69.6^\circ\text{N} - 74.3^\circ\text{N}$; $21^\circ\text{W} - 31^\circ\text{W}$; blue box 1 in Fig. 1), centre west ($69.3^\circ\text{N} - 72.5^\circ\text{N}$; $49^\circ\text{W} - 57^\circ\text{W}$; blue box 2), south ($59.5^\circ\text{N} - 63.3^\circ\text{N}$; $41^\circ\text{W} - 51^\circ\text{W}$; blue box 3) and north ($80.5^\circ\text{N} - 83^\circ\text{N}$; $42^\circ\text{W} - 62^\circ\text{W}$; blue box 4), and for the entire GrIS at both resolutions as well as the difference between 1-km and 11-km; (bottom) same for the ice-covered area (km^2).




Fig. 3. Correlation to elevation of annual mean a) total precipitation (solid and liquid), b) runoff, c) SMB, d) sublimation, e) melt and f) drifting snow erosion modelled by RACMO2.3 and calculated on the 11-km grid for the period 1958-2015.




Fig. 4. (a) Elevation dependent downscaling procedure: local estimate of a daily SMB components regression to elevation on the RACMO2.3 grid at 11-km. (b) Scheme of a tilted plane as described in the GIMP DEM at 1-km.







Fig. 5. Summer 2011 time series of daily, ice sheet integrated runoff difference (Gt/day) between the downscaled product at 1-km, using a minimum threshold of 3 to 9 regression points (legend), and the RACMO2.3 model at 11-km.



(a) Modelled SMB at 11 km



(b) Downscaled SMB at 1 km version v0.2



(c) Corrected SMB at 1 km version v1.0

Fig. 6. Comparison of SMB measurements collected at 213 sites with (a) modelled SMB from RACMO2.3 at 11-km; (b) downscaled SMB at 1-km (v0.2) and (c) corrected downscaled SMB at 1-km (v1.0). The red stars correspond to PROMICE station QAS_L located in southern Greenland (61.03°N, 46.85°W, 310 m.a.s.l). The red dashed line represents the regression including all measurements using a perpendicular fit.




Fig. 7. Annual mean observed (red dots) and downscaled (blue dots, v1.0) SMB for 8 selected transects in the GrIS ablation zone (mWE/yr). Name and locations of these transects (Fig. 1) are listed at the top of each graph. Graphs also list the number of sites used for each transect, linear SMB-to-elevation regression retrieved from observations and downscaled (v1.0) data in mmWE/yr per m, RMSE and mean bias.




Fig. 8. Comparison of accumulation observations collected at 182 sites with modelled SMB from RACMO2.3 at 11-km (red) and downscaled SMB v0.2 at 1-km (blue) in mWE/yr. Note that bias correction has not yet been applied.




Fig. 9. Centre east: a) Ice sheet mask in RACMO2.3 at 11-km (red) and in the down-sampled GIMP DEM at 1-km (orange) (blue box 1 in Fig. 1), and the mask of disconnected glaciers and ice caps at 1-km (blue); average (1958-2015) annual mean b) total precipitation, c) runoff and d) SMB (mmWE/yr) modelled by RACMO2.3 at 11km; e) elevation bias (m) between 1-km and 11-km resolutions. Figures f), g), h) represent annual mean total precipitation, runoff and SMB downscaled to 1-km using elevation dependence only (v0.2). Figure i) shows the bare ice albedo bias between MODIS measurements at 1-km (2000-2015) and RACMO2.3 at 11-km (2001-2010). Figures j), k) and l) are similar to f), g) and h) but incorporate the bare ice albedo and precipitation corrections (v1.0).




Fig. 10. Centre west: same as Fig. 9 but for central west Greenland (blue box 2 in Fig. 1).




Fig. 11. South: same as Fig. 9 but for south Greenland (blue box 3 in Fig. 1). The yellow dot in a) locates station QAS_L.




Fig. 12. North: same as Fig. 9 but for north Greenland (blue box 4 in Fig. 1). The green line in a) shows the grounded ice mask at 1-km. The yellow dot in a) locates the Petermann glacier site settled on a floating ice tongue.




Fig. 13. Annual mean model SMB RMSE (model vs. observations) of the 11-km SMB field in RACMO2.3 (red dots), the downscaled SMB dataset v0.2 (green dots) and v1.0 (blue dots) as a function of observed SMB (395 observations). Modelled SMB is grouped in 0.5 mWE/yr bins except for the first bin, which ranges from -6.00 to -3.75 mWE/yr. Numbers indicate the amount of observations used in each bin.