



Brief communication: Organochlorine pesticides in an archived firn core from Law Dome, East Antarctica

5 Marie Bigot*¹, Mark A.J. Curran^{2,3}, Andrew D. Moy^{2,3}, Derek C.G. Muir⁴, Darryl W. Hawker⁵, Roger
Cropp⁵, Camilla F. Teixeira⁴ and Susan Bengtson Nash¹

¹ Environmental Futures Research Institute, Griffith University, 170 Kessels Rd, Nathan, QLD 4111, Australia

² Australian Antarctic Division, 203 Channel Highway, Kingston, TAS 7050, Australia

10 ³ Antarctic Climate and Ecosystems Cooperative Research Centre, University of Tasmania, Private Bag 80, Hobart TAS
7001, Australia.

⁴ Aquatic Contaminants Research Division, Environment Canada, 867 Lakeshore Rd, Burlington ON, L7R 4A6 Canada

⁵ School of Environment, Griffith University, 170 Kessels Rd, Nathan, QLD 4111, Australia

Correspondence to: Marie Bigot (marie.bigot@griffithuni.edu.au)

Abstract. Organochlorine pesticides (OCPs) were, for the first time, quantified in archived firn cores from the East Antarctic
15 ice sheet representative of 1945-1957 C.E. and 1958-1967 C.E. The core sections were melted under high purity nitrogen
atmosphere and the melt water analysed. Methods allowed quantification of hexachlorocyclohexanes, heptachlor, *trans*-
chlordane, dieldrin and endrin. While the core presented evidence of nominal modern-use chemical contamination,
indicating handling and/or storage contamination, legacy OCP concentrations and deposition rates reported are orders of
20 magnitude lower than those from Arctic regions, lending support for their validity. The study further provides a description
of equipment used and suggests adapted methods to overcome logistical challenges associated with trace organic
contaminant detection in Polar Regions.

1 Introduction

Persistent Organic Pollutants (POPs) are ubiquitous, toxic, anthropogenic substances that have been widely used in
agriculture and manufacturing industries since the 1930s. A variety of organochlorine pesticides (OCPs) have been listed as
25 POPs under the Stockholm Convention, adopted by the United Nations Environment Program to ban, or severely restrict, the
use of these chemicals and hereby minimise the environmental and human health hazards that they pose. By definition, POPs
are resistant to environmental degradation processes, allowing them to undergo long-range environmental transport to
remote Polar Regions where cold conditions serve to restrict further movement.

30 Unlike seasonal sea-ice that persists for only limited periods of time, ice sheets are deposited over thousands of years. This
makes them a rich archive for the study of past atmospheric composition and climate variations. Levels of inorganic
chemical markers such as sodium, potassium, chloride, nitrate or sulphate have previously been successfully determined in
glaciology studies (Legrand et al., 1984). Glacial ice cores also hold temporal information regarding historical organic



pollution events, however their application for this purpose is far less prevalent in the documented literature. Inherent incompatibilities exist between traditional glaciology research methods and the requirements for robust POP quantification in this environmental compartment. Nevertheless, the work by Wang et al. (2008), for example, showed the potential of POPs in this regard by finding a strong correlation between historical usage of POPs in India and POP levels in ice cores of
5 corresponding deposition ages from Mt Everest. To our knowledge, only one study has determined OCP concentrations in glacial ice/firn from the Arctic (Isaksson et al., 2003) while none, to date, exist from the Antarctic. Other studies in Polar Regions have focused on other chemicals of interest (i.e. brominated flame retardants, polycyclic aromatic hydrocarbons, current use pesticides and polychlorinated biphenyls) and/or other frozen matrix (i.e. snow) (e.g. Fuoco et al., 2012;Gregor et al., 1995;Hermanson et al., 2010;Kang et al., 2012;Zhang et al., 2013;Peters et al., 1995).

10

In the current global climate context, accelerated melting and subsequent retreat of Antarctic ice shelves are frequently reported. As the ice compartment represents a reservoir of historically deposited POPs, quantification of concentrations in Antarctic continental ice is particularly relevant in order to predict the possible future POP re-emission into the atmosphere and oceans through ice-melt. This secondary input process has previously been demonstrated for melting alpine glaciers in
15 Switzerland (Bogdal et al., 2009). Under this scenario, global efforts to moderate polar ecosystems' exposure to these potentially hazardous compounds would be compromised.

20

This work investigates the potential for a well-studied archived Antarctic firn core to elucidate historical deposition rates of OCPs. It was designed as a proof of concept to examine the implications of mismatched sampling priorities between
classical glaciology programs and POP research, representing an important step for the assessment of the value of firn core archives world-wide for the progression of Polar contaminants research. Further, the work addresses some of the challenges associated with the sampling of this matrix for the trace levels of POPs generally observed in Antarctica and describes purpose built equipment for overcoming some of these challenges.

2 Materials and methods

25

2.1 Firn core sourcing

For this study, we accessed the historical firn core from a long-established glaciology study site located in East Antarctica: Dome Summit South (DSS), Law Dome. The 1196 m DSS ice core was collected in 1988-1993 and the entire records spans approximately 80 kyr B.P. (van Ommen et al., 2004). The DSS ice core chronology is derived from a combination of direct layer counting (van Ommen et al., 2004;Plummer et al., 2012;Roberts et al., 2015) and age ties to other records (van Ommen
30 et al., 2004). This ice core has provided a wealth of atmospheric circulation, ambient temperature, snow accumulation, and climate proxy records for climate reconstructions (Souney et al., 2002;Plummer et al., 2012;Roberts et al., 2015). Coring methods employed are presented elsewhere (Morgan et al., 1997). Following collection, the core was cut to one meter



lengths on site, stored in double plastic bags at -18°C , and archived in large cold storage facilities in Tasmania, Australia, until analysis.

There is an inherent mismatch between standard glaciology and POP sampling procedures, particularly in terms of the volumes of matrix required for robust analysis, the type of materials used during sampling and long term storage requirements. Nevertheless, given the massive logistical investment to sample ice in remote Polar environments, and the rich source of ancillary information accompanying the well-studied DSS core, access to the remaining firn core represented an opportunity to examine the implications of method incompatibilities and assess whether useful information could be derived from this, and therefore similar valuable archives, world-wide.

Sample volumes recently used for determination of OCPs in other Antarctic frozen matrices such as continental surface snow (Kang et al., 2012) and sea-ice (Dickhut et al., 2005) were 500 mL and 95.5 to 132 L, respectively. In the Arctic, Isaksson et al. (2003) used a total of 48 metres of glacial ice/firn core (of unknown diameter) separated into nine individual samples. No other previous study on OCPs has been published for continental firn cores in Antarctica, hence the chemical burden and consequent volume of firn required for confident detection was largely unknown. We therefore adopted a conservative approach by utilising just two large volume samples from a known period as opposed to a higher number of samples of smaller volume. The DSS firn core was selected as it yielded large volumes for the relevant deposition time periods of interest, enabling guidance on chemical deposition rates for future investigations. It was separated into batches of approximately 10 metre length each, representing two early OCP deposition periods: 1945-1957 C.E. (35.4 to 45.6 m depth, sample A) and 1958-1967 C.E. (26.3 to 35.4 m depth, sample B). Density of these firn sections had previously been estimated between 0.63 to $0.74\text{ g}\cdot\text{cm}^{-3}$ (for core depth between 21.5 to 47.6 m) (Morgan et al., 1997) and dating had been completed by direct layer counting.

2.2 Sampling procedures

Firn core samples were minimally exposed to laboratory ambient air, and whole portions were melted together in a clean stainless steel unit, purpose designed and built for POP analysis (Figure 1). Melted sample volumes were 144 and 133 L for sample A and B, respectively. The ice-melting unit, derived from a design by Gustafsson et al. (2005), allowed complete isolation of the sample and melting under high purity nitrogen atmosphere (Figure 1). The unit was fitted with a drain connected to a modified Kiel in-situ Pump (KISP, Aimes GmbH, Germany), previously used for the purpose of trace organics sampling in water (Petrick et al., 1996). This set-up was designed to provide a means of *in-situ* sample extraction for field based snow, ice and water campaigns. The ice-melting unit is fully transportable for direct sample loading and sealing in the field. The unit can then be brought to the operational Polar research station for access to nitrogen gas, water and electrical power to complete sample extraction.



Melt-water was pumped directly from the melting unit through a pre-furnaced (12h at 450°C) glass fiber filter (GFF, 142 mm, 0.7 µm, Whatman, England) and a stainless steel cartridge filled with approximately 150 mL of pre-cleaned Amberlite XAD-2 resin (Supelpak™-2, Sigma-Aldrich Co. LLC.) in series at a flow rate of 150 mL·min⁻¹. Full sampling material preparation methods and analytical procedures for filters and XAD have been described elsewhere (Bigot et al., 2016).
5 Sample extracts and blanks were analysed by Australian Laboratory Services (ALS Global, Burlington, ON, Canada) for OCPs using gas chromatography high resolution mass spectrometry (GC-HRMS) following standard protocols based on EPA method 1699.

2.3 Quality Assurance / Quality Control

The samples were handled in a clean room (positively pressured, high efficiency particulate and carbon filtered air) during
10 extraction at Environment Canada (Canadian Centre for Inland Waters (CCIW), Burlington, ON, Canada). Three blanks (XAD and filters) were treated in the same manner as the samples: exposed to ambient air during changing of sampling materials in the laboratory, transported together, and extracted and analysed in parallel with samples. All sample values reported are blank corrected using the average of three blanks, providing a means of correcting for possible contamination during laboratory, storage and/or transport of the samples.

15 True blanks (i.e. similar volume of archive samples representative of a deposition period pre-OCP production) were not available. In order to evaluate possible contamination, modern usage POPs, namely polybrominated diphenyl ethers (PBDEs), were used as markers of contamination resulting from the sampling, processing and/or storage of firn cores. PBDEs are commercial flame retarding compounds widely produced since the 1970's. Their concentrations were determined
20 in the final extracts using low resolution GC-MS at CCIW/Environment Canada.

Method detection limits (MDLs) were calculated for each compound as three times the standard deviation of the laboratory blanks. Recovery rates were evaluated via spiking sample materials with 200 µL of a solution containing δ-HCH and ¹³C₈ Mirex (10 pg·µL⁻¹ each), and BDE-71 (9.6 pg·µL⁻¹). XAD cartridges were spiked prior to sampling and filters were spiked
25 prior to extraction. Labelled target analytes were also added by ALS Global and responses were used to correct for losses during clean-up and instrument drift. A list of recoveries for all surrogate standards is available in the supplementary information. Results originally expressed in pg·L⁻¹ were corrected to estimate the mean deposition per year (pg·cm⁻²·yr⁻¹) using the basal area of the core as deposition surface, thus allowing comparison of the two time periods studied.

3 Results and discussion

30 The relatively large volumes of firn analysed in this study enabled the quantification of the legacy OCPs α- and γ-HCH, heptachlor, trans-chlordane, dieldrin, and endrin in the dissolved fraction of the melt water of both samples (Figure 2). These



compounds are likely to be deposited on snow after long range atmospheric transport from source regions within the Southern Hemisphere. Their presence have recently been reported in seawater and/or air close to the coast of East Antarctica (Bigot et al., 2016). Both samples, however, also contained traces of our storage and handling contamination markers, PBDEs, indicating that sampling and/or storage conditions of the archived firn cores introduced organic contamination.

5 While no direct link can be established between PBDE levels introduced and possible contamination of the firn core by legacy OCPs, the presence of the PBDE markers necessitate caution in the interpretation of results. Notwithstanding this, the firn deposition rates were below $0.8 \text{ pg}\cdot\text{cm}^{-2}\cdot\text{yr}^{-1}$ for α - and γ -HCH, heptachlor, trans-chlordane and endrin, while dieldrin had deposition rates of up to $4 \text{ pg}\cdot\text{cm}^{-2}\cdot\text{yr}^{-1}$. Previous studies reported data in melt-water concentrations rather than deposition rates. The deposition rates that we report here are derived from melt-water concentrations of < 60.1 and $310 \text{ pg}\cdot\text{L}^{-1}$,

10 respectively, which are 10 to 1000 fold lower than corresponding OCP levels reported for glacial ice/firn cores from the Arctic (Hermanson et al., 2010; Isaksson et al., 2003). Data reported from other environmental matrices, i.e. seawater and air, show a 10 to 100 fold decrease between Arctic and Antarctic concentrations of HCHs, thus lending support for the validity of the OCP results obtained.

15 The high sample volumes required for this analysis limit the temporal resolution of results and therefore the detail with which we are able to investigate accumulation history of OCPs. The samples represent firn from Law Dome over consecutive 13 and 10-year periods, respectively. Observed differences between the two time periods may reflect different usage patterns but may also be a function of the physicochemical properties of individual pesticides. Given OCPs are mainly transported to Polar Regions via atmospheric transport (Wania and Mackay, 1996), the Law Dome area would have been

20 impacted by air originating from the Southern Hemisphere. Access to detailed historical usage of OCPs in this part of the world is limited, however, most legacy OCPs were first used in the 1940's or early 1950's, e.g. in Australia and Africa (Harrison, 1997; Wandiga, 2001). The concentrations reported represent deposition of OCPs when their usage was at or near their maximum, before the first controls on the uses of these compounds were implemented in the 1970's. As a result, it gives an insight on possible maximum concentrations in Antarctic firn that may remobilise from melt water. Dieldrin and γ -

25 HCH show a distinct increase in their concentrations in sample B, spanning the period (1958 to 1967 C.E.), compared to those in Sample A (1957 – 1945 C.E.) (Figure 2). This could reflect an increase in Southern Hemisphere usage of these two compounds between 1945 and 1967 C.E. Alternatively, the compounds may have started to deposit later than 1945, in which case sample A's results may be affected by dilution of the deposited burden by uncontaminated sections of the core.

30 HCHs are the most commonly studied OCPs in the Antarctic marine abiotic environment. The relative contributions of each isomer ($\gamma > \alpha$) observed in the continental firn samples from Law Dome are consistent with reported Antarctic sea-ice and atmospheric profiles (Dickhut et al., 2005). All OCPs detected in this study have also been recently reported in atmospheric and oceanic samples in the same sector of Antarctica (Bigot et al., 2016). Some of these OCPs, heptachlor and dieldrin in



particular, are currently found in Australian air samples at concentrations amongst the highest reported globally (Wang et al., 2015).

5 Dieldrin was detected at the highest levels in both dissolved and particulate fractions of the melted firn samples analysed in the present study. It is a versatile cyclodiene insecticide that has been largely used around the world for treatment of soil for plantations, as well as for control of various pests. Its usage was restricted in the 1970s, but its use was banned only in 2004, under the Stockholm Convention. In the dissolved fraction, we report an average concentration of dieldrin of 222 and 310 $\text{pg}\cdot\text{L}^{-1}$ in sample A and B, respectively. The dissolved fraction accumulated in firn cores may, if released due to rapid continental ice melting, be available for uptake by marine organisms, and present a toxicological threat to the Antarctic food web. Dieldrin has only rarely been reported in Antarctica's marine environment (Bengtson Nash et al., 2008), yet it is currently one of the most prevalent OCPs found in this region (Bigot et al., 2016). Our findings indicate that ice-melt is likely to result in particularly high dieldrin enrichment compared to other OCPs found in the firn samples. Interestingly, it is also the only compound detected on particles contained in melt-water of the archived ice core at 3.81 and 7.05 $\text{pg}\cdot\text{L}^{-1}$ for sample A and B, respectively, which suggests it may also be transported on atmospheric aerosols that could originate from wind eroded soils for example. Dieldrin was the only OCP compound detected on aerosols sampled in adjacent areas of the Southern Ocean (Bigot et al., 2016).

This proof of concept study determined OCP concentrations that reached Antarctica's ice shelf during their early usage. Melted water volumes of 144 and 133 L for each sample allowed quantification of legacy OCPs that have previously been reported in the vicinity of Antarctica, however results should be considered as maximum possible values given possible contamination introduced during sampling and/or storage. Results for other OCPs were below MDLs, although higher volumes may facilitate quantification. It is important to note that the concentrations reported in this study may be representative of what is expected to be the highest deposition period and therefore volumes employed may be insufficient to detect lower concentrations in more recent parts of firn cores.

25 **4 Conclusion**

Results show that legacy OCPs could have accumulated in the East Antarctic ice shelf from as early as 70 years ago. Although East Antarctica currently exhibits a slight positive mass balance of its ice shelf as opposed to the rest of the continent (Sasgen et al., 2013), driven by climate conditions in the Pacific (Meehl et al., 2016), the potential for OCPs to remobilise into the atmosphere and oceans raises the need for development of POP research in the cryosphere. While we found evidence of nominal PBDE contamination throughout the core, OCP concentrations and deposition rates reported are orders of magnitude lower than those from Arctic regions. This appears realistic and therefore suggests that our results can be taken as first indications of OCP contamination at Law Dome, East Antarctica, through this time period.



Standard glaciology sampling procedures for inorganic analytes (e.g. metals), climate tracers and those for POPs are not necessarily compatible, particularly in terms of the volumes required for analysis, and methods used for collection, sampling and storage. Given the potential value of POP quantification in the Polar cryosphere, we advocate pro-active sampling in collaboration with glaciology programs to ensure sufficient volumes of dedicated samples according to POP work QA/QC procedures. The ice-melting unit used for the present work was designed to be transportable and offer a means of in-situ melting and pumping of large volumes of ice. It provides a cost-effective solution to the logistical challenges of transporting these volumes of ice and also reduces risks for contamination during storage and transport. These in-situ melting techniques have previously been successful in the Arctic (Gustafsson et al., 2005; Isaksson et al., 2003) and should be implemented when dealing with trace POP analysis in Antarctica. In lieu of such support, however, glaciology archives may provide useful indications for remote locations from which information is still lacking.

Acknowledgements. This study was funded by Australian Research Council Discovery project DP140100018 and the Australian Antarctic Division (AAD) provided funding and the DSS firn core samples (AAS project # 4061). Marie Bigot acknowledges a PhD scholarship from Griffith University. The authors would like to thank the field teams that collected samples in 1988 - 1993, as well as Meredith Nation, Lauren Wise, Debbie Lang and all personnel that provided assistance at the Australian Antarctic Division. We also thank Whitney Davis and Ron MacLeod at ALS Global for support provided.

References

- Bengtson Nash, S., Poulsen, A., Kawaguchi, S., Vetter, W., and Schlabach, M.: Persistent organohalogen contaminant burdens in Antarctic krill (*Euphausia superba*) from the eastern Antarctic sector: A baseline study, *Sci. Total Environ.*, 407, 304-314, doi: 10.1016/j.scitotenv.2008.08.034, 2008.
- Bigot, M., Muir, D., Hawker, D., Cropp, R., Dachs, J., Teixeira, C., and Bengtson Nash, S.: Air-seawater exchange of organochlorine pesticides in the Southern Ocean between Australia and Antarctica, *Environ. Sci. Technol.*, doi: 10.1021/acs.est.6b01970, 2016.
- Bogdal, C., Schmid, P., Zennegg, M., Anselmetti, F. S., Scheringer, M., and Hungerbühler, K.: Blast from the past: melting glaciers as a relevant source for persistent organic pollutants, *Environ. Sci. Technol.*, 43, 8173-8177, doi: 10.1021/es901628x, 2009.
- Dickhut, R. M., Cincinelli, A., Cochran, M., and Ducklow, H. W.: Atmospheric concentrations and air-water flux of organochlorine pesticides along the western Antarctic Peninsula, *Environ. Sci. Technol.*, 39, 465-470, doi: 10.1021/es048648p, 2005.
- Fuoco, R., Giannarelli, S., Onor, M., Ghimenti, S., Abete, C., Termine, M., and Francesconi, S.: A snow/firn four-century record of polycyclic aromatic hydrocarbons (PAHs) and polychlorobiphenyls (PCBs) at Talos Dome (Antarctica), *Microchem. J.*, 105, 133-141, doi: 10.1021/es048648p, 2012.
- Gregor, D., Peters, A., Teixeira, C., Jones, N., and Spencer, C.: The historical residue trend of PCBs in the Agassiz ice cap, Ellesmere Island, Canada, *Sci. Total Environ.*, 160, 117-126, doi: 10.1016/0048-9697(95)04349-6, 1995.
- Gustafsson, Ö., Andersson, P., Axelman, J., Bucheli, T. D., Kömp, P., McLachlan, M. S., Sobek, A., and Thörngren, J. O.: Observations of the PCB distribution within and in-between ice, snow, ice-rafted debris, ice-interstitial water, and seawater




- in the Barents Sea marginal ice zone and the North Pole area, *Sci. Total Environ.*, 342, 261-279, doi: 10.1016/j.scitotenv.2004.12.044, 2005.
- Harrison, S.: Organochlorines in Australia, Subregional Awareness Raising Workshop on Persistent Organic Pollutants (POPs), Bangkok, Thailand, 1997,
- 5 Hermanson, M. H., Isaksson, E., Forsström, S., Teixeira, C., Muir, D. C., Pohjola, V. A., and van de Wal, R. S.: Deposition history of brominated flame retardant compounds in an ice core from Holtedahlfonna, Svalbard, Norway, *Environ. Sci. Technol.*, 44, 7405-7410, doi: 10.1021/es1016608, 2010.
- Isaksson, E., Hermanson, M., Hicks, S., Igarashi, M., Kamiyama, K., Moore, J., Motoyama, H., Muir, D., Pohjola, V., and Vaikmäe, R.: Ice cores from Svalbard—useful archives of past climate and pollution history, *Phys. Chem. Earth Pt A/B/C*,
- 10 28, 1217-1228, doi: 10.1016/j.pce.2003.08.053, 2003.
- Kang, J.-H., Son, M.-H., Do Hur, S., Hong, S., Motoyama, H., Fukui, K., and Chang, Y.-S.: Deposition of organochlorine pesticides into the surface snow of East Antarctica, *Sci. Total Environ.*, 433, 290-295, doi: 10.1016/j.scitotenv.2012.06.037, 2012.
- Legrand, M., De Angelis, M., and Delmas, R.: Ion chromatographic determination of common ions at ultratrace levels in Antarctic snow and ice, *Anal. Chim. Acta*, 156, 181-192, doi: 10.1016/S0003-2670(00)85549-X, 1984.
- 15 Meehl, G. A., Arblaster, J. M., Bitz, C. M., Chung, C. T., and Teng, H.: Antarctic sea-ice expansion between 2000 and 2014 driven by tropical Pacific decadal climate variability, *Nature Geoscience*, doi: 10.1038/ngeo2751, 2016.
- Morgan, V., Wookey, C., Li, J., Van Ommen, T., Skinner, W., and Fitzpatrick, M.: Site information and initial results from deep ice drilling on Law Dome, Antarctica, *J. Glaciol.*, 43, 3-10, doi: 10.3198/1997JoG43-143-3-10, 1997.
- 20 Peters, A. J., Gregor, D. J., Teixeira, C. F., Jones, N. P., and Spencer, C.: The recent depositional trend of polycyclic aromatic hydrocarbons and elemental carbon to the Agassiz Ice Cap, Ellesmere Island, Canada, *Sci. Total Environ.*, 160, 167-179, doi: 10.1016/0048-9697(95)04354-4, 1995.
- Petrick, G., Schulz-Bull, D. E., Martens, V., Scholz, K., and Duinker, J. C.: An in-situ filtration/extraction system for the recovery of trace organics in solution and on particles tested in deep ocean water, *Mar. Chem.*, 54, 97-105, doi: 25 10.1016/0304-4203(96)00029-1, 1996.
- Plummer, C. T., Curran, M. A. J., van Ommen, T. D., Rasmussen, S. O., Moy, A. D., Vance, T. R., Clausen, H. B., Vinther, B. M., and Mayewski, P. A.: An independently dated 2000-yr volcanic record from Law Dome, East Antarctica, including a new perspective on the dating of the 1450s CE eruption of Kuwae, Vanuatu, *Climate of the Past*, 8, 1929-1940, doi: 10.5194/cp-8-1929-2012, 2012.
- 30 Roberts, J., Plummer, C., Vance, T., van Ommen, T., Moy, A., Poynter, S., Treverrow, A., Curran, M., and George, S.: A 2000-year annual record of snow accumulation rates for Law Dome, East Antarctica, *Climate of the Past*, 11, 697-707, doi: 10.5194/cp-11-697-2015, 2015.
- Sasgen, I., Konrad, H., Ivins, E., Van den Broeke, M., Bamber, J., Martinec, Z., and Klemann, V.: Antarctic ice-mass balance 2003 to 2012: regional reanalysis of GRACE satellite gravimetry measurements with improved estimate of glacial-isostatic adjustment based on GPS uplift rates, *The Cryosphere*, 7, 1499-1512, doi: 10.5194/tc-7-1499-2013, 2013.
- 35 Souney, J. M., Mayewski, P. A., Goodwin, I. D., Meeker, L. D., Morgan, V., Curran, M. A., van Ommen, T. D., and Palmer, A. S.: A 700-year record of atmospheric circulation developed from the Law Dome ice core, East Antarctica, *J. Geophys. Res. Atmos.*, 107, ACL 1-1-ACL 1-9, doi: 10.1029/2002JD002104, 2002.
- van Ommen, T. D., Morgan, V., and Curran, M. A.: Deglacial and holocene changes in accumulation at Law Dome, East Antarctica, *Annals of Glaciology*, 39, 359-365, doi: 10.3189/172756404781814221, 2004.
- 40 Wandiga, S. O.: Use and distribution of organochlorine pesticides. The future in Africa, *Pure Appl. Chem*, 73, 1147-1155, 2001.
- Wang, X., Kennedy, K., Powell, J., Keywood, M., Gillett, R., Thai, P., Bridgen, P., Broomhall, S., Paxman, C., Wania, F., and Mueller, J. F.: Spatial distribution of selected persistent organic pollutants (POPs) in Australia's atmosphere, *Env. Sci. Process. Impact.*, 17, 525-532, doi: 10.1039/C4EM00594E, 2015.
- 45 Wang, X. p., Xu, B. q., Kang, S. c., Cong, Z. y., and Yao, T. d.: The historical residue trends of DDT, hexachlorocyclohexanes and polycyclic aromatic hydrocarbons in an ice core from Mt. Everest, central Himalayas, China, *Atmos. Environ.*, 42, 6699-6709, doi: 10.1016/j.atmosenv.2008.04.035, 2008.
- Wania, F., and Mackay, D.: Tracking the distribution of persistent organic pollutants, *Environ. Sci. Technol.*, 30, 390-396, 50 doi: 10.1021/es962399q, 1996.



Zhang, X., Meyer, T., Muir, D. C., Teixeira, C., Wang, X., and Wania, F.: Atmospheric deposition of current use pesticides in the Arctic: Snow core records from the Devon Island Ice Cap, Nunavut, Canada, *Env. Sci. Process. Impact.*, 15, 2304-2311, doi: 10.1039/C3EM00433C, 2013.

5



10

Figure 1: Cross section diagram of the ice-melting unit used for this work




Figure 2: Deposition rates of 6 legacy OCPs at the Law Dome site in Antarctica between 1945 and 1957 (Sample A) and 1958 to 1967 (Sample B). Note: results should be considered as maximum possible values.