

Spatiotemporal Variability of Snow Depth across the Eurasian Continent from 1966 to 2012

Xinyue Zhong^{1, 3}, Tingjun Zhang², Shichang Kang^{3, 4}, Kang Wang⁵, Lei Zheng⁶,
Yuantao Hu², Huijuan Wang²

¹ Key Laboratory of Remote Sensing of Gansu Province, Cold and Arid Regions Environmental and Engineering Research Institute, Chinese Academy of Sciences (CAS), Lanzhou 730000, China

² Key Laboratory of Western China's Environmental Systems (Ministry of Education), College of Earth and Environmental Sciences, Lanzhou University, Lanzhou 730000, China

³ State Key Laboratory of Cryospheric Science, Cold and Arid Regions Environmental and Engineering Research Institute, CAS, Lanzhou 730000, China


⁴ CAS Center for Excellence in Tibetan Plateau Earth Sciences, Beijing 100101, China

⁵ Institute of Arctic and Alpine Research, University of Colorado Boulder, Boulder, Colorado, 80309, USA

⁶ Chinese Antarctic Center of Surveying and Mapping, Wuhan University, Wuhan 430079, China

Correspondence to: T. Zhang (tjzhang@lzu.edu.cn)

ABSTRACT

Snow depth is one of key physical parameters for understanding land surface energy balance, soil thermal regimes, water cycles, as well as assessing water resources from local community to regional industrial water supply. Data and knowledge on snow in general and snow depth/snow water equivalent in particular are prerequisites for climate change studies and local/regional development planning. Past studies by  in-situ data are mostly site-specific, while data from satellite remote sensing may cover a large area or in global scale, uncertainties remain large, even misleading. In this study, spatiotemporal change and variability in snow depth was investigated using long-term (1966-2012) ground-based measurements from 1814 stations across the Eurasian continent. Spatially, mean snow depths of >20 cm were recorded in north-eastern European Russia,

1 the Yenisey River basin, Kamchatka Peninsula, and Sakhalin. Annual mean and maximum snow
2 depth increased significantly from 1966 through 2012. Seasonally, monthly snow depth decreased
3 in autumn and increased in winter and spring over the study period. Regionally, snow depth
4 significantly increased in areas north of 50°N. Compared with air temperature, snowfall had more
5 influence on snow depth and snow water equivalent during November through March across the
6 former Soviet Union. This study provides a baseline for snow depth climatology and changes,
7 which were significant in climate system changes over the Eurasian continent.

8

1 Introduction

Snow depth, snow water equivalent (SWE) and snow density are all important parameters for water resource assessment, hydrological and climate model inputs and validation (Dressler et al., 2006; Lazar and Williams, 2008; Nayak et al., 2010).

Changes in snow cover, including snow depth and snow area extent, serve as an indicator of climate change because of their interactions and feedbacks with surface energy and moisture fluxes, hydrological processes, and atmospheric and oceanic circulations (Brown and Goodison, 1996; Armstrong and Brown, 2008; King et al., 2008). Changes in snow depth could have dramatic impacts on weather and climate through the surface energy balance (Sturm et al., 2001), soil temperature and frozen ground (Zhang, 2005), spring runoff, water supply, and human activity (AMAP, 2011).

During winter, the average maximum terrestrial snow cover is approximately $47 \times 10^6 \text{ km}^2$ over the Northern Hemisphere land surfaces (Robinson et al., 1993; IGOS, 2007). A large fraction of the Eurasian continent is covered by snow during the winter season, and some areas are covered by snow for more than half a year. There are long-term snow measurements and observations across the Eurasian continent with the first snow depth record dating back to 1881 in Latvia (Armstrong, 2001). These measurements provide valuable data and information for snow cover phenology and snow cover change detection. Many studies on snow depth have focused on local and regional scales over Russia (Ye et al., 1998; Kitaev et al., 2005; Bulygina et al., 2009, 2011; Brasnett, 1999) and on the Tibetan Plateau (TP) (Li and Mi, 1983; Ma and Qin, 2012), which have revealed the significant regional characteristics in the changes in snow depth. Annual mean snow depth has increased in northern Eurasia and the Arctic during the last 70 years (Ye et al., 1998; Kitaev et al., 2005; Callaghan et al., 2011a; Liston and Hiemstra, 2011) and showed large regional differences (Bulygina et al., 2009, 2011; Ma and Qin, 2012; Stuefer et al., 2013; Terzago et al., 2014). Changes in snow depth are primarily affected by air temperature and precipitation. Ye et al. (1998) and Kitaev et al. (2005) showed that higher air temperatures caused an increase in snowfall in winter from 1936 through 1995, and thus, greater snow depth

1 was observed in northern Eurasia in response to global warming. Furthermore, the
2 snow depth distribution and variation are controlled by terrain (i.e., elevation, slope,
3 aspect, and roughness) and vegetation (Lehning et al., 2011; Gr̃newald et al., 2014;
4 Revuelto et al., 2014; Rees et al., 2014; Dickerson-Lange et al., 2015). Snow depth is
5 closely related to synoptic-scale atmospheric circulation indices such as the North
6 Atlantic Oscillation/Arctic Oscillation (NAO/AO). For example, Kitaev et al. (2002)
7 reported that the NAO index was positively related to snow depth in the northern part
8 of the East European Plain of Russia and over western Siberia from 1966 to 1990;
9 however, the NAO index was negatively correlated with snow depth in most southern
10 regions of northern Eurasia. You et al. (2011) demonstrated that there was a positive
11 relationship between snow depth and the winter AO/NAO index and between snow
12 depth and Ni ño-3 region sea surface temperature (SST) on the eastern and central TP
13 from 1961 through 2005. However, most snow depth studies are at regional scale,
14 information of snow depth at continental scale is required over the Eurasian continent.

15 To increase the spatial coverage of snow depth, researchers have used different
16 instruments (e.g., LIDAR, airborne laser scanning (ALS), and unmanned aerial
17 systems (UASs)) (Hopkinson et al., 2004; Gr̃newald et al., 2013; B̃hler et al., 2016)
18 or developed and/or improved passive microwave snow algorithms (Foster et al.,
19 1997; Derksen et al., 2003; Grippaa et al., 2004; Che et al., 2016). Although snow
20 depth and snow water equivalent obtained by satellite remote sensing could mitigate
21 regional deficiency of the in situ snow depth observations, they have low spatial
22 resolution (25×25 km), and the accuracy is always affected by underlying surface
23 conditions and algorithms. Using ground-based snow depth measurements over the
24 Eurasian continent against snow depth obtained from passive microwave satellite
25 remote sensing, Zheng et al. (2015) found that the mean percentage error was greater
26 than 50% and can be up to approximately 200%. Utilization of snow depth obtained
27 from satellite remote sensing has large uncertainties and is impractical. Apart from
28 remote sensing, numerical modeling is often used to obtain accurate and
29 spatially-complete fields of snow depth and/ or snow water equivalent (SWE) (Liston
30 and Hiemstra, 2011; Terzago et al., 2014; Wei and Dong, 2015). However, remote

1 sensing data with coarse-scale measurement is an important input parameter that
2 affects simulation accuracy and does not provide a sufficient time series length.
3 Spatial interpolation is a common method for estimates in areas with devoid data.
4 However, uncertainty and potential bias in spatial interpolation can be introduced due
5 to specific algorithms especially in complex terrain areas. In addition, data acquisition
6 from large airborne equipment or aerial systems is costly and strict data use
7 limitations apply. Ground-based measurements provide currently available and
8 accurate snow depth over long time-series, which are critical data and information for
9 investigating snow depth climatology and variability and can provide the data base for
10 the verifications of remote sensing and model simulation.

11 The objective of this study is to develop a climatology and investigate the
12 variability of snow depth over the Eurasian continent from 1966 to 2012. In addition,
13 we analyse the effects of topography and climate factors (i.e., air temperature and
14 snowfall) on the changes in snow depth over the study area. This study is unique in
15 snow cover analysis using the most comprehensive daily snow depth observational
16 network at continental scale over Eurasia. The dataset and methodology are described
17 in Section 2 with the results, discussion, and conclusions presented in Sections 3, 4,
18 and 5, respectively.

19

20 **2 Data and Methodology**

21 The data used in this study include daily snow depth, snow water equivalent
22 (SWE), air temperature and precipitation. Measurements of daily snow depth were
23 conducted at 1103 meteorological stations over the Eurasian continent from 1881 to
24 2013 (Table 1). Snow depth was measured once a day at meteorological stations using
25 a graduated stake installed at a fixed point location within the station or by a wooden
26 ruler. Snow depth was measured using the same method across the Eurasian continent
27 since the meteorological observation standard was established by the former Union of
28 Soviet Socialist Republics (USSR) and followed by all of the former USSR republics,
29 Mongolia and China. Snow depth is one of the standard elements to be measured on a
30 daily basis (WMO, 1996). Historical snow course data over the former USSR from

1966 to 2011 were also used in this study. Snow course data include routine snow surveys performed throughout the accumulation season (every ten days) and during the snowmelt period (every five days) over the former USSR. Snow surveys were conducted over 1-2 km-long transects in both forest and open terrain around each station. Snow depth was measured every 10 m in the forest and every 20 m in open terrain (Bulygina et al. 2011).

SWE is an important parameter that is often used in water resource evaluation and hydroclimate studies. SWE was measured using a snow tube every 100 m along the 0.5-1.0 km courses and every 200 m along the 2 km course (Bulygina et al., 2011).


Daily air temperature was measured using a thermometer, which was placed at a height of 1.5 m above the ground surface in an instrument shelter at the meteorological station (WMO, 1996). The air temperature measurement was accurate to 0.1 °C. Air temperature was measured four times a day at 0200, 0800, 1400, and 2000 local time. The daily mean air temperature was calculated by a simple arithmetic average of the four measurements, whereas the monthly mean was based on the daily mean and the annual mean was based on the monthly mean. Precipitation was gathered and measured by a precipitation gauge and was reported with a 0.1-mm precision (Groisman and Rankova, 2001). The original precipitation data were not corrected by considering the gauge undercatch. Daily precipitation was partitioned into a solid and liquid fraction based on daily mean temperature (Brown, 2000). The solid fraction of precipitation, S_{rat} , was estimated by

$$S_{rat} = \begin{cases} 1.0 & \text{for } T_{mean} \leq -2.0^{\circ}C, \\ 0.0 & \text{for } T_{mean} \geq +2.0^{\circ}C, \\ 1.0 - 0.25(T_{mean} + 2.0) & \text{for } -2.0^{\circ}C < T_{mean} < +2.0^{\circ}C. \end{cases} \quad (1)$$

where T_{mean} is the mean daily air temperature (°C).

Snow depth and SWE at each station were determined as the average value of a series of measurements in each snow course survey (Bulygina et al., 2011). In individual measurements, both random and systematic errors inevitably occur (Kuusisto, 1984). To minimize these errors, a quality control of meteorological data was automatically undertaken prior to the datasets being stored at the Russian

1 Research Institute for Hydrometeorological Information-World Data Center
2 (RIHMI-WDC) (Veselov, 2002) and the National Meteorological Information Center
3 (NMIC) of China Meteorological Administration (Ma and Qin, 2012). We
4 implemented additional quality control using the following requirements: (1) To
5 ensure snow depth stability, at a given location, a month with less than 15 days of
6 snow depth measurements was deleted. (2) Stations with sudden and steep changes in
7 snow depth were eliminated from the list. (3) The World Meteorological Organization
8 common approach to calculate anomalies is based on a 30-years climate normal
9 period (IPCC, 2013). In our study, 1971-2000 was used as the normal period. To
10 ensure data continuity, stations with less than 20-years data during the 1971-2000
11 period were excluded. (4) At each station, we eliminated data points that exceeded
12 two standard deviations from their long-term (1971-2000) mean. After these four
13 steps of snow depth quality control, we used data from 1814 stations to investigate the
14 climatology and variability of snow depth over the Eurasian continent (Fig. 1 and
15 Table 1).

16 We defined a snow year starting from July 1st of a current year through June 30th
17 of the following year to capture the entire seasonal snow cycle. Procedures and
18 techniques for measuring snow depth may have changed over the course  station
19 history. Consequently, snow depth data may not be homogeneous in the time series
20 over the period of the record. Fortunately, there was no change in the procedure and
21 technique of snow depth measurements since 1965 in Russia and the other countries
22 in this study (Bulygina et al., 2009). Therefore, in this study, we chose to use snow
23 depth data from 1966 to 2012. The following variables were calculated for each
24 station:

25 (1) Monthly mean snow depth: in this study, we defined a snow cover day with
26 snow depth equal to or greater than 0 cm according to the standard method for
27 deriving monthly mean snow depth based on the World Meteorological Organization
28 (WMO) climatological products (Ma and Qin, 2012). According to the quality control,
29 months having more than 15 days with snow data were used. The monthly mean snow
30 depth was computed as an arithmetic sum of daily snow depth divided by the number

of days with snow on the ground within each month;

(2) Annual mean snow depth: an annual mean snow depth was calculated as an arithmetic sum of the monthly mean snow depth divided by the number of available snow months within each snow year. The annual mean snow depth was averaged for stations with more than 20 snow years during the 1966-2012 period;

(3) Annual mean maximum snow depth: an annual mean maximum snow depth was determined from the maximum daily snow depth in each snow year. It was calculated using the average value of the annual maximum snow depth from stations with more than 20 years of data during the 1966-2012 period.

Anomalies of monthly, annual mean, and annual mean maximum snow depth from their long-term (1971-2000) records were calculated for each station across the Eurasian continent. Composite time series of monthly and annual anomalies were obtained by using all of the available station data across the study area.


Wavelet analysis was performed to reveal the long-term low-frequency variations in snow depth over the entire study area. A wavelet is a wave-like oscillation with an amplitude that begins at 0, increases, and then decreases back to 0 (Graps, 1995). We applied a discrete wavelet transform, excluded the high-frequency components and then used the inverse transform to reconstruct the lower frequency signal. Any trend analysis is an approximate and simple approach to obtain what has occurred on average during the study period. A linear trend analysis provides an average rate of this change. The linear trend analysis is also a useful approximation when systematic low-frequency variations emerge even though there is a nonlinearity (Folland and Karl, 2001; Groisman et al., 2006). The linear trend coefficient of snow depth was calculated to represent the rate of change at each station. The Student's t-test was used to assess statistical significance of the slope in the linear regression analysis and the partial correlation coefficients, and a confidence level above 95% was considered significant in our study. The Durbin-Watson test was used to detect serial correlation of data in the time series, and the Cochran-Orcutt test was used to correct the serial correlation. Then, the serial correlations of the new data were rechecked and recalculated trends in the time series of the new data. The methods and test results

were described in the appendix.

3 Results

3.1 Climatology of Snow Depth

Distributions of long-term mean snow depth indicated a strong latitudinal zonality. Generally, snow depth increased with latitude northward across the Eurasian continent (Fig. 2). A maximum annual mean snow depth of 106.3 cm was observed west of the Yenisey River (dark blue circle) (Fig. 2a). In contrast, the minimum values (~0.01 cm) were observed in some areas south of the Yangtze River in China (small grey circles).

Annual mean snow depth for most areas in Russia  >10 cm. Snow depths were even greater in the north-eastern part of European Russia, the Yenisey River basin, the Kamchatka Peninsula, and Sakhalin with snow depths of >40 cm. Regions with the smallest annual mean snow depth (<5 cm) were located in the eastern and western areas of the Caucasus Mountains. Snow depth in other areas of the former USSR was ~2-10 cm, but shallow snow depths (no more than 1 cm) were observed in some southern regions of Central Asia. The annual average snow depth in the central Mongolian Plateau was lower than that in the northern areas with values of no more than 5 cm. Snow depth was >3 cm in the northern part of the Tianshan Mountains, Northeast China, and some regions of the southwestern TP. In the Altay Mountains and some areas of the north-eastern Inner Mongolia Plateau, annual mean snow depths were >5 cm.

Annual mean maximum snow depth (Fig. 2b) showed a similar spatial distribution pattern compared to the annual mean snow depth pattern. The maximum value was approximately 201.8 cm in snow depth. For the majority of Russia, the maximum snow depth was >40 cm. The regions with maximum snow depths (exceeding 80 cm) were in the north-eastern regions of European Russia, the northern part of the West Siberian Plain, the Yenisey River basin, the Kamchatka Peninsula, and Sakhalin; however, along the coast of the Caspian Sea, the maximum snow depth was <10 cm. Most of the rest of the former USSR had a maximum depth of >10 cm,

except for some regions of the Ukraine and Uzbekistan. Maximum snow depth was >10 cm in northern Mongolia and decreased to 6–10 cm when moving south to central and eastern Mongolia. Maximum snow depths were higher over the northern part of the Xinjiang Autonomous Region of China, Northeast China, and eastern and southwestern TP, which were mostly greater than 10 cm and even greater than 20 cm in some areas. For the remaining regions of China, the maximum snow depths were relatively small and mostly less than 10 cm.

In the autumn months (September to November), the snow depth was shallow (Figs. 3a-c). Monthly mean snow depth was <20 cm in most areas of European Russia and south of Siberia but ranged from ~20 cm to 40 cm in northern Siberia and the Russian Far East in November (Fig. 3c). Moving southward, the monthly mean snow depth was less than 5 cm north of Mongolia and across China. From December to February, the snow depth increased and the snow cover extent expanded significantly (Figs. 3d-f). Monthly snow depth values were >20 cm over the former USSR. Monthly mean snow depth was still <1 cm for the majority of China, except the northern Xinjiang Autonomous Region of China, Northeast China, and south-western TP where snow depth exceeded 10 cm. The snow depth was even more than 20 cm in some places of the Altai Mountains. In spring (March through May), snow cover areas decreased significantly (Figs. 3g-i), which was mainly because of snow disappearance in the majority of China. However, the monthly mean snow depth still exceeded 20 cm in most areas of Russia. Snow cover areas and snow depth gradually decreased in April and May. Snow cover was observed only in Russia and in the TP in June (Fig. 3j).

3.2 Variability of Snow Depth

There were long-term significant increasing trends in both annual mean snow depth and maximum snow depth from 1966 to 2012 over the Eurasian continent. Mean annual snow depth increased at a rate of approximately 0.2 cm decade⁻¹, whereas annual mean maximum snow depth increased at a rate of approximately 0.6 cm decade⁻¹ (Fig. 4). Both annual mean snow depth and maximum snow depth

1 exhibited a similar pattern of changes over the four decades, although the amplitude
2 of maximum snow depth anomaly (approximately ± 2 cm) was much larger than that
3 of the mean snow depth anomaly (approximately ± 1 cm). From the mid-1960s to the
4 early 1970s, annual mean snow depth decreased slightly then increased until the early
5 2000s and then decreased sharply until 2012 (Fig. 4a). Maximum snow depth
6 decreased by 2.5 cm from the mid-1960s through the early 1970s (Fig. 4b). There was
7 a sharp increase of approximately 3 to 4 cm in the maximum snow depth during the
8 1970s and then there was a large fluctuation without a significant trend from the late
9 1970s to the early 1990s. The maximum snow depth increased again from the early
10 1990s through the early 2010s.

11 Monthly snow depth changed significantly across the Eurasian continent from
12 1966 through 2012 (Fig. 5). Snow depth decreased in October at a rate of
13 approximately -0.1 cm decade⁻¹ (Fig. 5a), and there were no significant trends in
14 November and December with large inter-annual variations (Fig. 5b-c). From January
15 through April, snow depth showed statistically increasing trends with rates between
16 0.3 cm decade⁻¹ and 0.6 cm decade⁻¹ (Fig. 5d-g). Overall, snow depth decreased or
17 there was no change in autumn and increased in winter and spring with large
18 inter-annual variations over the study period.

19 Figure 6 shows the spatial distributions of linear trend coefficients of annual
20 mean snow depth and maximum snow depth for each station during 1966-2012 with p
21 ≤ 0.05 . The significant increasing trends (blue circles) of annual mean snow depth
22 occurred in European Russia, south of Siberia and the Russian Far East, the northern
23 Xinjiang Autonomous Region of China, and Northeast China (Fig. 6a). In contrast,
24 decreasing trends (red circles) were detected in western European Russia, some
25 regions of Siberia, north of the Russian Far East, and some regions to the south of
26 40°N in China. Over the entire Eurasian continent, the most significant linear trends
27 in annual mean snow depth were observed in regions north of 50°N , which indicated
28 that the increasing rate of annual mean snow depth was greater in higher latitude
29 regions.

30 In October and November, there were few stations with significant increasing

trends in snow depth ($P \leq 0.05$) (Figs. 7a and b). The increasing trends were mainly observed in most areas across the Eurasian continent in October although the magnitudes were generally small. Over November, the increasing trends in snow depth only appeared in Siberia and the Russian Far East, whereas decreasing trends occurred in monthly mean snow depth over eastern European Russia, the southern West Siberian Plain, and the northeast Russian Far East.

In winter months (December-February), there was a gradual expansion in areas with increasing trends in monthly mean snow depth variation with $P \leq 0.05$ (Figs. 7c–e), and this mainly occurred in eastern European Russia, southern Siberia, the northern Xinjiang Autonomous Region of China, and Northeast China. In contrast, significant decreasing trends were observed in northern and western European Russia and were scattered in Siberia, the northeast Russian Far East, and northern China.

From March to May, the number of stations with significant changes ($P \leq 0.05$) in monthly mean snow depth decreased, especially in May because of snow melt (only 78 stations) (Figs. 7f–h). Changes in monthly mean snow depth were consistent with the trends in winter over the former USSR, but more stations with decreasing trends were found in southern Siberia. There were few stations with statistically significant trends in snow depth across China; for these stations, monthly snow depths tended to decrease at most stations. Compared with regions south of 50°N, changes in monthly mean snow depth were more significant over regions north of 50°N.



4 Discussion

4.1 Comparisons with previous results

Studies on changes in snow depth have received much attention over different regions across Eurasian continent. This study, for the first time, investigated changes in snow depth using ground-based data and information over the region as a whole. Ma and Qin (2012) investigated changes in snow depth across China over period from 1957 to 2009. We found that the climatology (1966–2012) of snow depth from this study was basically consistent with the results from Ma and Qin (2012) over China. In terms of changes in snow depth, both studies showed increase in snow depth

1 but with slight difference in magnitude. This may be caused by using a different
2 number of stations and covering different study periods. Over northern Eurasia,
3 Kitaev et al. (2005) and Bulygina et al. (2011) investigated snow depth and its change.
4 The long-term (1966-2012) mean snow depth from this study was approximately 5-10
5 cm higher than the results from Kitaev et al. (2005) and Bulygina et al. (2011) over
6 northern Eurasia. These discrepancies may result from differences in the time frame
7 of data collection, the number of stations, calculation methods, and data quality
8 control. For example, Kitaev et al. (2005) investigated historical changes in snow
9 depth spanning 65 years from 1936 to 2000, while this study covered 47 years from
10 1966 through 2010. In this study, we intentionally did not use the earlier (1936-1965)
11 data due primarily to data quality. The earlier Russian snow depth data were
12 discontinuous and did not meet the data quality control requirements used in this
13 study. Historical changes of the hydrometeorological station locations were also a
14 critical reason for deleting many stations from the study. Based on results from this
15 study, we believe that snow depth data in early years (prior to 1965) may be
16 questionable and changes in snow depth prior to 1965 over Russia need further
17 in-depth investigation.

18 Ye et al. (1998) found that historical winter snow depth increased in northern
19 Russia (1.86 cm yr^{-1}) and decreased in southern Russia at a rate of -0.23 cm yr^{-1}
20 during 1936-1983 (Ye et al., 1998). Results from this study were essentially consistent
21 with Ye et al. (1998) in northern Russia; however, in southern Siberia where snow
22 depth increased at a rate of 0.42 cm yr^{-1} during the period from 1966 to 2012. We
23 believe that the difference is mainly due to the time periods covered by the two
24 studies.

25 Liston and Hiemstra (2011) conducted snow depth assimilation using the
26 SnowModel. Results from the SnowModel assimilations in general agree well with
27 ground-based measurements. For example, both observations from this study and
28 assimilations with the SnowModel (Liston and Hiemstra, 2011) presented that the
29 peak snow depth and SWE occurred more in the western portion of northern Eurasia
30 than the eastern portion of the Russian Far East. This may be primarily because the

SnowModel input data included ground-based measured air temperature, precipitation, wind conditions and in part snow depth. However, results from CMIP5 (Coupled Model Intercomparison Project Phase 5, Terzago et al., 2014; Wei and Dong, 2015) overestimated snow depth over the TP and underestimated in forest regions. This implies that large uncertainties currently still exist in modeling snow depth.

4.2 Impact of Topography on Snow Depth

Topography is an important factor affecting the climatology of snow depth and is the main reason accounting for the inhomogeneity of data (Grünewald and Lehning, 2011, 2013; Grünewald et al., 2014). To explore the effects of complex terrain on snow depth over Eurasia, we conducted a linear regression analysis of the annual mean snow depth with latitude, elevation and continentality (Fig. 8). Snow depth was positively correlated with latitude, i.e., snow depth generally increased with latitude (Fig. 8a). The increased rate of snow depth was approximately 0.81 cm per 1°N across the Eurasian continent. A closer relationship between latitude and snow depth was found in regions north of 40°N where snow cover was relatively stable with the number of annual mean continuous snow cover days at more than 30 (Zhang and Zhong, 2014).

There was a negative correlation between snow depth and elevation across the Eurasian continent (Fig. 8b); with every 100 m increase in elevation, snow depth decreased by ~0.5 cm ($P \leq 0.05$). Annual mean snow depth was less than 1 cm in most areas, with an elevation greater than 2000 m because a snow depth of 0 cm was used to calculate the mean snow depth. Therefore, although the TP is at a high elevation, the shallow snow depth in this area resulted in a generally negative correlation between snow depth and elevation across the Eurasian continent. Snow depths were averaged to 200 m elevation bands and then discussed the relationship between elevation level for the former USSR and China. Snow depths were deeper in the lower elevation bands between 0 and 600 m across the former USSR (Fig. 8c). However, there were shallow snow accumulation between 600 and 1000 m due to most accumulation areas located in forest. Then snow depth was followed by a significant positive trend and

1 reached a peak. Snow depths represented marked decrease in the highest elevation
2 band (2600~2900 m). There were only two stations in this band and more snow
3 accumulation difference between the two stations because of terrain and climate
4 factors. Snow depths were deeper in three elevation bands across China: 200~1000 m,
5 1600~1800 m and 2400~2600 m. Greater snow accumulation were attributed to heavy
6 snowfall and severe cold in these regions. An increasing trend of snow depth
7 presented in the higher elevations above 2600 m on the TP.

8 There was a statistically significant positive relationship between snow depth and
9 continentality over the Eurasian continent ($r=0.1$, $P\leq 0.05$, Fig. 8d). This indicated
10 that the continentality may be an important driving factor of snow depth
11 distribution over Eurasia, especially on the TP. Although previous studies showed that
12 the TP's largest snow accumulation occurred in winter, precipitation during winter
13 months was the smallest of the year (Ma, 2008). This was mainly due to the majority
14 of annual precipitation that occurs during the summer monsoon season on the TP,
15 which causes much less precipitation during the winter half year (or the snow
16 accumulation season).

18 4.3 Impact of Climate Factors on Snow Accumulation

19 In addition to the terrain factors, variations in snow depth are closely related to
20 climate variability. To examine the relationship between snow depth and climatic
21 factors, we calculated the long-term mean snow depth, air temperature and snowfall
22 of 386 stations from November through March across the former USSR (Fig. 9). The
23 period (snow cover years) spanned from 1966 through 2009 using available data.
24 Snow depth significantly decreases with increasing air temperature ($P\leq 0.05$) but the
25 Goodness of Fit of the relationship was only 16% (Fig. 9a). Compared with air
26 temperature, snowfall exhibited a strong relationship with snow depth (Fig. 9b). The
27 mean snow depth was less than 20 cm at most stations with an accumulated snowfall
28 of <50 mm from November through March. Snow depth increased with an increase in
29 accumulated snowfall, and the thickest snow depth of approximately 120 cm had a
30 maximum cumulative snowfall of approximately 350 mm.

Compared with the long-term inter-annual trends in change in snow depth, air temperature and snowfall, the variabilities in snow depth was mainly affected by the changes in snowfall. Overall, the trends in long-term air temperature, snowfall and snow depth displayed increasing trends from November to March (Fig. 10). This was because the increased precipitation fell as snow in cold areas where the increased temperature was still below freezing (Ye et al., 1998; Kitaev et al., 2005). Warmer air led to a greater supply of moisture for snowfall and hence the snow accumulation still increased (Ye et al., 1998). Significant increasing snowfall can explain the sudden drop in bulk snow density from the mid-1990s through the early 2000s (Zhong et al., 2014): increasing snowfall should decrease the density of the surface snowpack, which lowered the entire density of the snowpack. In addition, there were similar inter-annual variations in snowfall and heavy snowfall. This indicated that extreme snowfall events may be the main cause of the increase in snow depth.

The partial correlation coefficients between snow accumulation, air temperature and snowfall were calculated to discuss the spatial relationship between them (Fig. 11). A significant negative correlation ($p \leq 0.05$) between snow depth and air temperature was present in most areas of European Russia and southern Siberia (Fig. 11a). The stations with negative effects of air temperature on snow depth were fewer, and there were no statistically significant correlations in northern Siberia (Fig. 11b). This was because there was no obvious effect of increasing temperature on snow depth when the air temperature was below 0 °C, which occurred in most areas of Siberia from December through March.

Compared with previous studies, the sensitivity of snow depth to air temperature and precipitation for each station showed regional differences (Fallot et al., 1997; Park et al., 2013). The amount of snowfall can be affected by climate change and lead to differences in snow depth at different times (Ye et al., 1998; Kitaev et al., 2005; Ma and Qin, 2012). We found that there was a significant ($p \leq 0.05$) negative relationship between snow depth and air temperature in southern Siberia but not in northern Siberia. In addition to air temperature and precipitation, atmospheric circulation was a key factor affecting snow depth change (Cohen, 2011; Zhao et al., 2013; Ye et al.,

2015). Those factors above and related uncertainties may explain the regional and temporal differences in long-term mean snow depth and snow depth change.

Snow cover extent and snow cover duration decreased in response to climate change (Bulygina et al., 2009; Brown and Robinson, 2011; IPCC, 2013; Xu et al., 2017), however, snow accumulation increased significantly with in situ data over Eurasia. Our study showed that heavy snowfall may be the main reason for snow thickening, and atmospheric circulation was also an important factor.

4.4 Potential Effects of the Variations in Snow Depth

Snow depth is an important factor of controlling the ground thermal regime (Goodrich, 1982; Zhang et al., 1996; Zhang, 2005; Ling and Zhang, 2005; Park et al., 2014). Research has shown that thin snow cover resulted in a cooler soil surface, whereas thick snow cover led to a warmer soil surface (Kudryavtsev, 1992). Frauenfeld et al. (2004) indicated that the maximum snow depth by the end of winter had a significant influence on the active layer depth in the following summer. Snow depth was responsible for 50% or more of the changes in soil temperature at a depth of 3.6 m in north-eastern Siberia from 1901-2009 (Park et al., 2014). Results from this study indicated that snow depth significantly decreased on the TP and increased in Siberia. Although it is not clear what is the role (cooling or warming) of snow cover on soil thermal region on the TP, the decrease in snow depth would reduce the warming effect, offsetting the increase in permafrost temperatures (Zhang, 2012). Over Siberia, increase in snow depth would further increase permafrost temperatures (Zhang et al., 2001, 2005; Park et al., 2014), enhancing permafrost degradation over the region.

Snow cover has an important impact on the hydrological cycle (AMAP, 2011). Spring floods are generated by melting snow, and freshwater derives are from snowmelt in some snow-dominated basins (Barnett et al., 2005). Increasing snow depth may lead to frequent spring floods in northern Xinjiang and snow accumulation reduction can result in freshwater shortage on the TP. Furthermore, snow interacts with vegetation and in turn vegetation affects snow cover accumulation, redistribution

and the vertical profile in forests or shrubs (Hedstrom and Pomeroy, 1998; Pomeroy et al., 2006). Snow also influences plant growth, high snow depth with more water amount can increase soil moisture and promote vegetation productivity (Peng et al., 2010). Therefore, increasing snow depths could contribute to forest growth in northern Eurasia and north-eastern China.

5 Conclusions

In this study, daily snow depth and snow course data from 1814 stations were used to investigate spatial and temporal changes in annual mean snow depth and maximum snow depth over the Eurasian continent for the period from 1966 to 2012. Our results demonstrate that greater long-term average snow depth was observed in north-eastern European Russia, the Yenisey River basin, the Kamchatka Peninsula, and Sakhalin. In contrast, the shallowest snow depths were recorded in China, except for the northern Xinjiang Autonomous Region of China, Northeast China, and in some regions of the southwestern TP.

There were statistically significant trends in variations in long-term snow depth over the entire Eurasian continent. A similar increasing pattern of changes was exhibited in both annual snow depth and maximum snow depth, although the amplitude of the maximum snow depth anomaly was much larger than the equivalent value for mean snow depth. Monthly snow depth in autumn presented a decreasing trend, whereas there were increasing trends in the variations in snow depth during winter and spring, especially during the period of the mid-1980s through the 2000s.

Significant increasing trends in snow depth were detected in the eastern regions of European Russia, southern Siberia, the Russian Far East, the northern areas of the Xinjiang Autonomous Region of China, and north-eastern China. Decreasing linear trends were observed in most western areas of European Russia, some regions of southern Siberia, the north-eastern Russian Far East and most areas in the southern 40°N across China.

Compared with elevation, latitude played a more important role in snow depth climatology. Variations in snow depth were explained by air temperature and snowfall

1 in most areas of European Russia and some regions of southern Siberia and the effects
2 of the two factors on SWE only appeared in some of these areas; however, snowfall
3 especially heavy snowfall was the main driving force of the variance of snow depth
4 and SWE in the former USSR.
5

Appendix A: Analysis of serial correlation

In this research, the Kolmogorov-Smirnov (K-S) test was used to determine whether snow depth data followed a normal distribution. The results showed that all station data followed a normal distribution (such as annual mean snow depth for all stations, Fig. A1). We used ordinary linear regression (OLR) to detect trends in changes in snow depth. Failure to consider the serial correlation of data could lead to erroneous results when detecting the trends in a time series of snow depth, which is mainly because the probability of detecting false trends would be increased (Westerhead et al, 1998; Storch, 1999; Khaliq et al., 2009). To avoid this situation, we used the Durbin-Watson test to check the serial correlation (Neter et al., 1989; Tao et al., 2008):

$$d = \frac{\sum_{t=2}^n (e_t - e_{t-1})^2}{\sum_{t=1}^n e_t^2} \quad (A1)$$

where e_t was the residual estimated by the OLR, and t was the number of observations. d_l was the lower critical value, and d_u was the upper critical value, which could be obtained through the Durbin-Watson statistic table. If $d_u \leq d \leq 4 - d_u$, a serial correlation was absent; if $d \leq d_l$ or $d \geq 4 - d_l$, a serial correlation was present.

We used the Cochrane-Orcutt method to correct the variable if the serial correlation was present (Neter et al., 1989; Tao et al., 2008):

$$X'_t = X_t - \rho X_{t-1} \quad (A2)$$

$$Y'_t = Y_t - \rho Y_{t-1} \quad (A3)$$

where X' was the corrected year, Y' was the corrected anomaly in time series of snow depth for each station in this research, and the autocorrelation coefficient ρ was replaced by its estimate value r :

$$r = \frac{\sum_{t=2}^n e_{t-1} e_t}{\sum_{t=2}^n e_{t-1}^2} \quad (A4)$$

Then, the Durbin-Watson test was used to check the serial correlation of the new snow depth anomalies, and recalculated the trends in the time series of new data.

The Durbin-Watson test results show that there were no serial correlations in the

inter-annual trends in annual mean snow depth, maximum snow depth and monthly mean snow depth for all of the composite data ($d_u \leq d \leq 4 - d_u$) (Table A1). However, the serial correlation was present in some stations when we calculated the linear trend of annual snow depth, maximum depth and monthly mean snow depth for each station. The percentage of the stations with a serial correlation for annual snow depth and maximum depth were 18% and 21%, respectively. In the monthly test, the smallest proportion appeared in October at approximately 11%; the largest percentage of these stations for all of the stations was found in February and was up to 21%. Then, the Cochrane-Orcutt method was used to correct the variables and re-estimated the trends in snow depth for these station (Fig. 6-7 in the text). Using the Dikson site (73.5° N, 80.4° E, 42 m a.s.l.) as an example, the serial correlation was present when the trend in annual mean snow depth was calculated. Compared with the corrected result, the variance of the previous OLR statistic was overestimated, and annual mean snow depth increased at the rate of 0.113 cm yr⁻¹ (Table A2). The corrected result indicated that the variation of inter-annual mean snow depth was not significant ($P' > 0.05$). The serial correlation cannot be ignored for detecting trends in a time series of snow cover variables, which possibly invalidates the statistical test on slopes if this variable is not dealt with.

Acknowledgements. We express our gratitude to the researchers who assembled and digitized the snow depth data at meteorological stations and snow surveys across the Eurasian continent over a period of >40 years. This work was funded by the National Key Scientific Research Program of China (2013CBA01802), the Open Foundation from the State Key Laboratory of Cryospheric Sciences (SKLCS-OP-2016-12), the Project for Incubation of Specialists in Glaciology and Geocryology of the National Natural Science Foundation of China (J1210003/ J0109), and the Foundation for Excellent Youth Scholar of Cold and Arid Research Environmental and Engineering Research Institute, Chinese Academy of Sciences.

References

- AMAP: Snow, Water, Ice and Permafrost in the Arctic (SWIPA), Oslo, Norway: Arctic Monitoring and Assessment Programme (AMAP), 553pp, 2011.
- Armstrong, R. L.: Historical Soviet daily snow depth, version 2 (HSDSD). Boulder, Colorado: National Snow and Ice Data Center. CD-ROM. 2001.
- Armstrong, R. L. and Brown, R.: Introduction, in: Snow and climate: Physical processes, surface energy exchange and modeling, edited by: R. L. Armstrong, and Brun, E., Cambridge University Press, Cambridge, UK, 1-11, 2008.
- Barnett, T.P., Adam, J.C., and Lettenmaier, D.P.: Potential impact of a warming climate on water availability in snow-dominated regions, *Nature*, 438, 303-309, 2005.
- Brasnett, B.: A global analysis of snow depth for numerical weather prediction, *J. Appl. Meteorol.*, 38, 726-740, 1999.
- Brown, R. D.: Northern Hemisphere snow cover variability and change, 1915-97, *J. Climate*, 13, 2339-2355, 2000.
- Brown, R. D. and Goodison, B. E.: Interannual variability in reconstructed Canadian snow cover, 1915-1992, *J. Climate*, 9, 1299-1318, 1996.
- Brown, R. D. and Robinson, D. A.: Northern Hemisphere spring snow cover variability and change over 1922-2010 including an assessment of uncertainty, *The Cryosphere*, 5, 219-229, 2011.
- Bulygina O. N., Razuvaev, V. N., and Korshunova, N. N.: Changes in snow cover over Northern Eurasia in the last few decades, *Environ. Res. Lett.*, 4, 045026, doi:10.1088/1748-9326/4/4/045026, 2009.
- Bulygina, O. N., Groisman, P. Y., Razuvaev, V. N., and Korshunova, N. N.: Changes in snow cover characteristics over Northern Eurasia since 1966, *Environ. Res. Lett.*, 6, 045204, doi:10.1088/1748-9326/6/4/045204, 2011.
- Callaghan, T. V., Johansson, M., Brown, R. D., Groisman, P. Ya., Labba, N., and Radionov, V.: The changing face of Arctic snow cover: A synthesis of observed and projected changes, *Ambio*, 40, 17-31. doi:10.1007/s13280-011-0212-y, 2011.
- Che, T., Dai, L., Zheng, X., Li X., and Zhao, K.: Estimation of snow depth from passive microwave brightness temperature data in forest regions of northeast China, *Remote Sens.*

1 Environ., 183, 334-349, 2016.

2 Cohen, J.: Eurasian snow cover variability and links with stratosphere-troposphere coupling and
3 their potential use in seasonal to decadal climate predictions, US National Oceanic and
4 Atmospheric Administration, Climate test bed joint seminar series, NCEP, Camp Springs,
5 Maryland, 6pp, 2011.

6 Cohen, J. L., Furtado, J. C., Barlow, M. A., Alexeev, V. A., and Cherry, J. E.: Arctic warming,
7 increasing snow cover and widespread boreal winter cooling, Environ. Res. Lett., 7, 014007,
8 2012.

9 Derksen, C., Walker, A., and Goodison, B.: A comparison of 18 winter seasons of in situ and
10 passive microwave-derived snow water equivalent estimates in Western Canada, Remote
11 Sens. Environ., 88, 271-282, 2003.

12 Dickerson-Lange, S.E., Lutz, J.A., Martin, K.A., Raleigh, M.S., Gersonde, R., and Lundquist, J.D.:
13 Evaluating observational methods to quantify snow duration under diverse forest canopies,
14 Water Resour. Res., 51, 1203-1224, 2015.

15 Dressler, K.A., Leavesley, G.H., Bales, R.C., Fassnacht, S.R.: Evaluation of gridded snow water
16 equivalent and satellite snow cover products for mountain basins in a hydrologic model.
17 Hydrol. Process., 20, 673-688, 2006.

18 Fallot, J., Barry, B. G., and Hoogstrate, D.: Variations of mean cold season temperature,
19 precipitation and snow depths during the last 100 years in the former Soviet Union (FSU),
20 Hydrol. Sci. J., 42, 301-327, 1997.

21 Folland, C.K. and Karl, T.R.: Observed climate variability and change. Climate Change 2001: The
22 Scientific Basis, edited by: J. T. Houghton et al., Cambridge University Press, Cambridge,
23 UK, 99–181, 2001.

24 Foster, J.L., Chang, A.T.C., and Hall, D.K.: Comparison of snow mass estimates from a prototype
25 passive microwave snow algorithm, Remote Sens. Environ., 62, 132-142, 1997.

26 Goodrich, L.E.: The influence of snow cover on the ground thermal regime, Can. Geotech. J., 19,
27 421-432, 1982.

28 Ghatak, D., Frei, A., Gong, G., Stroeve, J., and Robinson, D.: On the emergence of an Arctic
29 amplification signal in terrestrial Arctic snow extent, J. Geophys. Res., 115, D24105, 2010.

30 Graps, A.: An introduction to wavelets, Computational Science & Engineering, IEEE, 2, 50-61,

1 1995.

2 Grippaa, M., Mognarda, N., Le Toana, T., and Josberger E.G.: Siberia snow depth climatology
3 derived from SSM/I data using a combined dynamic and static algorithm, *Remote Sens.*
4 *Environ.*, 93, 30-41, 2004.

5 Groisman, P.Y. and Rankova, E.Y.: Precipitation trends over the Russian permafrost-free zone:
6 removing the artifacts of pre-processing, *Int. J. Climatol.*, 21, 657-678, 2001.

7 Groisman, P., Knight, R., Razuvaev, V., Bulygina, O., and Karl, T.: State of the Ground:
8 Climatology and Changes during the Past 69 Years over Northern Eurasia for a Rarely Used
9 Measure of Snow Cover and Frozen Land, *J. Climate*, 19, 4933–4955, 2006.

10 Gr̈unewald, T. and Lehning, M.: Altitudinal dependency of snow amounts in two small alpine
11 catchments: can catchment-wide snow amounts be estimated via single snow or precipitation
12 stations? *Ann. Glaciol.*, 52, 153-158, 2011.

13 Gr̈unewald, T. and Lehning, M.: Can a point measurement represent the snow depth in its vicinity?
14 A comparison of areal snow depth measurements with selected index sites, *Proceedings of*
15 *the International Snow Science Workshop*, 27 October 7 to 11, Grenoble, France, 69-72,
16 2013.

17 Gr̈unewald, T., B̈uhler, Y., and Lehning, M.: Elevation dependency of mountain snow depth, *The*
18 *Cryosphere*, 8, 2381-2394, 2014.

19 Hedstrom, N.R. and Pimeroy, J.W.: Measurement and modelling of snow interception in the boreal
20 forest, *Hydrol. Process.*, 12, 1611-1625, 1998.

21 Integrated Global Observing Strategy (IGOS): Cryosphere Theme Report - For the monitoring of
22 our environment from space and from earth, WMO/TD-No.1405, World Meteorological
23 Organization, Geneva, 100 pp., 2007.

24 IPCC: Climate change 2013: The physical science basis. Contribution of working group I to the
25 fifth assessment report of the intergovernmental panel on climate change, edited by: T.F.
26 Stocker et al., Cambridge University Press, Cambridge, UK, and New York, USA, 1535 pp,
27 2013.

28 Ji, Z., and Kang, S.: Projection of snow cover changes over China under RCP scenarios, *Clim.*
29 *Dyn.*, 41: 589-600, 2013.

30 Khaliq, M.N., Ouarda, T.B.M.J., Gachon, P., Sushama, L., and St-Hilaire, A.: Indentification of

1 hydrological trends in the presence of serial and cross correlations: A review of selected
2 methods and their application to annual flow regimes of Canadian rivers, *J. Hydrol.*, 368,
3 117-130, 2009.

4 King, J. C., Pomeroy, J. W., Gray, D. M., Fierz, C., Fohn, P., Harding, R. J., Jordan, R. E., Martin,
5 E., and Pluss, C.: Snow-atmosphere energy and mass balance, in: *Snow and Climate:
6 Physical Processes, Surface Energy Exchange and Modeling*, edited by: Armstrong, R. L. and
7 Brun, E., Cambridge University Press, Cambridge, UK, 70–124, 2008.

8 Kitaev, L., Kislov, A., Krenke, A., Razuvaev, V., Martuganov, R., and Konstantinov, I.: The snow
9 cover characteristics of northern Eurasia and their relationship to climatic parameters, *Boreal
10 Environ. Res.*, 7, 437-445, 2002.

11 Kitaev, L., Førlund, E., Razuvaev, V., Tveito, O. E., and Krueger, O.: Distribution of snow cover
12 over Northern Eurasia, *Nord. Hydrol.*, 36, 311-319, 2005.

13 Kuusisto, E.: Snow accumulation and snowmelt in Finland, *Publications of the Water Research
14 Institute*, v. 55, Valtion painatuskeskus, Helsinki, 149 pp., 1984.

15 Lazar, B. and Williams, M.: Climate change in western ski areas: potential changes in the timing
16 of wet avalanches and snow quality for the Aspen ski area in the years 2030 and 2100, *Cold
17 Reg. Sci. Technol.*, 51, 219–228, 2008.

18 Lehning, M., Grönwald, T., and Schirmer, M.: Mountain snow distribution governed by an
19 altitudinal gradient and terrain roughness, *Geophys. Res. Lett.*, 38, L19504, 2011.

20 Li, P. and Mi, D.: Distribution of snow cover in China, *J. Glaciol. Geocryol.*, 5, 9-18, 1983 (in
21 Chinese with English abstract).

22 Ling, F. and Zhang, T.: Modeling the effect of variations in snowpack-disappearance date on
23 surface-energy balance on the Alaskan north slope, *Arct. Antarct. Alp. Res.*, 37, 483-489,
24 2005.

25 Liston, G.E., and Hiemstra, C.A.: The changing cryosphere: Pan-Arctic snow trends (1979-2009),
26 *J. Climate*, 24, 5691-5712, 2011.

27 Liu, J., Curry, J., Wang, H., Song, M., and Horton, R.: Impact of declining Arctic sea ice on winter
28 snowfall, *Proc. Natl. Acad. Sci. U.S.A.*, 109, 4074-4079, 2012.

29 Ma, L.: Spatiotemporal variation in snow cover and the relationship between snow cover and
30 atmospheric circulation in the Tibetan Plateau in recent 50 years, PhD dissertation, Graduate

1 University of Chinese Academy of Sciences, 156pp, 2008.

2 Ma, L. and Qin, D.: Spatial-temporal characteristics of observed key parameters for snow cover in
3 China during 1957-2009, *J. Glaciol. Geocryol.*, 34, 1-11, 2012 (in Chinese with English
4 abstract).

5 Ministry of Environment Province of British Columbia (MEPBC): Snow survey sampling guide,
6 Surface Water Section, Water Management Branch, Ministry of Environment Province of
7 British Columbia, 1981.

8 Nayak, A., Marks, D., Chandler, D.G., and Seyfried, M.: Long-term snow, climate, and streamflow
9 trends at the Reynolds Creek experimental watershed, Owyhee Mountains, Idaho, United
10 States. *Water Resour. Res.*, 46, W06519. doi: 10.1029/2008WR007525, 2010.

11 Neter, J., Wasserman, W., and Kutner, M.H.: Applied linear regression model, Boston, IRWIN,
12 1989.

13 Park, H., Walsh, J. E., Kim, Y., Nakai, T., and Ohata, T.: The role of declining Arctic sea ice in
14 recent decreasing terrestrial Arctic snow depths, *Polar Sci.*, 7, 174-187, 2013.

15 Park, H., Fedorov, A.N., Zheleznyak, M.N., Konstantinov, P.Y., and Walsh, J.E.: Effect of snow
16 cover on pan-Arctic permafrost thermal regimes, *Clim. Dyn.*, 44, 2873-2895, 2015.

17 Peng, S., Piao, S., Ciais, P., Fang, J., and Wang, X.: Change in winter snow depth and its impacts
18 on vegetation in China, *Global Change Biol.*, 16, 3004-3013, 2010.

19 Pomeroy, J.W., Bewley, D.S., Essery, R.L.H., Hedstrom, N.R., Link, T., Granger, R.J., Sicart, J.-E.,
20 Ellis, C.R., and Janowicz, J.R.: Shrub tundra snowmelt, *Hydrol. Process.*, 20, 923-941, 2006.

21 Rees, A., English, M., Derksen, C., Toose, P., and Silis, A.: Observations of late winter Canadian
22 tundra snow cover properties, *Hydrol. Process.*, 28, 3962-3977, 2014.

23 Revuelto, J., López-Moreno, J.I., Azorin-Molina, C., and Vicente-Serrano, S.M.: Topographic
24 control of snowpack distribution in a small catchment in the central Spanish Pyrenees: intra-
25 and inter-annual persistence, *The Cryosphere*, 8, 1989-2006, 2014.

26 Robinson, D. A., Dewey, K. F., and Heim, R. R.: Global snow cover monitoring: An update, *B.*
27 *Am. Meteorol. Soc.*, 74, 1689-1696, 1993.

28 Storch, H.V.: Misuses of Statistical Analysis in Climate Research, in: *Analysis of Climate*
29 *Variability*, edited by: Storch, H.V. and Navarra, A.: Springer Press, Berlin Heidelberg,
30 Germany, 11-26, 1999.

1 Stuefer, S., Kane, D.L., and Liston, G.E.: In situ snow water equivalent observations in the US
2 Arctic, *Hydrol. Res.*, 44, 21-34, 2013.

3 Sturm, M., McFadden, J. P., Liston, G. E., Chapin III, F. S., Racine, C. H., and Holmgren, J.:
4 Snow-shrub interactions in Arctic tundra: A hypothesis with climatic implications. *J. Climate*,
5 14, 336-344, 2001.

6 Tao, J., Zhang, X., Tao, J., and Shen, Q.: The checking and removing of the autocorrelation in
7 climatic time series, *Journal of Applied Meteorological Science*, 19, 47-52, 2008.

8 Terzago, S., Hardenberg, J. Palazzi, E., and Provenzale, A.: Snowpack changes in the Hindu
9 Kush-Karakoram-Himalaya from CMIP5 global climate models, *J. Hydrometeorol.*, 15,
10 2293-2313, 2014.

11 Veselov, V. M.: PC archives of the State Data Holding and technology of their organization,
12 Proceedings of the RIHMI-WDC (Russian Research Institute for Hydrometeorological
13 Information-World Data Center), 170, 16-30, 2002 (in Russian).

14 Weatherhead, E.C., Reinsel, G.C., Tiao, G.C., Meng, X., Choi, D., Cheang, W., Keller, T., DeLuisi,
15 J., Wuebbles, D.J., Kerr, J.B., Miller, A.J., Oltmans, S.J., and Frederick, J.E.: Factors
16 affecting the detection of trends: Statistical considerations and applications to environmental
17 data, *J. Geophys. Res.*, 103(D14), 17149-17161, 1998.

18 Wei, Z. and Dong, W.: Assessment of Simulations of Snow Depth in the Qinghai-Tibetan Plateau
19 Using CMIP5 Multi-Models, *Arct. Antarc. Alp. Res.*, 47, 611-625, 2015.

20 WMO: Guide to meteorological instruments and methods of observation, WMO-No.8, Geneva,
21 Switzerland, 1996.

22 Xu, W., Ma, L., Ma, M., Zhang, H., and Yuan, W.: Spatial-temporal variability of snow cover and
23 depth in the Qinghai-Tibetan Plateau, *J. Climate*, 30, 1521-1533, 2017.

24 Ye, H. C., Cho, H. R., and Gustafson, P. E.: The changes in Russian winter snow accumulation
25 during 1936-83 and its spatial patterns, *J. Climate*, 11, 856-863, 1998.

26 Ye, K., Wu, R., and Liu Y.: Interdecadal change of Eurasian snow, surface temperature, and
27 atmospheric circulation in the late 1980s, *J. Geophys. Res. Atmos.*, 120, 2738-2753, 2015.

28 You, Q., Kang, S., Ren, G., Fraedrich, K., Pepin, N., Yan, Y., and Ma L.: Observed changes in
29 snow depth and number of snow days in the eastern and central Tibetan Plateau, *Clim. Res.*,
30 46, 171-183, 2011.

1 Zhang, T.: Influence of the seasonal snow cover on the ground thermal regime: An overview, *Rev.*
2 *Geophys.*, 43, RG4002, doi: 10.1029/2004RG000157, 2005.

3 Zhang, T.: Progress in global permafrost and climate change studies, *Quaternary Sciences*, 32,
4 27-38, 2012 (in Chinese with English abstract).

5 Zhang, T., Osterkamp, T. E., and Stamnes, K.: Influence of the depth hoar layer of the seasonal
6 snow cover on the ground thermal regime, *Water Resour. Res.*, 32, 2075-2086, 1996.

7 Zhang, T., Barry, R.G., Gilichinsky, D., Bykhovets, S.S., Sorokovikov, V.A., and Ye, J.: An
8 amplified signal of climatic change in soil temperatures during the last century at Irkutsk,
9 Russia, *Clim. Change*, 49, 41-76, 2001.

10 Zhang, T. and Zhong, X.: Classification and regionalization of the seasonal snow cover across the
11 Eurasian continent, *J. Glaciol. Geocryol.*, 36, 481-490, 2014 (in Chinese with English
12 abstract).

13 Zhao, H., Higuchi, K., Waller, J., Auld, H., and Mote, T.: The impacts of the PNA and NAO on
14 annual maximum snowpack over southern Canada during 1979-2009, *Int. J. Climatol.*, 33,
15 388-395, 2013.

16 Zheng, L., Zhang, T., Che, T., Zhong, X., and Wang, K.: Evaluation of snow depth products
17 derived from passive microwave satellite remote sensing data using ground-based snow
18 measurements, *Remote Sensing Technology and Application*, 30, 413-423, 2015 (in Chinese
19 with English abstract).

20 Zhong, X., Zhang, T., and Wang, K.: Snow density climatology across the former USSR, *The*
21 *Cryosphere*, 8, 785-799, 2014.

22

Tables and Figures

Table 1. Sources of snow depth data

Dataset	Spatial distribution	Number of stations	Source
Daily snow depth	the former USSR	586	Russian Research Institute for Hydrometeorological Information-World Data Center (RIHMI-WDC) National Snow and Ice Data Center (NSIDC), University of Colorado at Boulder
	China	492	National Meteorological Information Center (NMIC) of the China Meteorological Administration
	Mongolia	25	NSIDC
Snow depth from snow courses	the former USSR	1044	RIHMI-WDC, NSIDC
Snow water equivalent (SWE)	the former USSR	386	RIHMI-WDC
Daily air temperature and precipitation	the former USSR	386	RIHMI-WDC

Table A1. Trends in snow depths with the Durbin-Watson test across Eurasia during 1966-2012

	d_1	d_u	d	$slope^*$	P^*
Mean	1.3034	1.3871	1.6435	0.02	0.0016
Maximum	1.3034	1.3871	1.8824	0.06	0.0004
October	1.3034	1.3871	2.1377	-0.01	0.0069
November	1.4872	1.5739	2.3667	0.00	0.7408
December	1.4872	1.5739	1.9684	0.02	0.0793
January	1.3034	1.3871	1.6326	0.04	0.0014
February	1.3034	1.3871	1.8469	0.06	0.0000
March	1.3034	1.3871	1.9874	0.06	0.0003
April	1.3034	1.3871	1.6754	0.03	0.0187
May	1.4872	1.5739	2.0703	0.00	0.5811

*: slope was the trend of changes in snow depth, the unit was cm yr^{-1} ; P was the confidence level.

Table A2. Trends in annual mean snow depth with the Durbin-Watson test for the Dikson site during 1966-2012

ID	d_1	d_u	d	$slope$	P	d'_1	d'_u	d'	$slope'^*$	P'^*
----	-------	-------	-----	---------	-----	--------	--------	------	------------	--------

	20674	1.3034	1.3871	1.2856	0.113	0.016	1.4872	1.5739	2.0249	0.0942	0.055
1	*: slope' was the corrected trend of changes in snow depth, the unit was cm yr ⁻¹ ; P' was the corrected confidence										
2	level.										
3											




Figure 1. Geographical locations of meteorological and snow course stations across the Eurasian continent. The red circles represent stations where snow depth was measured at both meteorological stations and snow course surveys, the green circles show stations where snow depth was measured at snow surveys only, and the blue circles show stations where snow depth was measured at meteorological stations only. The abbreviations of countries represented separately: ARM-Armenia, AZE-Azerbaijan, BLR-Belarus, EST-Estonia, GEO-Georgia, KAZ-Kazakhstan, KGZ-Kyrgyzstan, LTU-Lithuania, LVA-Latvia, MDA-Moldova, TJK-Tajikistan, TKM-Turkmenistan, UKR- Ukraine, UZB-Uzbekistan.






Figure 2. Annual mean snow depth (a) and maximum snow depth (b) across the Eurasian continent (cm) during 1966-2012.



1



1
2 **Figure 3.** Monthly mean snow depth (from September to June) (cm) across the Eurasian continent
3 (cm) during 1966-2012. (a) September, (b) October, (c) November, (d) December, (e) January, (f)
4 February, (g) March, (h) April, (i) May, (j) June.






Figure 4. Composite of inter-annual variation of annual mean snow depth (a) and maximum snow depth (b) from 1966 through 2012 with respect to the 1971-2000 mean across the Eurasian continent. The line with dots is the anomaly of snow depth; the thick curve represents the smoothed curve using wavelet analysis; the thick line presents a linear regression trend.



1


2 **Figure 5.** Composites of inter-annual variation of monthly mean snow depth (from October to

1 May) from 1966 through 2012 with respect to the 1971-2000 mean across the Eurasian continent.
2 (a) October, (b) November, (c) December, (d) January, (e) February, (f) March, (g) April, (h) May.
3 The line with dots is the anomaly of snow depth; the thick curve represents the smoothed curve
4 using wavelet analysis; the thick line presents a linear regression trend. Linear regression was only
5 shown when the rate of change was at the 95% level.
6



7
8 **Figure 6.** Spatial distribution of linear trend coefficients (cm yr⁻¹) of annual mean snow depth (a)
9 and maximum snow depth (b) for each station in 1966-2012. The rate of change was at the 95%
10 level. Red circles represent a decreasing trend, and blue circles represent an increasing trend.

1



2




Figure 7. Spatial distributions of linear trend coefficients (cm yr^{-1}) of monthly mean snow depth (from October to May) during 1966 to 2012. (a)October, (b) November, (c) December, (d) January, (e) February, (f) March, (g) April, (h) May. The rate of change was at the 95% level. Red circles represent a decreasing trend, and blue circles represent an increasing trend.





Figure 8. The relationship between annual mean snow depth and latitude (a), elevation (b and c) and continentality (d) for all stations across the Eurasian continent during 1966-2012. Asterisks show the mean snow depth of each station; the thick line is a linear regression trend.



1
2 **Figure 9.** The relationships among annual mean snow depth, air temperature and snowfall for 386
3 stations from November through March during 1966-2009 over the USSR. The thick line is a
4 linear regression trend.




Figure 10. Composite of inter-annual variation of annual mean air temperature (a), annual snowfall (b), annual heavy snowfall (c) and annual snow depth (d) from November through March during 1966-2009 with respect to the 1971-2000 mean across the former USSR. The line with dots is the composite of the annual means; the thick curve represents the smoothed curve using wavelet analysis; the thick line presents a linear regression trend.




Figure 11. Spatial distributions of partial correlation coefficients of snow depth and air temperature (a), snow depth and snowfall (b), SWE and air temperature (c), SWE and snowfall from November through March during 1966-2009 across the former USSR. The coefficients reaching to 0.95 confidence level are displayed. Red circles represent a negative relationship, and blue circles indicate a positive relationship.




Figure A1. Normal distribution test of annual mean snow depth for all station by K-S test.