

Interactive comment on “Brief Communication: Glaciers in the Hunza Catchment (Karakoram) are in balance since the 1970s” by Tobias Bolch et al.

Anonymous Referee #1

Received and published: 14 October 2016

General comments:

This study utilizes digital elevation models extracted from satellite imagery (historical Hexagon and modern SRTM and ASTER) to calculate a regional geodetic mass balance for glaciers in the Hunza River basin, Karakoram region, using DEM differencing. The authors show that given the uncertainties of the methodology, the regional geodetic mass balance is not statistically different from zero change, consistent with previous mass balance studies on shorter more recent timescales. Their results suggest that the so-called "Karakoram anomaly" is not limited to the past ~ 15 years, but extends back to at least 1973. This is the first study using elevation differences to confirm this finding over a several-decade timespan, which supports previous studies showing no significant changes in debris cover or glacier area in the Karakoram over similar (1970's-present) time periods. Overall, it is a nice paper, and is ready for publication

[Printer-friendly version](#)

[Discussion paper](#)



after a few minor additions.

A table of values showing the standard deviation of mean elevation change between the ASTER and SRTM DEMs for assumed stable (non-glacier) terrain is needed to better assess the relative vertical accuracy of the DEMs.

Regarding the satellite imagery datasets, a paragraph, table, or figure to clearly show which DEMs are being subtracted from one another for each given time period, i.e. SRTM minus Hexagon for 1973-1999, and ASTER minus SRTM for 1999-2009. This would serve to clarify the methods section significantly.

Since the primary motivation of the paper is to extend the geodetic mass balance record further back in time, I would recommend an additional calculation of the full timespan (1973 - 2009) mass balance. This would also serve to validate the 1973-1999 and 1999-2009 mass balances, and remove the significant uncertainty regarding SRTM penetration into the ice.

The equations used for estimating uncertainty lean toward the more conservative side (i.e. large error bars). For example, linearly adding up the errors in Eq. 3 instead of adding in quadrature, which assumes that the error components in Eq. 3 are completely correlated with one another. The authors should make clear in the conclusion of the manuscript - results show no statistical difference from zero change, given the somewhat large/conservative uncertainties used with the DEM differencing method.

Specific comments:

P3 L6 Were any glaciers covered only partially by scenes from different years? If so, it may be best to use a weighted mean (weighted by the percentage of a glacier's area covered by each scene).

P3 L8 It is still somewhat unclear to me how the Cartosat-1 data is being used. I assume the authors compute Cartosat minus SRTM, then compare to ASTER minus SRTM in order to check consistency between the datasets. This should be further

[Printer-friendly version](#)[Discussion paper](#)

clarified in the text.

P2 L18 "assuming a full penetration of the radar beam into snow..." - regarding the ablation region. What about additional penetration into the ice itself, is this taken into account?

P2 L20 It would be useful here to refer to the later section (3.2) so the reader can easily find the discussion regarding void filling with the ASTER GDEM2 and associated uncertainties.

P3 L24 "All stereo images have been processed with a RMS of $< \sim 1.5$ pixels." Which aspect of the stereo photogrammetry is this referring to? Is this the reprojection error of triangulated ground control points after bundle adjustment, or something to do with the reseau grid distortion removal, or something else? A more detailed explanation is needed to interpret the meaning.

P3 L28 See previous comment regarding P3 L24

P3 L32 What kind of spatial trend corrections were made? Rotation, translation, or perhaps polynomial surface corrections... if so are they first order (linear), or higher order polynomials, or some other method?

P4 L12 Was the outlier threshold applied to both Hexagon and ASTER data, or to Hexagon only? If no outlier filtering was needed for the ASTER DEMs, this should be stated explicitly in the text.

P4 L17 It would be helpful to know the percentage of total pixels excluded (using the outlier threshold filter) for each glacier, to ensure that no large regions were interpolated using the ordinary kriging; otherwise unrealistic elevations could result. The text later discusses the percentage of voids in the SRTM data, but says nothing regarding the percentage of data gaps in the Hexagon data.

P6 L11 "we confirm for the first time using elevation differences..." Should specify: over this longer time period (because elevation differences have been used over shorter

[Printer-friendly version](#)[Discussion paper](#)

time periods in previous studies).

P6 L13 Going back to the previous comments regarding P4 L12 and P4 L17 - Since both time periods use different data sources and therefore contain differing amounts of data voids - a percentage of voids for each would help eliminate doubts regarding direct comparisons between the two time periods, which use SRTM - Hexagon for 1973-1999, and ASTER - SRTM for 1999-2009. Could a difference in data gaps/holes make a significant difference when comparing these datasets/time periods?

P6 L17 What is meant by "different surge stages in the two periods..."? What is different between the two periods regarding surges, the magnitude, timing, or something else? Or is the word "different" simply being used in a fashion equivalent to "separate"?

P8 L1 When calculating the mass budget with the non-void-filled version of SRTM for comparison, were the voids interpolated, or was the mass balance computed using only the volume change of existing pixels, then divided by the glacier area only covered by existing pixels? More details would be helpful.

Technical corrections:

P2 L7 complicate the

P3 L10 "The major advantage of this dataset is besides the high spatial resolution and also the 12 bit pixel depth" – strange wording. Should change to something like: "The major advantages of this dataset are the high spatial resolution and 12-bit pixel depth."

P7 L11 "... ASTER DEMs where lower..." change "where" to "were"

P8 L6 voids

P8 L4 versions

Interactive comment on The Cryosphere Discuss., doi:10.5194/tc-2016-197, 2016.

Printer-friendly version

Discussion paper

