

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25

Rapid Wastage of the Hazen Plateau Ice Caps, Northeastern Ellesmere Island, Nunavut, Canada

Mark C. Serreze¹, Bruce Raup², Carsten Braun³, Douglas R. Hardy⁴ and Raymond S. Bradley⁴

¹Department of Geography, and Cooperative Institute for Research in Environmental Sciences,
University of Colorado, Boulder Colorado, USA

²Cooperative Institute for Research in Environmental Sciences, University of Colorado, Boulder Colorado,
USA

³Geography and Regional Planning / Environmental Science, Westfield State University, Westfield,
Massachusetts, USA

⁴Department of Geosciences, University of Massachusetts, Amherst Massachusetts, USA

26

Abstract

27 Two pairs of small stagnant ice bodies on the Hazen Plateau of northeastern Ellesmere Island, the St.
28 Patrick Bay ice caps and the Murray and Simmons ice caps, are rapidly shrinking, and the remnants of
29 the St. Patrick Bay ice caps are likely to disappear entirely within the next five years. Vertical aerial
30 photographs of these Little Ice Age relics taken during August of 1959 show that the larger of the St.
31 Patrick Bay ice caps had an area of 7.48 km², and the smaller one 2.93 km²; the Murray and Simmons
32 ice caps covered 4.37 km² and 7.45 km² respectively. Outlines determined from ASTER satellite data for
33 July 2016 show that, compared to 1959, the larger and the smaller of the St. Patrick Bay ice caps had
34 both been reduced to only 5% of their former area, with the Murray and Simmons ice caps faring better
35 at 39% and 25%, likely reflecting their higher elevation. Consistent with findings from other glaciological
36 studies in the Queen Elizabeth Islands, ASTER imagery in conjunction with past GPS surveys documents a
37 strikingly rapid wastage of the St. Patrick Bay ice caps over the last 15 years. These two ice caps shrank
38 noticeably even between 2014 and 2015, apparently in direct response to the especially warm summer
39 of 2015 over northeastern Ellesmere Island. The well-documented recession patterns of the Hazen
40 Plateau ice caps over the last 55+ years offer an opportunity to examine the processes of plant
41 recolonization of polar landscapes.

42 **Keywords:** Arctic, ice caps, mass balance, Little Ice Age, Hazen Plateau, Ellesmere Island, ASTER

43

44

1. Introduction

45 The Hazen Plateau of northeastern Ellesmere Island, Nunavut, Canada, is a rolling upland, with
46 elevations rising from about 300 meters above sea level near Lake Hazen to over 1000 m along the
47 northeast coast of the island. The plateau is unglaciated with the exception of two pairs of small
48 stagnant ice caps - the unofficially-named St. Patrick Bay ice caps, and, 110 km to the southwest, the
49 Murray and Simmons ice caps (**Figure 1**). They are collectively referred to here as the Hazen Plateau ice
50 caps. As of 2001, the larger St. Patrick Bay ice cap ranged in elevation between about 880 m and 720 m
51 above sea level, with the smaller one spanning 820 m to 700 m. The Murray and Simmons ice caps lie in
52 higher terrain; in 2001, both fell between about 1100 m and 1000 m above sea level. The Hazen Plateau
53 ice caps are interpreted as forming and attaining their maximum extents during the Little Ice Age (LIA, c.
54 1600-1850) (Koerner, 1989). Like much of the Queen Elizabeth Islands, the Hazen Plateau is presently a
55 polar desert; annual precipitation is typically only 150-200 mm, with a late summer and early autumn
56 maximum (Serreze and Barry, 2014). Summer precipitation may be variously rain or snow. Summers are
57 very cool but variable; assessed as part of a multiyear glaciological study (Braun et al., 2004), the
58 average 10 m July air temperature at the Murray ice cap summit (1100 m) measured for the years 1999
59 through 2001, respectively, was 4.0°C, 0.2°C and 1.6°C.

60 This paper documents the behavior of the Hazen Plateau ice caps over the past 55+ years in the context
61 of other glaciological studies in the Canadian Arctic. The analysis is based on a combination of past
62 work using aerial photography, direct mass balance measurements from several field investigations, and
63 GPS surveys of ice cap areas collected as part of these investigations – along with new information on

64 ice cap areas using data at 15 m resolution from the ASTER (Advanced Spaceborne Thermal Emission
65 and Reflection Radiometer) instrument. ASTER flies onboard the NASA's Earth Observing System Terra
66 satellite, launched in December 1999. It provides reflectance at a 15 m resolution and is a key asset of
67 the international GLIMS initiative (Global Land Ice Measurements from Space) for mapping glacier
68 outlines (Raup et al., 2007; Kargel et al., 2014).

69 **2. Previous Work**

70 Table 1 lists all available direct mass balance estimates of the ice caps (in meters water equivalent, or
71 w.e.). Table 2 provides all available estimates of ice cap areas (km²). The first information on the St.
72 Patrick Bay ice caps that we are aware of is oblique aerial photographs taken in late July of 1947 as part
73 of the U.S. Operation Polaris Trimetregon Survey. These photographs show the ice caps standing out
74 prominently against the snow-free tundra surface. Vertical aerial photographs collected by the Canada
75 Department of Energy, Mines and Resources followed in August of 1959. These photographs show
76 prominent, exposed surface dirt layers and stratigraphic layering on the St. Patrick Bay ice caps, and the
77 Murray and Simmons ice caps are also bare of snow. From digitizing the 1959 photographs and
78 mapping the ice cap outlines, the larger of the St. Patrick Bay ice caps then had an area of 7.48 km², and
79 the smaller one 2.94 km². The Murray and Simmons ice caps covered, respectively 4.37 and 7.45 km²
80 (Serreze, 1985; Bradley and Serreze, 1987; Braun et al., 2004). We estimate that these areas are
81 accurate to within 5%.

82 In July of 1972, Canadian scientists H. Serson and J. A. Morrison surveyed the larger of the two St.
83 Patrick Bay ice caps. They landed by helicopter in foul weather to find the ice cap totally covered with
84 snow. They installed eight accumulation stakes along a roughly two kilometer transect partway across
85 the ice cap. The range in elevation along this transect was about 60 m, which compares to a range for
86 the entire ice cap of about 160 m. Later that same summer, on August 20-21, the ice cap was visited by
87 G. Hattersley-Smith and A. Davidson, who noted a "partial cover of winter snow all around the ice
88 margins for at least a kilometer" (Hattersley-Smith and Serson, 1973), in striking contrast with
89 conditions depicted in the August 1947 and 1959 aerial photographs. They concluded that while the ice
90 cap had been in decline (as suggested from the 1947 and 1959 photographs), by the early 1970s it had
91 returned to good health, "thickening slightly and extending its margins" (icy firn was observed atop the
92 dirty melt surface and a perennial snow cover extended beyond the ice cap margins). This is consistent
93 with a known shift towards cooler summers and increased precipitation over the eastern Canadian
94 Arctic (Bradley and Miller, 1972; Bradley and England, 1978). Hattersley Smith and Serson estimated a
95 mass balance for the 1971/1972 season of +0.14 m w.e.

96 In 1982 and 1983, the St. Patrick Bay ice caps were the focus of detailed energy and mass balance
97 investigations (Serreze, 1985; Bradley and Serreze, 1987; Serreze and Bradley, 1987). The stake network
98 was expanded on the larger St. Patrick Bay ice cap and several stakes were installed on the smaller one.
99 At the end of the 1982 field season in early August, the entire ice cap was bare ice with a well-developed
100 cryoconite surface. Assuming that the 1982 melt season had largely ended by early August (all visible
101 melt had stopped by the time that the field camp had been evacuated), the 1981/1982 mass balance for
102 the larger ice cap was estimated at -0.14 m w.e.. Given that more melt may have occurred, this is likely

103 a minimum estimate. Based on the stake line installed in 1972, Bradley and Serreze (1987) estimated
 104 that the overall mass balance for the period 1972-1982 was approximately -1.3 m w.e. (-0.14 a⁻¹). This
 105 result finds qualitative support in comparisons between the 1959 aerial photographs and subsequent
 106 vertical aerial photographs taken on 1 August 1978 showing that the larger and smaller of the ice caps
 107 had decreased in area by 7% and 11% over that interval. Like the 1959 photographs, the August 1978
 108 photographs revealed a snow-free plateau and bare ice with a prominent ablation surface. Aerial
 109 photographs taken four years earlier, on 4 August 1974, showed broadly similar conditions. As part of
 110 the St. Patrick Bay Project, a network of stakes installed on the Simmons ice cap in 1976 (Bradley and
 111 England, 1977) was re-surveyed on 11 July 1983. Of the 18 original stakes, only 6 could be located; the
 112 others were presumed to have melted out. Based on these sparse data, Bradley and Serreze (1987)
 113 estimated that over the period 1976-1983, the Simmons ice cap experienced a total mass loss of at least
 114 -0.49 meters w.e. (-0.08 a⁻¹). Collectively, these observations provided strong evidence that the period
 115 of recovery inferred by Hattersley-Smith and Serson (1973) was short-lived.

116 However, the summer of 1983 was cool, and the snow never completely melted off the surrounding
 117 tundra. The 1982/1983 annual mass balance for the larger St. Patrick Bay ice cap was estimated at
 118 +0.14 m w.e. , and given their higher elevation, it is reasonable to assume that the 1982/1983 balance
 119 year for the Simmons and Murray ice caps was also positive.

Table 1. Directly measured mass balances (meter water equivalent) of the Hazen Plateau ice caps. Where a value represents a multiyear record, the average annual value is shown in parentheses. Asterisks denote minimum estimates.

Balance year or period	Large St. Patrick Bay	Small St. Patrick Bay	Murray	Simmons
1971/72	+0.14 ¹	-----	-----	-----
1971/72-1981/82	-1.3 ² (-0.14) ²	-----	-----	-----
1975/76-1982/83	-----	-----	-----	*-0.49 (-0.08) ²
1981/82	*-0.14 ²	-----	-----	-----
1982/83	+0.14 ²	-----	-----	-----
1983/84-1997/98	-----	-----	-----	*-0.49 (-0.03) ⁴
1983/84-1999/00	*-1.01 (-0.06) ³	*-1.26 ³ (-0.07) ³	-----	-----
1998/99	-----	-----	-0.49 ³	-----
1999/00	-----	-----	-0.29 ³	-0.40 ³
2000/01	-----	-----	-0.47 ³	-0.52 ³
2001/02	-----	-----	-0.29 ³	-----

Sources: ¹Hattersley-Smith and Serson (1973); ²Bradley and Serreze (1987); ³Braun et al. (2004)

120

121 To our knowledge, there were no further visits to the Hazen Plateau ice caps until 1999, when C. Braun,
 122 D. Hardy and R. Bradley of the University of Massachusetts Amherst established a network of 11
 123 accumulation stakes on the Murray Ice Cap, which they further expanded in the year 2000. A new
 124 network of 15 stakes was established on the Simmons ice cap in 2000. Winter snow accumulation was

125 measured on both ice caps in late May of 1999 through 2001, and summer ablation was measured in
 126 late July and early August from 1999-2002. For the four years analyzed, 1999-2002, annual balances of
 127 both ice caps were negative in all years, ranging from -0.29 m w.e. (Murray ice cap in 2000) to -0.52 m
 128 w.e. (Simmons ice cap in 2001). In the summer of 2001, C. Braun and D. Hardy used portable GPS to
 129 survey the perimeter of all four ice caps. The larger and smaller of the St. Patrick Bay ice caps had shrunk
 130 to 62% and 59% of their 1959 areas, respectively. The Murray and Simmons ice caps had shrunk to 70%
 131 and 53% of their 1959 areas. Some of the accumulation stakes inserted into the larger St. Patrick Bay
 132 ice cap in 1982 and 1983 were located but all had melted out. Knowing how deep they had been
 133 originally inserted enabled a minimum estimate (-1.01 m w.e., -0.06 a⁻¹) of the mass loss between 1984
 134 and 2000. This is based on the mean remaining depth of stake insertion into the ice in 1983 and an
 135 assumed ice density of 900 kg m³ (Braun et al., 2004). In the late summer of 2003, C. Braun mapped the
 136 margins of the Murray and Simmons ice caps via portable GPS by holding the device out the window of a
 137 low-flying helicopter. The same approach was used to assess the ice cap margins in 2006, this time by
 138 University of Massachusetts graduate student T. Cook.

139 3. Updated History, 1959 to 2016

140 3.1 Ice Cap Areas

141 The use of ASTER in conjunction with the air photographs and GPS surveys enables a fairly detailed
 142 assessment of changes in ice cap areas from 1959 through the present. Clear-sky late summer (July or
 143 August) scenes of the St. Patrick Bay ice caps showing a strong brightness contrast between the ice and
 144 the bare, dark plateau surface, enabled manual mapping of the ice cap perimeters from ASTER for the
 145 years 2005, 2009, 2014, 2015 and 2016. For the Murray and Simmons ice caps, ASTER estimates were
 146 obtained for 2001, 2007 and 2016. For 2001, areas of the Murray and Simmons were available from
 147 both ASTER and the surface-based GPS surveys. Considering the GPS surveys for this year as ground
 148 truth, the ASTER areas for this year are accurate to within 1% for the Murray ice cap and 3% for the
 149 Simmons ice cap. It is assumed that this is representative of the accuracy of area mapping from ASTER
 150 for the other years.

151 As of July 2016, and Murray and Simmons ice caps cover 39% and 25% of the areas in 1959 based on the
 152 aerial photographs. By sharp contrast, both of the St. Patrick Bay ice caps in 2016 cover only 5% of their
 153 former areas, and have been reduced to ice patches, with the smaller ice body now covering only 0.15
 154 km².

	Larger St. Patrick Bay Ice Cap Area and % of 1959		Smaller St. Patrick Bay Ice Cap Area and % of 1959		Murray Ice Cap Area and % of 1959		Simmons Ice Cap Area and % of 1959	
1959	7.48 ¹	100%	2.94 ¹	100%	4.37 ¹	100%	7.45 ¹	100%
1978	6.69 ¹	89%	2.74 ¹	93%	-----	-----	-----	-----
1999	-----	-----	-----	-----	3.28 ²	75%	-----	-----
2000	-----	-----	-----	-----	3.15 ²	72%	-----	-----

2001	4.61 ²	62%	1.72 ²	58%	3.05 ² (3.08 ⁴)	70%	3.94 ² (3.83 ⁴)	53%
2003	-----	-----	-----	-----	2.91 ³	66%	3.31 ³	44%
2005	3.68 ⁴	49%	1.03 ⁴	35%	-----	-----	-----	-----
2006	-----	-----	-----	-----	2.86 ³	65%	3.19 ³	43%
2007	-----	-----	-----	-----	2.76 ⁴	63%	2.92 ⁴	39%
2009	2.54 ⁴	34%	0.63 ⁴	21%	-----	-----	-----	-----
2014	1.03 ⁴	14%	0.29 ⁴	10%	-----	-----	-----	-----
2015	0.52 ⁴	7%	0.18 ⁴	6%	-----	-----	-----	-----
2016	0.35 ⁴	5%	0.15 ⁴	5%	1.72 ⁴	39%	1.84 ⁴	25%
¹ Aerial photographs, ² Surface GPS surveys, <i>Braun et al. (2004)</i> , ³ GPS helicopter surveys, ⁴ ASTER								

155

156 Outlines of the St. Patrick Bay ice caps for 1959 from aerial photography, for 2001 from GPS surveys, and
 157 for 2014 and 2015 from ASTER are shown in **Figure 2**. The reductions in ice cap area are striking. Note
 158 the obvious shrinkage even between the years 2014 and 2015. Shrinkage of the Murray and Simmons
 159 ice caps is shown in **Figure 3**, based on outlines from 1959, 2001 and 2016. The shrinkage of these two
 160 ice caps is clearly evident, albeit less pronounced.

161 Using the area estimates through 2002 and extrapolating forward, Braun et al. (2004) suggested that the
 162 Hazen Plateau ice caps would disappear by the middle of the 21st century or soon thereafter, and that,
 163 given their larger size, the Simmons ice cap and the larger of the two St. Patrick Bay ice caps would be
 164 the last to go. However, based on data through 2016 and extrapolating forward (**Figure 4**), it now
 165 appears that both of the St. Patrick Bay ice caps will disappear around the year 2020.

166 From the analyses described above, and results from other glaciological investigations for the Canadian
 167 Arctic and the Arctic as a whole, the following conclusions are drawn:

- 168 • The Hazen Plateau ice caps are unlikely to be relics of the last glacial maximum, but rather likely
 169 formed during the Little Ice Age (LIA, c. 1600-1850) (Koerner, 1989). They may have retained
 170 their LIA extents through the first couple decades of the 20th century (Hattersley-Smith, 1969;
 171 Sharp et al., 2014), but have been in overall decline ever since. Braun et al. (2004) speculate on
 172 the basis of a mapped lichen trim line that the Murray ice cap may have attained a maximum LIA
 173 extent of about 9.6 km², over twice the mapped 1959 area of 4.35 km². Similar trim lines were
 174 observed around the other three ice caps and although not mapped in detail, strongly point to
 175 much more extensive ice cover during the LIA. To place these findings in a broader context, for
 176 the Queen Elizabeth Islands as a whole, trim lines based on high-resolution satellite imagery
 177 point to a 37% reduction in perennial snow and ice extent between the LIA maximum extent and
 178 the year 1960. Over the lower-lying central and western islands, a complete removal of
 179 perennial snow and ice occurred by 1960 (Wolken et al., 2008).
- 180 • From the 1960s through part of the 1970s, the ice caps may have experienced a period of
 181 reduced loss or occasional growth (1971/1972, 1982/1983) in response to cooling. This basic
 182 pattern likely holds for monitored Canadian Arctic glaciers and ice caps as a whole (Bradley and

183 Miller, 1972; Hattersley-Smith and Serson, 1973; Ommanney, 1977; Bradley and England, 1978;
184 Braun et al., 2004; Sharp et al., 2014).

185 • Since then, apart from occasional years such as 1982/1983, annual mass balances of the four ice
186 caps have been persistently negative (Braun et al., 2004). This is in turn consistent with the
187 broader pattern of reductions in mass and area of Arctic glaciers and ice caps (Dowdeswell et
188 al., 1997; Dyurgerov and Meier, 1997; Arendt et al., 2002; Koerner, 2005; Sharp et al., 2011;
189 Fisher et al., 2012; Sharp et al., 2014; Mortimer et al., 2016). It is also consistent with a negative
190 mass balance of the Greenland Ice Sheet since at least the 1990s (Shepard et al., 2012).

191 Mass balance summaries for four monitored glaciers and ice caps in the Canadian Arctic (Devon Ice
192 Cap, Meighan Ice Cap, Melville South Ice Cap and the White Glacier) are provided as part of the
193 American Meteorological Society (AMS) State of the Climate reports. As assessed over the period
194 1980 through 2010, all four have had negative average annual mass balances, ranging from -0.15 m
195 w.e. for the Devon Ice Cap to -0.29 w.e. for the Melville South Ice Cap (AMS, 2016). Cumulative
196 changes in regional total stored water for the period 2003 through 2015 based on gravimetric data
197 from the GRACE mission (Gravity Recovery and Climate Experiment) are qualitatively consistent with
198 these mass balance measurements (AMS, 2016). Based on ice core data, Fisher et al. (2012)
199 document rapid acceleration of ice cap melt rates of over the last few decades across the entire
200 Canadian Arctic; the large reductions in area of the Hazen Plateau ice caps, in particular the lower-
201 elevation St. Patrick Bay ice caps, is consistent with this finding. However, reflecting variable climate
202 conditions, annual balances are also quite variable. For example, for the 2013/2014 balance year
203 (the most recent data available), the White Glacier had a strongly negative balance (-0.42 m w.e.)
204 while the small Meighan Ice Cap actually gained mass (+0.06) (AMS, 2016). Sharp et al. (2014) show
205 that while the larger ice bodies in the Canadian Arctic have seen the largest losses in mass, the
206 smaller masses have lost a larger proportion on their areas. This is also consistent with the behavior
207 of the Hazen Plateau ice caps. Below we examine variability in climate conditions over the Hazen
208 Plateau, and links to mass balance and area changes.

209 3.2 Associated Climate Conditions

210 The annual mass balance of low-accumulation ice caps and glaciers in the Canadian High Arctic is known
211 to be primarily governed by summer warmth rather than winter accumulation (e.g., Bradley and
212 England, 1978; Koerner, 2005). To place the behavior of the Hazen Plateau ice caps in a climate context,
213 use is made of summer-averaged (June through August) 850 hectopascal (hPa) temperature anomalies
214 from the radiosonde record at Alert, located on the northeast coast of Ellesmere Island (Figure 1) along
215 with estimated summer temperature anomalies for the LIA. The Alert radiosonde record extends back
216 to 1957. We use monthly mean records contained in the Integrated Global Radiosonde Archive (IGRA,
217 Durre et al., 2006), based on daily 00 and 12 UTC soundings. Summer averages (J,J,A) were eliminated if
218 based on fewer than 70 values. The 850 hPa level is about 1400 m above sea level for a standard
219 atmosphere, hence roughly 600-700 m above the surface in the vicinity of the St. Patrick Bay ice caps
220 and 300-400 m above the surface in the vicinity of the Murray and Simmons ice caps. While arguably it
221 might be better to look at the 925 hPa level as it is closer to the plateau surface, this level has many

222 missing values in the IGRA record. The time series of anomalies, computed with respect to the standard
223 averaging period 1981-2010, follows in **Figure 5**.

224 Kaufman et al. (2010) took advantage of a variety of proxy sources (e.g., tree rings, ice cores, lake cores)
225 to assemble a record of Arctic summer surface temperature anomalies that extend back 2000 years.
226 From their analysis, LIA summer Arctic temperatures anomalies averaged around -0.6°C with respect to
227 a 1961-1990 reference period. The 1961-1990 summer mean of -3.2°C from the radiosonde data
228 compares to a mean of -2.6°C for 1981-2000. The latter period is hence about 0.6°C warmer. With the
229 assumption that (a) temperature anomalies at 850 hPa have been similar to those at the surface
230 (supported by Sharp et al. (2011) in their analysis of Canadian Arctic ice caps but possibly complicated by
231 the temperature inversion structure), and (b) LIA conditions over the Hazen Plateau were at least
232 broadly similar to those for the Arctic as a whole, these results imply that Arctic LIA temperature
233 anomalies were about -1.2°C relative to a 1981-2000 baseline. This estimated LIA temperature anomaly
234 is shown in Figure 5 as a dashed line.

235 If it is also accepted that the ice caps were broadly in equilibrium with average LIA summer
236 temperatures, Figure 5 suggests generally strong negative annual balances from the beginning of the
237 record through the early 1960s. This was followed by smaller negative and occasionally positive annual
238 balances from the middle of the 1960s through about 2000, and a preponderance of strong negative
239 balances from the beginning of the century through the present. For comparison with the radiosonde
240 record, we also examined 850 hPa summer temperatures over the Hazen Plateau from the National
241 Centers for Environmental Prediction/National Center for Atmospheric Research (NCEP/NCAR)
242 reanalysis (Kalnay et al., 1996) which extend back to 1948. Given that the Alert radiosonde data are
243 assimilated into the reanalysis, it follows that the radiosonde and NCEP/NCAR time series look similar
244 for the period of overlap. The NCEP/NCAR records suggest that the period 1948-1956 not covered by
245 the IGRA record was warm overall with mostly positive anomalies relative to the 1981-2000 baseline.
246 The cooling between the late 1940s through the middle 1960s broadly corresponds to the cooling over
247 the eastern Canadian Arctic such as discussed by Bradley and Miller (1972), Bradley and England (1978)
248 and other studies. The time series of decadal mean summer temperatures at the 700 hPa level for the
249 major glaciated regions of the Canadian Arctic presented by Sharp et al. (2011) based on the
250 NCEP/NCAR reanalysis (their Figure 9.3) is also consistent with the pattern shown in Figure 5.

251 An examination of selected individual years is instructive. Based on summer 1957 temperatures, the
252 1956/1957 annual balance must have been strongly negative. The same can be said for 1959/1960 and
253 1962/1963. By sharp contrast, the summer of 1972, when Hattersley-Smith and Serson (1973) visited
254 the ice caps and remarked upon the extensive August snow cover over the plateau and estimated a
255 positive balance of $+0.14$ m w.e. (for the 1971/1972 season), was the coldest in the radiosonde record,
256 and about 2°C below the estimated LIA average. There is also a clear contrast between 1982 (a known
257 negative annual balance year for the St. Patrick Bay ice caps) and 1983 (a known positive balance year,
258 with summer 850 hPa temperatures slightly below the LIA average). Given the low temperatures for the
259 summer of 1992, which followed the 1991 eruption of Mt. Pinatubo, the balance for 1991/1992 was
260 likely positive. In the sense that summer temperatures were above the estimated LIA average, negative
261 balances for the Murray ice caps for 1998/1999 through 2001/2002 (Braun et al., 2004) are all

262 consistent with Figure 5. Note however that the largest negative balance of -0.49 m w.e. for 1999/2000
263 corresponds to the coldest of the four summers, arguing for influences of local effects on summer
264 temperature or perhaps a low winter accumulation.

265 Regarding the summer of 2013, the obvious exception to the pattern of recent warm years, the ASTER
266 data and daily images from the Moderate Resolution Imaging Spectroradiometer (MODIS) show
267 extensive cloud cover through the summer, making it difficult to determine whether the snow cover
268 ever entirely cleared off the plateau. It is likely, however, that the 2012/2013 balance year was positive
269 for the Hazen Plateau ice caps - the Devon Ice Cap, Meighen Ice Cap and the White Glacier all gained
270 mass. Only the Melville South Ice Cap, lying well to the west, had a negative balance (AMS, 2014).
271 Consistent with this view, **Figure 6** shows that summer (J,J,A) averaged 850 hPa temperature anomalies
272 over the Queen Elizabeth Islands from the NCEP/NCAR reanalysis were about 2°C below the 1981-2010
273 baseline in the area centered over Axel Heiberg and Ellesmere islands. This reflects the influence of an
274 unusually-deep circumpolar vortex at the 500 hPa level, centered just south of the Pole along about
275 90°W longitude. By sharp contrast, the notable area reduction of the St. Patrick Bay ice caps between
276 August 2014 and 2015 aligns with the very warm summer of 2015, essentially tied with 1957 as the
277 highest in the record. From **Figure 7**, July 2015 temperatures at the 850 hPa level from the NCEP/NCAR
278 reanalysis were 3-4°C above the 1981-2010 baseline over most of northeastern Ellesmere Island. Mass
279 balance estimates for monitored glaciers in the Queen Elizabeth Islands for the 2014/2015 season that
280 would provide context were not available us at the time that this paper came to press.

281 **4. Conclusions**

282 Regarding accelerating wastage of the St. Patrick Bay ice caps since the dawn of the 21st century, the
283 outsized warming of the Arctic in recent decades compared to the rest of the Northern Hemisphere
284 (termed Arctic Amplification), is overall most strongly expressed during the cold season, and is not
285 nearly as prominent in summer (Serreze and Barry, 2011). Nevertheless, from the NASA Goddard
286 Institute for Space Sciences (GISS) analysis (<http://data.giss.nasa.gov/gistemp/>), the trend in July surface
287 air temperatures over Northeastern Ellesmere Island over the period 1960-2015 is about 2°C (expressed
288 as a total change) which stands out compared to the rest of the Arctic. On the basis of a satellite-
289 derived record (from MODIS) of summer land surface temperatures, the more recent period of 2000 to
290 2015 has seen an average warming rate over the Queen Elizabeth Islands of 0.06°C per year, or a total of
291 nearly 1.0°C, most of this occurring between 2005 and 2012 (Mortimer et al., 2016). They associate this
292 warming with increasingly negative mass balances for glaciers and ice caps in the region. However,
293 conditions over the Hazen Plateau are highly variable, and the summers of 1957, 1960 and 1963 were
294 almost as warm as those seen in 2015, and the summer of 2013 was quite cool, very likely resulting in a
295 positive balance for 2012/2013.

296 Rapid wastage of the St. Patrick Bay ice caps over the past 15 years likely also reflects a reduction in
297 summer albedo, as dirt layers become progressively exposed and accumulate at the surface. During the
298 1982 and 1983 field campaigns, it was observed that summer precipitation over the ice caps was
299 typically in the form of snow, temporarily increasing the surface albedo and adding some mass. The
300 frequency of summer snowfall has likely declined in the (generally) sharply warming climate over the

301 past 15 years. Also, as suggested from the prominent decline in the area of the larger St. Patrick Bay ice
302 cap between 2014 and 2015, when there is an especially warm summer, the thin collar of ice at the ice
303 cap margins (a feature evident in field observations) will be prone to completely melting. The less
304 pronounced area reduction of the Murray and Simmons ice caps must partly be due to their higher
305 elevation and relatively cooler summer conditions. However, the elevation difference is only about 200-
306 300 m, which argues that the stronger response of the St. Patrick Bay ice caps to warming may also be
307 related to ice thickness. Regional differences in the temperature lapse rate (notably the temperature
308 inversion structure) could also be involved.

309 It is possible that the Hazen Plateau caps could see some temporary recovery given the large natural
310 variability in the Arctic. However, as noted by Alt (1978) and Bradley and England (1978), for stagnant
311 ice caps such as these, all it takes is one warm summer to erase any accumulated mass gains of a
312 previous decade. Assessing variability and trends in Arctic precipitation is notoriously difficult, but as
313 evaluated over the period 1950-2007, annual precipitation has generally increased across Canada, and
314 especially across Northern Canada. For example, at station Eureka in central Ellesmere Island (see
315 Figure 1), annual precipitation appears to have increased by at least 40% (Zhang et al., 2008). Trends
316 over the plateau are not known, but this suggests that, if anything, precipitation changes are helping to
317 buffer the ice caps from summer mass loss.

318 Paradoxically, perhaps, loss of the Hazen Plateau ice caps may open new research opportunities. As they
319 recede, plant remains are exposed that can be dated and used to better understand the past climate
320 history of the region. From radiocarbon dates on rooted tundra plants exposed by receding cold-based
321 ice caps on Baffin Island – given that the plants are killed when the snowline drops below the collection
322 sites – Miller et al. (2013) were able to construct a record of summer temperatures over Arctic Canada
323 for the past 5000 years. La Farge et al. (2013) discovered that ice loss in Sverdrup Pass, Ellesmere Island,
324 has exposed nearly intact plant communities for which radiocarbon dates point to entombment during
325 the LIA. They also found that these recently-exposed, subglacial bryophytes can regenerate, which may
326 have important implications for recolonization of polar landscapes. The area surrounding the receding
327 Hazen Plateau ice caps provides a unique opportunity to examine this process of recolonization in the
328 High Arctic, as the rates of ice recession are now well-documented for the last 55+ years (Table 1).

329 **Author Contribution:** M. Serreze led the overall effort. C. Braun, D. Hardy, and R.S. Bradley provided
330 GPS data and historical documents. B. Raup analyzed the ASTER data. All authors contributed to the
331 writing.

332 **Data Availability:** Radiosonde data for station Alert are available at from the Integrated Global
333 Radiosonde Archive (<ftp://ftp.ncdc.noaa.gov/pub/data/igra/>). ASTER data can be obtained through the
334 NASA Land Processes DAAC (<https://lpdaac.usgs.gov/>).

335 **Acknowledgements:** This study was supported by the University of Colorado Boulder, the NASA Snow
336 and Ice DAAC award NNG13HQ033 to the University of Colorado, and NSF Award 9819362 to the
337 University of Massachusetts.

338 **Competing Interests:** None

339

References

- 340 Alt, B.: Synoptic climate controls of mass balance variations on Devon Island Ice Cap, *Arc. Alp. Res.*, 10,
341 61-80, 1978.
- 342
- 343 American Meteorological Society: State of the Climate in 2014, J.S. Blunden and D.S. Arndt (eds.),
344 Special Supplement to the Bulletin of the American Meteorological Society, 96, 267 p.
- 345
- 346 American Meteorological Society: State of the Climate in 2015, J.S. Blunden and D.S. Arndt (eds.),
347 Special Supplement to the Bulletin of the American Meteorological Society, 97, 275 p.
- 348
- 349 Arendt, A. A., Echelmeyer, K. A., Harrison, W. D., et al.: Rapid wastage of Alaska Glaciers and their
350 contribution to rising sea level, *Science*, 297, 382-385, 2002.
- 351
- 352 Bradley, R. S., and England, J.: The Simmonds Ice Cap, in: R.S. Bradley and J. England, eds., *Past Glacial*
353 *Activity in the High Arctic*, University of Massachusetts Amherst, Department of Geology and
354 *Geography*, 177-182, 1977.
- 355 Bradley, R. S., and England, J.: Recent climatic fluctuations of the Canadian High Arctic and their
356 significance for glaciology, *Arc. Alp. Res.*, 10, 715-731, 1978.
- 357 Bradley, R. J., and Miller, G. H.: Recent climatic change and increased glacierization in the eastern
358 Canadian Arctic, *Nature*, 237, 385-387, 1972.
- 359 Bradley, R. S., and Serreze, M. C.: Mass balance of two high Arctic plateau ice caps, *J. Glaciol.*, 33, 123-
360 128, 1987.
- 361 Braun, C., Hardy, D. R., and Bradley, R. S.: Mass balance and area changes of four high Arctic plateau ice
362 caps, *Geografiska Annaler*, 86A, 43-52, 2004.
- 363 Dowdeswell, J. A., Hagen, J. O., Bjornsson, H., Glazovsky, A. F., Harrison, W. D., Holmlund, P., et al.: The
364 mass balance of circum-Arctic glaciers and recent climate change, *Quat. Res.*, 48, 1-14, 1997.
- 365 Durre, I., Vose, R. S., and Wuetz, D.B.: Overview of the integrated global radiosonde archive, *J. Clim.*,
366 19, 53-68, 2006.
- 367 Dyurgerov, M. B., and Meier, M. F.: Year-to-year fluctuation of global mass balance of small glaciers and
368 their contribution to sea level changes, *Arc. Alp. Res.*, 29, 392-402, 1997.
- 369 Fisher, D., Zheng, J., Burgess, D., Zdanowicz, C., Kinnard, C., Sharp, M., Bourgeois, J.: Recent melt rates of
370 Canadian arctic ice caps are the highest in four millennia, *Global and Planetary Change*, 84-85,
371 3-7, doi: 10.1016/j.gloplacha.2011.06.005, 2012.
- 372 Gardner, A. S., Moholdt, G., Wouters, B., Wolken, G. J., Burgess, D. O., Sharp, M. J., Cogley, J. G., Braun,
373 C., and Labine, C.: Sharp acceleration of mass loss from Canadian Arctic Archipelago glaciers and
374 ice caps, *Nature*, 473, 357-360, 2011.

- 375 Hattersley-Smith, G.: Glacial features of Tanquary Fiord and adjoining areas of Ellesmere Island, N.W.T.,
376 J. Glaciol., 8, 23-50, 1969.
- 377 Hattersley-Smith, G., and Serson, H.: Reconnaissance of a small ice cap near St. Patrick Bay, Robeson
378 Channel, Northern Ellesmere Island, Canada, J. Glaciol., 12, 417-421, 1973.
- 379 Kalnay, E., Kanamitsu, M., Kistler, R., et al.: The NCEP/NCAR 40-year re-analysis project, Bull. Amer. Met.
380 Soc., 77, 437-471, 1996.
- 381 Kargel, J. S., Leonard, G. J., Bishop, M. P., Kaab, A., and Raup, B. (Eds.): Global Land Ice Measurements
382 from Space, Springer-Praxis, 33 chapters, 876 pages. ISBN: 978-3-540-79817-0, 2014.
- 383 Kaufman, D. S., Schneider, D. P., McKay, N. P., et al.: Recent warming reverses long-term Arctic cooling,
384 *Science*, 325, 1236-1239, doi: 10.1126/science.1173983, 2009.
- 385 Koerner, R. M.: Queen Elizabeth Island Glaciers: In: Fulton, R.J. (ed.), Quaternary Geology of Canada and
386 Greenland. Geology of Canada (No. 1), Geological Survey of Canada, 464-477, 1989.
- 387 Koerner, R. M.: Mass balance of glaciers in the Queen Elizabeth Islands, Nunavut, Canada, Ann. Glaciol.,
388 42, 417-423, doi: 10.3189/172756405781813122, 2005.
- 389 La Farge, C., Williams, K. H., and England, J. H.: Regeneration of Little Ice Age bryophytes emerging from
390 a polar glaciers with implications of totipotency in extreme environments, Proc. Nat. Acad. Sci.,
391 110, 9839-9844, do: 10.1072/pnas.1304199110, 2013.
- 392 Miller, G. H., Lehman, S. J., Refsnider, K. A., Southon, J. R., and Zhong, Y.: Unprecedented recent
393 summer warmth in Arctic Canada, Geophys. Res. Lett., 40, 5745-5751,
394 doi:10.1002/2013GL0571888, 2013.
- 395 Mortimer, C. A., Sharp, M., and Wouters, B.: Glacier surface temperatures in the Canadian high Arctic,
396 2000-2015, J. Glaciol., doi: 10.1017/jog.2016.80, 2016.
- 397 Ommanney, C. S. L.: Quadrennial Report to the Permanent Service on the Fluctuations of Glaciers on
398 Canadian Glacier Variations and Mass Balance Changes, Fisheries and Environment Canada,
399 Inland Waters Directorate, Glaciology Divisions, Ottawa, 1977.
- 400 Raup, B. H., Racoviteanu, A., Khalsa, S. J. S., Helm, C., Armstrong, R., and Arnaud, Y.: The GLIMS
401 geospatial glacier database: a new tool for studying glacier change, Glob. Planet. Change, 56,
402 101-110, doi:10.1016/j.gloplacha.2006.07.018, 2007.
- 403 Sharp, M., Burgess, D.O., Cawkwell, F., et al.: Remote sensing of recent glacier changes in the Canadian
404 Arctic, Chapter 9 in: J.S. Kargel et al. (eds.), Global Land Ice Measurements from Space, Springer
405 Praxis Books, 205-228, doi: 10.1007/978-3-540-79818-7_9, 2014.

406 Sharp, M., Burgess, D. O., Cogley, J. G., Ecclestone, M., Labine, C., and Wolken, G. J.: Extreme melt on
407 Canada's Arctic ice caps in the 21st century, *Geophys. Res. Lett.*, 38, L11501,
408 doi:10.1029/2011GL047381, 2011.

409 Serreze, M. C.: Topoclimatic Investigations of a Small, Sub-Polar Ice Cap with Implications for
410 Glacierization, Masters Thesis, Department of Geology and Geography, University of
411 Massachusetts, Amherst, 201 p., 1985.

412 Serreze, M. C., and Barry, R. G.: Processes and impacts of Arctic amplification: A research synthesis,
413 *Glob. Plan. Change*, 77, 85-96, 2011.

414 Serreze, M. C. ,and Barry, R. G.: The Arctic Climate System, Second Edition, Cambridge University Press,
415 404 p., 2014.

416 Serreze, M. C.,and Bradley, R. S.: Radiation and cloud observations on a high Arctic plateau ice cap, *J.*
417 *Glaciol.*, 33, 162-168, 1987.


418 Shepherd, A., Ivins, E. R., Valentina, G. A., et al.: A reconciled estimate of ice-sheet mass balance,
419 *Science*, 338, doi: 10.1126/science.1228102, 2012.

420 Wolken, G.J., Sharp, M.L., and England, J.H.: Changes in late-Neoglacial perennial snow/ice extent and
421 equilibrium-line altitudes in the Queen Elizabeth Islands, Arctic Canada, *The Holocene*, 18, 615-
422 627, 2008.

423 Zhang, X., Brown, R., Vincent, L., Skinner, W., Feng, Y., and Mekis, E.: Canadian climate trends, 1950-
424 2007. Canadian Biodiversity: Ecosystem Status and Trends 2010, Technical Thematic Report No.
425 5. Canadian Councils of Resource Ministers. Ottawa, ON. iv + 21 p.,
426 <http://www.biodivcanada.ca/default.asp?lang=En&n=137E1147-0>, 2011.

427


428



429

430 Figure 1. The location of the St. Patrick Bay (STPBIC), Murray (MIC) and Simmons (SIC) ice caps. The
431 inset map shows Ellesmere Island (EL), Axel Heiberg Island (AHI), Greenland (GL), the Arctic Ocean (AO)
432 and station Alerts (ALR) and Eureka (EUR). Use is made of 850 hPa temperature data from the Alert
433 radiosonde record and precipitation records from Eureka.


434



435

436 Figure 2: Outlines of the St. Patrick Bay ice caps based on aerial photography from August 1959, GPS
 437 surveys conducted during August 2001, and for August of 2014 and 2015 from ASTER. The base image is
 438 from August 2015.

439




440

441 Figure 3: Outlines of the Murray and Simmons ice caps based on aerial photography from August 1959,
442 GPS surveys conducted during August 2001, and for July 2016 from ASTER. The base image is from
443 August 2016.

444

445


446




447

448 Figure 4. Time history of ice cap areas and projected times of disappearance.

449




450
 451 Figure 5. Temperature anomalies at the 850 hPa level ($^{\circ}\text{C}$) over the period 1957-2016 (referenced to the
 452 period 1981-2010) from the Alert radiosonde record. The horizontal black line shows the estimated
 453 summer average Arctic temperature anomaly for the LIA relative to 1981-2010 (-1.2°C). Also shown are
 454 the annual mass balance estimates for the larger St. Patrick Bay (SBP) ice cap for the 1981/1982 and
 455 1982/1983 balance years, and for the Murray Ice Cap (M) for the 1999/2000, 2000/2001, 2001/2002
 456 and 2002/2003 balance years (in m w.e.).
 457



458

459

460 Figure 6. Summer (J,J,A) 2013 air temperature anomalies at the 850 hPa level from the NCEP/NCAR
461 reanalysis relative to a 1981-2010 baseline.



462

463

464 Figure 7. July 2015 air temperature anomalies at the 850 hPa level from the NCEP/NCAR reanalysis
465 relative to a 1981-2010 baseline.

466

467

468

469

470