

Innsbruck, 21/03/2017

Dear Jean-Luis Tison,

Thank you for your useful comments and suggestions. I made all the changes required (see list of changes attached) and I also decided to drop the section 5.5. as suggested. As a consequence also the title and the abstract changed slightly (see marked up version attached below).

Finally, I would like to thank you for your patience and efforts as Editor and for giving us the possibility to publish our work in The Cryosphere.

I wish you all the best for your trip to Antarctica!

Sincerely,
Daniela Festi



LIST OF CHANGES

Editors comments in red.

Authors reply in black.

PAGE 2

L236-39: Drop these lines. This remains a weakness of the paper. I would like you to delete it.

See further⁵¹²

Authors: Done.

PAGE 5

L12-13..and in related figure: I would choose another color of these inversions in the graph to make these pop up clearly

Authors: Done. We used empty squares. Fig. capture has been updated as well.

PAGE 6

L25: delete "." at end of sentence

Authors: Done.

PAGE 7

L 2: blank missing before "provide"

Authors: Done.

L 4: change to: "the new method shows clear improvements at various levels"

Authors: Done.

L8: Change to: "For example, in the year 2005..."

Authors: Done.

L16: delete "etc."

Authors: Done.

L31replace "pollen dating" by "pollen analysis"

Authors: Done.

L34 delete sign before "This"

Authors: Done.

L36 delete sign before "the modelling"

Authors: Done.

L36 and 37Change to: "the modelling approach appears globally robust following validation..."

Authors: Done.

L37 to 39This is important. It deserves another insert plot in Figure 3, of Eismodel vs. Field validation

Authors: Since it was very hard to insert a further graph into Fig 3 we added a Figure 4 containing the 2 scatter plots (the one previously included in Fig 3 (comparison EISmodel and pollen dates) and the one here required.

PAGE 8

L3: Change to: "Such efficiency, however, cannot.."

Authors: Done.

L4 and 5: This is making the "depth-to-date" application more problematic then.. I think it should be mentioned

Authors: We added a sentence. I agree that if there is such a delay it must be considered, quantified and integrated in the model.

L11 and 12: change to: "helps in in the interpretation...inversions (specific color dots in Figure 2b).

Authors: Done.

L 13: Change to: " can potentially be of two origins: a) there might be...artefact.."

Authors: Done.

L 15: Change to: "accuracy; b) on the other hand, "

Authors: Done.

L17: Change to: ".. and snow: redistribution of..., and mixing..."

Authors: Done.

L20: Change to: "..in section 5.3"

Authors: Done.

L36: There are discordances between the depths of "ice lenses" in figure 2d and "meltlayers" in Figure 3! Needs consistency!

Authors: They are not discordances. In temperate firn ice lenses are not expected to form only in correspondence of meltlayers but also as a result of meltwater percolating from the surface and refreezing over internal discontinuities of the snowpack, such as sun crusts or wind crusts.

L41: delete "completely"

Authors: Done.

PAGE 9

L7:Also discordances between depths of "ice lenses" and "meltlayers" Figure 2 and 3

Authors: See comment above. Also in this case the water seems to have percolated till the winter layers forming a thick ice lens.

L7:delete "-" in front of "A further.."

Authors: This seems to be a deleted blank. See now Page 7 L 29.

L14change to: "..where these authors.."

Authors: Done.

L19Change to: "However, in the same study, no evidnece.."

Authors: Done.

L23delete "-" in front of "These"

Authors: This was also a deleted blank. See now in P8 L4.

L27Change to: "..2010) similar values being reported.."

Authors: Done.

L28Change to: "could show..."

Authors: Done.

L29 replace "dislocation" by "transport"

Authors: Done.

PAGE 10

L3: spelling: "recognized"

Authors: Done.

L22:change to: "..in order to support it."

Authors: Done.

L36 and 37:But then, if that is the case, one can wonder how far down in the deep ice core you will be able to decipher "sub-seasonal" behaviour?...Also, changes between glacial and interglacials are likely to involve more drastic changes than some 10 days of shift in the flowering timing!.. The whole speciation could change!...This is not convincing and I would

delete lines 32 to 37... I would also add in the previous sentence that the feasibility of the approach in deep ice cores also relies on potential regional shifts in pollen speciation.

Authors: We deleted those lines and added the sentences suggested (and the corresponding citations from Reference lists).

PAGE 10 and 11

All 5.5 : This remains a very weak section of the paper!.. The $T^\circ/\delta D$ relationship is really blurred and has nothing to do with what would be expected of a clean GMWL!.. It is therefore very difficult to interpret, and, to me, brings no added value. It is simply pushing your application too far!.. That section is not convincing and should be dropped.

Authors: Deleted.

PAGE 11

L23: Change to: "...source, with negligible source-sink lag, the existence.."

Authors: Done.

L29: Change to: "...and can also take place ...Alternatively, data from the closest.."

Authors: Done.

L32: Change to: ".. As witnessed by the recent launch..."

Authors: Done.

L33: Change to: "..Theoretically, our approach could be applied to deeper ice cores, if the sampling resolution...high and if no significant shift occurred in the pollen specification. In such cases our method could ... model by documenting the loss of seasonal signal with depth in the core."

Authors: Done.

L41replace: "types" by "conditions"

Authors: Done.

PAGE 12

L1 to 3: Too weak. Drop this.

Authors: Done.

L8: Change to: ".. In this paleo-climatic archive under "borderline" environmental conditions for proxies reliability."

Authors: Done.

FIGURES

Figure 2, caption "Number and thickness of ice lenses": it is not clear if this is total thickness or individual thickness.. Surely not all ice lenses at a given depth have the same thickness (?)

Authors: it refers to mean individual thickness of lenses per sample (added in capture).

Figure 4: axes Change to: "Mass balance from modelling" and "Mass balance from Pollen"

Authors: Done. Fig. 4 has been renumbered as Figure 5.

Figure 5: Delete.

Authors: Former figure 5 has been deleted.

Linking pollen deposition and, snow accumulation ~~and isotopic composition~~ on the Alto dell'Ortles glacier (South Tyrol, Italy) for sub-seasonal dating of a firn temperate core

5

Daniela Festi¹, Luca Carturan², Werner Kofler¹, Giancarlo dalla Fontana², Fabrizio de Blasi², Federico Cazorzi³, Edith Bucher⁴, Volkmar Mair⁵, Paolo Gabrielli^{6,7}, Klaus Oegg¹

¹Institute of Botany, University of Innsbruck, Sternwartestraße 15, A-6020 Innsbruck, Austria

10 ²Department of Land, Environment, Agriculture and Forestry, University of Padova, Viale dell'Università 16, 35020 Legnaro (PD) Italy

³ Department of Agriculture and Environmental Sciences, University of Udine, via delle Scienze 208, 33100 Udine, Italy

⁴ Autonome Provinz Bozen Südtirol, Landesagentur für Umwelt, Biologisches Labor, Unterbergstraße 2, 39055 Leifers, BZ, Italy

15 ⁵ Autonome Provinz Bozen Südtirol, Amt für Geologie und Baustoffprüfung, Eggentalerstraße 48, 39053 Kardaun (BZ), Italy

⁶Byrd Polar and Climate Research Center, The Ohio State University, 1090 Carmack Road, Columbus, Ohio 43210-1002, USA

⁷School of Earth Sciences, 275 Mendenhall Laboratory, The Ohio State University, 125 South Oval 8 Mall, Columbus, OH 43210, USA

20 *Correspondence to:* Daniela Festi Daniela.Festi@uibk.ac.at

Abstract. Dating of ice cores from temperate non-polar glaciers is challenging and often problematic. Yet, a proper timescale is essential for a correct interpretation of the proxies measured in the cores. Here, we introduce a new method developed to obtain a sub-seasonal timescale relying on statistically measured similarities between pollen spectra obtained from core samples, and daily airborne pollen monitoring samples collected in the same area. This approach was developed on a 10 m
25 core retrieved from the temperate firn portion of Alto dell'Ortles glacier (Eastern Italian Alps), for which a five-year annual/seasonal timescale already exists. The aim was to considerably improve this timescale, reaching the highest possible temporal resolution, and testing the efficiency and limits of pollen as a chronological tool. A test of the new timescale was performed by comparing our results to the output (date of layer formation) of the mass balance model EISModel, during the period encompassed by the timescale. The correspondence of the results supports the new sub-seasonal timescale based on
30 pollen analysis. This comparison also allows to draw important conclusions on the post depositional effects of meltwater percolation on the pollen content of the firn core, as well as on the climatic interpretation of the pollen signal. ~~Finally, we provide an example of useful application of the timescale related to the direct comparison of measured meteorological parameters and the stable isotopes composition of the core.~~

35 **Keywords:** cryopalynology, ice core, chronology, paleoclimatology

1. Introduction

Ice core dating is crucial for the interpretation of the paleo-climatic and paleo-environmental proxies contained in glacial archives. When combined with the detection of absolute temporal horizons, annual layer counting is the most accurate

technique to date ice cores (Thompson et al., 2013). However, low snow accumulation and/or post-deposition effects (e.g. meltwater percolation) hamper the detection of annual layers, especially in temperate glaciers where annual signals are most often smoothed (Eichler et al., 2001). In addition, no proxies had so far a level of temporal precision that allows studying past seasonal changes in detail, while a sub-seasonal temporal resolution would be desirable to reconstruct these changes in detail.

Several studies conducted on glaciers worldwide have proven that pollen (Nakazawa et al., 2011, 2005, 2004, 2015; Santibañez et al., 2008; Uetake et al., 2006) and stable isotopes (Gabrielli et al., 2008; Haeberli et al., 1983; Thompson, 1980; Dansgaard, 1964; Vareschi, 1934) are valuable proxies to detect seasonality in ice cores. However, annual layer counting from oxygen and hydrogen isotopes ratios is a far more common chronological tool than the palynological approach, the potential of which remains largely unexplored. The reason for the scarce use of cryopalynology has functional and conceptual basis. The main issue is that pollen analyses requires a minimum ice sample size of up to 1 L (e.g., Burogois, 2000), which is problematic to obtain because sample volume from ice cores is very limited, especially when working at high resolution. Yet, studies from the Altai (Nakazawa et al., 2015, 2011, 2005 and 2004) and the European Alps (Festi et al., 2015; Bortenschlager, 1970a and b; Vareschi et al., 1934) suggest that this is still a limit for clean samples obtained from polar ice caps, whereas in low and middle latitude glaciers, the minimum sample size can be reduced to 10-30 mL (Festi et al., 2015; Nakazawa et al., 2011 and 2005), thanks to the proximity of the source vegetation and the consequent much greater pollen deposition. An additional limit is that pollen analyses are time-consuming and work-intensive, because they imply manual identification and quantification of pollen grains. In order to overcome this latter issue Nakazawa et al., (2011, 2005 and 2004) adopted a simplified approach, focusing on three main taxa (Pinaceae, Betulaceae and *Artemisia*) that are representative of the three corresponding flowering seasons (spring, early summer, late summer). By doing so, they were able to detect seasonal changes in ice cores from the Altai (Nakazawa et al., 2015, 2012, 2006 and 2005). Finally, there is little knowledge about the effects of percolating water on the palynological signal and only two studies directly addressed this issue, obtaining contrasting results (Ewing et al., 2014; Nakazawa and Suzuki, 2008).

In Festi et al. (2015) we developed an efficient method to detect seasonality employing a 10 m shallow core extracted in June 2009 from the Alto dell'Ortles glacier (South Tyrol, Italy) (Fig. 1) and we improved the existing timescale based on the isotopic composition of the core (Gabrielli et al., 2010). In Festi et al. (2015) we conducted accurate taxonomical identification and implemented a statistical approach consisting in performing a principal component analysis (PCA) on pollen concentration values, and extracting the three principal components (PC) indicative of the three flowering seasons. As each PC summarizes the seasonal information of the pollen assemblage, score values of PC indicate seasonal/annual patterns and enable the identification of seasonal and annual firn layers (Fig.2 a). In Festi et al. (2015) we also discussed that the main pollen input on the glacier likely comes from the near valleys, as the pollen spectra from the ice samples and from valley floor at Solda's monitoring station (Fig. 1) are very similar.

The present study uses the palynological data discussed in Festi et al. (2015) to develop a new refined and innovative pollen-based method to date ice core samples at a sub-seasonal resolution. The aim was to improve the chronology enhancing resolution, dating efficiency, coherence of seasonal patterns (i.e. overlapping of components).

Combined with mass balance modelling, the new high resolution results also provide new insights into the processes controlling the formation and preservation of the palynological signal in firn and ice. ~~Finally, we give an example on the potential application of the new pollen-based timescale by applying it to the isotopic composition of the core, hereby performing a direct comparison between meteorological parameters (i.e. temperature) and isotopes measured in the core, allowing to gain a new insight on the processes of formation of the isotopic signal.~~

2. Study site

The Alto dell'Ortles is the highest glacier of South Tyrol (Italy) in the Eastern European Alps (46° 30' 32'' N, 10° 32' 41'' E) (Fig. 1), ranging in altitude between 3018 and 3905 m a.s.l. and covering an area of 1.07 km². The maximum glacier thickness is about 75 m (Gabrielli et al., 2012) and encompasses the last ~ 7 kyr (Gabrielli et al., 2016). In its upper part the glacier is polythermal, with temperate firn and cold ice underneath (Gabrielli et al., 2012). The local climate is dry and continental, characterized by a mean annual precipitation of 800-950 mm y⁻¹ at the valley floor in Solda (Adler, 2015). This study focuses on the uppermost 10 meters of firn accumulated on the glacier from spring 2005 to June 2009, that have been retrieved using a lightweight hand auger at 3830 m a.s.l.. Our previous research (Festi et al., 2015; Gabrielli et al., 2010), proved that the seasonal/annual signature of pollen and stable isotopes is well preserved in these shallow firn layers. Figure 2(a) shows the formerly published timescale (Festi et al., 2015), which is based on the palynological analyses of 103 continuous ~10 cm samples along the core.

In this study, we also use air temperature data recorded by a standalone data logger placed on the Mt. Ortles at 3835 m a.s.l., as well as meteorological (air temperature and precipitation) and airborne pollen data collected at the meteorological station of Solda (Festi et al., 2015). The station is located 4.5 km northeast of the Mt. Ortles at an altitude of 1850 m a.s.l. (Fig. 1).

3. Methods

This section describes i) the novel pollen-based method “from depth-to-day” developed to obtain a high resolution timescale for the 2009 Mt. Ortles shallow firn core, and ii) the approach used to obtain a core layer dating by using a mass balance model (EISModel), which serves as a comparison for the newly developed palynological timescale.

3.1. High resolution pollen-based timescale: the depth-to-day method

Details on the composition of the Mt. Ortles and Solda's pollen assemblages are reported in Festi et al. (2015). Here, we use these data to develop an enhanced resolved chronology at sub-seasonal timescale. The extensive pollen analyses carried out on the 103 samples obtained from the 10 m core provide a high diversity of 64 pollen types, including the main pollen types characterizing the forest vegetation in the region (e.g., *Pinus* sp., *Picea*, *Fagus*, *Corylus*, *Betula*, *Fraxinus*, *Quercus*), as well as herbs (e.g. Poaceae, Chenopodiaceae, Asteraceae, Cannabaceae). The Ortles pollen assemblages proved to be representative for the regional vegetation, and to be comparable with airborne assemblages recorded at the Solda aerobiological station during the years 2008 to 2010 (Festi et al 2015). Solda's airborne pollen data provided crucial daily information about the timing of local flowering of different plant taxa and of the daily changes in airborne pollen concentration for the years 2008 to 2010. Every plant in the region releases pollen during a certain period of the calendar year, and this is repeated on an annual cycle. However, the onset of flowering may differ by several days (1 to 7 depending on the species; Festi et al., 2015) from year to year, due to different weather conditions.

Solda's airborne pollen samples are characterized by their specific pollen content on a specific day of the year (DOY), while each of the 10 cm sequential Mt. Ortles samples is characterized by its pollen assemblage at a specific depth in the firn core. The three-year palynological dataset of the Solda's aerobiological station has been considered as a representative calibration data-set to define the flowering DOY for in the entire period covered by the 10 m Ortles firn core (2005 to 2009). For every 10 cm sequential Mt. Ortles sample, the three Solda's airborne samples with the highest similarity (one per each year of the three years Solda dataset) were selected, using the Jaccard similarity index (Jaccard, 1901). This index was chosen because it is asymmetrical and hence avoids the double zero problem, i.e. it excludes similarity in case of absence of a pollen type from pairs of compared samples. The Jaccard similarity index was calculated with the SPSS software obtaining a matrix of similarity indices. Indices typically presented values scaled from zero to one: the higher the value, the greater the similarity between two samples. The lower boundary for the Jaccard index was set at 0.5 to ensure high similarity and avoid possible mismatches.

Three potential DOYs were obtained for every Mt. Ortles sample. For each sample, the average DOY and uncertainty (one sigma) have been calculated (Table 1, Fig. 2 (b)). Mt. Ortles samples having pollen concentrations reflecting “winter” season (<0.5 grains ml⁻¹), were excluded from the analyses, as Solda’s airborne station observes only pollen distributions during the vegetation period (March - October). Therefore, the added value of the sub-seasonal dating method mostly lies in spring and summer months, i.e. the ablation season on glaciers. This time matching between flowering and ablation is one of the reasons why we implemented this procedure.

In summary, by coupling the Mt. Ortles firn samples with the most statistically similar assemblage of the Solda’s airborne samples, we establish a link between pollen deposition at a specific sample depth on Alto dell’Ortles and a specific DOY. This “space-for-time” substitution (depth-to-day) by pollen is the key concept of the new dating technique developed. This method is based on the assumption that there is no time lag between the flowering in Solda and the pollen deposition on the glacier, thanks to the efficient uplift of pollen grains by thermic wind (Barry and Chorley, 2009). Depth values were converted to water equivalent values to enable comparison with EISModel results, using a polynomial function fitted to the 2009 firn core density sampled at 10 cm depth intervals. Samples are ordered (Table1) according to their increasing depths from the top of the core, and increasing w.e. from the bottom of the core.

3.2. EISModel

Here we briefly describe the essential parts of the mass balance model used in this study (Carturan et al., 2012a). The cumulated mass balance from 2005 to 2009 at the coring site was calculated using EISModel (Carturan et al., 2012a; Cazorzi and Dalla Fontana, 1996). Before being included in the model, the raw meteorological data recorded on Mt. Ortles and at Solda, were checked and validated against other meteorological weather stations located in the proximity of Mt. Ortles (Madriccio at 2825 m and Cima Beltovo at 3328 m). The precipitation data recorded at Solda were corrected for gauge undercatch errors, using the procedure described in Carturan et al. (2012b).

The mass balance model simulates accumulation and melt processes at hourly time steps. Snow accumulation was calculated from the precipitation data recorded at Solda, extrapolated to the elevation of the study site using a Precipitation Linear Increase Factor *PLIF* (% km⁻¹), to account for the increase of precipitation with altitude. Combining Solda’s precipitation with measurements of snow water equivalent performed in the area of Madriccio weather station at the end of winter 2013 and 2014, at an altitude of 2700-2800 m (Carturan L., unpublished data), has enabled the calculation of an average *PLIF* of 50% km⁻¹, which was assumed to be valid also for the period 2005-2009. Effects from preferential snow deposition, sublimation and erosion by wind were not taken into account in extrapolating precipitations at the core site, assuming they compensate each other. In absence of direct observations, this is a reasonable assumption and its possible effects are discussed further on in the paper. Internal accumulation due to refreezing of percolating water was not calculated because it is negligible in temperate-firn layers (March and Trabant, 1997; Cogley et al., 2011; Zemp et al., 2013)

Ablation was calculated by means of an enhanced temperature-index approach, using the clear-sky shortwave radiation computed from a Digital Elevation Model (LiDAR survey of 2005, provided by the Province of Bolzano) as a distributed morpho-energetic index (Carturan et al., 2012a). The melt rate MLT_t (mm h⁻¹) was calculated for each hour (t) as follows:

$$MLT_t = RTMF \cdot CSR_t(1 - \alpha_t) \cdot T_t \quad (1)$$

where T_t (°C) is the air temperature, CSR_t (W m⁻²) is the clear sky shortwave radiation and α_t is the surface albedo (calculated in function of T_t , Carturan et al., 2012a). *RTMF* is a calibration coefficient called Radiation-Temperature Melt Factor (mm h⁻¹°C⁻¹W⁻¹ m²). The T_t at the study site was calculated from the air temperature measured at Solda, applying a monthly-variable lapse rate ranging from -4.6°C km⁻¹ in November to -6.8°C km⁻¹ in July (mean value = -5.9°C km⁻¹), as calculated between Solda and the air temperature logger placed at the core site at 3835 m on Alto dell’Ortles in the period from October

2011 to May 2014. The RTMF value was obtained by using mass balance measurements (snow depth soundings and density measurements in snow pits) carried out at the core site on June 12 and August 31, 2009.

For each snow layer deposited (i.e. the water equivalent that accumulates at the surface of the snowpack during an hourly time step), the model provides its time and date of formation as well as the air temperature during its deposition.

4. Results

4.1. Pollen based timescale

Results of depth-to-day match of firn and Solda's samples are shown in Table 1 and Fig. 2(b). Each date illustrates the time period encompassed by the sample, with the amplitude of the error bars indicating the number of days potentially included in each sample. The surface sample was dated to the 6th of June 2009, in good agreement with the fact that the core was extracted in the first half of June. The dates cluster in 5 groups representing 5 vegetation periods, and are distributed in chronological order within each year. Few inversions are present (i.e., samples at 78 cm w.e. in 2006; 175 cm w.e. in 2007; 324 cm w.e. in 2008; 434, 448 and 452 cm in 2009), but the estimated dates are generally within the uncertainty of adjacent samples. Such inversions are reported in Table 1, as they potentially represent disturbances in the snow deposition (see discussion paragraph 5.2). We observe a substantial difference in the pollen distribution patterns in the snow, both within and among flowering seasons. Flowering seasons stand out distinctively as layers with high pollen concentration, variable inter-annual thickness and seasonality patterns. According to their thickness and vertical distribution of pollen, the flowering years cluster in two groups: 2005 and 2006 vs. 2007, 2008 and 2009. In details, the flowering seasons of 2007 and 2008 correspond to very thick firn layers (76 and 65 cm w.e. respectively), into which pollen is distributed with a clear seasonal pattern. In contrast, the flowering seasons of 2005 and 2006 are characterized by: i) a significantly lower firn thickness (12 and 34 cm w.e. respectively), ii) the occurrence of a thin lower layer with a distinct spring pollen content, and iii) a thin upper layer containing mixed spring/summer pollen. On the contrary, the non-flowering seasons (October to February) are clearly visible as firn layers which are free (or nearly free) of pollen and present significant differences in thickness. Winter 2007/08 is the thinnest, with only 33 cm w.e., followed by 2006/07 with 44 cm w.e., 2005/06 with 60 cm w.e. and finally 2008/09 with 91 cm w.e.:

4.2. EISModel output

The cumulated mass balance simulated by the model shows high variability of accumulation and melt rates during the period from 2005 to 2009 (Fig. 3). In particular, there is a strong difference between the first two years (2005 and 2006), during which snow accumulation mostly occurred in late summer after weeks with high ablation, and the last three years (from 2007 to 2009), characterized by higher accumulation rates and a much lower summer ablation. In 2005, the ablation removed snow layers accumulated between the 9th April and the 31st of July. The same happened in 2006 to the snow layers accumulated between the end of March and the end of July.

Given that the precipitation regime in the Mt. Ortles area is characterized by a winter minimum and a summer maximum (Schwarb, 2000), corresponding maxima and minima in the snow accumulation rate are expected at the study site, in absence of significant ablation. This behaviour was indeed modelled in the hydrological years 2006 - 2007 and 2007 - 2008. In 2008 - 2009 the modelled accumulation rate was high also during winter, as a result of abundant winter precipitation in the Italian Alps.

5. Discussion

5.1. Improvements in the pollen timescale

The depth-to-day method and the PCs approach (Festi et al., 2015) provide converging evidences that the core encompasses 5 vegetation periods, that can be assigned to specific years by counting back from the core's extraction date in June 2009 (Fig. 2, panels (a) and (b)). However, the new method ~~shows presents clear advantages improvements~~ at ~~different various~~ levels: i) it provides a sub-seasonal time resolved dating; ii) it estimates the timespan encompassed by each dated sample (represented by the amplitude of the error); and iii) it resolves the main issue of the PCs method, namely extracting a coherent chronological succession when PCs overlap, e.g. in years 2005 and 2006. ~~For example, in~~ the year 2005 the overlapping of principal components (PCs) makes the interpretation of the seasonality problematic. On the contrary, the depth-to-day method provides three distinctive dates in chronological order, encompassing a time period from April to July (Fig 2(b)). A similar situation is found in the year 2006, which mainly corresponds to an overlapping of PCs between 80 and 85 cm w.e. of depth. This overlapping follows a very low rise in the spring component and precedes an equally low increase in the late-summer component. The depth-to day method provides a better chronological succession of the samples, by dating the first sample of 2006 to March, a second cluster of three samples to May-beginning of June (corresponding to the PCs overlapping) and the final sample to the mid-end of June. The central cluster of dates includes also an inversion, likely indicating disturbances in the pollen and/or in the snow deposition, possibly caused by melting or wind erosion and ~~rededposition redidposition etc.~~ (see also section 5.3). Also in 2007 the succession of dates provides a more straightforward timescale than the components do, solving the case of overlapping components between 160-180 cm w.e. of depth. The ability of the depth-to-day method to produce a detailed dating for layers otherwise characterized by the PCs overlapping suggests that this method is more efficient in detecting changes in the pollen assemblage. In general, component overlapping can be expected because the flowering of the different plants occurs in a continuum, without pauses between spring, early-summer and late summer season, and therefore without sharp changes in the airborne pollen assemblage. The only break in the flowering is represented by the autumn and winter, as in this period there is no local pollen production, leading to firn layers with no, or sporadic pollen occurrence. In addition, the PC overlapping may also result from sampling two flowering season in a single snow/firn sample. In summary, the day-to-depth method significantly improved the timescale by Festi et al. (2015), providing a clear and intuitive representation of the chronological succession, showing a better efficiency in detecting changes in pollen assemblages that allows establishing a sub-seasonal resolution timescale, and providing the estimation of the number of days included in each dated sample.

5.2. Comparison of the pollen and modelled timescales

Application of the “depth-to-day” pollen concept and mass balance modelling provides two independent dating of the snow and firn layers of the 10 m firn core drilled in June 2009. Layer dating obtained with EISModel and pollen ~~analysis dating~~ are in very good agreement (Pearson correlation coefficient $r=0.99$; $p<0.01$) as shown in Figure ~~34~~. The chronological development of snow accumulation as modelled by EISModel matches very well with the absolute dates provided by the pollen-based timescale (Fig. 3 ~~and 4~~). This can be partially due to the compensating effects of a small overestimation of the modelled ablation during summer and the higher accumulation during winter (Fig. ~~54~~). The most likely cause of such behaviour is the simplified modelling of snow accumulation, which does not account for redistribution processes. Nevertheless, ~~the~~ the modelling approach ~~appears looks globally~~ robust ~~following, as revealed by~~ validation tests carried out in the period from August 2008 to August 2013. Without parameter adjustments, the EISModel reliably calculated the mass balance measured in 10 different dates at the core site (+10% cumulated error in 5 years, 23 cm w.e. RMSE for single dates).

The two methods provide converging evidence that two distinct types of warm seasons occurred during the investigated period, i.e. summers with high ablation in 2005 and 2006, and summers with high accumulation in 2007 and 2008 (Fig. 3 and ~~54~~). The high agreement between pollen dating and EISModel results further confirms that in the Ortles region pollen grains are very efficiently transported upward from the vegetation source to the glacier with no/negligible time lag. This is due to the vicinity of the vegetation to the ice body and to the steep altitudinal gradient (Fig.1). Such efficiency, ~~however~~, cannot be

assumed to be a common feature for every glacier-periglacial environment. In fact, Nakazawa et al. (2015) observed a significant delay (about one month) between pollen production and its deposition on a glacier in the Mongolian Altai. In this context, the application of the depth-to-day method can be problematic, however if the time-lag between pollen production and deposition is known it can be integrated in the dating model.

Comparing the detailed chronological distributions obtained by the palynological methods and by the EISModel, ~~is of helps~~ in the interpretation of problematic features such as the pollen timescale inversions (specific color dots in Fig.2b). Theoretically, chronological inversions can potentially ~~be of two have different~~ origins: a) there statistical and/or physical. ~~They~~ might be a statistical artefacts due to the limited time period (three years) covered by the available data from the aerobiological station of Solda. A longer training period for our palynological method would probably have enabled a better dating accuracy; ~~b) On~~ other hand, there can also be physical processes that have altered the original deposition of pollen and snow; ~~for example the~~ redistribution of surface snow layers caused by wind erosion and redistribution, and ~~the~~ mixing of pollen in melting layers. The high-altitude location of the Ortles core site favors strong wind redistribution because it is not sheltered by the surrounding relief, and because the low temperature keeps the snow dry (and therefore mobilizable) for long periods, also in spring and summer. The effects from surface melt and water percolation are discussed in ~~the following~~ section 5.3.

5.3. Melt water effect on the pollen signal

For the year 2005 and 2006 the EISModel calculations indicate the occurrence of strong summer ablation, leading to the removal of spring and early summer layers for a thickness of about 30 cm w.e. The occurrence of a 0.45 cm thick ice lens at the bottom of the core, and a thick 6.3 cm lens at 50 cm w.e. (Fig 2d) are indications of water percolation and refreezing due to summer melting. Despite water percolation, dating of the 2005 samples provided dates which are in chronological order, with no evidence of pollen displacement. In the 2006 sequence, the only hint for a minor displacement of the pollen is a cluster of three samples dated to May-June, which presents small inversions. While it is hard to assign the origin of such inversions to either percolating water or a limited training set for the depth-to-day method, a minor displacement of the pollen within these layers cannot be ~~completely~~ excluded. On the contrary, a major displacement can be excluded because there is no evidence of pollen displacement downward to the early spring layers in both 2005 and 2006, as well as in the 2005/06 winter strata, despite the fact that ablation reached 30 cm w.e. in summer 2006. In fact, pollen concentration in this winter layer remains below 0.5 pollens mL⁻¹. This value is similar to winter pollen concentration values of the hydrological years characterized by lower summer ablation (i.e. 2007 and 2008), and it is in agreement with the low winter pollen accumulation reported from the western Alps (Haeberli et al., 1983). A further indication of water percolation and refreezing comes from a 9 cm ice lens detected at 150 cm w.e., corresponding to the first datable sample of the year 2007. According to EISModel calculations, summer ablation in 2007 was lower than in 2005 and 2006, and there are no clear evidences of downward pollen transportation. In 2007 the dates are, indeed, in chronological order and include inversion of minor entity. Based on these results, it is possible to infer that during periods with high summer ablation, as during the warm seasons 2005 and 2006, pollen grains are not transported by percolation to lower layers, but mainly concentrate into surface layers similarly to dust and debris (Gabielli et al., 2014).

Our results on the resilience of pollen to percolating water also agree with observations by Nakazawa and Suzuki (2008) in a snowpack study on the Norikura Highlands (1590 m asl) in Japan, where ~~the authors~~ found that during melt phases pollen concentrates on the surface of the snow and is not transported downward by meltwater. This outcome supports the idea that pollen grains are too big (5-200 µm) to be easily displaced by percolation within snow or firn (Nakazawa et al., 2004). Nakazawa et al (2015) in their study on the Potanin Glacier (Mongolian Altai) postulated that the spring pollen layers could be enriched in pollen as a result of heavy summer melting. Also in this case, However, in the same study, no evidence of pollen transport was found in the winter layers. In contrast, Ewing et al. (2014) obtained different results in a laboratory experiment,

simulating post-depositional processes with different glacier snowpack conditions using Styrofoam coolers (60 x 30 x 30 cm) filled with natural winter snow accumulation. In this study, they observed a major vertical displacement of pollen grains concluding that meltwater highly affects pollen distribution.

These divergent conclusions on the effect of percolating water on pollen grains point to the fact that more specific studies on this phenomenon are needed, as it is likely influenced by laboratory design conditions, the natural interplay of local micro-climatic conditions, physical characteristics of snow and firn, and possibly pollen grain size and shape. For example, on the Alto dell'Ortles, fresh winter snow usually accumulates in windy conditions and has a density of 300 kg m⁻³ (Gabrielli et al., 2010) and similar values are reported by Nakazawa et al (2012, 2015). In contrast, snow deposited at other sites with low wind conditions could show a lower density (50-70 kg m⁻³) and higher porosity, which would facilitate the vertical ~~dislocation-transport~~ of pollen grains. For the Alto dell'Ortles glacier, a major displacement of pollen grains by meltwater percolation can be ruled out during the studied time period. The few pollen grains retrieved in the cold season layers are likely associated to the occasional input of redeposited regional pollen grains or long-distance transport of pollen brought by windy events, which could carry pollen from the near Mediterranean region, where the flowering season is longer. A local input of redeposited pollen during winter is unlikely at the study site, because during the cold season the local atmospheric boundary layer generally lies below 2000 m a.s.l., trapping pollen and pollutants in the lowermost layers of the troposphere (Gabrielli et al., 2011).

The combined use of the pollen and EISModel timescale further corroborates the finding by Festi et al. (2015), which improved the dating of the 2009 shallow core obtained from chemistry-based seasonality (Gabrielli et al., 2010) that is more likely to be affected by meltwater percolation. Therefore, an approach that combines at least two of these methods turns out to be a more reliable approach for dating firn cores from temperate glaciers in the Alps. This may be valid also for cores retrieved from other ice bodies located in an environmental setting similar to the European Alps, where the vegetation is close to the glaciers and leads to abundant pollen deposition.

5.4. The potential of pollen for qualitative climatic reconstruction

The comparison between pollen content of sequential firn samples and cumulative mass balance modelling (Fig. 3 and 54), led us to argue that pollen has a good potential not only for dating, but also for inferring the impact of past climatic conditions in firn and ice cores at seasonal resolution, as already recognized by Nakazawa et al (2015). Based on the pollen content in the Mt. Ortles strata, three main types of pollen assemblages can be identified and correlated with the corresponding seasonal climatic conditions inferred by EISModel: i) thin pollen rich layers with fairly clear date order but PCs overlapping (i.e. 2005 and 2006). EISModel indicates that such layers are the result of intense summer ablation thus pointing to warm and dry summer periods; ii) thick layers with significant pollen concentration and well-distinguished succession of dates (i.e. 2007 and 2008). For these layers EISModel shows a spring/summer snow deposition characterized by minimal melting, generated by abundant precipitation and low temperatures; iii) thick layers with no (or nearly absent) pollen, representing snow deposition during the autumn and winter seasons.

Figure 4Figure 5 shows the correlation between the inferred summer and winter mass balance according to the pollen dates and EISModel. The pollen based seasonal snow accumulation has been defined for each flowering year (summer in graph) as the depth in w.e. encompassed between the first and last datable sample of each year, while winter (non-flowering season) mass balance has been calculated as the w.e. accumulated between consecutive summer (flowering seasons). For a direct comparison, the same dates for each season were used for EISModel. Figure 4Figure 5 shows that the thickness of the warm/cold season layers dated by pollen analyses and modelled by EISModel gives a fairly good qualitative indication of the amount of seasonal precipitation: summer 2005 and 2006 stand out for the low accumulation in w.e. with both methods; on the contrary summer 2007 and 2008 are inferred by both methods as high accumulation summers. Similarly, among the cold seasons winter 2008-09 stands out for its higher w.e. thickness correlated to exceptionally high precipitation.

Undoubtedly, this is only the first step towards a qualitative climatic interpretation of the pollen signal and further investigations on longer climate series and pollen sequences from glacier cores are required in order to support this evidence. Further studies are also required to detect as many combinations of pollen assemblage types-ice layers as possible, and to further correlate them with the corresponding conditions of formation. In fact, there are several other possible combinations, e.g. thin layers with high pollen concentration formed during dry (but not particularly hot) summer periods, or thick layers with high pollen content but inconsistent or mixed sequence of seasonal components, deriving from relatively low winter accumulation and possible blending of two or more years. Our results suggest that, combining the classical stable isotope method with palynological analyses does not only enhance the accuracy of the ice core dating, but also offers the potential to provide a calibration set to obtain qualitative paleo-climatic information at seasonal resolution. Clearly, the feasibility of this approach in deeper ice cores depends on the amount of ice available for pollen analyses, the resolution achievable during sampling and the condition of no delay between pollen production and deposition. When applying this qualitative method to deeper ice cores a thinning of the layers by compression must also be taken into account while comparing ice or firn accumulated in different epochs. ~~Finally, the feasibility of the approach in deep ice cores also relies on potential regional shifts in pollen speciation. Long-term studies on the phenology and historic series of pollen monitoring encompassing the last 35 years show that the shift of the onset/end of the blooming season due to temperature trends are on average of 10 days in Europe (Menzel & Fabian 1999, Nature) as well as in the Ortles region (Bortenschlager & Borthenschlager 2007, Grana). We therefore suggest that this difference is not particularly relevant when dating a deep core, as this value is surely smaller than the number of days encompassed in one deep core sample.~~

5.5. Application of the pollen-based timescale

~~The application of the new detailed sub-seasonal timescale to other climatic proxies contained in ice cores (e.g. stable isotopes, major ions, dust, etc.) enables a direct comparison with meteorological variables and opens novel possibilities for a better understanding of processes related to the deposition and preservation of specific proxies. An example of a possible application is provided in Fig. 5(a), where the timescale has been applied to the δD values obtained from the same shallow firn core retrieved on Alto dell'Ortles. For this purpose, a complete timescale was calculated for the core, in order to date all the snow and firn samples and to plot their isotopic composition as a function of time. For samples belonging to the flowering seasons, absolute dates obtained with the depth to day pollen method were used (Table 1 and Fig. 2) and inversions were also included in the timescale. For the cold season layers, in absence of direct measurements of snow accumulation, a simple linear regression of depth vs. time was calculated. By applying this timescale to the δD values, a direct comparison between isotopes and mean daily temperature data as recorded in Solda became possible. This is supposed to be meaningful as there is a very high correlation ($r = 0.95$) between daily air temperature measured at Solda and on the Mt. Ortles shallow core drilling site, during the period from 2011-2014. Figure 5 shows a good correlation (Pearson; $r = 0.642$; $p < 0.001$) between the mean daily temperature and the measured isotopic composition, which is however not stationary as it is affected by the smoothing of δD in the deepest firn layers (Gabrielli et al., 2010) because of meltwater percolation from the upper layers and partial melting of the snow accumulated during the 2005 and 2006 warm seasons. In contrast, the years 2007 and 2008 show a distinct δD summer peak, likely linked to the above mentioned higher accumulation and lower ablation during those summers. In conclusion, this is an example that clearly highlights the utility of combining different approaches for investigating the involved physical processes and for interpreting the proxy data contained in paleo-climatic archives.~~

6. Conclusions

In this study, we have proposed a sub-seasonal timescale for the 10 m Mt. Ortles firn core. The day-to-depth method significantly improved the timescale by Festi et al. (2015), providing a clear and intuitive representation of the chronological

succession, proving a better efficiency in detecting changes in pollen assemblages that allows establishing a sub-seasonal resolution timescale, and providing the estimation of the number of days included in each dated sample. The method can be applied to all types of glaciers, regardless of their thermic state, provided the proximity of the pollen source, with negligible source –sink lag, the existence of flowering seasons (typical of mid-latitudes) or a clear contrast between a flowering and non-flowering season (typical of the tropics and equator due to the alternation of dry and humid season), and the support of modern pollen monitoring data. We also show that a three- years training set of pollen monitoring is sufficient to provide meaningful comparison with glacier samples. This consideration becomes particularly relevant when working in glaciated regions that are not covered by the pollen allergy network, which is the main source of modern pollen data, e.g. World Allergy Organisation-WAO (worldallergy.org), European Aeroallergen Network-EAN (ean.pollenifo.eu), etc. In this case, a three years monitoring can be included in the cryopalynological study and can also take place also after the coring. ~~In a~~Alternatively, data from the closest stations in the region can be used and a correction for the delay between pollen production and deposition can be applied to the dating model. The method is particularly relevant at the moment because of the increasing interest in ice cores from non-polar areas, as ~~testified~~witnedded by the recent launch of the “Protecting ice memory” project, which aims at creating the first archive of glacial ice for future generations. ~~In fact~~Theoretically, our approach ~~couldan~~ be applied to date deeper ice cores, if sampling resolution is sufficiently high and if no significant shift occurred in the source vegetation. In such ~~cases~~cores our method ~~coulan~~ make a significant contribution to the chronological model also by documentingeteeting the loss of seasonal signal with depth in the core the transition from annual to non annual layering in the deeper ice.

The combined use of a mass balance model and pollen-based dating methodology brings compelling evidence that on the Alto dell’Ortles glacier pollen grains are resilient to downward transport by percolating meltwater, also in the case of strong melting as in 2005 and 2006. The independent dating of firn layers by mass balance modelling and pollen match well, and highlight detectable intra-seasonal and inter-annual differences of high altitude snow accumulation rates on Mt. Ortles. More specifically, we found evidence of peculiar types of pollen distribution in firn layers, that may be related to well defined weather types-conditions (e.g. warm-dry, warm-humid or cold-humid weather). These results reveal the good potential of pollen for inferring past climatic conditions at a sub-seasonal resolution in ice core records. ~~Finally, we show that a sub-seasonal timescale is valuable because it can be used to investigate the relationship between climatic proxies contained in ice cores and measured meteorological variables for sites where this relationship has not been studied on site~~. Future studies could focus on joined fieldwork observations of meteorological variables (temperature, snow, wind, etc.) with measurements of snow accumulation/ablation, pollen accumulation, signature of isotopes and other chemical species as well as on monitoring of post depositional effects (snow redistribution, water percolation, sublimation) on the proxies in order to directly test the assumptions made in the present study. This type of approach opens the possibility of gaining new insights in this paleo-climatic archive under “borderline” environmental conditions for proxies reliability. ~~to the physical processes involved in the formation and preservation of the signal stored in this paleo-climatic archive.~~

Author contributions

D. Festi, W. Kofler and K. OeggI performed palynological analyses on the Ortles samples and developed the pollen based timescale. E. Bucher performed the pollen analyses at the aerobiological station of Solda. L. Carturan, F. Cazorzi and F. de Blasi processed the raw meteorological data and mass balance measurements, and carried out the mass balance simulations using EISModel. P. Gabrielli planned the logistic of the field campaign, drilled the firn core and coordinated the processing of the samples. D. Festi, L. Carturan, P. Gabrielli and K. OeggI prepared the manuscript with contributions from all co-authors.

Acknowledgements

We thank the Autonome Provinz Bozen – Südtirol, Abteilung Bildungsförderung, Universität und Forschung for financial support to the project PAMOGIS (Pollen Analyses of the Mt.Ortles Ice Samples). This work is a contribution to the “Ortles

project”- a program supported by NSF Awards #1060115 and #1461422 to The Ohio State University, the Fire protection and civil division of the Autonomous Province of Bolzano in collaboration with the Forest division of the Autonomous Province of Bolzano and the National Park of Stelvio. This is the Ortles project publication no. 8. We acknowledge the collaboration in the various phases of the 2009 and following years field operations of Hanspeter Staffler, Michela Munari and Roberto Dinale (Fire Protection and Civil Division of the Autonomous Province of Bolzano) Ludwig Noessig, Elmar Wolfsgruber and Claudio Carraro (Ufficio geologia e prove materiali of the Autonomous Province of Bolzano), Marc Zebisch and Philipp Rastner (EURAC), Jacopo Gabrieli (IDPA CNR-Venice), Roberto Seppi (University of Pavia), Thomas Zanoner (University of Padova), Karl Krainer (University of Innsbruck), Paul Vallelonga (University of Copenhagen), Michele Lanzinger, Matteo Cattadori and Roberto Filippi (Museo Tridentino di Scienze Naturali) and Silvia Forti (Istituto di Cultura le Marcelline). We thank Michele Bertò for assistance in designing Fig 2. For the logistic support, we are grateful to Toni Stocker (Alpine guides of Solda) and the helicopter company Airway. This is also BPCRC contribution no. 1562. We thank two anonymous reviewers for their comments. We are grateful to Jean-Luis Tison for his useful remarks and suggestions that helped improve the original manuscript.

References

Adler, S. : Das Klima von Tirol-Südtirol-Belluno: 1981-2010; Vergangenheit-Gegenwart-Zukunft. Zentralanstalt für Meteorologie und Geodynamik; Südtirol Abteilung Brand- und Zivilschutz, Bozen, 102, 2015.

Barry, R. G., and Chorley, R. J.: Atmosphere, weather and climate. Routledge, London and New York, 9th edition, 2009.

~~Bortenschlager, S. and Bortenschlager I.: Altering airborne pollen concentration due to Global Warming. A comparative analysis of airborne pollen records from Innsbruck and Obergurgl (Austria) fro the period 1989–2001. Grana, 44, 172–180, DOI: 10.1080/00173130410005582, 2005.~~

Bortenschlager, S.: Neue Pollenanalytische Untersuchungen von Gletschereis und Gletschernahen Mooren in den Ostalpen. Zeitschrift für Gletscherkunde und Glaziologie, 6 (1-2), 107-118, 1970 (a).

Bortenschlager, S.: Probleme und Ergebnisse der Untersuchung von Pollenspektren im Hochgebirge (Kesselwandferner, Öztaler Alpen, 3240 m). Mitteilungen der Ostalpin-Dinarischen Sektion der Internationalen Vereinigung für Vegetationskunde, 10(2), 5–9, 1970 (b).

Bourgeois, J.C.: Seasonal and inter-annual pollen variability in snow layers of arctic ice caps. Rev. Paleobot. Palyno., 108(1), 17–36, doi:10.1016/S0034-6667(99)00031-7, 2000.

Carturan, L., Cazorzi, F., and Dalla Fontana, G.: Distributed mass-balance modelling on two neighboring glaciers in Ortles-Cevedale, Italy, from 2004 to 2009, J. Glaciol., 58, 467-486, 2012 (a).

Carturan, L., Dalla Fontana, G., and Borga, M.: Estimation of winter precipitation in a high-altitude catchment of the Eastern Italian Alps: validation by means of glacier mass balance observations, Geogr. Fis. Din. Quat., 35, 37-48, 2012 (b).

Cazorzi, F. and Dalla Fontana, G.: Snowmelt modeling by combining air temperature and a distributed radiation index, J. Hydrol., 181, 169–187, 1996.

Cogley, J.G., Hock, R., Rasmussen, L.A., Arendt, A., Bauder, A., Braithwaite, R.J., Jansson, P., Kaser, G., Möller, M., Nicholson, L., Zemp, M.: Glossary of Glacier Mass Balance and Related Terms, IHP-VII Technical Documents in Hydrology

- No. 86, IACS Contribution No. 2, Paris, UNESCO-IHP, 114 pp., 2011. Dansgaard, W.: Stable isotopes in precipitation. *Tellus*, 16(4), 436–468, doi: 10.1111/j.2153-3490.1964.tb00181.x, 1964.
- Eichler, A., Schwikowski, M., and Gäggeler, H. W.: Meltwater-induced relocation of chemical species in Alpine firn, *Tellus B*, 53B, 192-203, 2001.
- 5 Ewing, M.E., Reese C.A., and Nolan, M.S.: The potential effects of percolating snowmelt on palynological records from firn and glacier ice. *J. Glaciol.*, 60, 661-669, <http://dx.doi.org/10.3189/2014JoG13J158>, 2014.
- Festi, D., Kofler, W., Bucher, E., Carturan, L., Mair, V., Gabrielli, P., Oeggl, K.: A novel pollen-based method to detect seasonality in ice cores: a case study from the Ortles glacier, South Tyrol, Italy. *Journal of Glaciology*, 61 (229), 815-824, <http://dx.doi.org/10.3189/2015JoG14J236>, 2015.
- 10 Gabrieli, J., Carturan, L., Gabrielli, P., Kehrwald, N., Turetta, C., Cozzi, G., Spolaor, A., Dinale, R., Staffler, H., Seppi, R., dalla Fontana, G., Thompson, L., and Barbante, C.: Impact of Po Valley emissions on the highest glacier of the Eastern European Alps, *Atmos. Chem. Phys.*, 11, 8087-8102, doi:10.5194/acp-11-8087-2011, 2011.
- Gabrielli, P., Barbante, C., Bertagna, G., Bertó, M., Binder, D., Carton, A., Carturan, L., Cazorzi, F., Cozzi, G., Dalla Fontana, G., Davis, M., De Blasi, F., Dinale, R., Dragà, G., Dreossi, G., Festi, D., Frezzotti, M., Gabrieli, J., Galos, S., Ginot, P., 15 Heidenwolf, P., Jenk, T. M., Kehrwald, N., Kenny, D., Magand, O., Mair, V., Mikhalenko, V., Lin, P. N., Oeggl, K., Piffer, G., Rinaldi, M., Schotterer, U., Schwikowski, M., Seppi, R., Spolaor, A., Stenni, B., Tonidandel, D., Uglietti, C., Zagorodnov, V., Zanon, T., and Zennaro, P.: Age of the Mt. Ortles ice cores, the Tyrolean Iceman and glaciation of the highest summit of South Tyrol since the Northern Hemisphere Climatic Optimum, *The Cryosphere*, 10, 2779-2797, doi:10.5194/tc-10-2779-2016, 2016.
- 20 Gabrielli, P., Hardy, D. R., Kehrwald, N., Davis, M., Cozzi, G., Turetta, C., Barbante, C., and Thompson, L. G.: Deglaciated areas of Kilimanjaro as a source of volcanic trace elements deposited on the ice cap during the late Holocene, *Quat. Sci. Rev.*, 93, 1-10, 2014.
- Gabrielli, P., Barbante, C., Carturan, L., Cozzi, G., Dalla Fontana, G., Dinale, R., Draga, G., Gabrieli, J., Kehrwald, N., Mair, V., Mikhalenko, V.N., Piffer, G., Rinaldi, M., Seppi, R., Spolaor, A., Thompson, L.G. and Tonidandel, D.: Discovery of cold 25 ice in a new drilling site in the Eastern European Alps. *Geogr. Fis. Dinam. Quat.*, 101–105, doi:10.4461/GFDQ.2012.35.10, 2012.
- Gabrielli, P., Carturan, L., Gabrieli, J., Dinale, R., Krainer, K., Hausmann, H., Davis, M., Zagorodnov, V., Seppi, R. and Barbante, C. Atmospheric warming threatens the untapped glacial archive of Ortles mountain, South Tyrol. *J. Glaciol.*, 56(199), 843–853, <http://dx.doi.org/10.3189/002214310794457263>, 2010.
- 30 Gabrielli, P., Barbante, C., Plane, J.M.C., Boutron, C.F., Jaffrezo, J.L., Mather, T.A., Stenni, B., Gaspari, V., Cozzi, G., Ferrari, C., Cescon, P.: Siderophile metal fallout to Greenland from the 1991 winter eruption of Hekla (Iceland) and during the global atmospheric perturbation of Pinatubo. *Chem. Geol.* 255, 78 – 86, doi:10.1016/j.chemgeo.2008.06.012, 2008
- Haeberli, W., Schotterer, U., Wagenbach, D., Haberli-Schwiter, H. and Bortenschlager, S.: Accumulation characteristics on a cold, high -Alpine firn saddle from a snow-pit study on Colle Gnifetti, Monte Rosa, Swiss Alps. *J. Glaciol.*, 29 (102), 260– 35 271, 1983.

- Jaccard, P. : Étude comparative de la distribution florale dans une portion des Alpes et des Jura. Bulletin de la Société Vaudoise des Sciences Naturelles 37: 547–579, 1901.
- March, R.S., Trabant, D.C.: Mass balance, meteorological, ice motion, surface altitude, and runoff data at Gulkana Glacier, Alaska, 1993 balance year. USGS Water-Resources Investigations Report, 96–4299, 30 pp. 1997
- 5 ~~Menzel, A. and Fabian, P.: Growing season extended in Europe. Nature, 397, 659, 1999.~~
- Nakazawa, F.: Application of pollen analysis to dating of ice cores from lower-latitude glaciers. J. Geophys. Res., 109 (F4), F04001, doi:10.1029/2004JF000125, 2004.
- Nakazawa, F., Fujita, K., Takeuchi, N., Fujiki, T., Uetake, J., Aizen, V. and Nakawo, M.: Dating of seasonal snow/firn accumulation layers using pollen analysis. J. Glaciol., 51 (174), 483–490, doi: 10.3189/172756505781829179, 2005.
- 10 Nakazawa, F., Konya, K., Kadota, T., Ohata, T.: Reconstruction of the depositional environment upstream of Potanin Glacier, Mongolian Altai, from pollen analysis. Environ. Res. Lett. 7, 2012 035402, doi: 10.1088/1748-9326/7/3/035402, 2012.
- Nakazawa, F., Konya, K., Kadota, T., Ohata, T.: Depositional and summer snow melting features in 2007–2011 on the upstream side of Potanin Glacier, Mongolian Altai, reconstructed by pollen and oxygen isotope analysis. Environ. Earth Sci., 74, 1851–1859 DOI 10.1007/s12665-015-4436-1, 2015.
- 15 Nakazawa, F., Miyake, T., Fujita, K., Takeuchi, N., Uetake, J., Fujiki, T., Aizen, V. and Nakawo, M.: Establishing the Timing of Chemical Deposition Events on Belukha Glacier, Altai Mountains, Russia, Using Pollen Analysis. Arctic, Antarctic, and Alpine Research, 43 (1), 66–72, doi: 10.1657/1938-4246-43.1.66, 2011.
- Nakazawa, F. and Suzuki, K.: The alteration in the pollen concentration peak in a melting snow cover. Bull. Glaciol. Res. 25, 1–7, 2008.
- 20 Santibañez, P., Kohshima, S., Scheihing, R., Jaramillo, J., Shiraiwa, T., Matoba, S., Kanda, D., Labarca, P., Casassa, G.: Glacier mass balance interpreted from biological analysis of firn cores in the Chilean lake district. J. Glaciol., 54(186), 452–462, doi: <http://dx.doi.org/10.3189/002214308785837101>, 2008.
- Schwarb, M.: The alpine precipitation climate. Evaluation of a high-resolution analysis scheme using comprehensive rain-gauge data, Doctoral dissertation, Diss., Naturwissenschaften ETH Zürich, Nr. 13911, 2000.
- 25 Thompson, L.G.: Glaciological investigations of the tropical Quelccaya ice cap, Peru. J. Glaciol. 25(91), 69–84, <http://dx.doi.org/10.3198/1974JoG25-91-69-84>, 1980.
- Thompson, L.G.: Glaciological investigations of the tropical Quelccaya ice cap, Peru. J. Glaciol. 25(91), 69–84, <http://dx.doi.org/10.3198/1974JoG25-91-69-84>, 1980.
- Thompson, L.G., Mosley-Thompson, E., Davis, M.E., Zagorodnov, V.S., Howat, I.M., Mikhalev, V.N., Lin P.-N.
- 30 Annually resolved ice core records of tropical climate variability over the past ~1800 years. Science, 340(6135), 945–950, doi:10.1126/science.1234210, 2013.
- Uetake, J., Kohshima, S., Nakazawa, F., Suzuki, K., Kohno, M., Kameda, T., Arkhipov, S. and Fujii, Y.: Biological ice-core analysis of Sofiyskiy glacier in the Russian Altai. Ann. of Glaciol., 43(1), 70–78, doi: 10.3189/172756406781811925, 2006.

Vareschi, V.: Pollenanalysen aus Gletschereis. Ber. Geobot. Inst. Rübel, 81–99, 1934.

Zemp, M., Thibert, E., Huss, M., Stumm, D., Rolstad Denby, C., Nuth, C., Nussbaumer, S.U., Moholdt, G., Mercer, A., Mayer, C., Joerg, P.C., Jansson, P., Hynek, B., Fischer, A., Escher-Vetter, H., Elvehøy, H., Andreassen, L.M.: Reanalysing glacier mass balance measurement series. *The Cryosphere*, 7, 1227–1245, doi:10.5194/tc-7-1227-2013, 2013.

5





Figure 1 Geographical setting of Mt. Ortles and of the meteorological station of Solda. b) Close up of the Alto dell'Ortles Glacier, where the 2009 shallow firn core was retrieved (figure adapted from Festi et al., 2015).






Figure 2 (a) Mt. Orles firn core 2009 timescale (Festi et al., 2015) based on Principal Component Analyses. As every principal component (PC) condenses the seasonal information of the pollen spectrum, scores values should to be interpreted as follows: a sample (w.e. depth down) presenting high component scores values for a specific PC is characterized by a pollen content reflecting predominantly the season corresponding to that particular PC. (b) Mean date (day of the year ± 1 sigma) obtained by using the new method developed in this paper: empty squares represent inversions. (c) Pollen concentration in the Mt. Orles 2009 firn core. (d) Number and mean thickness (colour scale) of ice lenses per sample.




Figure 3 Comparison between cumulated mass balance modelled by EISMModel and obtained by pollen dating. Alternating grey/white bars have a six months duration and roughly indicate the warm/cold season. Horizontal dash lines indicate melt layers.





Figure 4 Mass balance modelled by EISModel compared with a) pollen dating in the period from 2005 to 2009, and b) mass balance measurements in the validation period from 2009 to 2013.




Figure 4-5 Comparison of the winter and summer mass balance as obtained by pollen dating and modelling (EISModel).




Figure 5 Isotopic variations in the Mt. Ortes 2009 core plotted with the new timescale (this work: pollen dates in red, interpolated dates in blue) (a) compared with the air temperature measured at Solda during the same period (b); Correlation diagram of Ortes isotopic composition (δD) and Solda's mean daily temperature.

5

10

Table 1. Results of the depth to date matching obtained by similarity analyses between Mt. Ortes snow samples and pollen monitoring data from Solda from the years 2008 to 2010. DOY= day of the year. Samples are ordered according to their depth in the core.

Depth (cm)	Water Equivalent (cm)	2008 (DOY)	2009 (DOY)	2010 (DOY)	Mean (DOY)	Date (dd.mm.yy)	SD (\pm days)
10	459	157	157	-	157.0	06.06.2009	0
20	456	159	155	154	156.0	05.06.2009	3
30	452	163	157	187	169.0	18.06.2009	16
40	448	166	159	183	169.3	17.06.2009	12
50	445	-	130	151	140.5	21.05.2009	15
60	441	132	129	145	135.3	15.05.2009	9
70	437	132	137	145	138.0	18.05.2009	7
79.5	434	132	104	116	117.3	27.04.2009	14
99	427	129	129	146	134.7	15.05.2009	12
110.5	422	132	104	113	116.3	26.04.2009	12
120.5	418	111	99	98	102.7	13.04.2009	10
130.5	415	111	99	98	102.7	14.04.2009	14

355.5	324	219	231	230	226.7	13.08.2008	7
365.5	319	229	-	249	239.0	26.08.2008	14
372.5	316	225	233	230	229.3	16.08.2008	4
389.5	309	178	231	201	203.3	21.07.2008	27
399.5	305	211	157	162	176.7	25.06.2008	30
410.5	300	193	166	162	173.7	22.06.2008	17
419.5	296	149	159	163	157.0	05.06.2008	7
452.5	281	152	159	166	159.0	07.06.2008	7
462.5	277	149	159	145	151.0	30.05.2008	7
472.5	272	147	142	157	148.7	28.05.2008	8
492.5	263	132	141	146	139.7	19.05.2008	7
502.5	259	132	131	146	136.3	15.05.2008	8
572.5	226	233	252	-	242.5	30.08.2007	13
604.5	211	212	-	-	211.0	30.07.2007	0
613.5	207	211	191	199	200.3	19.07.2007	10
623.5	202	194	191	191	192.0	11.07.2007	2
633.5	197	164	159	159	160.7	09.06.2007	3
643.5	192	173	157	159	163.0	12.06.2007	9
679.5	175	-	127	154	140.5	21.05.2007	19
689.5	170	-	-	146	146.0	26.05.2007	0
699.5	165	132	138	145	138.3	18.05.2007	7
708.5	160	129	131	154	138.0	18.05.2007	14
728.5	150	132	128	145	135.0	15.05.2007	9
815.5	106	167	157	172	165	14.06.2006	8
854.5	85	134	141	130	135.0	15.05.2006	6
864.5	80	132	138	145	138.3	18.05.2006	7
868.5	78	148	139	169	152.0	01.06.2006	15
878.5	72	-	77	63	70.0	11.03.2006	10
988.5	12	201	162	160	174.3	23.06.2005	23
998.5	7	132	131	145	136.0	16.05.2005	8
1010.5	0	134	104	107	115.0	25.04.2005	17
