

Lisbon, 30 April 2017

Dear Dr Marcia Phillips,

Thank you very much for the careful review, comments and insights on our manuscript on GST and permafrost in the High Atlas. We have tried to answer your requests in this new version of the manuscript and we think that with the new informations, the text is more solid and we hope that it can be accepted for publication in the special issue of The Cryosphere.

Please find below a full copy of your comments and questions, which we have intercalated with our replies.

We have submitted both the clean version of the new manuscript, as well as a version with « track changes » as a supplement file.

Best wishes,

Gonçalo Vieira, Carla Mora and Ali Faleh

### **Comments by the Editor and replies**

Prof. E. Serrano (Reviewer 1) commented that the analysis of the landforms was poor - and although you describe some of them, it is impossible for the reader to locate these, as there is no overview of them. A simple geomorphological map, an aerial photograph or even an excerpt from Google Earth with labelled landforms would be useful. This is particularly important in this setting, because the distribution of possible permafrost does not appear to be linked to an elevational gradient – but rather to morphological terrain characteristics. The process of ice genesis at the base of steep slopes due to mass wasting (avalanches, rockfall) needs to be discussed further and will be mentioned again below.

*Reply : We have prepared a geomorphological sketch map, now included as figure 2, allowing to better discuss on the geomorphological features and their relation with measured data. Using the new map, we have also improved the description of the study area in section 2, including more information on mass wasting processes. The map also identifies the main scarps and differentiates between talus slopes, debris-mantled surfaces, typical of the higher reaches of the study area and the areas with deformation features.*

Dr. B. Staub (Reviewer 2) suggested carrying out a GIS analysis to investigate potential solar radiation, slope angles and terrain curvature. This would allow interesting comparisons with GIS analyses carried out in other mountain regions, e.g. (Kenner and Magnusson, 2017) and would help to determine whether the presence of permafrost is possible or not. From a purely morphological point of view, a simple GIS analysis would allow to determine locations where snow avalanches and rock fall can be deposited – thus pointing towards potential locations of buried ice at the foot of slopes (where air temperature regimes, solar radiation and GST have less influence on the distribution of permafrost than the presence of massive ice buried under a layer of rock debris).

*Reply : We have added a GIS analysis of potential radiation and slope angle as suggested and included it in the methodology, results and discussion. The results serve essentially as reference, since due to the scarce number of sites with GST data, we cannot realistically approach any modelling of relationships between measurements and GIS variables. The discussion on the possible presence of buried-ice has been added to the manuscript, but our observations do not provide clear support to this hypothesis. There are no clear avalanche tracks, nor gullies and the slope above T3 is not the pronest to snow avalanching, the same having been confirmed by our guide. There isn't also any evidence of thermokarst showing degradation of buried ice, nor any spring in June-July close to the arcuate features. The guide indicated that the snow accumulation present in the area results essentially from wind blown snow and we have advanced with the hypothesis of this being a form for ice to enter the talus.*

Reviewer 2 also suggested including freezing and thawing degree-days – this would be a further interesting way of comparing the GST regime at the 4 sites and of making first steps towards the development of High Atlas-specific GST indices – if they have any relation to permafrost distribution at all here. You state that you have these values and I suggest you include them. The extremely high summer GSTs at site T3 and the low ones in winter indicating possible permafrost at this site are worth further reflection and discussion.

*Reply : We have calculated the freezing and thawing degree-days and included them in Table 2. The results are presented in p.5, line 5-12. and also included the discussion, which is now substantially different.*

Although you adapted your previous statements and analyses regarding the comparisons of weather data measured at Menara and on Toubkal, you did not attempt any of the other approaches suggested by Reviewer 2, such as land surface temperature data analysis, which would provide an interesting comparison with your point data and should at least be mentioned in the outlook.

*Reply : We sincerely do not think that remote sensing LST will bring significant insight into assessing permafrost distribution, since LST are collected in the early morning, suffer from significant shadowing effects and are essentially an instantaneous snapshot of GST. Such GSTs vary enormously and very possibly will not correlate well with MAGST. Furthermore LST will only work during the snow free season and in that part of the year, GSTs are normally quite high and show no relation with MAGST. We therefore decided that there is no significant added-value in including this approach in the Outlook section.*

As mentioned above, an important point I suggest you include and discuss is the potential distribution of avalanche snow and of rock fall debris (see Figure 11 for example), based on slope angles. Buried avalanche deposits may be the main / only type of permafrost remaining in this region – independent of elevation but dependent on slope angle and snow / rock debris availability. The ice may be very old. Current avalanche deposits give clues to where snow may have been buried in the past. In the Swiss Alps we have for example observed excess ice at the base of slopes at low elevations (and hence not included on the Swiss permafrost distribution map), and buried under a thick layer of talus (see e.g. [www.permos.ch](http://www.permos.ch) – the Flüelapass site is one of these). Lambiel & Pieracci (2008) and (Scapozza et al., 2011) also provide such examples. You observed creeping features, lobate deposits and ridges/furrows at T3, which is at the base of a steep slope. These indicate that there is/was ice buried here – and if it is still

present, it is a much more likely explanation for the low GSTs in winter than intra-talus ventilation. You should carefully consider what effects buried ground ice would have on the ground thermal regime. The presence of buried ice would also be of interest regarding future water resources in the High Atlas – an important aspect for further research in this arid area.

*Reply : We have further discussed the avalanching hypothesis (was already mentioned in the previous version) and accomodated the buried-ice genesis hypothesis relating it to rockfalls, in order to keep it as an open possibility. We have also improved the description of the arcuate features, which are quite incipient. However, with the current data, we cannot go much further in the discussion, than to present and discuss the various hypothesis, and this is what we did in several parts of the manuscript.*

### **Some open questions:**

Did you notice any cold air fluxes at the base of the slope during very hot days in summer (pointing towards ventilation)?

*Reply : No, we didn't. As we say in the text, the talus is not openwork, but matrix-supported.*

Are there any springs emanating from the lobes at T3 or elsewhere?

*Reply : No, the area was dry during the two visits and we only found water in a small creek in the valley floor away from T3. This water should have been essentially derived from snow melt.*

Were spring temperatures measured?

*Reply : No. We did not find springs close to the talus.*

Have the local guides noticed a change in the occurrence of mass wasting?

*Reply : We haven't collected information on this issue, but is a good point, which we will adress in person with the guides in the future, since by email it is difficult to obtain accurate information.*

### **Further questions**

Title: I suggest you add '...the possible presence of permafrost...'

*Reply : Done.*

p. 3, line 17: is the atmosphere-soil interaction the major controlling factor on the ground thermal regime here? It may be one of the controlling factors – but has less relevance if you consider the presence of buried ice.

*Reply : We have modified the sentence to « The detailed analysis of the GST provides a good insight on the atmosphere-soil interaction, as one of the major controlling factors on the ground thermal regime.” This phrasing opens up the possibility for other factors such as buried ice, that, although possible, is not supported by surficial evidence.*

Study area: mention present/past snow avalanche activity

*Reply : information about avalanches is absent for the Ikhibi South valley, although we have found references to avalanching in the northwest facing slopes on the other side of the ridge. However, there the slope is steeper and there are avalanche corridors. After consulting with our guide, he mentioned that avalanches do not occur at the T3 monitoring site and that snow accumulation is mainly due to sheltering and wind drift. We added the following sentence in p. 4, line 30 : « Oral information from local guides confirms that during the cold season snow accumulates due to wind redistribution from the col and that avalanches do not occur at the site.»*

Methods: how were the temperature loggers calibrated? Ice-water mixture? Active layer zero curtain?

*Reply : The active layer zero-curtain was used to calibrate the loggers, with actual differences found to be below the sensor error (from 0.02 to 0.08 °C). We have added a sentence in p5, line 15 : «The data was corrected using the freezing zero-curtain, having resulted in correction from -0.02 to -0.08 °C.»*

Methods: remove the last sentence (iButtons would for example have been a cheap and practical solution – and why would there be a higher probability of disappearance at high altitude sites??)

*Reply : We have deleted the sentence. As a note, we have decided to use Hobo TidBit, 5x more expensive than iButtons, due to their better accuracy, resolution and due to being more robust. The mention to the disappearance was meant to air temperature loggers, since one i-button installed at the top of Toubkal was stolen and only those installed close to shops or huts made it through the year.*

3.2, p. 6, line 5: define the difference between 'snow' and 'significant snow' – and how is it determined?

*Reply: we have added the following sentence: "The later showed a homogeneous spectral signal of the snow surface across the areas surrounding the sensor sites, while the "snow" class showed still some spectral mixture, but clearly the presence of snow cover."*

Figure 4: where is the extrapolated data for D. Toubkal? (Figure 12?)

*Reply : Given the comments of the referees, we decided to remove the extrapolated data and to present only the measured data. However, we forgot to delete the mention to that in the caption. It is now fixed.*

Figure 9: please label the axes in English (months)

*Reply : OK. Done.*

What are your thoughts on the usefulness and quality of the remote sensing snow cover data?

*Reply : The type of remote sensing imagery we have used is of limited application, but here it was meant to provide a confirmation of the periods with snow cover at the different sites. Since it is an area with absent data on snow cover, we think this was an adequate way to approach the question. It also provides significant information on the overall patterns of snow covers and melting, with important information emphasising on the fast melt of snow in the ridges and south facing slopes, contrary to shady slopes and high valley floors.*

Conclusions: please make these more concise and add a separate section: Outlook.

*Reply : Done. We have shortened the conclusions and added Outlook as section 7.*

# New observations indicate the possible presence of permafrost in North Africa (Djebel Toubkal, High Atlas, Morocco)

Gonalo Vieira<sup>1</sup>, Carla Mora<sup>1</sup>, Ali Faleh<sup>2</sup>

1) Centre for Geographical Studies, IGOT, Universidade de Lisboa, Portugal.

2) Universit  Sidi Mohammed Ben Abdellah, F s, Morocco.

Correspondence to: Gonalo Vieira (vieira@campus.ul.pt)

**Abstract.** Relict and present-day periglacial activity have been reported in the literature for the upper reaches of the High Atlas mountains, the highest range in North Africa (Djebel Toubkal – 4,167 m a.s.l.). Lobate features in the Irhzer Ikbi South at 3,800 m a.s.l. have been previously interpreted as an active rock glacier, but no measurements of ground or air temperatures are known to exist for the area. In order to assess on the possible presence of permafrost, we analyse data from June 2015 to June 2016 from two air temperature sites at 2,370 and 3,200 m a.s.l., and from four ground surface temperature (GST) sites at 3,200, 3,815, 3,980 and 4,160 m a.s.l. allowing to characterize conditions along an altitudinal gradient along the Oued Ihghyghaye valley to the summit of the Djebel Toubkal. GST were collected at 1-hour intervals and the presence of snow cover at the monitoring sites was validated using Landsat-8 and Sentinel-2 imagery. Two field visits allowed for logger installation and collection and for assessing the geomorphological features in the area. The results show that snow plays a major role on the thermal regime of the shallow ground, inducing important spatial variability. The lowest site at 3,210 m showed a regime characterized by frequent freeze-thaw cycles during the cold season but with a small number of days of snow. When snow settled, the ground remained isothermal at 0  C and, the thermal regime indicating the absence of permafrost. The highest sites at 3,980 and 4,160 m a.s.l. showed very frequent freeze-thaw cycles and a small influence of the snow cover on GST, reflecting the lack of snow accumulation due to the wind-exposed settings in a ridge and in the summit plateau. The site located at 3,815 m in the Irhzer Ikbi South valley showed a stable thermal regime from December to March with GST varying from -4.5 to -6  C, under a continuous snow cover. The site’s location in a concave setting favours wind-driven snow accumulation and lower incoming solar radiation due to the shadowing effect of a southwards ridge, favouring the maintenance of a thick snow pack. The stable and low GST are interpreted as a strong indicator of the probable presence of permafrost at this site, an interpretation supported by the presence of lobate and arcuate features in the talus deposits. These results are still a first approach and observations through geophysics and boreholes are foreseen. This is the first time that probable permafrost is reported from temperature observations in the mountains of North Africa.

## 1. Introduction

Permafrost occurrence in arid and semi-arid mountains plays a significant environmental role due to its influence on hydrology. Frozen ground induces refreezing of rain and snowmelt and acts as a subsurface water reservoir able to support streamflow even during the dry season. Its influence can therefore be significant for ecosystems and biodiversity, and in some mountain areas, permafrost may show impacts for agriculture and grazing. However, permafrost research in high remote mountain regions is still in its early stages (see Rangecroft et al., 2013). In fast changing and sensitive mountain environments, permafrost niches can also provide special conditions allowing for the occurrence of biological refugia, which may include endemisms and extremophiles of scientific significance (Hu et al., 2015; Jansson and Taş, 2014). Ice-rich permafrost and ground ice may also allow for environmental reconstruction (Lacelle and Vasil'chuk, 2013). Permafrost also plays a significant role for geomorphological dynamics in mountains, with a number of specific associated landforms and hazards linked to its warming and consequent thaw, such as rock falls and landslides (Haeberli et al., 2010).

The climatic importance of permafrost and the active layer has lead to their classification as Essential Climate Variable (ECV 9) by the Global Climate Observing System of the World Meteorological Organization (Smith and Brown, 2009). The International Permafrost Association maintains the Global Terrestrial Network for Permafrost (GTN-P), which includes over 1074 boreholes, but with only 31 sites in mountain permafrost settings (Biskaborn et al., 2015), which are still poorly assessed regions (Gruber and Haeberli, 2009).

Contemporary permafrost occurrence is known in the Western Mediterranean region but is mostly constrained to small areas at high altitude or shady sites. In the Pyrenees, several active rock glaciers are present and their distribution suggests that the lower limit of permafrost is at about 2,630-2,700 m a.s.l. (Oliva et al., 2016b; Serrano et al., 2009). Geophysical and temperature observations compiled from several authors by Serrano et al. (2009) suggest that continuous permafrost occurs above 3,000-3,100 m a.s.l. In the Sierra Nevada (37° 03' N, 03° 19' W) an isolated patch of permafrost and a very small rock glacier lobe occurs in the Veleta cirque in a north facing cirque at 3,150 m a.s.l., protected by a steep rockwall and reflecting relict conditions associated with buried ice (Tanarro et al., 2001). The mean annual ground temperature for 1998/99-2008/09 in a shallow borehole in the rock glacier was 0.6 °C at 0.05 m depth and -1.4 °C at 1.5 m (Salvador Franch et al., 2011). A deep borehole in the summit area at the Veleta Peak (3,380 m a.s.l.) shows that permafrost is absent and mean annual ground temperatures were 3.2 °C at 0.6 m and 2 °C at 20 m depth (Oliva et al., 2016a).

Periglacial features are widespread in the High Atlas (Hughes et al., 2011) having been described for the Central High Atlas by Couvreur (1966), who has reported active solifluction above 2,200 m a.s.l. The same author states that permafrost is absent in that part of the High Atlas and notes that there is a strong lithological control on the types of periglacial features that occur. For the Western High Atlas, Chardon and Riser (1981) have pushed the limit of frost activity towards 2,500 m and consider that frost action dominates the morphogenesis above 3,000 m. Robinson and Williams (1992) on a study on sandstone weathering report frequent air temperature minima from -10 to 0 °C in winter and as low as -20 °C, at 2,000 m a.s.l. The only landforms supporting a permafrost-related morphogenesis described in the literature are ~~some~~ rock glaciers reported for the

High Atlas by Dresch (1941), Wilche (1953) and Chardon and Riser (1981). Most of them are relict features and at least one case, the Arround rock mass near Imlil, has been recently re-interpreted as a very large rock slide (Hughes et al., 2014). The only reference that we have found for active permafrost-related landforms is Chardon and Riser (1981), who interpret lobate features in the Irhzer Ikbi South at 3,800 m a.s.l. as an active rock glacier. However, recent studies are missing and the literature lacks direct observations and quantitative data supporting the presence of permafrost. The only direct thermal observation of permafrost in the whole Africa, known to the authors, is from Mount Kilimanjaro, where permafrost has been reported at 5,785 m a.s.l. with temperatures of -0.03 °C at 3 m depth (Yoshikawa, 2013).

Given the climate change scenarios that the Mediterranean regions face, marked by warming and precipitation decrease (Giorgi and Lionello, 2007; Montanari, 2013; Simonneaux et al., 2015), permafrost should be close to the threshold of disappearance in most Mediterranean mountains. The subsurface nature of permafrost and the presence of a thawed surface layer in the warmer season (the active layer) strongly limit its identification, characterization and mapping, especially in remote mountain areas (Gruber and Haeberli, 2009).

The present research aims at contributing to solve the question of the presence of permafrost in North Africa and is an exploratory step towards an in-depth assessment aiming at the characterization and modelling of permafrost in the High Atlas.

For such an initial assessment, we have installed a set of ground surface temperature (GST) and air temperature data loggers across an altitudinal gradient from 3,200 m to the summit of the Djebel Toubkal in order to characterize the ground temperature regime and heat exchange at the ground-atmosphere interface. The detailed analysis of the GST provides a good insight on the atmosphere-soil interaction, as one of the major controlling factors on the ground thermal regime.

## 2. Study Area

The Djebel Toubkal is located in the Western High Atlas (31° 4' N, 7° 55' W) and is the highest mountain in North Africa with 4,167 m a.s.l. (Figure 1). The Atlas Mountains comprise a series of ranges and plateaus extending from southwest Morocco to northern Tunisia across more than 2,400 km (Mark and Osmaston, 2008). In Morocco, the Atlas Mountains comprise, from north to south: the Middle Atlas (Djebel Bou Naceur, 3,340 m), the High Atlas (Djebel Toubkal, 4,167 m) and the Anti-Atlas (Djebel Sirwa, 3,304 m). The High and Middle Atlas are intracontinental fold-thrust belts located in the foreland of the Rif (Arbolea et al., 2004). The three major massifs in the High Atlas are, from west to east: the Djebel Toubkal Massif, the Irhil M'Goun Massif (4,071 m) and the Djebel Ayachi (3,751 m).

The climate in the High Atlas is marked by the influence of the Atlantic Ocean to the West, the Mediterranean Sea to the North and the Sahara Desert to the South, resulting in a semi-arid to arid climate (Knippertz et al., 2003; Marchane et al., 2015). The rainy season lasts from November to April and the dry season coincides with the summer, reflecting the Mediterranean style of the climate (N'da et al., 2016). Annual rainfall exceeds 600 mm above 700 m, with summer precipitation being mostly convective. Boudhar et al. (2014) report an average of 520 mm of annual precipitation for the period of 1989 to 2010 in Oukaimeden at 3,200 m elevation. Snow is present from November to April/May in the highest parts of the mountains, but



with irregular regimes (Badri et al., 1994; Peyron, 1980) and is rarely continuous at mid-altitude, with events of snowfall and subsequent melt sometimes happening within one week. However, in the highest reaches, snow cover lasts for several weeks to months (Boudhar et al., 2009). Snowmelt contributes to 15-50% of the stream flow in the Tensift catchment, playing a significant role for irrigation (Boudhar et al., 2009). The low atmospheric humidity and typically subfreezing temperatures above 3,000 m favour losses by sublimation, which can account up to 44% of snow ablation, while at lower altitudes melting prevails (Schulz and de Jong, 2004). Seemingly, the only perennial snow patch in North Africa, occurs in the northern cliffs of the Tazaghart plateau (3,980 m a.s.l.), close to the Toubkal. This feature is described in various recent papers and was identified by Dresch (1941) together with other periglacial features (see Hughes, 2014). Its presence may be related to the high snow feeding area in the plateau above, together with the shelter effect of the steep north-exposed cliff face.

The present study was conducted in the upper reaches of the Oued Ihghyghaye valley, between the marabout of Sidi-Chamharouch and the summit of Djebel Toubkal (Figure 1). The lithology of the study area is composed by Precambrian volcanics, such as Piroxene-bearing doleritic basalts and megaporphyric basalts of the Sidi Chamharouch formation (Zahour et al., 2016) and andesites in the Djebel Toubkal (Cheggour, 2008; Rauh, 1952; Ros et al., 2000). The area shows a typical alpine relief with sharp crests rising above 3,500 m and long deep valleys, with the upper catchments showing evidence of Late Pleistocene glaciation with frequent landforms such as roches mouttonnées and moraines (Chardon and Riser, 1981; Hannah et al., 2016; Hughes et al., 2011; Hughes and Woodward, 2008; Mark and Osmaston, 2008; de Martonne, 1924). Extensive talus slopes and debris cones, together with widespread evidence of frost shattering mark the landscape above 3,000 m.

The detailed study area for ground surface temperatures develops between the Neltner refuge and the Toubkal summit along the Irhzer Ikhibi-South valley, which is the main climbing route. The valley with about 1,400 m in length is a hanging tributary of the Oued Ihghyghayene valley and rises southeastwards of the Neltner refuge above a rock knob with ~~numerous~~ glacier polished outcrops at 3,350-3,400 m. Up valley from the knob, the floor shows a steep longitudinal gradient and is filled by accumulations of large boulders ~~rsly accumulations, grading-passing towards the east~~ in-to the distal parts of ~~the~~ talus slopes, which are matrix-supported accumulations of decimetric to metric angular clasts (figure 2), ~~matrix-supported. These deposits are formed by large boulders in the south slope and formed by smaller boulders in the north slope. At ci. 3,800 m~~ The deposits show small lobate forms and incipient arcuate ridges-ramparts and furrows at ci. 3,800 m suggesting supporting an active periglacial dynamics, either by solifluction or permafrost creep. Chardon & Riser (1998) ~~81~~ interpret this sector as an active rock glacier, ~~but no steep front occurs and no debris body is clearly defined and therefore we do not support this interpretation. No water springs have been observed near the talus slope. The~~ Slopes surrounding the Irhzer Ikhibi-South valley shows dissimetric slopes, with 15-25° prevailing in the north slope and 20 to 40° prevailing in the south slope, valley headwalls showing 30-40°, with scarps and free faces occurring at 3,800-3,900 m, except in the north slope. Above the scarps, slope angle is 15-30° and the surfaces are mantled with angular clasts and boulders, with a similar situation in ~~are steep, with angular taluses with free faces in the rock knob area, along the south slope and in the Toubkal face. Toubkal's west col and most of the slope~~ the north slope of the Irhzer Ikhibi-South of valley are debris mantled. The south ridge rises above 3,900 m causing

~~a significant shadowing effect during the winter in~~ the valley floor ~~in winter~~. A snow patch occurs frequently until June in the ~~Irhzer Ikhibi South~~ valley, especially in the Toubkal slope and may be partly responsible for the debris ramparts, ~~first~~ described ~~above by Chardon & Riser (1981) that occur at 3,800 m. Oral information from local guides confirms that during the cold season snow accumulates due to wind redistribution from the col and that avalanches do not occur at the site.~~ Google Earth  
5 imagery allows identifying around the Toubkal numerous debris-mantled slopes and taluses with flow-like lineaments, suggesting creep, and small rock glacier-like features are also identifiable. At the col of the Irhzer Ikhibi Nord, ~~just~~ north of the Toubkal, at 3,900 m, solifluction lobes are present.

### 3. Methods

#### 3.1 Air and ground surface temperature monitoring

10 Air temperature, relative humidity and ground surface temperature data loggers were installed in June 2015 from Sidi Chamharouch (2, 370 m) to Djebel Toubkal (4,160 m a.s.l.) across an altitudinal transect aiming at an hourly characterization of the soil and air climate for 2015-16. For air temperature and relative humidity, we used Hobo ProV2 loggers, with an accuracy of  $\pm 0.2$  °C, installed in radiation shields at ci. 2 m height. One was installed close to a shop in Sidi Chamharouch (2,370 m) and the other near the Neltner refuge of the Club Alpin Français de Casablanca (3,210 m). Both sites were surveyed  
15 by local partners. A minilogger ibutton DS-1922L was installed at the summit of Djebel Toubkal (4,167 m), hidden in a shady location in the iron trig that stands at the top. However, this logger disappeared and data was lost.  
For GST, single channel miniloggers Hobo TidBit with an accuracy of  $\pm 0.2$  °C were glued to the lower face of a 15x15x0.2 cm high diffusivity steel plate that maximizes contact with the soil particles, when buried at 2-3 cm depth (see Ferreira et al. 2016). To check for drifts on temperature accuracy after retrieval, the loggers were tested under various temperature settings  
20 (-20 °C to 39 °C) and showed average differences under 0.1 °C, which is well within sensor error. ~~The data was corrected using the freezing zero-curtain, having resulted in correction from -0.02 to -0.08 °C.~~ Four of such plates were used between the Neltner refuge and the summit of Djebel Toubkal. The sites were ~~chosen selected in order~~ to characterize the altitude control on GST and were installed along the main climbing route, ~~with care in order to avoid stepping and surfacing due to the large number of climbers.~~ All sites were installed in stony silty-sandy soils, matrix-supported, in gentle slope positions, but where  
25 locally the terrain was relatively flat. Neltner (NLT) was installed above the refuge in a boulderly diamicton. Toubkal 3 (T3) was installed in boulderly diamicton in a valley position, where snow accumulation is favoured. Toubkal 2 (T2) was installed in a debris-~~covered-mantled~~ slope, a few meters below a ridge crest. Toubkal 1 (T1) was installed in the debris-~~covered-mantled~~ surface of the Djbel Toubkal summit plateau, about 100 m from the summit, to avoid the proximity of climbers. Given the known high spatial variability of GST in complex mountain settings (see Gubler et al., 2011) the interpretation of the results  
30 must be conducted with care and this is also the reason why we tried to limit the soil differences between sites. Details are provided in Table 1 and figure ~~32~~.

Temperatures were recorded hourly from 16 June 2015 to 16 July 2016 with the objective of having a whole year of data, centered in the cold season. All measures presented are derived from hourly data. ~~No further loggers were installed due to funding limitations and high probability of disappearance at high altitude sites.~~

3.2 Potential solar radiation modelling

Slope angle and potential solar radiation was modelled using the algorithms of SAGA GIS 4.0, using a standard lumped atmospheric transmittance effect of 70% and the ASTER Global Digital Elevation Model version 2 from NASA, with a pixel size of 30 m. The total potential solar radiation (direct + diffuse) was used for the whole year and also for July in order to assess the influence of solar input in the ground temperature maxima.

3.3 Remote sensing characterization of the snow cover

Although ground surface temperature regimes allow for identifying the presence of snow cover with high degree of confidence, for this study it was essential to demonstrate that snow played a major role on GST and that it was present at some of the sites. For characterizing the snow cover we used 18 scenes from Landsat 8 OLI and 2 scenes from Sentinel 2-A collected between from 16/09/2015 and to 14/06/2016. Landsat scenes were obtained at 16-day intervals, at 10 AM local time (USGS, 2016) and only one scene showed partial cloud cover. Sentinel-2 scenes at 10:30 AM complement the series and allow confirming the results obtained with Landsat. Pixel size is 30 m for Landsat 8 and 10 m for Sentinel 2 (ESA, 2015). We have used full resolution georeferenced visible colour composites provided by USGS at EarthExplorer. The images allowed to identify the general snow cover conditions at the monitoring sites. For the interpretation, caution was necessary since the spatial resolution of the imagery is much larger than the microscale variability that may affect the monitoring sites. The snow conditions at each site were classified by visual inspection of the imagery as « no snow », « possible snow/snow margin », « snow » and « significant snow ». The later showed a homogeneous spectral signal of the snow surface across the areas surrounding the sensor sites, while the “snow” class showed still some spectral mixture, but clearly the presence of snow cover. The classification was done on-screen in QGIS using an overlay of elevation contours and the coordinates of the monitoring sites for better accuracy. Differences between snow and cloud cover were easily identifiable and clouds were rare.

3.4 Climate series and extrapolation

To assess the climatic representativity of the period of June 2015 to July 2016, a long-term climate series of temperature and precipitation from a nearby meteorological station is needed. This allows comparing monthly records with the reference series and better frame the study period and discussing the results. However, the High Atlas has no long-term meteorological stations and the regional network is very sparse. The only long-term meteorological data available are from Marrakesh (Menara) in the plains north of the mountain range at 468 m a.s.l. and about 65 km from the study site, or from Ouarzazate, in the southern piedmont, at 1,153 m a.s.l, but in a very dry setting. Middel, located 350 km to the east of the Toubkal Massif at 1,515 m a.s.l. shows very incomplete data. Given the more complete data series of Menara we used it for framing the climate characteristics

Formatada: Inglês (Reino Unido)

Formatada: Normal

Formatada: Inglês (Reino Unido)

Formatada: Inglês (Reino Unido)

Formatada: Inglês (Reino Unido)

Formatada: Inglês (Reino Unido)

Formatada: Inglês (Reino Unido)

of the study period in a time series. The data was obtained from the Custom Monthly Summaries of the Global Historical Climate Network (NCDC) for 1977-2016. Before 1977 there are several gaps in the series.

4. Results

4.1 Climate characteristics of the study period

5 Climate records from Marrakesh (Menara – Figure 43) for June 2015 to March 2016 (no data available afterwards) show that mean monthly temperature from June to December was close to the median, but January was extremely warm, with a value (15.2 °C) close to the maximum (15.5 °C) of the period 1977-2015 and well above the 3rd quartile (13.4 °C). February with 14.9 °C was between the median and the 3rd quartile and March showed a mean monthly temperature close to the 1st quartile, with 15.4 °C. Precipitation showed very high values in August and September (close to the maximum), decreasing afterwards, with 10 November to January as very dry months, below the 1st quartile and close to the minimum. February showed precipitation close to the median and March, close to the 3rd quartile. The study period was therefore initially characterized by a wet summer, followed by a dry autumn and early winter, which coincided with a very warm January, followed by a very cold and wet March.

The mean monthly air temperatures measured at Sidi Chamharouch and Neltner showed similar variation as Menara, with the 15 exceptions of February and March, which were colder in the mountain massif (figure 54). The lowest mean air temperature was recorded at Neltner with -1.2 °C in February 2016 (3.8 °C in Sidi Chamharouch) and the highest was July 2015 with 14.0 °C in Neltner (20.9 °C in Sidi Chamharouch). The mean annual air temperature (MAAT) for July 2015 to June 2016 was 6.2 °C at Neltner and 11.1 °C at Sidi Chamharouch, which results in a lapse rate of -0.59 °C.100 m<sup>-1</sup>, a value close to the -0.56 °C.100 m<sup>-1</sup> calculated by Boudhar et al. (2009) from weather station data from Saada (411 m) and Oukaïmeden (2,760 m) for 20 1998-2005. The extrapolation of the lapse rate obtained for our two stations to the summit results in a MAAT of 0.6 °C.

Reports from local guides indicate that the winter of 2015/16 was anomalous for snow conditions, with a very late onset of the snow pack in mid-February. This is confirmed by the remote sensing data (figure 56), which shows snow in early October 2015 affecting the Toubkal massif with a peak in the scene of 18/10/2015, then decreasing progressively until early January when a short duration cover shows up, then melting again until 7 February. It was only between 7 and 23 February that 25 significant snowfall occurred covering the whole study area. The snow cover remained in the valley floors and concave areas until 11 April, but quickly melted from the ridges and south facing slopes. In mid-May another large snow fall event took place, but snow melted quickly in two weeks and by mid-June was completely gone from the study area.

4.2 Ground Surface Temperatures

30 Mean monthly ground surface temperatures at the Toubkal massif showed a similar annual regime at the 4 sites (figure 76), with the warmest month in June 2016 showing values from 13.5 °C at T2 to 17.7 °C at NLT. The cold season showed mean

monthly temperatures below 0 °C from November to March at the 3 higher sites, with NLT still showing a positive average in November. T3 showed stable mean GST below -5 °C from December to March, while T1 showed the minimum in February with -4.5 °C.

The daily absolute temperature maxima and minima follow in general the regime of the monthly means, except at T3 and evidence very high amplitudes. A plateau in the curve for the daily maxima is observable from April to September, with T3 showing the highest values reaching 39.3 and 40.5 °C in June and July 2016, respectively. At the same site, the stabilization of the absolute maxima below -4.4 °C in January and February 2016, as well as the sudden rise from 0 °C in March to 31.3 °C in April, are noteworthy. The mean annual GST (MAGST) showed a decrease with altitude with values of 6.02 °C at NLT, 3.2 °C at T3, 3.4-3 °C at T2 and 2.8 °C at T1, the summit of Toubkal (figure 87 and table 2). Degree-days of freezing were extremely varied, with the minimum at NLT with 110, the maximum at T3 with 779, and 325 and 499 at T2 and T1 respectively (table 2). Degree-days of thawing showed very high values, ranging from 1536 at T1 to 2317 at NLT, with a general decrease with altitude. Annual potential incoming solar radiation (figure 9) showed much higher values at the interfluvial sites (T1 – 2432 and T2 – 2631 kWh.m<sup>-2</sup>) than at the valley sites (NLT – 1703 and T3 – 1892 kWh.m<sup>-2</sup>), while July values were much more homogeneous ranging from 223 kWh.m<sup>-2</sup> at NLT to 262 kWh.m<sup>-2</sup> at T1.

Four altitudinal patterns of mean monthly GST occurred in the study period (figure 108). Group 1 includes the warm season, with June, July and August and shows decreasing GST with altitude, with T2 slightly colder than T1. Group 2 integrates the transition months, with May, September and October showing a similar continuous decrease of GST with altitude and a rate of ci. -0.47 °C.100 m<sup>-1</sup>. Group 3 includes the cold season, with January, February, March, November and December and shows a general decreasing of GST with altitude, but with T3 being the coldest site. Finally, April 2016 shows up as an outlier, with an inverted rate with altitude, but with T3 as the warmest site (Group 4).

The hourly records of GST allow for a more accurate analysis of the conditions influencing the monthly means and to assess the environmental controls on the ground thermal regime (figure 119). The first striking characteristic of the GST at the 4 sites is the large diurnal thermal amplitude range, especially from May to September with averages of 11.4 to 12 °C, except at T3, with 19.7 °C. Maximum amplitudes were from 19.2 to 22.5 °C, except at T3, with 35 °C.

The cold season is clearly defined in the GST, lasting from mid-October to the end of April, with May being a transition month. Differences in GST regimes are clear during the cold season. NLT showed a long period from mid-October to mid-February with small thermal amplitude and numerous freeze-thaw cycles. Afterwards, temperatures remained stable just below freezing until mid-April. At T3 the cold season was marked by subfreezing temperatures from mid-October to mid-April, with temperatures decreasing regularly until mid-December after a zero-curtain effect lasting about 2 weeks in late-October. Then, a stable value starting at around -4.5 °C and decreasing irregularly to about -6 °C occurred, as situation which lasted until late March, when temperatures increased quickly and then stabilized for about 10 days with a zero-curtain effect. During this period the diurnal amplitudes of GST at T3 were typically between 0.4 and 1.0 °C, with an average of 0.8 °C. The GST regime is especially significant in this paper and will be analysed in more detail below. The two upper sites, T2 and T1, show very similar GST regimes during the cold season, with the main differences being the number of freeze-thaw cycles which is larger

Formatada: Não Realce

Formatada: Superior à linha

Formatada: Superior à linha

Formatada: Superior à linha

Formatada: Superior à linha

at T2, with lower maxima during the cold season at T1. Both sites show a short zero-curtain effect in late-October, simultaneously to T1. GST daily amplitudes from mid-December to mid-March averaged 5.2 °C at T2 and 4.8 °C at T1 and were variable at NLT.

After an increase in GST in April 2016, May showed a significant cooling that lasted for 8 days in NLT and 19 days at T3.

- 5 This cold event resulted in a stabilization of GST at 0.5 °C at NLT, at 0.2 °C at T3 and at 0 °C at T1 and in a cold but unstable regime at T2.

## 5. Discussion

- 10 MAGST in the study area varied between 6.0 and 2.8 °C, values which are high when compared to mid-latitude mountains with permafrost. There is a clear altitudinal control on the MAGST but site specific factors, such as high snow accumulation and lower annual potential solar radiation, affect the spatial pattern and lower the temperatures as occurs at T3. Consequently, this site showed 779 degree-days of freezing (DDf) in contrast with 325 and 499 DDf at the ridge and summit sites. The effect of increased solar radiation at ridge and summit sites, which show very high values of potential radiation, together with the wind exposition and lack of snow, are also causes for the freeze-thaw regimes observed even during the cold season. The low cloudiness of the mountain range, as demonstrated by the high number of cloudfree remote sensing scenes, emphasises this effect. Potential solar radiation in the summer is high, but homogeneous between sites and do not explain the differences in diurnal thermal regimes, with very high maxima, especially at T3. This fact is noteworthy in the extreme value of 1,936 thawing degree-days (DDt) at the later site, when compared to 1,536-1,542 DDt at the ridge and summit sites. The cause to this difference must be in ground thermophysical properties, as well as to the concave and wind sheltered location at T3, receiving increased radiation from the valley slopes.
- 15
- 20 The hourly GST data ~~across the altitudinal gradient in the Toubkal Massif allows allow~~ for a ~~first detailed~~ insight into the ground thermal regime of the High Atlas periglacial zone. The comparison of the snow cover at different altitudes derived from the remote sensing imagery with the GST regimes confirms that the presence of the snow pack is the cause for stable temperatures close or below 0 °C during the cold season at the different sites (figure ~~65~~). This fact is clear both for the period from late October to mid-April, but also for the short cooling event in May. The effect is well known and has been shown for
- 25 different regions. While a thin and compacted snow layer a few centimeters thick allows for fast heat transfer between the ground and the atmosphere, and in some cases, even to an increase in ground cooling due to high albedo and high ice thermal diffusivity, a thick snow pack acts as buffer between the ground and the atmosphere. If snow is thick enough, the thermal wave will be delayed in the ground (Goodrich, 1982; Staub and Delaloye, 2016; Williams and Smith, 1989). If the ground is unfrozen at depth, heat will flow towards the snow pack and the ground surface temperature will be controlled by phase change at the
- 30 snow interface, generating near 0 °C isothermal regimes (Vieira et al., 2003) - the so-called zero-curtain effect (Outcalt et al., 1990). On the other hand, if the ground is colder than the snow pack, it will generate a heat sink at depth, inducing a decrease

in GST, which are function of the insulating capacity of the snow pack, that also depends on forcing induced by the atmosphere at the snow-air interface (see Haeberli and Patzelt, 1982; Ishikawa, 2002).

From the four monitored sites, Site T3 shows a remarkable regime with relatively stable GST at ci. -5.8 °C from December 2015 to March 2016. The small temperature range reflects the insulating effect of snow cover, but ~~still~~ a small diurnal ~~heat transfer effect~~ wave is identifiable in the GST with daily amplitudes of up to 1 °C. As an example, the detailed examination of the hourly temperatures at T3 from 4 to 20 December 2015 shows GST consistently below estimated air temperatures, except for a few hours in 11 and 12 December (figure 14.2). This situation occurred all over the snow cover period. ~~It is also noticeable~~ Data also shows that the thermal wave is delayed in the snowpack for about 5-6 hours with the resulting GST curve being much smoother than air temperatures.

Ishikawa (2003) reports two sites with GST regimes very close to the one of T3 in the Hidaka Mountains (Hokkaido, Japan), located in low altitude openwork boulder deposits, favouring cold air drainage and funneling in winter. The author has classified this type of settings as the extrazonal permafrost zone. Lambiel and Pieracci (2008) also report a ~~very~~ similar thermal regime to T3 at the base of a talus slope at low altitude in the Western Swiss Alps, reporting GST in the end of the winter below -5 °C as an indicator of the probable presence of permafrost. Values of spring GST below thick snow covers under -3 °C are frequently used in the literature as an indication of the probable presence of permafrost, an approach derived from the widely used Bottom Temperature of Snow (BTS) method (Hoelzle, 1992; Lewkowicz and Ednie, 2004). Numerous cases with sporadic permafrost developing in talus slopes have been reported and the cooling process below coarse debris is well-known (e.g. Delaloye et al., 2003; Delaloye and Lambiel, 2005; Gadek, 2012; Sawada et al., 2003). Gadek and Kędzia (2008) in a study for the Tatra Mountains (Poland and Slovakia), where different GST regimes were analysed, associated sites with steadily decreasing temperatures during the winter to increasing snow cover, and where GST dropped to below about -5 °C, permafrost was found. Such sites were not open work, which limited cold air flow through the boulders, a situation similar to T3 in the Toubkal. Furthermore, small variations in GST through the cold season indicated that the snow insulation was not perfect. In those situations, ground cooling was attributed to the concave locations with cold air flowing over the surface, which cooled the snow and ground. This process acted together with the low summer solar irradiation to promote cooling, with a larger influence on low GST than snow thickness or elevation. Other authors have indicated snow redistribution by avalanches as an important factor promoting cooler ground and permafrost formation at the base of slopes, since snow lasts longer at those localities, mitigating ground surface warming (Haeberli, 1975). Onaca et al. (2015) indicate this as an important effect for the maintenance of permafrost in the Retezat Mountains (Romania), where permafrost occurs at sites with MAGST below 0 °C, BTS lower than -3 °C and ground freezing index higher than 600 degree-days. Snow avalanche and rock and debris fall events have also been shown to be responsible for burial of snow and ice at the base of talus slopes, even at low altitude (see for example (Scapozza et al., 2011)). Such buried-ice may be preserved outside the climate boundary of permafrost and act as heat sinks promoting lower GST and are therefore another hypothesis for the conditions observed at T3.

Following the literature synthesis made above, the low and stable GST measured at T3 (3,815 m) ~~are-is~~ a strong indicator of probable permafrost occurrence. However, MAGST is relatively high, with 3.2 °C resulting from the very high summer GST, likely caused by the concave setting and thermophysical properties of the soil. At the site, the surface material is a boulderly matrix-supported deposit, with a pebbly-sandy matrix and no visible voids. This limits the interpretation of a talus-ventilation related origin for the low GST and suggests conditions close to those described by Gądek and Kędzia (2008) in the Tatra Mountains and also by Onaca et al. (2015) to the Tatras. Further supporting the importance of the effect of the long lasting snow cover, the area at T3 has shown to be one of the latest sites with snow melt. Local guides also indicate that wind redeposition of snow is an important mechanism for snow accumulation at the site. The geomorphic setting at the lower part of a talus favours the possibility of It is also possible that the deposit is open work at depth, which could help explain the low temperatures, but this hypothesis lacks verification. the presence of ice burial by rockfall and avalanching, an hypothesis than should be tested, for example, by Electrical Resistivity Tomography surveying and/or refraction seismics (see (Hauck et al., 2007)). The possibility of the deposit to be open work at depth, which could help explain the low temperatures, cannot be excluded, but lacks verification.

The GST regimes measured at Neltner (3,200 m) show clearly that permafrost is absent at the site. The snow pack settled only in mid-February and temperatures become very stable close to 0 °C, which shows that there is no heat sink at depth, but rather a warmer unfrozen ground. Snow lasted about a week longer than at T3, possibly due to local effects, such as shadowing or thicker snow accumulation associated with snow drift.

The upper sites at the ridge (T2 – 3,980 m) and summit of Toubkal (T3 – 4,160 m) show small events of stable GST as a reaction to the scarce number of snow fall episodes, but long lasting zero-curtains are not visible after the initial one in late October, which was synchronous at T1, T2 and T3 (figure 112). After that episode, GST shows irregular curves with ranges of ~1 to 8 °C reflecting an absent or very thin and compact snow layer with high conductivity. The lack of zero-curtain effects and the frequent freeze-thaw cycles at T2 indicate the absence of snow and a very dry soil. At both sites, the data suggest that the ground remains frozen below the surface during the cold season, however the GST regime analysis does not allow assessing on the presence of permafrost.

The very high daily GST ranges during the warm season are explained by the high insolation of the Mediterranean high mountain, together with the scarce moisture and rocky nature of the soil. We have no data to explain with safety the very high GST at T3, but they should be function of local differences in soil thermophysical properties, together with the concave setting of the site, receiving more reflected and emitted radiation from the surrounding slopes and also due to a wind shelter effect. Both T2 and T1 are convex and very wind exposed sites, which may explain lower maxima than at T3.

The probable presence of permafrost at site T3 is not unexpected, since the valley shows landforms typical of mountain permafrost, such as lobate deposits and poorly developed transverse ridges and furrows in the lower part of the scree slope (figure 134), a fact that has been also described by Chardon and Riser (1981). The high altitude and sheltered location favouring wind driven snow accumulation and longer lasting snow cover, together with the talus setting, also favours ice-burial either by snow deposition (dry snow into the debris voids) or rockfalls. This would led to the formation of interstitial ice that would

Código de campo alterado



facilitate the creep of the frozen talus and the formation of solifluction and creep features Amplified cooling over the snow surface due to the concave setting and cold air drainage, as shown for the Tatra mountains, may also be a controlling process in ground cooling. -The longer lasting snow can also provide moisture that would refreeze at depth when percolating in the debris.

will also contribute to increased water availability in the soil surface, promoting refreezing within the soil, with the possibility for the presence of interstitial ice. This would facilitate the creep of the frozen talus and the formation of solifluction and creep features that are detectable.

The extrapolation of the long-terms records from Menara to the summit of Djebel Toubkal using a lapse rate of  $-0.59\text{ }^{\circ}\text{C}.100\text{ m}^{-1}$ , a similar approach to Hannah et al. (2016) for paleoequilibrium line altitude estimation in the region allows for an insight into the climate sensitivity of the high reaches of the Toubkal Massif. The results should be interpreted with care, but it is worth noting that the extrapolation of the warming trend observed in Marrakesh, represents a gradual shift towards positive MAAT at the summit (Figure 124). This suggests that the probable permafrost sites in the High Atlas are possibly at risk of disappearing if the trend is to be continued.

## 6. Conclusions

The analysis of ~~hourly ground surface temperatures~~ GST from June 2015 to July 2016 across an altitudinal transect in the Toubkal Massif provided, for the first time, data for the high altitude periglacial domain of the High Atlas ~~Mountains~~. The period was marked by high temperatures in December and January, with very low precipitation from November to January, ~~which caused a the~~ late onset of the winter snow cover ~~when compared to other years~~. GST from 3,210 to 4,160 m a.s.l. showed two ~~very~~-contrasted periods: a hot season from late-May to late-September, and a long cold season, from mid-October to mid-April. The ~~hot season-former~~ showed positive air temperatures at all sites and was marked by very high daily temperature amplitudes, with maxima reaching  $40.1\text{ }^{\circ}\text{C}$  at Toubkal 3. This regime was controlled by the high solar radiation inputs and by the dryness of the soil. The cold season was marked by ~~either~~ subfreezing GST or by frequent freeze-thaw cycles, depending on snow conditions. Neltner in a valley floor at 3,210 m showed frequent freeze-thaw cycles during the start of the cold season, until a heavy snow fall event in mid-February, inducing GST to stabilize close to  $0\text{ }^{\circ}\text{C}$ . The high altitude sites (T2 ~~(-3,964)~~ and T1 ~~(-4,160)~~ m) located in wind swept areas showed subfreezing temperatures with frequent freeze-thaw cycles illustrating the lack of an insulating snow cover. These conditions prevailed during the whole cold season, with mean monthly GST below  $0\text{ }^{\circ}\text{C}$  from November to March.

~~The~~ The monitoring site Toubkal 3, ~~mini~~logger located in ~~Irhzer Ikhibi-South a valley in the lower section of a talus slope~~ at 3,820 m a.s.l. showed ~~the most~~ remarkable GST regime ~~during the cold season~~. After an early onset of a stable snow pack in mid-October, GST decreased regularly until mid-December and then showed minor oscillations around ci.  $-5.8\text{ }^{\circ}\text{C}$ . These conditions were only interrupted in late-March with snow melt, ~~and were followed by a zero curtain effect in April~~.

The ~~GST low-GST at regime Toubkal 3 at the site with values around -5 °C with low thermal amplitude, under the snow pack, shows supports~~ the likely presence of permafrost, ~~which agrees with the presence of poorly developed arcuate ridges and furrows in the area.~~ The site is in a valley ~~section~~ with long lasting snow cover ~~due to wind accumulation and -moderate potential incoming solar radiation and in the distal part of a talus, a setting which is known to favour lower ground temperatures. The~~  
5 ~~data and observations are still scarce and do not allow for conclusions on the origin of the lower temperatures at the site, nor at assessing their spatial significance. Possibly, the low GST are a result of the joint effects of snow, cold air flow within the talus and over the snow surface, as well as from the presence of interstitial/buried ice derived from rockfall events, faeing west, but close to a slope that suffers the shadowing effect of the ridge located southwards. The boulderly surface of the lower talus shows poorly developed ridges and furrows, which agree with the presence of permafrost.~~

10 ~~These results presented here~~ should be interpreted with care and although the data suggests the presence of permafrost, for a more accurate assessment, more observations are needed. ~~The possible presence of permafrost at 3,800 m in the Djebel Toubkal massif, to be confirmed, could become spatially significant, since such altitudes are frequent in the High Atlas, not only in the Toubkal region, but also further east, in the M'Goun massif. For the continuation of this research, forthcoming studies should target at: i. Electrical resistivity surveying to identify possible anomalies and the presence of ice, ii. Installing a larger number~~  
15 ~~of GST loggers at different settings around T3, iii) Installing an air temperature logger at T3 to assess the possible shadowing effect on air temperature and GST cooling, and iv) installation of a borehole for monitoring temperatures.~~

## 7. Outlook

~~-For the continuation of this research, forthcoming studies should target at: i. Electrical resistivity surveying to identify possible anomalies and the presence of ice, ii. Installing a larger number of GST loggers at different settings around T3, iii) Installing~~  
20 ~~an air temperature logger at T3 to assess the possible shadowing effect on air temperature and GST cooling, and iv) installation of a borehole for monitoring temperatures.~~

~~The possible presence of permafrost at 3,800 m in the Djebel Toubkal massif, to be confirmed, could become spatially significant, since such altitudes are frequent in the High Atlas, not only in the Toubkal region, but also further east, in the M'Goun massif. The presence of permafrost would also need to be assessed for: ice-content and consequent impacts for~~

25 hydrology (at least at a local level), thermal state and spatial distribution at the regional scale, ecological significance (vegetation communities, endemisms, refugia), possible presence of extremophiles and possible analysis of permafrost ice as a paleoenvironmental archive. Given the warming trend as shown by the Menara data-series, future climate scenarios and the very sensitive setting of the upper reaches of the High Atlas, just above 0 °C MAAT, further research at an interdisciplinary level is needed, since the possible permafrost remnants could quickly disappear facing climate change. And this can be the last  
30 permafrost remnant in North Africa.

Formatada: Cabeçalho 1

## Acknowledgements

Special thanks are due to Philip Hughes, who provided valuable advice on routes and logistics for the Toubkal massif and information on the geomorphology of the region. Warmest thanks to Sebastião Vieira, who participated in the field season of 2015 and helped with the setting of the loggers, for his fantastic company and patience. Mustapha Asquarray (Dar Assarou) from Imlil provided excellent guiding services and support in the Toubkal region, with Ibrahim ait Tadrart Buff and Mohammed Boyikd providing mule transport support. To the Ait Elkadi family guards of the Neltner Hut and to Hsin Nsliman at Sidi Chamharouch Cafe, we thank for the surveillance of the air temperature/RH loggers. Hassan Michalou is thanked for helping with the air temperature minilogger at the summit of Toubkal. The manuscript benefited from the comments of Enrique Serrano and Benno Staub, which conducted the review and to whom we thank. This research benefited from support of the bilateral project COLDATLAS – « Does permafrost occurs in the high mountains of North Africa? » funded by the FCT/CNRST (Portugal-Morocco), with partial funding from the CEG/IGOT - Universidade de Lisboa.

## References

- Arbolea, M. L., Teixell, A., Charroud, M. and Julivert, M.: A structural transect through the High and Middle Atlas of Morocco, *J. African Earth Sci.*, 39(3–5), 319–327, doi:10.1016/j.jafrearsci.2004.07.036, 2004.
- Badri, W., Gauquelin, T., Minet, J. and Savoie, J. M.: Données météorologiques nouvelles sur le massif de l'â€™ Oukaimeden (2570 m , Haut Atlas de Marrakech , Maroc ) : un exemple de climat de haute montagne méditerranéenne ., *Publ. l'Association Int. Climatol.*, 7, 190–198, 1994.
- Biskaborn, B. K., Lanckman, J.-P., Lantuit, H., Elger, K., Streletskiy, D. a., Cable, W. L. and Romanovsky, V. E.: The Global Terrestrial Network for Permafrost Database: metadata statistics and prospective analysis on future permafrost temperature and active layer depth monitoring site distribution, *Earth Syst. Sci. Data Discuss.*, 8(1), 279–315, doi:10.5194/essdd-8-279-2015, 2015.
- Boudhar, A., Hanich, L., Boulet, G., Duchemin, B., Berjamy, B. and Chehbouni, A.: Evaluation of the Snowmelt Runoff Model in the Moroccan High Atlas Mountains using two snow-cover estimates, *Hydrol. Sci. J.*, 54(March 2015), 1094–1113, doi:10.1623/hysj.54.6.1094, 2009.
- Boudhar, A., Boulet, G., Hanich, L., Sicart, J. E. and Chehbouni, A.: Energy fluxes and melt rate of a seasonal snow cover in the Moroccan High Atlas, *Hydrol. Sci. J.*, 6667(August 2015), 141217125340005, doi:10.1080/02626667.2014.965173, 2014.
- Chardon, M. and Riser, J.: Formes et processus géomorphologiques dans le Haut-Atlas marocain, *Rev. Géographie Alp.*, 69, 561–582, 1981.
- Cheggour, A.: Mesures de l'érosion hydrique à différentes échelles spatiales dans un bassin versant montagneux semi-aride et spatialisation par des S.I.G.: Application au bassin versant de la Rhéraya, Haut Atlas, Maroc, Cady Ayyad, Marrakesh., 2008.
- Couvreur, G.: Les formations périglaciaires du Haut Atlas central marocain, *Rev. Géographie du Maroc*, 10, 47–50, 1966.

- Delaloye, R. and Lambiel, C.: Evidence of winter ascending air circulation throughout talus slopes and rock glaciers situated in the lower belt of alpine discontinuous permafrost (Swiss Alps), *Nor. Geogr. Tidsskr. - Nor. J. Geogr.*, 59(2), 194–203, doi:10.1080/00291950510020673, 2005.
- Delaloye, R., Reynard, E., Lambiel, C., Marescot, L. and Monnet, R.: Thermal anomaly in a cold scree slope (Creux du Van, Switzerland), *Proc. Eighth Int. Conf. Permafrost, Zürich, Switzerland.*, 175–180 [online] Available from: <http://onlinelibrary.wiley.com/doi/10.1002/cbdv.200490137/abstract>, 2003.
- Dresch, J.: Recherches sur l'évolution du relief dans le Massif Central du Grand Atlas le Haouz et le Sous, Arrault et Cie, Maitres Imprimeurs, Tours., 1941.
- ESA: SENTINEL-2 User Handbook, , (1), 64, doi:GMES-S1OP-EOPG-TN-13-0001, 2015.
- Ferreira, A., Vieira, G., Ramos, M. and Nieuwendam, A.: Ground temperature and permafrost distribution in Hurd Peninsula (Livingston Island, Maritime Antarctic): An assessment using freezing indexes and TTOP modelling, *CATENA*, doi:10.1016/j.catena.2016.08.027, 2016.
- Gadek, B.: Debris slopes ventilation in the periglacial zone of the Tatra Mountains (Poland and Slovakia): The indicators, *Cold Reg. Sci. Technol.*, 74–75, 1–10, doi:10.1016/j.coldregions.2012.01.007, 2012.
- Gądek, B. and Kędzia, S.: Winter ground surface temperature regimes in the zone of sporadic discontinuous permafrost, Tatra Mountains (Poland and Slovakia), *Permafr. Periglac. Process.*, 19(3), 315–321, doi:10.1002/ppp.623, 2008.
- Giorgi, F. and Lionello, P.: Climate change projections for the Mediterranean region, , doi:10.1016/j.gloplacha.2007.09.005, 2007.
- Goodrich, L. E.: The influence of snow cover on the ground thermal regime, *Can. Geotech. J.*, 19(4), 421–432, doi:10.1139/t82-047, 1982.
- Gruber, S. and Haeberli, W.: Mountain Permafrost, in *Permafrost Soils*, pp. 33–44., 2009.
- Gubler, S., Fiddes, J., Keller, M. and Gruber, S.: Scale-dependent measurement and analysis of ground surface temperature variability in alpine terrain, *Cryosph.*, 5(2), 431–443, doi:10.5194/tc-5-431-2011, 2011.
- Haeberli, W.: Untersuchungen zur Verbreitung von Permafrost zwischen Fluelapass und Piz Grialetsch (GR), *Mitteilungen der VAW, ETH Zurich*, 17, 1975.
- Haeberli, W. and Patzelt, G.: Permafrostkartierung im gebiet der Hochebenkar-blockgletscher, Obergurgl, Otztaler Alpen, *Zeitschrift für Gletscherkd. und Glazialgeol.*, 2(18), 127–150, 1982.
- Haeberli, W., Noetzli, J., Arenson, L., Delaloye, R., Gärtner-Roer, I., Gruber, S., Isaksen, K., Kneisel, C., Krautblatter, M. and Phillips, M.: Mountain permafrost: development and challenges of a young research field, *J. Glaciol.*, 56(200), 1043–1058, doi:10.3189/002214311796406121, 2010.
- Hannah, G., Hughes, P. D. and Gibbard, P. L.: Pleistocene plateau ice fields in the High Atlas, Morocco, in *Quaternary glaciation in the Mediterranean mountains*, vol. 433, edited by P. D. Hughes and J. C. Woodward, p. SP433.12, Geological Society of London., 2016.
- Hauck, C., Vieira, G., Gruber, S., Blanco, J. and Ramos, M.: Geophysical identification of permafrost in Livingston Island,

- Maritime Antarctica, *J. Geophys. Res. Earth Surf.*, 112(2), 1–15, doi:10.1029/2006JF000544, 2007.
- Hoelzle, M.: Permafrost occurrence from BTS measurements and climatic parameters in the eastern Swiss Alps, *Permafr. Periglac. Process.*, 3(2), 143–147, doi:10.1002/ppp.3430030212, 1992.
- Hu, W., Zhang, Q., Tian, T., Cheng, G., An, L. and Feng, H.: The microbial diversity, distribution, and ecology of permafrost in China: a review, *Extremophiles*, 19(4), 693–705, doi:10.1007/s00792-015-0749-y, 2015.
- Hughes, P. and Woodward, J.: Timing of glaciation in the Mediterranean mountains during the last cold stage, *J. Quat. Sci.*, 23, 575–588, doi:10.1002/jqs, 2008.
- Hughes, P. D.: Little Ice Age glaciers in the Mediterranean mountains, *Méditerranée*, 122, 63–79, doi:10.4000/mediterranee.7146, 2014.
- Hughes, P. D., Fenton, C. R. and Gibbard, P. L.: Quaternary Glaciations of the Atlas Mountains, North Africa., 2011.
- Hughes, P. D., Fink, D., Fletcher, W. J. and Hannah, G.: Catastrophic rock avalanches in a glaciated valley of the High Atlas, Morocco: <sup>10</sup>Be exposure ages reveal a 4.5 ka seismic event, *Geol. Soc. Am. Bull.*, doi:10.1130/B30894.1, 2014.
- Ishikawa, M.: Thermal regimes at the snow – ground interface and their implications for permafrost investigation, , 1264, 1–16, 2002.
- Ishikawa, M.: Thermal regimes at the snow-ground interface and their implications for permafrost investigation, *Geomorphology*, 52(1–2), 105–120, doi:10.1016/S0169-555X(02)00251-9, 2003.
- Jansson, J. K. and Taş, N.: The microbial ecology of permafrost, *Nat. Rev. Microbiol.*, 12(6), 414–425, doi:10.1038/nrmicro3262, 2014.
- Knippertz, P., Christoph, M. and Speth, P.: Long-term precipitation variability in Morocco and the link to the large-scale circulation in recent and future climates, *Meteorol. Atmos. Phys.*, 83(1–2), 67–88, doi:10.1007/s00703-002-0561-y, 2003.
- Lacelle, D. and Vasil'chuk, Y. K.: Recent Progress (2007-2012) in Permafrost Isotope Geochemistry, *Permafr. Periglac. Process.*, 24(2), 138–145, doi:10.1002/ppp.1768, 2013.
- Lambiel, C. and Pieracci, K.: Permafrost distribution in talus slopes located within the alpine periglacial belt, Swiss Alps, *Permafr. Periglac. Process.*, 19(3), 293–304, doi:10.1002/ppp.624, 2008.
- Lewkowicz, A. G. and Ednie, M.: Probability mapping of mountain permafrost using the BTS method, Wolf Creek, Yukon Territory, Canada, *Permafr. Periglac. Process.*, 15(1), 67–80, doi:10.1002/ppp.480, 2004.
- Marchane, A., Jarlan, L., Hanich, L., Boudhar, A., Gascoin, S., Tavernier, A., Filali, N., Le Page, M., Hagolle, O. and Berjamy, B.: Assessment of daily MODIS snow cover products to monitor snow cover dynamics over the Moroccan Atlas mountain range, *Remote Sens. Environ.*, 160, 72–86, doi:10.1016/j.rse.2015.01.002, 2015.
- Mark, B. G. and Osmaston, H. A.: Quaternary glaciation in Africa: Key chronologies and climatic implications, *J. Quat. Sci.*, 23(6–7), 589–608, doi:10.1002/jqs.1222, 2008.
- de Martonne, E.: Les formes glaciaires sur le versant nord du Haut Atlas, *Ann. Geogr.*, 33(183), 296–302, 1924.
- Montanari, B.: The Future of Agriculture in the High Atlas Mountains of Morocco: The Need to Integrate Traditional Ecological Knowledge, in *The Future of Mountain Agriculture*, pp. 51–72, Springer Berlin Heidelberg, Berlin, Heidelberg.,

- 2013.
- N'da, A. B., Bouchaou, L., Reichert, B., Hanich, L., Ait Brahim, Y., Chehbouni, A., Beraaouz, E. H. and Michelot, J.-L.: Isotopic signatures for the assessment of snow water resources in the Moroccan high Atlas mountains: contribution to surface and groundwater recharge, *Environ. Earth Sci.*, 75(9), 755, doi:10.1007/s12665-016-5566-9, 2016.
- 5 Oliva, M., Gómez-Ortiz, A., Salvador-Franch, F., Salvà-Catarineu, M., Palacios, D., Tanarro, L., Ramos, M., Pereira, P. and Ruiz-Fernández, J.: Inexistence of permafrost at the top of the Veleta peak (Sierra Nevada, Spain), *Sci. Total Environ.*, 550(April), 484–494, doi:10.1016/j.scitotenv.2016.01.150, 2016a.
- Oliva, M., Serrano, E., Omez-Ortiz, A. G., Gonz Alez-Amuchastegui, M. J., Nieuwendam, A., Palacios, D., Erez-Alberti, A. P., Pellitero-Ondicol, R., Ruiz-Fern Andez H, J., Valc Arcel, M., Vieira, G. and Antoniades, D.: Spatial and temporal variability of perglaciation of the Iberian Peninsula, , doi:10.1016/j.quascirev.2016.02.017, 2016b.
- 10 Onaca, A., Ardelean, A. C., Urdea, P., Ardelean, F. and Sîrbu, F.: Detection of mountain permafrost by combining conventional geophysical methods and thermal monitoring in the Retezat Mountains, Romania, *Cold Reg. Sci. Technol.*, 119(August), 111–123, doi:10.1016/j.coldregions.2015.08.001, 2015.
- Outcalt, S. I., Nelson, F. E. and Hinkel, K. M.: The zero???curtain effect: Heat and mass transfer across an isothermal region in freezing soil, *Water Resour. Res.*, 26(7), 1509–1516, doi:10.1029/WR026i007p01509, 1990.
- 15 Peyron, M.: Les chutes de neige dans l'Atlas marocain, *Rev. géographie Alp.*, 68(3), 237–254, doi:10.3406/rga.1980.2203, 1980.
- Rangecroft, S., Harrison, S., Anderson, K., Magrath, J., Castel, A. P. and Pacheco, P.: Climate change and water resources in arid mountains: An example from the bolivian andes, *Ambio*, 42(7), 852–863, doi:10.1007/s13280-013-0430-6, 2013.
- 20 Rauh, W.: *Vegetationsstudien im Hohen Atlas un dessen Vorland*, Springer-Verlag., 1952.
- Robinson, D. A. and Williams, R. B. G.: Sandstone weathering in the High Atlas, Morocco, *Zeitschrift fur Geomorphol.*, 36(4), 413–429, 1992.
- Ros, R. M., Cano, M. J., Muñoz, J. and Guerra, J.: Contribution to the bryophyte flora of Morocco: the Jbel Toubkal, *J. Bryol.*, (22), 283–289, 2000.
- 25 Salvador Franch, F., Gómez Ortiz, A., Salvà Catarineu, M. and Palacios Estremera, D.: Caracterización térmica de la capa activa de un glaciar rocoso en medio periglaciario de alta montaña mediterránea : El ejemplo del Corral del Veleta (Sierra Nevada, España), *Cuad. Investig. Geográfica*, 37(2), 25, doi:10.18172/cig.1255, 2011.
- Sawada, Y., Ishikawa, M. and Ono, Y.: Thermal regime of sporadic permafrost in a block slope on Mt. Nishi-Nupukaushinupuri, Hokkaido Island, Northern Japan, *Geomorphology*, 52(1–2), 121–130, doi:10.1016/S0169-555X(02)00252-0, 2003.
- 30 Scapozza, C., Lambiel, C., Baron, L., Marescot, L. and Reynard, E.: Internal structure and permafrost distribution in two alpine periglacial talus slopes, Valais, Swiss Alps, *Geomorphology*, 132(3–4), 208–221, doi:10.1016/j.geomorph.2011.05.010, 2011.
- Schulz, O. and de Jong, C.: Snowmelt and sublimation: field experiments and modelling in the High Atlas Mountains of Morocco, *Hydrol. Earth Syst. Sci.*, 8, 1076–1089, doi:10.5194/hess-8-1076-2004, 2004.

Serrano, E., Morales, C., González-Trueba, J. and Martín, R.: Cartografía del permafrost de montaña en los Pirineos españoles, Finisterra. Rev. Port. Geogr., 44(87), 45–54 [online] Available from: [https://www.redib.org/recursos/Record/oai\\_articulo665186-cartografia-permafrost-montana-pirineos-espanoles](https://www.redib.org/recursos/Record/oai_articulo665186-cartografia-permafrost-montana-pirineos-espanoles), 2009.

Simonneaux, V., Cheggour, A., Deschamps, C., Mouillot, F., Cerdan, O. and Le Bissonnais, Y.: Land use and climate change effects on soil erosion in a semi-arid mountainous watershed (High Atlas, Morocco), J. Arid Environ., 122, 64–75, doi:10.1016/j.jaridenv.2015.06.002, 2015.

Smith, S. and Brown, J.: Essential Climate Variables: Permafrost and seasonally frozen ground, GTOS, 62, 22, 2009.

Staub, B. and Delaloye, R.: Using Near-Surface Ground Temperature Data to Derive Snow Insulation and Melt Indices for Mountain Permafrost Applications, Permafr. Periglac. Process., 248(March 2015), 237–248, doi:10.1002/ppp.1890, 2016.

Tanarro, L. M., Hoelzle, M., García, A., Ramos, M., Gruber, S., Gómez, A., Piquer, M. and Palacios, D.: Permafrost distribution modelling in the mountains of the Mediterranean: Corral del Veleto, Sierra Nevada, Spain, Nor. Geogr. Tidsskr. - Nor. J. Geogr., 55(4), 253–260, doi:10.1080/00291950152746612, 2001.

USGS: Landsat 8 (L8) Data users handbook., 2016.

Vieira, G., Mora, C. and Ramos, M.: Ground temperature regimes and geomorphological implications in a Mediterranean mountain ( Serra da Estrela , Portugal ), Geomorphology, 52, 57–72, doi:10.1016/S0169-555X(02)00248-9, 2003.

Wilche, K.: Klimamorphologische und talgeschichtliche studien im M’Goungebiet, Mitt. Der Geogr. Ges. Wien, 95, 4–41, 1953.


Williams, P. J. and Smith, M. W.: The Frozen Earth, Cambridge Univ Press., 1989.

Yoshikawa, K.: Africa, Kilimanjaro, in Permafrost in our time, edited by K. Yoshikawa, pp. 238–239, University of Alaska Fairbanks, Valencia, California., 2013.

Zahour, G., Hadi, H. El, Tahiri, A., Zerhouni, Y., Alikouss, S., Zahour, R. and Reddad, A.: The Late Neoproterozoic Continental Tholeiitic Basalts of the Toubkal Inlier (Western High-Atlas, Morocco): A Post-Pan-African Rifting Witness in the Northern Margin of the West African Craton, Open J. Ecol., 6(8), 509–516, doi:10.4236/oje.2016.68048, 2016.


Sites	Alt (m)	Model	Variables	Interval (h)
Sidi Chamharouch	2,370	Hobo Prov2	AT, HR	1
Neltner (NLT)	3,200	Hobo Prov2, TidBit	AT, HR, GST	1
Toubkal 3 (T3)	3,815	TidBit	GST	1
Toubkal 2 (T2)	3,980	TidBit	GST	1
Toubkal 1 (T1)	4,160	Ibutton, TidBit	AT, GST	4,1

**Table 1 – Temperature data loggers installed in the High Atlas. Variables: GST -Ground surface temperature, AT - Air temperature, HR - Relative humidity.**




5 Figure 1 – Location and topography of the Toubkal Massif study area. Yellow circles are the sites of the data loggers. Contour equidistance is 50 m.





5 **Figure 2: Geomorphological sketch map of the Toubkal Massif.**

**Formatada:** Tipo de letra: Negrito



**Figure 23:** Location of the ground surface temperature loggers: a. Neltner (3,200 m), b. Toubkal 3 (3,815 m), c. Toubkal 2 (3,980 m), d. Toubkal 1 (4,160 m).




Figure 43: Statistics of monthly temperature and precipitation at Menara (Marrakesh) from 1977 to 2015 and records of the study period (July 2015 – March 2016). Source: NCDC/GHCN.






Figure 54: Mean monthly air temperatures from July 2015 to June 2016 in Marrakesh (Menara) and, in the two study sites Sidi Chamharouch (2,370 m) and Neltner (3,210 m) and extrapolated to the summit of Djebel Toubkal (4,167 m a.s.l.).



**Figure 65:** Satellite true colour composite scenes from Landsat-8 (USGS) and Sentinel-2 (ESA) used for assessing the snow cover at the monitoring sites from September 2015 to June 2016. Light blue indicates snow cover and brown and green indicate snow free terrain. Satellite imagery obtained from USGS EarthExplorer.



**Figure 67:** Monthly ground surface and air temperatures in the Djebel Toubkal massif from June 2015 to July 2016. Extremes are absolute monthly maximum and minimum temperatures.





Figure 78: Mean annual ground surface temperatures vs altitude for the 4 monitored sites in the Djebel Toubkal massif.






**Figure 9: Distribution of annual potential solar radiation in the Toubkal massif, with indication of the ground surface temperature monitoring sites.**

Formatada: Normal, Esquerda


Formatada: Tipo de letra: Negrito





**Figure 108:** Mean monthly ground surface temperatures at the 4 study sites in the Toubkal Massif from July 2015 to June 2016.





**Figure 119:** Ground surface temperature regimes at the 4 sites in the Toubkal Massif and air temperature at Neltner. The plot shows hourly data. Symbols indicate the snow cover conditions at the date of satellite scenes: - no snow, + possible snow/snow margin, ++ snow, +++ - significant snow, ? - Uncertainty in classification.





Figure 102: Example of the hourly GST regime at Toubkal 3 and comparison with air temperatures measured at Neltner and extrapolated to T3 using the observed lapse rate of  $-0.59^{\circ}\text{C}.100\text{ m}^{-1}$ .

5



Figure 143: The Irhzer Ikhibi south valley, where the datalogger Toubkal 3 was installed. The arrows indicate arcuate boulder ridges and furrows in the talus slope.



**Figure 124:** Estimated mean annual air temperatures at the Djebel Toubkal (4,167 m asl) obtained from extrapolation of temperatures from Menara (Marrakesh) using a lapse rate of  $-0.59^{\circ}\text{C} \cdot 100 \text{ m}^{-1}$ .