

We very much thank the reviewer for your valuable comments and suggestions. Answers and explanations to all detailed questions and annotations raised by the reviewer is provided in the following. **Marked-up manuscript version was add at the end.**

(RC: Reviewer comments; AC: Author comments).

Reviewer comments

Specific Comments

RC: I found this new version of the manuscript greatly improved with respect to the first one. But I think that further work is needed. The data presented in this work are so abundant and rich that in order to prepare a balanced paper also a great effort in terms of interpretation and data-analysis are required. This is the weak side of the manuscript. But I am sure that with a little bit more work it will be possible to draw interesting conclusions and evidences. In general, I suggest to the authors to shorten a lot of sections. Reading the manuscript sometimes it seems to read a phd-thesis, not a draft of a paper. The space given to references and to their explanations is simply too much. Please consider to resume the most important information taken from literature in a more concise way, not presenting each single reference with an entire sentence, which describe it. This is particularly true for chapter 3.2, but also other sections of the work are similar.

AC: *We tried to simplify the text by removing some parts. We also split the text into more chapters making reading easier.*

New chapters:

3.1 Dating of the ice core

3.2 Glaciochemical records

3.3 Principal component analysis

3.4 Intraseasonal concentration fluxes variability

3.5 Interannual atmospheric variability

3.6 Atmospheric transport to Mount Johns ice core site

3.7 Relationships between atmospheric circulation, temperature, and sea ice concentration

RC: Moving to the content I have some doubts about the interpretation. What is lacking is a comprehensive analysis of all the data. At the present state, the discussion about elemental fractions, PCA and cluster is completely separated from back trajectories

analysis and from the meteorological considerations. TO draw robust conclusions putting all the pieces together is necessary.

AC: *We added a new chapter, “3.7 Relationships between atmospheric circulation, temperature, and sea ice concentration“, where all results are discussed together and in a clearer way.*

RC: Some details about the analysis of chemical data. At first the equation system presented to distinguish the ss and nss fractions of Al and Na is not convincing. In the equations, you show that nss Na depends on nss Al and that ss Al depends on ss Na. Without setting, a reference how is it possible to solve this system? Sodium is a function of Aluminum and vice versa, I really could't understand. You need to take an assumption if you want to solve the problem. I think that the only reasonable possible one is to assume that Al is only crustal and that nss Al is null. Assuming this point it will be possible to distinguish Na ss and nss fractions.

AC: *Yes, we were assuming Al as crustal origin only; the sentence was rewritten making it clear. The text was changed to: “In this study, we assume that Al is only from crustal origin (total Al is equal to nssAl).”*

RC: About PCA. I don't really agree with your interpretation of PC. In addition I think that with such good results their interpretation should be deepened. My idea about PCA is:

-PC1 is a sort of impurity index. We see that each element is positively correlated with this PC. This means that when the atmospheric load is high (high impurity content) PC1 show high values. At the opposite when impurity concentration reaches minima values PC1 is low. It would be interesting to show the record of PC1 along depth (or age) and see if presents seasonal values, I guess so.

-PC2 and PC3 are related to dust vs marine aerosol deposition. Here we see that Ba, Fe, Ti, Al are positively correlated, while other elements related to marine aerosol as K and Na are negatively correlated. These components distinguish the two fractions. Probably if we look further in deep we could say that PC2 highlights aluminosilicate (Al is the most important element), while PC3 highlights the siliclastic fraction of mineral dust (Fe and Ti).

-PC4 seems related to carbonate content, being Ca and Sr major constituents of CaCo₃. I suggest to the authors to better develop this part, studying the trends related to each

component along the core and trying to understand the different seasonal cycles. This could help also for the successive part dedicated to the analysis of atmospheric transport and pathways.

AC: We agree. The section is being rewritten and improved. We agreed that some interpretations were superficial and we are expanding on these issues.

RC: Another important problem concerns the use of elemental concentration instead of fluxes. Since this is a well-dated core, with a good seasonal signal, I think that it would be worth to convert all concentrations in fluxes. In this way, it would be possible to highlight the real changes of deposition, excluding the effect associated to snow dilution. What is said here is that 3 major periods are recognized in terms of deposition, related to high and low concentration. Maybe such an effect is only related to changes of snow precipitation. Using fluxes such problems would be completely excluded and final interpretation would be improved.

AC: Both concentration and flux presents similar variability, this indicates that the concentrations are independent of accumulation rate and that the variability exhibited in the record is likely not a function of changes in snow accumulation. Anyway, now we are presented the fluxes instead elemental concentrations, in the text.

Technical Comments

RC: 102 and 108: class100, add ISO5

AC: Done

RC: 110: change to This process is important to guarantee a complete dissolution of particulate and non-soluble elemental fractions.

AC: Done

RC: 119: applied to, not as

AC: Done

RC: 122: (such event occurred in less than 1% of the samples); minor, not less

AC: Done

RC: 180-182: Al is not chosen because of its abundance. Al is chosen because in addition to being abundant (and easy to determine) is also quite stable and immobile from a chemical perspective. Thus, it can be reliably used as crustal tracer. (See for example Nesbitt and Young 1982)

AC: *The text was changed to: "Aluminum was used as the reference element in this work because it is a good proxy of continental dust"*

RC: 190: see my comments about Na and Al

AC: *Done*

RC: 217: I don't understand which those percentages are related to. In Tab S3 it seems that global volcanic deposition is more important than the local one, but in the text you state the opposite. Is it wrong or am I missing something?

AC: *The values were in reverse order. My mistake.*

RC: 249: specify that while the peak of species associated to marine aerosol is mainly related to winter sea ice peak, the peak of dust deposition is related mainly to atmospheric circulation patterns. In coastal sites such dynamics are greatly influenced by local factors, for this reason the dust peak doesn't occur in the same season in whole Antarctica, each site presents its peculiarities.

AC: *The section is being rewritten and improved.*

RC: 259: associated to

AC: *Done*

A 125-year record of climate and chemistry variability at the Pine Island Glacier ice divide, Antarctica

Franciele Schwanck¹, Jefferson C. Simões¹, Michael Handley², Paul A. Mayewski², Jeffrey D. Auger², Ronaldo T. Bernardo¹, Francisco E. Aquino¹

¹Centro Polar e Climático, Universidade Federal do Rio Grande do Sul (UFRGS), Porto Alegre, 91540-000, Brazil

²Climate Change Institute, University of Maine, Orono, 04469, United States

Correspondence to: Franciele Schwanck (franschwanck@gmail.com)

Abstract. The Mount Johns (MJ) ice core (79°55'S; 94°23'W) was drilled near the Pine Island Glacier ice divide on the West Antarctic Ice Sheet during the 2008–2009 austral summer, to a depth of 92.26 m. The upper 45 m of the record covers approximately 125 years (1883–2008) showing marked seasonal variability. Trace element concentrations in 2,137 samples were determined using inductively coupled plasma mass spectrometry. In this study, we reconstruct mineral dust and sea salt aerosol transport and investigate the influence of climate variables on the elemental concentrations to the MJ site. The ice core record reflects changes in emissions as well as atmospheric circulation and transport processes. Our trajectory analysis shows distinct seasonality, with strong westerly transport in the winter months and a secondary northeasterly transport in the summer. During summer months, the trajectories present slow-moving (short) transport and are more locally influenced than in other seasons. Finally, our reanalysis correlations with trace element suggest that marine derived trace element concentrations are strongly influenced by sea ice concentration and sea surface temperature anomalies. The results show that seasonal elemental concentration maxima in sea-salt elements correlate well with the sea ice concentration winter maxima in the West Amundsen and Ross Seas. Lastly, we observed an increased concentration of marine aerosols when sea surface temperature decreased.

1 Introduction

The West Antarctic Ice Sheet (WAIS) is more susceptible to marine influences than East Antarctica Ice Sheet (EAIS). The lower average elevation of the WAIS compared to EAIS, 1100 m and 3000 m, respectively, (Bedmap 2 project data; Fretwell et al., 2013) facilitates the advection of air masses toward the interior of the continent, thereby directly contributing to the ice sheet's surface mass balance through precipitation (Nicolas and Bromwich, 2011). During recent decades, rapid changes have occurred in the WAIS sector, including flow velocity acceleration, retraction of ice streams, and mass loss (Pritchard et al., 2012). These changes influence the global climate through their contributions to sea level rise (Pritchard et al. 2009; Shepherd et al. 2012) and deep ocean circulations (Holland and Kwok, 2012). WAIS contains sufficient water to raise the global sea level by over 3 m (Bamber et al., 2009; Fretwell et al., 2013).

Major and minor impurities in polar ice originate from sources such as oceans, landmasses, volcanism, biogenic activity, biomass burning, and anthropogenic inputs (Legrand and Mayewski, 1997; Planchon et al., 2002; Vallelonga et al., 2004; Weller et al., 2008; Dixon et al., 2013; Schwanck et al., 2016). These impurities are transported over long distances to the polar regions through the troposphere (Petit and Delmonte, 2009) and stratosphere, e. g. volcanic emissions (Krinner et al., 2010). Sea salt and mineral dust can be used to reconstruct climate conditions as well as atmospheric transport patterns (Albani et al., 2012;

Chewings et al., 2014). These chemical elements are strongly influenced by the climate conditions in the source region, and the depositional record provides important information about cyclone activity, wind intensity (Koffman et al., 2014a), sea ice conditions (Criscitiello et al., 2014), and aridity and vegetation cover (McConnell et al., 2007).

Marine aerosol concentrations are strongly linked to cyclone frequency and intensity that provides high wind speeds over the ocean surface, with the aerosols deposited along the storm track (Fischer et al., 2004). Another primary source of trace elements is mineral dust. It is transported from arid, continental regions such as Australia (Revel-Rolland et al., 2006; Li et al., 2008), South America (Delmonte et al., 2010; Li et al., 2010), and probably New Zealand (Neff and Bertler, 2015) to the Antarctic. Once entrained into the atmosphere, small grains (0.1 to 5 μm) can be transported over long distances (Gaiero et al., 2007; Mahowald et al., 2014) through advection before they are deposited on the Antarctic snow surface (Prospero et al., 2002; Delmonte et al., 2013).

Here, we present a 125-year record of eleven trace elements (Al, Ba, Ca, Fe, K, Mg, Mn, Na, S, Sr, and Ti) from the Mount Johns (MJ) ice core in the WAIS. We focus on the influence of atmospheric circulation, surface temperature, and sea ice concentration on the transportation of mineral dust and sea salt aerosol to the MJ site. Correlations were made between trace element concentrations and the European Centre for Medium Range Weather Forecasts (ECMWF) Reanalysis Interim (ERA-Interim; Dee et al., 2011) climate variables. Modeled air mass trajectories were also employed to explore atmospheric transport, independent of deposition processes, from 1979 to 2008.

2 Methods

2.1 Site description and core collection

Presently, Pine Island Glacier (PIG) is responsible for 20 % of the total ice discharge from the WAIS (Rignot et al., 2008). Accelerated thinning observed since the 1980s is directly linked to enhanced sub-ice-shelf melting, which is induced by the recent alteration of Circumpolar Deep Water (Pritchard et al., 2012; Steig et al., 2012; Favier et al., 2014). Sea level pressure and geopotential height anomalies, associated with increased strength of the circumpolar westerlies (Steig et al., 2012), favor reduced sea ice extent in the Amundsen and Bellingshausen Seas (Schneider et al., 2011) and the advection of warm air onto the continent (Steig et al., 2009; Ding et al., 2011). In this context, both atmospheric and oceanic variability are important for determining the response of the WAIS over long time-scales.

In this study, we use an ice core drilled at the nunatak Mount Johns and near the Pine Island Glacier divide (Fig. 1) to reconstruct mineral dust and marine aerosol transport and the influence of climate variables on the elemental concentrations. The MJ ice core (79°55'28" S, 94°23'18" W, and 91.20 m depth) was recovered in the austral summer of 2008/2009. The ice thickness at the ice core site reaches 2,115 m (determined from the Bedmap 2 project data; Fretwell et al., 2013). The average accumulation rate for the period is 0.21 m water equivalent per year (based on calculated field measurements of snow and ice density) and the mean surface temperature (measured at a depth of 12 m) is -33°C, measured using a calibrated platinum probe.

Drilling was performed using the Fast Electromechanical Lightweight Ice Coring System (FELICS) (Ginot et al., 2002). The MJ core (8.5 cm diameter) was cut into sections of approximately 1 m in length, packed

in polyethylene bags and then stored in high-density Styrofoam boxes and transported by air to Punta Arenas, Chile. Then, it was sent frozen to the Climate Change Institute (CCI) at the University of Maine, USA, where it was sub-sampled and analyzed using Inductively Coupled Plasma Sector Field Mass Spectrometry (ICP-SFMS).

2.2 Laboratory analyses

Decontamination of the MJ ice core was carried out in a certified cold room **ISO 5** with temperature below -20°C . Prior to melting, core ends and breaks were manually scraped using a clean ceramic knife to reduce potential contamination from drilling or other sources (procedure performed according Tao et al., 2001 and described in Schwanck et al., 2016). The core was melted using a continuous ice core melter system developed by CCI researchers (details in Osterberg et al. 2006). This system uses fraction collectors to gather discrete, high resolution, continuous and co-registered meltwater samples. Our samples were collected into acid-cleaned (Optima HNO_3), low density polyethylene (LDPE) vials and acidified to 1 % with double-distilled HNO_3 . All sample handling was conducted in an **ISO 5** clean room under a laminar flow High Efficiency Particle Air (HEPA) bench. The samples were allowed to react with the acid (at room temperature) at least one month before analysis. **This process is important to guarantee a complete dissolution of particulate and non-soluble elemental fractions** (following studies Rhodes et al., 2011 and Koffman et al., 2014b).

Trace element concentrations in 2,137 discrete samples, corresponding to the upper 45 m of the MJ ice core, were determined using the CCI Thermo Electron Element2 ICP-SFMS coupled to an ESI model SC-4 autosampler. Working conditions and measurement parameters are described in Table S1 (Supporting Information). The ICP-SFMS is calibrated daily with five standards that bracket the expected sample concentration range. Although there is no certified reference material for trace elements in polar snow and ice, the analyzed samples were certified with water reference material (SLRS-4, National Research Council Canada, Ottawa, Canada) to ensure the concentrations were within the certification range, confirming the accuracy of this method (details in Osterberg et al., 2006).

Samples of de-ionized water, or “blanks”, were prepared, treated, and analyzed using the same method **applied to** snow samples. The Method Detection Limits (MDL) were defined as three times the standard deviation of blank samples (10 blank samples were used). Concentrations below the MDL were replaced with the MDL values, which occurred in a few cases for Al, Ba, Ca, Fe, Mn, Sr, and Ti (**minor** than 1 %). Blank concentrations and MDLs were either similar or less than published values using comparable methods and instruments (Table S2 – Supporting Information).

Water isotope analyses were performed at the Centro Polar e Climatico, Brazil, Isotopes Lab using a Picarro L2130-i wavelength-scanned cavity ring-down spectroscopy (WS-CRDS) instrument (Picarro Inc., USA). The samples collected at the CCI were sent frozen to Brazil and melted on the day of analysis. Aliquots of water were filled in 2 ml glass vials and sealed with polytetrafluorethylene/silicone caps. The vials were then placed in a PAL COMBI-HTCxt autosampler (CTC Analytics AG, Switzerland) connected to the Picarro L2130-i for δD and $\delta^{18}\text{O}$. Reproducibility of measurements is typically 0.9 ‰ for δD and 0.2 ‰ for $\delta^{18}\text{O}$.

2.3 Meteorological data analyses

To explore possible sources of the observed trace elements in aerosols, air mass backward trajectories were simulated for 1000 m above ground level over the MJ ice core site. Trajectory simulations were made using the Hybrid Single-Particle Lagrangian Integrated Trajectory (HySPLIT) model, developed by the NOAA Air Resources Laboratory (Draxler et al., 2010) in conjunction with the global reanalysis datasets from the National Centers for Environmental Prediction (NCEP) and the National Center for Atmospheric Research (NCAR), known as NCEP/NCAR reanalysis model (NCEP1) (Kalnay et al., 1996; Kistler et al., 2001). Despite limitations prior to the satellite era (1979), the NCEP/NCAR reanalysis model represents a useful tool for understanding the climate of the Southern Hemisphere since 1979 (Bromwich and Fogt, 2004) and has been used with success in Antarctic back-trajectory modeling (e.g. Sinclair et al., 2010; Dixon et al., 2011; Markle et al., 2012).

Five-day (120 hr), 3-D back-trajectories were created from the MJ site at 00:00 UTC daily from January 1979 to December 2008 (a total of 10,655 trajectories). We have tested the effect of model initiation heights at 500 m, 1000 m, and 1500 m on our trajectories and confirmed they are spatially consistent. The initial heights of 500 m are affected by surface topography, which is imperfectly represented in the reanalysis model (Dee et al., 2011). For this reason, we have chosen the 1000 m level as the initial height condition for our back trajectories. At this altitude, orographic influences are minimized while the trajectories are sufficiently close to the terrain to be dynamically linked to the surface wind field (Sinclair et al., 2010). The five day simulation is an appropriate time-length when considering the maximum lifetime transport (10 days) of small size (0.1 – 2.5 μm) fractions of mineral dust and other aerosols, while transport of large particles (> 2.5 μm) is likely restricted to the first several days (Albani et al., 2012). Cataldo et al. (2013) measured the size of dust particles in a separate ice core drilled near MJ and found that the mean dust size within the core ranges from 1.2 μm to 2.4 μm .

In order to obtain information about airflow patterns at the MJ site, a cluster analysis was applied to a database of individual trajectories (10,655 daily trajectories). The HySPLIT model's cluster analysis algorithm groups trajectories by minimizing the spatial variability between trajectories within some defined number of clusters (Draxler, 1999). For the trajectories presented here, it is determined that five clusters are sufficient to capture seasonal variability during the 1979 to 2008 period.

Bracegirdle and Marshall (2012) determined that ERA-Interim was the most accurate of six reanalysis models over Antarctica when compared against surface and midtropospheric pressure and temperature observations. ERA-Interim was thus utilized to provide annual mean 2-m air temperature (T_{2m}), sea surface temperature (SST), and sea ice concentration (SIC) from 1979 to 2008. ERA-Interim outputs were obtained from the ECMWF Data Server (<http://apps.ecmwf.int/datasets/>) at a resolution of 1.5°. Spatial correlations were performed between MJ ice core element data and ERA-Interim climate variables.

3 Results and discussion

3.1 Dating of the ice core

The aerosol records show strong seasonal variations due to seasonality in aerosol sourced and transport efficiency, and these seasonal signals can be used to detect annual layers (e.g. Sigl et al., 2016). A typical annual layer at MJ site is characterized by a maximum of sea-salt aerosol deposition (e.g. Na, Sr) during

austral winter, and a maximum of marine biogenic aerosol emission (e.g. S) in late austral summer. Manual interpretation of the data was done by multiple individuals to identify the individual layers. The CCI software package (Kurbatov et al., 2005) was also used to identify matching seasonal peaks from Ca, Na and Sr and the major historical volcanic eruptions. In this study, water isotopes were used to confirm the dating previously performed in Schwanck et al., 2016. The major historical volcanic events during this period, such as Pinatubo (1991), Agung (1963), Santa Maria (1902), and Krakatoa (1883) are identified by large peaks in S concentration and are used as absolute time horizons during timescale development (Fig. S1 – Supporting Information). On the basis of our dating, the upper 45 m of the MJ ice core covers the period from the 1883 to 2008 A. D. We estimate a maximum dating error of ± 1 year between the 1991 A.D. Pinatubo and 1963 A.D. Agung eruptions, and ± 2 years between the 1963 A.D. Agung and 1883 A.D. Krakatoa eruptions.

3.2 Glaciochemical records

Concentrations of 11 trace elements (Al, Ba, Ca, Fe, K, Mg, Mn, Na, S, Sr, and Ti) were measured in 2,137 discrete ice core samples. Table 1 shows a statistical summary of the trace element concentrations measured from the MJ ice core.

The soil and rock dust contribution for the measured trace elements were given by the crustal Enrichment Factor (EF_C) according to Eq. (1) (Osterberg, 2007):

$$EF_C = \frac{X_{ice}/Al_{ice}}{X_{ref}/Al_{ref}}, \quad (1)$$

where X_{ice} is the trace element concentration in the sample, Al_{ice} is the aluminum concentration in the sample, and X_{ref} and Al_{ref} are the trace element and the aluminum concentrations in the reference material, respectively. Aluminum was used as the reference element in this work because it is a good proxy of continental dust (Planchon et al., 2002). The mean elemental concentration used for reference is the average composition of the upper continental crust taken from the literature (Wedepohl, 1995).

Figure 2 shows trace element EF_C values during the austral summer and winter months in the MJ ice core. Elements with EF_C lower than 10 are considered to be non-enriched and predominantly have a crustal dust origin (Duce et al., 1975). EF_C higher than 10 indicates contributions from other sources, such as marine aerosol, volcanism, biogenic activity or anthropogenic emissions. Marine aerosol contributions were estimated using the oceanic enrichment factor (EF_O) according to Eq. (2) (Osterberg, 2007):

$$EF_O = \frac{X_{ice}/Na_{ice}}{X_{ref}/Na_{ref}}, \quad (2)$$

where X_{ice} is the trace element concentration in the sample, Na_{ice} is the Na concentration in the sample, and X_{ref} and Na_{ref} are the trace element and Na concentration in the reference material, respectively. Sodium is used as the reference element because it is the main sea salt constituent (Weller et al., 2008; Dixon et al., 2013). We used the average composition of ocean water (Lide, 2005) as a reference for the ocean elemental abundances.

Non-sea-salt sulfur ratios were calculated using Eq. (3) (Palmer et al., 2002; Becagli et al., 2005):

$$nssS = S - 0.084 \times ssNa, \quad (3)$$

where S is the total sulfur concentration on the sample, 0.084 is the mean S/Na ratio in seawater (Lide, 2005) and $ssNa$ is the Na actually derived from sea spray. Since some Na derives from continental dust, $nssNa$ was calculated using the equation system (4) reported below:

$$ssNa = Na - nssNa$$

$$nssNa = nssAl \times (Na/Al)_{crust} \quad (4)$$

$$Al = nssAl,$$

where the mean Na/Al ratio is 0.3315 in the crust (Wedepohl, 1995). In this study, we assume that Al is only from crustal origin (total Al is equal to $nssAl$). Wagenbach and others (1998) suggested that the seawater ratio at Antarctic coastal sites is variable because of fractionation processes (formation of frost flowers on fresh sea ice and precipitation of mirabilite at temperatures below $-8.2^{\circ}C$). It can be recognized by obtaining negative $nssS$ concentrations. The Mount Johns record shows a deficit of $nssS$ during the winter in a few years. However, it corresponds to less than 1% of the samples and we consider that the sulfur fractionation does not affect the ratio in the MJ area.

The sulfur contribution from marine phytoplankton activity (via atmospheric oxidation of dimethylsulfide produced by marine phytoplankton during summer months) was estimated from the methanesulfonate (MSA is a specific marker of marine biogenic emissions) concentration (following Udisti et al., 2016). About 30% of the $nssS$ concentration in the MJ area was considered from biogenic origin.

Approximately 10 to 15 % of the non-sea-salt sulfate/sulfur concentration in the Antarctic atmosphere originates from volcanic activity (Boutron and Paterson, 1986; Hur et al., 2007). We used the Hinkley et al. (1999) element/ S ratios to calculate inputs from the global mean volcanic quiescent degassing background for the element Mn and S (there is not data available for the other elements). Furthermore, we used the metal/ S ratios from the Mount Erebus ($77^{\circ}32'S$, $167^{\circ}10'E$) plume (Zreda-Gostynska et al., 1997) to represent local source contributions for Al , Ca , Fe , K , Mn , Na , S , and Ti . The first step to calculating the volcanic contribution is to remove the oceanic and crustal fraction of elements. We are left with the excess [excess = total - (oceanic + crustal)] elemental concentrations to calculate the local (3 % - 5 %) and global (10 % - 15 %) volcanic contributions (Table S3 - Supporting Information).

Only S and Mn show significant input of volcanic emissions with contributions ranging from 20 to 33 % from global volcanic sources and 3 to 5 % from regional emissions for mean excess elemental concentrations. The other elements (Al , Ca , Fe , K , Na , and Ti) presented less than 1 % of volcanic input. Due to lack of data in the literature, we did not calculate volcanic increments for the elements Ba , Mg , and Sr .

3.3 Principal component analysis

To evaluate the origin and transport of the trace elements determined in the MJ ice core, we applied the principal component analysis (PCA) in order to reveal linear relations existing in the chemical data set. For our study, OriginPro 2017 for Windows (OriginLab Corporation, USA), was utilized for the multivariate statistical analysis, and for descriptive and correlation analyses.

PCA is widely used to reduce data (Loska and Wiechuya, 2003; Gabrielli et al., 2008) and to extract a small number of latent factors (principal components, PCs) for analyzing relationships among the observed variables (Table S4 - Supporting Information). The PCA applied to the entire chemical data set resulted in

four PCs (Table 2) reporting 42.24 %, 13.27 %, 11.16 %, and 8.11% of the total variance of 11 trace elements.

All trace elements correlate positively on PC1 with Ba, K, Mg, Mn, Na, and Sr obtained the highest scores;

PC2 is positively dominated by Al, Fe, Mg, and Ti, and with Ca, K, Mn, Na, and Sr negatively correlated.

PC3 is positively dominated by Ba, Ca, Fe, and Ti while Al and Mg are negatively correlated. Finally, PC4

is positively dominated by Ca, Mg, and Sr, while S and Mn are negatively correlated.

PC1 appears to be an impurity index with each element correlated positively within the PC. When the

atmosphere load is high (high impurity content) PC1 shows high values. Conversely, when impurity

concentration reaches minimum values, PC1 is low (Figure 3). PC2 and PC3 suggest to be derived from

soil dust, as the elements that are positively correlated are the main components of soil and crustal.

However, the negatively correlated elements in PC2 point to marine aerosol deposition. Also, we can say

that PC2 highlights aluminosilicate (Al is the most important element), while PC3 highlights the siliciclastic

fraction of mineral dust (Fe and Ti). Last, PC4 is related to carbonate content, where Ca, Mg, and Sr are

major constituents of CaCO_3 .

Based on the analysis of crustal and marine enrichment factors and PCA we have classified the concentrations as predominantly crustal for the elements Al, Mg, and Ti while Na, Sr, and Ca are primarily sea salt derived elements. The elements Ba, Mn, Fe, and K show to be from mixed sources (mineral dust and sea salt aerosol). Furthermore, the S record has a considerable volcanic and biogenic input and Mn has an additional volcanic input. We acknowledge that Fe may have an additional contribution of biomass burning aerosol (Winton et al., 2016), but due to lack of data we will not address this issue here. Based on the low values of E_{Fc} and E_{Fo}, we consider that the presented concentrations are from natural origin and possible anthropogenic contributions to these elements would be insignificant in this area.

3.4 Intraseasonal concentration fluxes variability

Generally, trace element concentrations from sea salt aerosol observed in coastal and interior West Antarctic ice cores show a clear seasonal signal, with higher concentrations in austral winter and spring months (June to November) and lower concentrations in austral summer months (December-February) (Legrand and Mayewski, 1997; Sigl et al., 2016). Impurities from continental dust can peak in both the austral summer (Weller et al., 2008; Tuohy et al., 2015) or winter months (Hur et al., 2007), depending on site location. Dust deposition is mainly influenced by local factors and atmospheric circulation patterns.

Additionally, biogenic aerosols (e.g. sulfur) show peaks in summer months due to an increased phytoplankton activity (Weller et al., 2011). We found high concentrations in austral winter and low concentrations in austral summer for most of the elements analyzed, with the exception of sulfur, which presents peaks in the summer due to the biogenic contribution and of the elements with crustal influence (Al, Ba, Mg, and Ti) that present the highest concentrations in spring/summer.

To investigate if observed changes in chemistry are related to changes in accumulation rate, we calculate the annual deposition flux for all elements. The flux is calculated by multiplying the concentrations in the ice by the annual water-equivalent accumulation rate. The annual flux for all trace elements is shown in Fig. 4. Both concentration and flux presents similar variability (not shown), this indicates that the

concentrations are independent of accumulation rate and that the variability exhibited in the record is likely not a function of changes in snow accumulation.

The fluxes are highly variable down the length of the core. In particular, there are three distinct phases in the record: (i) between 1885 and around 1930 fluxes peak for Al, Ba, Fe, Mg, S, and Ti; (ii) between approximately 1930 and 1955 minimum fluxes can be observed for the above trace elements; and (iii) between approximately 1955 and 2008 a second increasing trend is observed for Al, Ba, Fe, Mg, S, and Ti.

The ice core record reflects changes in emissions as well as atmospheric circulation and transport processes. The El Niño-Southern Oscillation (ENSO) is the variation in southern Pacific sea surface temperature that induces regional-scale changes in atmospheric circulation in those latitudes. Several researchers have reported links between ENSO and aerosol deposition flux in Antarctica (Vance et al., 2013; Criscitiello et al., 2014). Below we will discuss in detail the influence of climate variables in the trace element fluxes presented.

3.5 Interannual atmospheric variability

Sea ice concentration trends are generally consistent with sea surface temperature (SST) trends, such that regions of increasing (decreasing) sea ice are nearly always found in an environment of decreasing (increasing) SST. However, winds can influence sea ice concentration in several ways, including atmospheric thermal advection, oceanic currents, and wind-driven dynamic transport (Holland and Kwok, 2012). Schneider et al. (2011) suggest that atmospheric circulation trends have influenced both the sea ice concentration and the temperature trends in the Pacific sector of the Antarctic.

Correlations were made between annual means of reanalysis variables (SIC, SST, and 2-m air temperature) from the ERA-Interim product and the studied trace element annual average concentrations from 1979–2008. Observed correlations suggest a relationship between sea ice concentration, sea ice temperature, and aerosol transport to the ice core site. Figure 5 shows that correlations between SIC and most of the analyzed elements are positive ($r > 0.55$; $p < 0.05$) in the Ross Sea and negative ($r > -0.45$; $p < 0.05$) in the Bellingshausen Sea, consistent with others authors (Simpkins et al., 2012; Turner et al., 2015a). Previous studies using satellite observations show a dipole structure in SIC with increasing sea ice in the Ross Sea (Stammerjohn et al., 2015) and decreasing in the Amundsen–Bellingshausen Seas (Holland and Kwok, 2012). This dipolar pattern is related to thermodynamic and dynamic forcing associated with variability in the pressure anomalies extending over the Amundsen Sea (Turner et al., 2015b).

The strongest positive correlations ($r = 0.65$; $p < 0.05$) are in the region between 180° and 140° W, suggesting that concentrations at the MJ site increase when SIC is high in the West Amundsen and Ross Seas. This may indicate the dominant source area of the marine aerosols to the MJ area. Some studies show that processes associated with sea ice formation (e.g. frost flower formation, brine production, and blowing snow released from sea ice surfaces) are the dominant source of sea salt aerosols for Antarctica (Rankin et al., 2002; Kaspari et al., 2005; Criscitiello et al., 2013). Aluminum and magnesium exhibit negative correlations ($r = -0.55$; $p < 0.05$) in the Ross Sea region and positive correlations ($r = 0.50$; $p < 0.05$) in the Antarctica Peninsula area. Aluminum in polar ice derives almost exclusively from crustal dust (McConnell et al., 2007), which explains the correlation difference between Al and other trace elements. However, Mg exhibits the same pattern observed for Al, showing a relationship between the two elements. As previously

presented in other studies, the nss fraction of Mg can be linked to insoluble silicate or soluble carbonate or evaporite (e.g., gypsum, halite) minerals derived from terrestrial sites (McConnell et al., 2007; Kreutz and Koffman, 2013), which would explain the strong correlation with Al found. The nssMg fraction measured in this work corresponds to approximately 72% of the total concentration. This negative correlation is also related to a different seasonality pattern, with mineral dust and marine aerosol peaks occurring in different periods of the year.

Figure 6 shows associations between SST and annual trace element concentrations. Negative correlations ($r > -0.70$; $p < 0.05$) between SST in the Ross, Amundsen, and Bellingshausen Seas with the trace element concentrations indicate increased transport of marine aerosols when the SST is cooler. Strong negative correlation ($r > -0.75$; $p < 0.05$) is also observed in the Weddell Sea and Indian Ocean. On the contrary, Al and Mg concentrations are positively correlated ($r > 0.65$; $p < 0.05$) with the Ross, Bellingshausen, and Amundsen Seas as well as the southern Indian Ocean SST indicating that Al and Mg concentrations increase when SST is high. Once more, Al and Mg exhibit a very strong relationship with each other and an inverse correlation to that presented by the other elements. In the case of these two elements, transport and deposition processes are associated with mineral dust seasonality pattern and influenced by warmer SST (usually associated with the spring/summer period) between the source and the MJ site.

The 1979–2008 correlations for T_{2m} are displayed in Fig. 7. Annual T_{2m} anomaly maps generally reveal weak, negative anomalies ($r > -0.45$; $p < 0.05$) over the Ross and West Amundsen Seas for the measured elements, while showing low, positive anomalies ($r > 0.45$; $p < 0.05$) for Al and Mg specifically.

Ligtenberg et al., (2013) show that snowfall and T_{2m} in the Antarctic continent appear to be linked: periods with increasing T_{2m} coincide with periods of increasing snowfall. Changes in T_{2m} show warmer and moister conditions extending considerably farther inland, consistent with enhanced air intrusions (Nicolas and Bromwich 2011). Changes in snow accumulation are also linked to the deepening of the Amundsen Sea Low (ASL), tropical SST, and large-scale atmospheric circulation (Thomas et al., 2015).

3.6 Atmospheric transport to Mount Johns ice core site

Atmospheric transport to Antarctica is dominated by the circumpolar westerly winds over the Southern Ocean and the permanent cyclone belt over the polar fronts (Hoskins and Hodges, 2005). The baroclinic zone, between 60–70°S, is a very active cyclone generating area due to the interaction of cold, dry air from the continent and relatively warmer, moist air from the Southern Ocean. The prevailing mid-latitude westerlies direct the cyclones, circulating around the Antarctic continent (King and Turner, 1997). The Antarctic plateau is dominated by a high-pressure (anticyclonic, counter-clockwise) and the wind regime is governed by katabatic winds (cold, dense air flowing downhill due to gravity).

The Amundsen Sea Low (ASL) is a major driver of West Antarctic climate variability (Turner et al., 2013). It is a mobile, climatological low-pressure located between 170–298° E and 80–60° S, in the Southern Pacific (Kreutz et al., 2000; Hosking et al., 2013). The depth and location of the low-pressure center affects the climatic conditions and the strength of the westerlies along the coastal regions and the interior of West Antarctica (Ding et al., 2011; Schneider et al., 2012).

Back-trajectory modeling has been used to understand the synoptic controls on precipitation and the transport of trace elements to the study site. Here, we calculated 5-day, backward trajectories using the

NOAA HySPLIT Model and clustered the trajectories over austral spring (September–November), summer (December–January), autumn (March–May), and winter (June–August) months. Daily simulations were generated during the 1979 to 2008 period. For the simulated trajectories, we have determined that five clusters are sufficient to capture the seasonal trajectory variability.

The MJ site receives the majority of air masses from the Amundsen Sea and, secondarily, from across the Antarctic Peninsula and Weddell Sea (Fig. 8). An additional source could be a continental local contribution. It is possible to identify two clusters with dominant westerly flow patterns ranging from fast (long) to slow-moving (short) depending on the season; one cluster includes fast-moving trajectories with strong cyclonic curvature around the Ross Ice Shelf (whole year); a group with direction that varies from westerly flow to northeasterly depending on the season, and a continental grouping that contains mainly katabatic flow paths from the interior. The clusters show that air masses circulate around the Antarctic continent until they are diverted to the interior, as synoptic storms, across the Amundsen and Bellingshausen Seas. In some cases, the air masses arrive on the WAIS after being diverted across the Weddell Sea and travelling over the Filchner-Ronne Ice Shelf.

The trajectories are classified into two groups: 1) oceanic influenced (blue), and 2) continental influenced (red). These classifications are defined by characteristics identified from the mean trajectories of each cluster, speed (proportional to trajectory length), source region, and pathway. The frequency distribution of cluster classes for the 1979–2008 period shows distinct seasonality between the austral summer and other seasons, with comparatively stronger, westerly transport in the cold months and a secondary, northeasterly transport in the warm months. On average, the oceanic group has a maximum, seasonal frequency in winter while the continental group peaks in winter/spring. During the summer, the trajectories generally are slow-moving (short) and are more locally influenced than in others seasons.

3.7 Relationships between atmospheric circulation, temperature, and sea ice concentration

The highest concentrations of trace elements in the winter are associated with air masses clustered within the oceanic trajectories. Low pressure centers over the Bellingshausen-Amundsen Sea are associated with cold temperatures and sea ice formation in the region (Bromwich et al., 2013; Criscitiello et al., 2014). Seasonal and interannual variability of the trace element concentrations are also likely due to wind speed and transport efficiency at the time of deposition. Hoskins and Hodges (2005) suggest that storminess over the Southern Ocean and the strength of inland transport, both of which are enhanced during winter, can explain the winter maxima deposition in the WAIS. The wind associated with the low pressure systems over the Bellingshausen-Amundsen Sea would facilitate the generation and transport of sea-salt aerosols from either an open ocean or a sea ice source (Bromwich et al., 2013; Criscitiello et al., 2014; Pasteris et al., 2014). The westward extent of trajectories from winter (maximum sea ice area) to spring (maximum sea ice retreat) is between 180 and 140°W, coincident with the area of greatest correlation between trace elements concentrations and SIC, SST, and T_{2m} . The mechanism for deposition of trace elements at MJ site is intrinsically linked with enhanced sea ice production in the Amundsen-Ross Sea. Thus, not only is there a clear mechanism for trace elements transport to the MJ site, but the factors governing this transport (for example, local wind conditions and temperature) are also related to the sea ice concentration itself.

These air masses also have the potential to capture impurities transported over the South Pacific from mid-latitude continental regions such as South America or Australia. Moreover, winds remobilize mineral dust from ice-free areas in the WAIS, for example trajectories crossing Marie Byrd Land, Ellsworth Land, the Antarctic Peninsula, and the Weddell Sea. Li et al. (2008) show that due to the prevailing westerlies, the distribution and deposition of dust has an eastward transport so that half of the Atlantic Ocean and Indian Ocean are influenced by South American dust and the Pacific region is influenced by Australian dust. Modeling studies of dust transport to Antarctica (Krinner et al., 2010) show that the annual mean concentration of dust in West Antarctica (particularly Marie Byrd Land), is mostly represented by dust originating from Australia. This is in agreement with Neff and Bertler (2015) and Tuohy et al. (2015) who show dust transportation from Australia and New Zealand to the South Pacific. As observed previously in other ice core sites from the WAIS (Dixon et al., 2011; Koffman et al., 2014a; Neff and Bertler, 2015), mineral dust reaching the MJ ice core site is unlikely to be associated with a single dust source due to mixing along-transport from presumed continental sources. Previous work at the MJ core site, Cataldo et al., (2013) found particle sizes ranging from 1.1 μm to 2.4 μm . Based on these data we assume that the concentrations are more influenced by remote continental sources than local sources. Although we do not disregard that local contributions can be a secondary source for dust.

Previous studies have also identified a teleconnection between the tropical Pacific Ocean and aerosol deposition in Antarctica (Vance et al., 2013; Criscitiello et al., 2014). SST anomalies under areas of strong tropical convection in the central tropical Pacific have a significant influence on the atmospheric circulation in the Bellingshausen-Amundsen Sea area through the generation of a large-scale, atmospheric wave train (Lachlan-Cope and Connolley, 2006; Ding et al., 2011). The wave train pattern is prominent in winter and spring months (Lachlan-Cope and Connolley, 2006). Since sea salt aerosols are deposited throughout the year, notably in winter, it is expected that tropical forcing and atmospheric Rossby waves would influence trace element concentrations. However, the elements of crustal origin presented different variability with maximum concentration in period of lower SIC and higher temperature (usually associated with the spring/summer). Mineral dust deposition at other locations in Antarctica also exhibits seasonal variability with a maximum in summer, e.g. Berkner Island (Bory et al., 2010), Windless Bight, McMurdo Sound (Dunbar et al., 2009), and South Pole (Legrand and Kirchner, 1988).

The strong, negative correlation between trace element concentrations and SST further demonstrate that concentrations contained in the MJ ice core record provide an indication of past variability. Based on significant, negative correlations with SST, and supported by annual variability of $\delta^{18}\text{O}/^{16}\text{O}$ (Fig. S2 – Supporting Information), we suggest that high concentrations observed before 1930 in our ice core record are directly related to cooler SST affecting atmospheric transport in the Amundsen-Ross Sea region. We assume that the low concentrations observed later (1930-1955) are consistent with a warming in the region. Schneider et al. (2008) show extreme positive anomalies (representative of West Antarctic surface temperature) during the 1936-1945 period. The authors interpreted these anomalies as indicative of strong teleconnections in part driven by the ENSO (1939–1942). The second increasing trend observed for Al, Ba, Fe, Mg, S, and Ti (between 1955 and 1992), when compared with $\delta^{18}\text{O}$ record, is also associated with a period of cooling in the MJ area or in the region where these aerosols are generated, Jones et al. (2016) show a cooling of SST in the Ross sector during the study time (1979-2014), this may be related to the

increase of concentrations, since our correlations have pointed this region as a source area of marine aerosols. This cooling is followed by a warming period into recent years in MJ record (associated with decreased concentrations of Al, Mg, and S). Therefore, it seems possible that different trace element concentrations respond to different forcing on different timescales.

The Antarctic climate system is strongly interconnected; regional and seasonal complexity as well as physical relationships between different climate variables must be taken into account to identify the mechanisms driving the production, transport, and deposition of trace elements in WAIS. While these results are promising, further work involving assimilation methods using ice core proxy records and climate simulations in order to best reconstruct the past state of the Antarctic atmospheric circulation is required.

4. Conclusions

Using high-resolution ICP-SFMS, several trace elements were measured in an ice core from WAIS and used to evaluate inputs from natural aerosol emission sources. It was found that natural contributions from mineral dust are important sources for Al, Mg, and Ti, while marine aerosols from the open sea and sea ice in the southern Pacific are important sources of Na, Sr, and Ca, over the MJ site. However, Ba, Fe, K, Mn, and S show to be from mixed sources (mineral dust and sea-salt aerosols). Additionally, S and Mn exhibit important volcanic contribution and S has a considerable biogenic input in the summer season. Utilizing back trajectories from HySPLIT, we identify two dominant air-mass trajectory clusters: marine and continental. The analysis shows distinct differences between seasons, relatively stronger, westerly transport during the winter months and a secondary, northeasterly transport during the summer months. During the summer, the trajectories present slow-moving (short) and are locally influenced when compared to other seasons. Based on particle size (1.1 μm to 2.4 μm) for MJ ice core and the air-mass trajectory, we conclude that the trace element concentrations are influenced by remote continental sources more so than local sources, although we do not ignore that local contributions can be a secondary source for mineral dust.

Our reanalysis-element correlations suggest that marine derived trace element concentrations are strongly influenced by SIC and SST anomalies. The results show that seasonal concentration maxima in sea-salt elements correlate well with the SIC winter maxima. The strongest correlations are in the region between 180° and 140° W, suggesting that elemental concentrations in the MJ site increase when SIC is high in the West Amundsen and Ross Seas. This may indicate the dominant source of the marine aerosols over the MJ site. Correlations between SST in the Ross, Amundsen and Bellingshausen Seas with the MJ site indicate increased transport of marine aerosols when the SST is relatively cooler. Previous studies indicate that wind associated with low pressure systems over the Bellingshausen-Amundsen Sea facilitate the generation and transport of sea-salt aerosols from either an open ocean or a sea ice source (Bromwich *et al.*, 2013; Criscitiello *et al.*, 2014; Pasteris *et al.*, 2014), as well as affect the mineral dust transport from circum-Antarctic continents to the Southern Ocean and WAIS (Neff and Bertler, 2015).

We show that both sources and transport of mineral dust and marine aerosols to West Antarctica are controlled by the climate variables in response to remote atmospheric forcing. While these results are promising, further work is needed to obtain a more detailed picture of past variability and its relationship with regional aerosol transport.

Acknowledgments

This research is part of the Brazilian Antarctic Program (PROANTAR) and was financed with funds from the National Council for Scientific and Technological Development (CNPq), project 407888/2013-6.

References

- Albani, S., Mahowald, N. M., Delmonte, B., Maggi, V. and Winckler, G.: Comparing modeled and observed changes in mineral dust transport and deposition to Antarctica between the Last Glacial Maximum and current climates, *Climate Dynamics*, 38(9-10), 1731-1755, 2012.
- Bamber, J. L., Riva, R. E. M., Vermeersen, B. L. A. and LeBrocq, A. M.: Reassessment of the potential sea-level rise from a collapse of the West Antarctic Ice Sheet, *Science*, 324, 901-903, doi: 10.1126/science.1169335, 2009.
- Becagli, S., Proposito, M., Benassai, S., Gragnani, R., Magand, O., Traversi, R. and Udisti, R.: Spatial distribution of biogenic sulphur compounds (MSA, nssSO_4^{2-}) in the northern Victoria Land-Dome C-Wilkes Land area, East Antarctica, *Annals of Glaciology*, 41, 23-31, 2005.
- Bory, A., Wolff, E., Mulvaney, R., Jagoutz, E., Wegner, A., Ruth, U. and Elderfield, H.: Multiple sources supply eolian mineral dust to the Atlantic sector of coastal Antarctica: Evidence from recent snow layers at the top of Berkner Island ice sheet, *Earth and Planetary Science Letters*, 291, 138-148, 2010.
- Boutron, C.F. and Patterson, C.C.: Lead concentration changes in Antarctic ice during the Wisconsin/Holocene transition, *Nature*, 323, 222–225, 1986.
- Bracegirdle, T. J. and Marshall, G. J.: The reliability of Antarctic tropospheric pressure and temperature in the latest global reanalyses, *Journal of Climate*, 25, 7138–7146, 2012.
- Bromwich, D. H. and Fogt, R. L.: Strong trends in the skill of the ERA-40 and NCEP/NCAR reanalyses in the high and middle latitudes of the Southern Hemisphere, 1958-2001, *Journal of Climate*, 17, 4603–4619, 2004.
- Bromwich, D. H., Nicolas, J. P., Monaghan, A. J., Lazzara, M. A., Keller, L. M., Weidner, G. A. and Wilson, A. B.: Central West Antarctica among the most rapidly warming regions on Earth, *Nature Geoscience*, 6, 139–145, 2013.
- Cataldo, M., Evangelista, H., Simões, J. C., Godoi, R. H. M., Simmonds, I., Hollanda, M. H., Wainer, I., Aquino, F. and Van Grieken, R.: Mineral dust variability in central West Antarctica associated with ozone depletion, *Atmospheric Chemistry and Physics*, 13, 2165-2175, 2013.
- Chewings, J. M., Atkins, C. B., Dunbar, G. B. and Golledge, N. R.: Aeolian sediment transport and deposition in a modern high-latitude glacial marine environment, *Sedimentology*, doi: 10.1111/sed.12108, 2014.
- Crisciatiello, A. S., Das, S. B., Evans, M. J., Frey, K. E., Conway, H., Joughin, I., Medley, B. and Steig, E.: Ice sheet record of recent sea-ice behavior and polynya variability in the Amundsen Sea, West Antarctica, *Journal of Geophysical Research: Oceans*, 118, 1-13, 2013.
- Crisciatiello, A. S., Das, S. B., Karnauskas, K. B., Evans, M. J., Frey, K. E., Joughin, I., Steig, E. J., McConnell, J. R. and Medley, B.: Tropical Pacific influence on the source and transport of marine aerosols to West Antarctica, *Journal of Climate*, 27, 1343-1363, 2014.
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., Van de Berg, L., Bidlot, J., Bormann, N.,

- Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, I., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., Rosnay, P. de, Tavolato, C., Thépaut, J.-N. and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Quarterly Journal of the Royal Meteorological Society*, 137, 553-597, 2011.
- Delmonte, B., Andersson P. S., Schöberg, H., Hansson, M., Petit, J.-R., Delmas, R., Gaiero, D. M., Maggi, V. and Frezzotti, M.: Geographic provenance of aeolian dust in East Antarctica during Pleistocene glaciations: preliminary results from Talos Dome and comparison with East Antarctic and new Andean ice core data, *Quaternary Science Reviews*, 29, 256-264, 2010.
- Delmonte, B., Baroni, C., Andersson, P. S., Narcisi, B., Salvatore, M. C., Petit, J.-R., Scarchilli, C., Frezzotti, M., Albani, S. and Maggi, V.: Modern and Holocene Aeolian dust variability from Talos Dome (northern Victoria Land) to the interior of the Antarctic ice sheet, *Quaternary Science Reviews*, 64, 76-89, 2013.
- Ding, Q., Steig, E. J., Battisti, D. S. and Küttel, M.: Winter warming in West Antarctica caused by central tropical Pacific warming, *Nature Geoscience*, 4, 398–403, doi: 10.1038/ngeo1129, 2011.
- Dixon, D. A., Mayewski, P. A., Goodwin, I. D., Marshall, G. J., Freeman, R., Maasch, K. A. and Sneed, S. B.: An ice-proxy for northerly air mass incursions into West Antarctica, *International Journal of Climatology*, 32, 1455–1465, doi: 10.1002/joc.2371, 2011.
- Dixon, D. A., Mayewski, P. A., Korotkikh, E., Sneed, S. B., Handley, M. J., Introne, D. S. and Scambos, T. A.: Variations in snow and firn chemistry along US ITASE traverses and the effect of surface glazing, *The Cryosphere*, 7, 515–535, 2013.
- Draxler, R. R.: HYSPLIT4 user's guide, technical memorandum, NOAA, Silver Spring, Md, 1999.
- Draxler, R. R., Ginoux, P., Stein, A. F.: An empirically derived emission algorithm for wind-blown dust, *Journal of Geophysical Research - Atmospheres*, 115 (D16212), doi: 10.1029/2009JD013167, 2010.
- Duce, R. A., Hoffman, G. L. and Zoller, W. H.: Atmospheric trace-metals at remote northern and southern hemisphere sites: pollution or natural? *Science*, 187(4171), 59–61, 1975.
- Dunbar, G. B., Bertler, N. A. N. and McKay, R. M.: Sediment flux through the McMurdo Ice Shelf in Windless Bight, Antarctica, *Global and Planetary Change*, 69, 87-93, 2009.
- Favier, L., Durand, G., Cornford, S. L., Gudmundsson, G. H., Gagliardini, O., Gillet-Chaulet, F., Zwinger, T., Payne, A. J. and Le Brocq, A. M.: Retreat of Pine Island Glacier controlled by marine ice-sheet instability, *Nature Climate Change*, 4, 117-121, doi: 10.1038/NCLIMATE2094, 2014.
- Fischer, H., Traufetter, F., Oerter, H., Weller, R. and Miller, H.: Prevalence of the Antarctic Circumpolar Wave over the last two millennia recorded in Dronning Maud Land ice, *Geophysical Research Letters*, 31, doi: 10.1029/2003GL019186, 2004.
- Fretwell, P., Pritchard, H. D., Vaughan, D. G., Bamber, J. L., Barrand, N. E., Bell, R., Bianchi, C., Bingham, R. G., Blankenship, D. D., Casassa, G., Catania, G., Callens, D., Conway, H., Cook, A. J., Corr, H. F. J., Damaske, D., Damm, V., Ferraccioli, F., Forsberg, R., Fujita, S., Gim, Y., Gogineni, P., Griggs, J. A., Hindmarsh, R. C. A., Holmlund, P., Holt, J. W., Jacobel, R. W., Jenkins, A., Jokat, W., Jordan, T., King, E. C., Kohler, J., Krabill, W., Riger-Kusk, M., Langley, K. A., Leitchenkov, G., Leuschen, C., Luyendyk, B. P., Matsuoka, K., Mouginot, J., Nitsche, F. O., Nogi, Y., Nost, O. A., Popov, S. V., Rignot, E., Rippin,

D. M., Rivera, A., Roberts, J., Ross, N., Siegert, M. J., Smith, A. M., Steinhage, D., Studinger, M., Sun, B., Tinto, B. K., Welch, B. C., Wilson, D., Young, D. A., Xiangbin, C. and Zirizzotti, A.: Bedmap2: improved ice bed, surface and thickness datasets for Antarctica, *The Cryosphere*, 7, 375-393, 2013.

Gabrielli, P., Cozzi, G., Torcini, S., Cescon, P. and Barbante, C.: Trace elements in winter snow of the Dolomites (Italy): A statistical study of natural and anthropogenic contributions, *Chemosphere*, 72, 1504-1509, 2008.

Gaiero, D. M., Brunet, F., Probst, J-L. and Depetris, P. J.: A uniform isotopic and chemical signature of dust exported from Patagonia: Rock sources and occurrence in Southern environments, *Chemical Geology*, 238, 107-120, 2007.

Ginot, P., Stampfli, F., Stampfli, D., Schwikowski, M. and Gäggeler, H. W.: FELICS, a new ice core drilling system for high-altitude glaciers, *Memoirs of National Institute of Polar Research, Special Issue*, 56, 38-48, 2002.

Hinkley, T. K., Lamothe, P. J., Wilson, S. A., Finnegan, D. L., Gerlach, T. M.: Metal emissions from Kilauea, and a suggested revision of the estimated worldwide metal output by quiescent degassing of volcanoes, *Earth and Planetary Science Letters*, 170, 315-325, 1999.

Holland, P. R. and Kwok, R.: Wind-driven trends in Antarctic sea-ice drift, *Nature Geoscience*, 5, 872-875, doi:10.1038/ngeo1627, 2012.

Hoskins, B. J. and Hodges, K. I.: A new perspective on Southern Hemisphere storm tracks, *Journal of Climate*, 18, 4108-4129, 2005.

Hosking, J. S., Orr, A., Marshall, G. J., Turner, J. and Phillips, T.: The influence of the Amundsen-Bellinghshausen Seas low on the climate of West Antarctica and its representation in coupled climate model simulations, *Journal of Climate*, 26, 6633-6648, 2013.

Hur, S., Cunde, X., Hong, S., Barbante, C., Gabrielli, P., Lee, K., Boutron, C. F. and Ming, Y.: Seasonal patterns of heavy metal deposition to the snow on Lambert Glacier basin, East Antarctica, *Atmospheric Environment*, 41, 8567-8578, 2007.

Jones, J. M., Gille, S. T., Goosse, H., Abram, N. J., Canziani, P. O., Charman, D. J., Clem, K. R., Crosta, X., Lavergne, C., Eisenman, I., England, M. H., Fogt, R. L., Frankcombe, L. M., Marshall, G. J., Masson-Delmotte, V., Morrison, A. K., Orsi, A. J., Raphael, M. N., Renwick, J. A., Schneider, D. P., Simpkins, G. R., Steig, E. J., Stenni, B., Swingedouw, D. and Vance, T. R.: Assessing recent trends in high-latitude Southern Hemisphere surface climate, *Nature Climate Change*, 6, 917-936, 2016.

Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha, S., White, G., Woollen, J., Zhu, Y., Chelliah, M., Ebisuzaki, W., Higgins, W., Janowiak, J., Mo, K. C., Ropelewski, C., Wang, J., Leetmaa, A., Reynolds, R., Jenne, R. and Joseph, D.: The NCEP/NCAR 40-year reanalysis project, *Bulletin of the American Meteorological Society*, 77(3), 437-471, 1996.

Kaspari, S., Dixon, D. A., Sneed, S. B. and Handley, M. J.: Sources and transport pathways of marine aerosol species into West Antarctica, *Annals of Glaciology*, 41(1), 1-9, 2005.

King, J. C. and Turner, J.: *Antarctic Meteorology and Climatology*, University Press Cambridge, Cambridge, 409 pp, 1997.

Kistler, R., Kalnay, E., Collins, W., Saha, S., White, G., Woollen, J., Chelliah, M., Ebisuzaki, W., Kanamitsu, M., Kousky, V., van den Dool, H., Jenne, R. and Fiorino, M.: The NCEP-NCAR 50-year

reanalysis: Monthly means CD-ROM and documentation, *Bulletin of American Meteorological Society*, 82, 247-267, 2001.

Koffman, B. G., Kreutz, K. J., Breton, D. J., Kane, E. J., Winski, D. A., Birkel, S. D., Kurbatov, A. V. and Handley, M. J.: Centennial-scale variability of the Southern Hemisphere westerly wind belt in the eastern Pacific over the past two millennia, *Climate of the Past*, 10, 1125-1144, doi: 10.5194/cp-10-1125-2014, 2014a.

Koffman, B. G., Handley, M. J., Osterberg, E. C., Wells, M. L. and Kreutz, K. J.: Dependence of ice-core relative trace-element concentration on acidification, *Journal of Glaciology*, 60(219), 103-112, 2014b.

Kreutz, K. J., Mayeswki, P. A., Pittalwala, I. I., Meeker, L. D., Twickler, M. S. and Whitlow, S. I.: Sea level pressure variability in the Amundsen Sea region inferred from a West Antarctic glaciochemical record, *Journal of Geophysical Research*, 105(D3), 4047-4059, 2000.

Kreutz, K. J. and Koffman, B. G.: Glaciochemistry, *Encyclopedia of Quaternary Science*, 2, 326-333, 2013.

Krinner, G., Petit, J.-R. and Delmonte, B.: Altitude of atmospheric tracer transport towards Antarctica in present and glacial climate, *Quaternary Science Reviews*, 29, 274-284, 2010.

Kurbatov, A. V., Mayewski, P. A. and Abdul Jawad, B. S.: Ice Core Dating Software for interactive dating of Ice Cores, *Eos, Transactions, American Geophysical Union. Fall Meeting Supplement*, 86(52), Abstract PP33C-1603, 2005.

Lachlan-Cope, T. and Connolley, W.: Teleconnections between the tropical Pacific and the Amundsen-Bellinghousen Sea: role of the El Niño/Southern Oscillation, *Journal of Geophysical Research*, 111 (D23101), 2006.

Legrand, M. and Kirchner, S.: Polar atmospheric circulation and chemistry of recent (1957-1983) South Polar precipitation, *Geophysical Research Letters*, 15(8), 879-992, 1988.

Legrand, M. and Mayewski, P.: Glaciochemistry of polar cores: A review, *Reviews of Geophysics*, 35, 219-243, 1997.

Li, F., Ginoux, P. and Ramaswamy, V.: Distribution, transport, and deposition of mineral dust in the Southern Ocean and Antarctica: Contribution of major sources, *Journal of Geophysical Research*, 113, doi: 10.1029/2007JD009190, 2008.

Li, F., Ginoux, P. and Ramaswamy, V.: Transport of Patagonian dust to Antarctica, *Journal of Geophysical Research*, 115, doi:10.1029/2009JD012356, 2010.

Lide, D. R.: Abundance of Elements in the Earth's crust and in the sea, In: Lide, D. R. ed. *CRC Handbook of Chemistry and Physics*, Internet Version 2005, Boca Raton, CRC Press, 14-17, 2005.

Ligtenberg, S. R. M., van de Berg, W. J., van den Broeke, M. R., Rae, J. G. L. and van Meijgaard, E.: Future surface mass balance of the Antarctic ice sheet and its influence on sea level change, simulated by a regional atmospheric climate model, *Climate Dynamics*, 41, 867-884, doi: 10.1007/s00382-013-1749-1, 2013.

Loska, K. and Wiechula, D.: Application of principal component analysis for the estimation of source of heavy metal contamination in surface sediments from the Rybnik Reservoir, *Chemosphere*, 51, 723-733, 2003.

Mahowald, N., Albani, S., Kok, J. F., Engelstaeder, S., Scanza, R., Ward, D. S. and Flanner, M. G.: The size distribution of desert dust aerosols and its impact on the Earth system, *Aeolian Research*, 15, 53-71, 2014.

Markle, B. R., Bertler, N. A. N., Sinclair, K. E. and Sneed, S. B.: Synoptic variability in the Ross Sea region, Antarctica, as seen from back-trajectory modeling and ice core analysis, *Journal of Geophysical Research*, 17, doi:10.1029/2011JD016437, 2012.

McConnell, J. R., Aristarain, A. J., Banta, J. R., Edwards, P. R. and Simões, J. C.: 20th-Century doubling in dust archived in an Antarctic Peninsula ice core parallels climate change and desertification in South America, *PNAS*, 104(14), 5743-5748, 2007.

Neff, P. D. and Bertler, N. A. N.: Trajectory modeling of modern dust transport to the Southern Ocean and Antarctica, *Journal of Geophysical Research: Atmospheres*, 120, doi: 10.1002/2015JD023304, 2015.

Nicolas, J. P. and Bromwich, D. H.: Climate of West Antarctica and influence of marine air intrusions, *Journal of Climate*, 24, 49-67, 2011.

Osterberg, E. C., Handley, M. J., Sneed, S. B., Mayewski, P. A. and Kreutz, K. J.: Continuous ice core melter system with discrete sampling for major ion, trace element, and stable isotope analysis, *Environmental Science and Technology*, 40 (10), 3355-3361, 2006.

Osterberg, E. C. North Pacific late Holocene climate variability and atmospheric composition, PhD Thesis, University of Maine, Orono, USA, 161 pp, 2007.

Palmer, A. S., Morgan, V. I., Curran, M. A. J., Van Ommen, T. D. and Mayewski, P. A.: Antarctic volcanic flux ratios from Law Dome ice cores, *Annals of Glaciology*, 35, 329-332, 2002.

Pasteris, D. R., McConnell, J. R., Das, S. B., Criscitiello, A. S., Evans, M. J., Maselli, O. J., Sigl, M. and Layman, L.: Seasonally resolved ice core records from West Antarctica indicate a sea ice source of sea-salt aerosol and a biomass burning source of ammonium, *Journal of Geophysical Research: Atmospheres*, 119 (14), 9168-9182, 2014.

Petit, J. R. and Delmonte, B.: A model for large glacial-interglacial climate-induced changes in dust and sea salt concentrations in deep ice cores (central Antarctica): palaeoclimatic implications and prospects for refining ice core chronologies, *Tellus*, 61B, 768-790, doi: 10.1111/j.1600-0889.2009.00437.x, 2009.

Planchon, F. A. M., Boutron, C. F., Barbante, C., Cozzi, G., Gaspari, V., Wolff, E. W., Ferrari, C. P. and Cescon, P.: Changes in heavy metals in Antarctic snow from Coats Land since the mid-19th to the late-20th century, *Earth and Planetary Science Letters*, 200(1-2), 207-222, 2002.

Pritchard, H. D., Arthern, R. J., Vaughan, D. G. and Edwards, L. A.: Extensive dynamic thinning on the margins of the Greenland and Antarctic ice sheets, *Nature*, 461, 971-975, doi:10.1038/nature08471, 2009.

Pritchard, H. D., Ligtenberg, S. R. M., Fricker, H. A., Vaughan, D. G., van den Broeke, M. R. and Padman, L.: Antarctic ice-sheet loss driven by basal melting of ice shelves, *Nature*, 484, 502-505, doi: 10.1038/nature10968, 2012.

Prospero, J. M., Ginoux, P., Torres, O., Nicholson, S. E. and Gill, T. E.: Environmental characterization of global sources of atmospheric soil dust identified with the Nimbus 7 total ozone mapping spectrometer (TOMS) absorbing aerosol product, *Reviews of Geophysics*, 40(1), doi: 10.1029/2000RG000095, 2002.

Rankin, A. M., Wolff, E. W. and Martin, S.: Frost flowers: Implications for tropospheric chemistry and ice core interpretation, *Journal of Geophysical Research*, 107(D23), doi: 10.1029/2002JD002492, 2002.

Revel-Rolland, M., Deckker, P. de, Delmonte, B., Hesse, P. P., Magee, J. W., Basile-Doelsch, I., Grousset, F. and Bosch, D.: Eastern Australia: A possible source of dust in East Antarctica interglacial ice, *Earth and Planetary Science Letters*, 249, 1-13, 2006.

Rhodes, R. H., Baker, J. A., Millet, M-A. and Bertler, N. A. N.: Experimental investigation of the effects of mineral dust on the reproducibility and accuracy of ice core trace element analyses, *Chemical Geology*, 286, 207-221, 2011.

Rignot, E., Bamber, J. L., Van den Broeke, M. R., Davis, C., Li, Y., Van de Berg, W. J. and Meijgaard, E. V.: Recent Antarctic ice mass loss from radar interferometry and regional climate modeling, *Nature Geoscience*, 1, 106-110, doi: 10.1038/ngeo102, 2008.

Schneider, D. P. and Steig, E. J.: Ice cores record significant 1940s Antarctic warmth related to tropical climate variability, *PNAS*, 105(34), 12154-12158, 2008.

Schneider, D. P., Deser, C. and Okumura, Y.: An assessment and interpretation of the observed warming of West Antarctica in the austral spring, *Climate Dynamics*, 38, 323-347, 2011.

Schneider, D. P., Okumura, Y. and Deser, C.: Observed Antarctic interannual climate variability and tropical linkages, *Journal of Climate*, 25, 4048-4066, 2012.

Schwanck, F., Simões, J. C., Handley, M., Mayewski, P. A., Bernardo, R. T. and Aquino, F. E.: Anomalous high arsenic concentration in a West Antarctic ice core and its relationship to copper mining in Chile, *Atmospheric Environment*, 125, 257-264, 2016.

Shepherd, A., Ivins, E. R., Geruo, A., Barletta, V. R., Bentley, M. J., Bettadpur, S., Briggs, K. H., Bromwich, D. H., Forsberg, R., Galin, N., Horwath, M., Jacobs, S., Joughin, I., King, M. A., Lenaerts, J. T. M., Li, J., Ligtenberg, S. R. M., Luckman, A., Luthcke, S. B., McMillan, M., Meister, R., Milne, G., Mouginot, J., Muir, A., Nicolas, J. P., Paden, J., Payne, A. J., Pritchard, H., Rignot, E., Rott, H., Sørensen, L. S., Scambos, T. A., Scheuchl, B., Schrama, E. J. O., Smith, B., Sundal, A. V., van Angelen, J. H., van de Berg, W. J., van den Broeke, M. R., Vaughan, D. G., Velicogna, I., Wahr, J., Whitehouse, P. L., Wingham, D. J., Yi, D., Young, D. and Zwally, J.: A reconciled estimate of ice-sheet mass balance, *Science*, 338, 1183-1189, 2012.

Sigl, M., Fudge, T. J., Winstrup, M., Cole-Dai, J., Ferris, D., McConnell, J. R., Taylor, K. C., Welten, K. C., Woodruff, T. E., Adolphi, F., Bisiaux, M., Brook, E. J., Buizert, C., Caffee, M. W., Dunbar, N. W., Edwards, R., Geng, L., Iverson, N., Koffman, B., Layman, L., Maselli, O. J., McGwire, K., Muscheler, R., Nishiizumi, K., Pasteris, D. R., Rhodes, R. H. and Sowers, T. A.: The WAIS Divide deep ice core WD2014 chronology – Part 2: Annual-layer counting (0–31 ka BP), *Climate of the Past*, 12, 769-786, doi:10.5194/cp-12-769-2016, 2016.

Simpkins, G. R., Ciasto, L. M., Thompson, D. W. J. and England, M. H.: Seasonal relationships between large-scale climate variability and Antarctic Sea Ice concentration, *Journal of Climate*, 25, 5451-5469, doi: 10.1175/JCLI-D-11-00367.1, 2012.

Sinclair, K. E., Bertler, N. A. N. and Trompeter W. J.: Synoptic controls on precipitation pathways and snow delivery to high-accumulation ice core sites in the Ross Sea region, Antarctica, *Journal of Geophysical Research*, 115(D22), 2010.

Stammerjohn, S. E., Maksym, T., Masson, R. A., Lowry, K. E., Arrigo, K. R., Yuan, X., Raphael, M., Randall-Goodwin, E., Sherrell, R. M. and Yager, P. L.: Seasonal sea ice changes in the Amundsen Sea,

Antarctica, over the period of 1979-2014, *Elementa Science of the Anthropocene*, doi: 10.12952/journal.elementa.000055, 2015.

Steig, E. J., Schneider, D. P., Rutherford, S. D., Mann, M. E., Comiso, J. C. and Shindell, D. T.: Warming of the Antarctic ice-sheet surface since the 1957 International Geophysical Year, *Nature*, 457, 459-462, doi: 10.1038/nature07669, 2009.

Steig, E. J., Ding, Q., Battisti, D. S. and Jenkins, A.: Tropical forcing of Circumpolar Deep Water Inflow and outlet glacier thinning in the Amundsen Sea Embayment, West Antarctica, *Annals of Glaciology*, 60(53), 19-28, doi: 10.3189/2012AoG60A110, 2012.

Tao, G., Yamada, R., Fujikawa, Y., Kudo, A., Zheng, F., Fisher, D. A., Koerner, R. A.: Determination of trace amounts of heavy metals in Arctic ice core samples using inductively coupled plasma mass spectrometry, *Talanta*, 55, 765-772, 2001.

Thomas, E. R., Hosking, J. S., Tuckwell, R. R., Warren, R. A. and Ludlow, E. C.: Twentieth century increase in snowfall in coastal West Antarctica, *Geophysical Research Letters*, 42, doi: 10.1002/2015GL065750, 2015.

Tuohy, A., Bertler, N., Neff, P., Edwards, R., Emanuelsson, D., Beers, T. and Mayewski, P.: Transport and deposition of heavy metals in the Ross Sea Region, Antarctica, *Journal of Geophysical Research: Atmospheres*, 120 (20), 10996-11011, 2015.

Turner, J., Phillips, T., Hosking, J. S., Marshall, G. J. and Orr, A.: The Amundsen Sea Low, *International Journal of Climatology*, 33, 1818-1829, 2013.

Turner, J., Hosking, J. S., Marshall, G. J., Phillips, T. and Bracegirdle, T. J.: Antarctic sea ice increase consistent with intrinsic variability of the Amundsen Sea Low, *Climate Dynamics*, doi: 10.1007/s00382-015-2708-9, 2015a.

Turner, J., Hosking, J. S., Bracegirdle, T. J., Marshall, G. J. and Phillips, T.: Recent changes in Antarctic Sea Ice, *Philosophical Transactions Royal Society A*, 373, 2015b.

Udisti, R., Bazzano, A., Becagli, S., Bolzacchini, E., Caiazzo, L., Cappelletti, D., Ferrero, L., Frosini, D., Giardi, F., Grotti, M., Lupi, A., Malandrino, M., Mazzola, M., Moroni, B., Severi, M., Traversi, R., Viola, A. and Vitale, V.: Sulfate coarse apportionment in the Ny-Ålesund (Svalbard Islands) Arctic aerosol, *Fis. Acc. Lincei*, 27(Suppl 1): 85. doi:10.1007/s12210-016-0517-7, 2016.

Vallelonga, P., Barbante, C., Cozzi, G., Gaspari, V., Candelone, J-P., Van de Velde, K., Morgan, V. I., Rosman, K. J. R., Boutron, C. F. and Cescon, P.: Elemental indicators of natural and anthropogenic aerosol inputs to Law Dome, Antarctica, *Annals of Glaciology*, 39, 169-174, 2004.

Vance, T. R., Van Ommen, T.D., Curran, M. A. J., Plummer, C. T. and Moy, A. D.: A millennial proxy record of ENSO and Eastern Australian rainfall from the Law Dome ice core, East Antarctica, *Journal of Climate*, 26, 710-725, 2013.

Wagenbach, D., Ducroz, F., Mulvaney, R., Keck, L., Minikin, A., Legrand, M., Hall, J. S. and Wolff, E. W.: Sea-salt aerosol in coastal Antarctic regions, *Journal of Geophysical Research*, 103(D9), 10961-10974, 1998.

Wedepohl, K. H.: The composition of the continental crust, *Geochimica et Cosmochimica Acta*, 59(7), 1217-1232, 1995.

Weller, R., Woltjen, J., Piel, C., Resenberg, R., Wagenbach, D., König-Langlo, G. and Kriews, M.: Seasonal variability of crustal and marine trace elements in the aerosol at Neumayer station, Antarctica, *Tellus*, 60B (5), 742-752, 2008.

Weller, R., Wagenbach, D., Legrand, M., Elsässer, C., Tian-Kunze, X. and König-Langlo, G.: Continuous 25-yr aerosol records at coastal Antarctica – I: inter-annual variability of ionic compounds and links to climate indices, *Tellus*, 63B (5), 901-919, 2011.

Winton, V. H. L., Edwards, R., Delmonte, B., Ellis, A., Andersson, P. S., Bowie, A., Bertler, N. A. N., Neff, P. and Tuohy, A.: Multiple sources of soluble atmospheric iron to Antarctic waters, *Global Biogeochemical Cycles*, 30, 421-437, 2016.

Zreda-Gostynska, G., Kyle, P. R., Finnegan, D. and Prestbo, K. M. Volcanic gas emissions from Mount Erebus and their impact on the Antarctic environment, *Journal of Geophysical Research*, 102, 15039-15055, 1997.

Table 1. Statistical summary of trace elements concentrations determined in MJ ice core.

Elements	Mean	Median	SD	Average Uncertainty	Min	Max	MDL*
Al (ng g ⁻¹)	2.64	2.46	1.65	0.03	0.30	22.77	0.30
Ba (pg g ⁻¹)	9.31	5.49	11.45	0.24	0.38	93.46	0.38
Ca (ng g ⁻¹)	3.01	1.88	3.24	0.07	0.09	21.22	0.09
Fe (ng g ⁻¹)	0.62	0.41	0.68	0.01	0.05	5.58	0.05
K (ng g ⁻¹)	1.78	0.94	3.39	0.07	0.06	39.54	0.06
Mg (ng g ⁻¹)	9.27	9.09	4.71	0.10	0.66	45.18	0.38
Mn (pg g ⁻¹)	28.10	11.99	61.60	1.34	0.94	783.79	0.94
Na (ng g ⁻¹)	21.91	12.64	33.10	0.69	0.21	381.61	0.21
S (ng g ⁻¹)	10.24	9.01	6.30	0.14	0.58	62.53	0.06
Sr (pg g ⁻¹)	23.21	17.74	17.85	0.38	0.74	117.94	0.74
Ti (pg g ⁻¹)	14.15	9.00	18.76	0.37	0.66	209.35	0.66

*MDL = Method Detection Limit.

Table 2. Principal component loadings for the Mount Johns trace elements concentrations (PCA loadings > 0.3 are shown in bold).

Element	Component			
	1	2	3	4
Al	0.24	0.59	-0.26	0.05
Ba	0.36	0.01	0.30	-0.07
Ca	0.22	-0.18	0.24	0.66
Fe	0.20	0.19	0.59	-0.02
K	0.32	-0.32	0.06	-0.17
Mg	0.32	0.28	-0.50	0.17
Mn	0.37	-0.18	0.03	-0.36

Na	0.37	-0.34	-0.19	-0.03
S	0.29	0.06	-0.10	-0.44
Sr	0.34	-0.17	-0.17	0.40
Ti	0.21	0.46	0.32	0.03
Percent of variance	42.24 %	13.27 %	11.16 %	8.11 %
Cumulative percent	42.24 %	55.51 %	66.67 %	74.77 %




Figure 1. Map of Antarctica showing the site of the MJ ice core site (red arrow) and locations of sites discussed in the text (figure adapted from the U.S. Geological Survey, <http://lima.usgs.gov/>).




Figure 2. (a) Mean element enrichment factors in reference to Earth's crust (EFc) and (b) oceanic composition (EFo) at the MJ site for austral summer (December to February) and austral winter (June to August).




Figure 3. Variability of PC1 values in relation to age for magnesium (black) and potassium (gray). The red line marks the limit between positive and negative values.




Figure 4. Annual fluxes of Al, Ba, Ca, Fe, K, Mg, Na, S, Sr, and Ti in the MJ ice core (Mn is not shown). Colored bands define three distinct phases in the record from 1883 to 2008.




Figure 5: Correlation of the 1979–2008 ERA-Interim reanalysis sea ice concentration versus annually-averaged trace element time series in the MJ ice core.




Figure 6: Correlation of the 1979–2008 ERA-Interim reanalysis sea surface temperature (SST) parameter versus the annually-averaged trace elements time series in the MJ ice core.




Figure 7: Correlation of the 1979–2008 ERA-Interim reanalysis 2-m air temperature parameter versus the annually-averaged trace elements time series in the MJ ice core.




Figure 8. HySPLIT seasonal clusters of daily 5 day back trajectories from 1979 to 2008 arriving at the Mount Johns ice core site, West Antarctica. Percentage of daily trajectories included in each cluster is indicated, number of daily trajectories for each season is indicated at the bottom right of each panel. Blue shaded area represent oceanic group clusters, while red shaded area show the continental group clusters. Trajectories calculated using the NOAA Hysplit Model (version 4.9).