

Heterogeneous glacier thinning patterns over the last 40 years in Langtang Himal

S. Ragettli^{1,2}, T. Bolch^{2,3} and F. Pellicciotti⁴

[1]{Institute of Environmental Engineering, ETH Zürich, Switzerland}

[2]{University of Zurich, Department of Geography, Zurich, Switzerland}

[3]{Institute for Cartography, Technische Universität Dresden, Dresden, Germany}

[4]{Northumbria University, Department of Geography, Newcastle upon Tyne, UK}

Correspondence to: S. Ragettli (ragettli@ifu.baug.ethz.ch)

Abstract

This study presents volume and mass changes of seven glaciers (five partially debris-covered, two debris-free) in the upper Langtang catchment in Nepal. We use a digital elevation model (DEM) from 1974 stereo Hexagon satellite data and seven DEMs derived from 2006-2015 stereo or tri-stereo satellite imagery (e.g. SPOT6/7). The availability of multiple independent DEM differences allows identifying a robust signal and narrowing down the uncertainty about recent volume changes. The volume changes calculated over several multi-year periods between 2006 and 2015 consistently indicate that glacier thinning has accelerated with respect to the period 1974-2006. We calculate an ensemble-mean elevation change rate of $-0.45 \pm 0.18 \text{ m a}^{-1}$ for 2006-2015, while for the period 1974-2006 we identify a rate of $-0.24 \pm 0.08 \text{ m a}^{-1}$. However, the behavior of glaciers in the study area is heterogeneous, and the presence or absence of debris does not seem to be a good predictor for mass balance trends. Debris-covered tongues have non-linear thinning profiles, and we show that recent accelerations in thinning correlate with the presence of supraglacial cliffs and lakes. At stagnating glacier areas near the glacier front, on the other hand, thinning rates decreased with time or remained constant. The April 2015 Nepal earthquake triggered large avalanches in the study catchment. Analysis of two post-earthquake DEMs revealed that the avalanche deposit volumes remaining six months after the earthquake are negligible in comparison to 2006-2015 elevation changes. However, the deposits compensate about 40% the mass loss of debris-covered tongues of one average year.

1 **1 Introduction**

2 Atmospheric warming has caused widespread recent glacier thinning and retreat in the
3 Himalayan region (Bolch et al., 2012). The impact of current and future glacier changes on
4 Himalayan hydrology and downstream water supply strongly depends on the rate of such
5 changes. However, planimetric and volumetric glacier changes are difficult to characterize
6 due to limited data availability, and many recent studies have highlighted the spatially
7 heterogeneous distribution of glacier wastage in the Himalayas (Fujita and Nuimura, 2011;
8 Bolch et al., 2012; Kääb et al., 2012). Prominent examples of current-day regional differences
9 in glacier evolution across the Hindu Kush–Karakoram–Himalaya (HKH) are the reported
10 positive glacier mass balances in the Pamir and Karakoram. Glaciers in the rest of the HKH
11 are thinning and receding (Bolch et al., 2012; Kääb et al., 2012; Gardelle et al., 2013). Across
12 regions, differences in recent glacier evolution can often be associated to differences in
13 climatic regimes (Fujita, 2008), particularly to the varying influence of the south Asian
14 monsoon and westerly disturbances (Yao et al., 2012). However, also within the same
15 climatic region the rate of glacier changes can be heterogeneous (Scherler et al., 2011b). A
16 main focus of current research is on the effect of supraglacial debris-cover on glacier response
17 to climate. Thick debris cover is a common feature in the HKH (Scherler et al., 2011b;
18 Racoviteanu et al., 2015) and a homogenous layer of thick debris effectively reduces melt
19 rates of underlying ice (e.g. Östrem, 1959; Mattson et al., 1993). However, the
20 characterization of debris-covered glacier response to climate is complicated by the frequent
21 occurrence of ice cliffs and supraglacial lakes. At exposed cliffs, melt rates are much higher
22 compared to the ice covered by a thick debris mantle (Sakai et al., 1998, 2002; Immerzeel et
23 al., 2014a; Steiner et al., 2015; Buri et al., 2016), and also at supraglacial ponds energy
24 absorption is several times larger than that at the surrounding debris-covered surface (Sakai et
25 al., 2000; Miles et al., 2016a). Recent large-scale geodetic studies based on remote sensing
26 have provided evidence that the present-day surface lowering rates of some debris-covered
27 areas in the HKH might be similar to those of debris-free areas even within the same
28 altitudinal range (Kääb et al., 2012; Nuimura et al., 2012; Gardelle et al., 2013), and surmise
29 this could be due to enhanced melt from exposed ice cliffs and supraglacial lakes. Several
30 detailed modelling studies on the other hand have provided evidence for a melt reducing
31 effect of debris at the glacier scale (e.g. Juen et al., 2014; Ragettli et al., 2015), and have
32 shown how supraglacial debris prolongs the response of the glacier to warming (Banerjee and
33 Shankar, 2013; Rowan et al., 2015). Discrepancies between the different conclusions may be

1 associated to glacier samples that are not comparable or to model uncertainties (particularly
2 regarding the representation of the effect of supraglacial cliffs and lakes on total melt).
3 Models can also provide actual melt rates while geodetic studies only provide glacier thinning
4 rates, which are affected by glacier emergence velocity.

5 Programs to monitor debris-covered glaciers have been initiated in the Karakorum (e.g.
6 Mayer et al., 2006; Mihalcea et al., 2006, 2008) and in the Central Himalaya (e.g. Pratap et
7 al., 2015; Ragettli et al., 2015). Yet, due to the logistical and financial constraints, long-term
8 mass balance measurements are basically inexistent in the HKH. To document changes in
9 debris-covered glacier thinning over time, declassified high-resolution reconnaissance
10 satellite data available from the 1960s and 1970s are an important source of information
11 (Bolch et al., 2008, 2011; Maurer and Rupper, 2015). However, a common problem of
12 previous multi-temporal geodetic studies is the relatively low statistical significance of
13 detected changes in glacier thinning over time. The uncertainties in digital elevation models
14 (DEMs) derived from optical data and mass change measurements by DEM differentiation in
15 the HKH arise from the difficult conditions for photogrammetric elevation analysis (due to
16 extreme topography, surfaces with low contrast like bright snow cover or cast shadows).
17 Radar derived DEMs provide more accurate results in snow covered areas but have even
18 higher problems in steep terrain due to the side looking geometry. Unknown penetration of
19 the radar beam into snow and ice is another shortcoming of DEMs derived from radar data.
20 Uncertainties in volume loss estimates are therefore usually higher than identified acceleration
21 in glacier thinning (Nuimura et al., 2012). The uncertainties are especially high over short
22 periods. For long periods with much larger absolute elevation changes, the effect of errors in
23 the DEMs weighs less and uncertainties in glacier volume changes are lower.

24 This study presents volume and mass changes of seven glaciers (five partially debris-covered,
25 two debris-free) in the upper Langtang catchment in Nepal. The aim of this study is to
26 determine changes in thinning rates with high confidence by considering multiple
27 independent DEM differences for short periods. For this we use seven DEMs derived from
28 2006-2015 stereo or tri-stereo satellite imagery and one DEM obtained from 1974 stereo
29 Hexagon satellite data. We obtain an ensemble of multi-annual elevation changes that
30 provides a range of plausible values for the period between October 2006 and October 2015.
31 We then assess if the elevation changes between different overlapping periods between 2006
32 and 2015 show similar characteristics. If this is the case, the ensemble of results can be used

1 to identify statistically significant changes in volume loss rates with respect to the longer
2 period 1974-2006. Three main research questions are then addressed. First, we assess if
3 overall thinning of glaciers in the region has accelerated. Second, we determine if spatial
4 thinning patterns have changed over time. To explain changes in thinning rates we derive a
5 number of glacier surface properties and glacier surface velocities. Third, we evaluate if there
6 are major differences between the response of debris-covered and debris-free glaciers in the
7 sample. Finally, we also look at the cryospheric impact of the April 2015 Nepal earthquake
8 (7.8 magnitude, epicenter approximately 80 km west of the Langtang Valley). The earthquake
9 devastated large parts of the Langtang catchment by triggering large avalanches (Kargel et al.,
10 2016). Two post-earthquake DEMs from May and October 2015 are used to quantify the
11 impact of the avalanche events on the mass balance of the debris-covered glacier tongues and
12 assess its significance in comparison to multi-annual volume changes.

13

14 **2 Study Site**

15 We analyze the seven largest glaciers in the Langtang Valley (Langtang, Langshisha,
16 Shalbachum, Lirung, Ghanna, Yala, Kimoshung), located in the monsoon-dominated Central
17 Himalaya in Nepal, approximately 50 km north of Kathmandu and 100 km west of the
18 Everest region. While Yala and Kimoshung Glaciers are debris-free glaciers, all other studied
19 glaciers have tongues that are almost entirely covered by supraglacial debris (Figure 1).
20 Langtang Glacier is the largest glacier in the valley with an area of 46.5 km² in 2006 (Table 1)
21 and a total length of approximately 18 km. The smallest glacier is Ghanna Glacier with an
22 area of 1.4 km².

23 Critical debris thicknesses leading to a reduction of melt rates are exceeded over most parts of
24 the debris-covered glacier area (Ragettli et al., 2015). Relatively thin debris appears only at
25 the transition zone between accumulation and ablation area. At Lirung, Shalbachum, Ghanna
26 and Langshisha Glaciers the upper margins of debris-covered sections are located at the foot
27 of steep cirques and icefalls, and transition zones are therefore very short. Ice cliffs and
28 supraglacial ponds increase the heterogeneity of glacier surface characteristics in the
29 Langtang Valley (Pellicciotti et al., 2015).

30 The ablation season of glaciers in the Langtang Valley lasts from April to September. The
31 monsoon season (mid June – September) is at the same time the warmest and the wettest
32 period of the year. Snow cover at the lower elevation of debris-covered glaciers is common

1 only in winter (December – March). However, outside the monsoon period precipitation is
2 limited and winters are rather dry (Collier and Immerzeel, 2015).

3

4 **3 Data and methods**

5 **3.1 Satellite imagery**

6 Multitemporal high-resolution data from different sensors are applied to assess glacier change
7 in the upper Langtang catchment. Each type of remote sensing data employed to calculate
8 glacier elevation changes is listed below. Spatial and radiometric resolutions and base to
9 height (b/h) ratios are provided in Table 2.

- 10 • The oldest data originate from Hexagon KH-9 stereo satellite images from November
11 1974 (Surazakov and Aizen, 2010; Pieczonka et al., 2013; Maurer and Rupper, 2015).
12 These are declassified images from a US reconnaissance satellite program (Burnett,
13 2012).
- 14 • Cartosat-1 is a remote sensing satellite built by the Indian Space Research
15 Organisation (Tiwari et al., 2008). We purchased radiometrically corrected along-track
16 stereo imagery (processed at level ‘ortho-kit’) of the upper Langtang catchment from
17 October 2006 and November 2009. Cartosat-1 data have been previously used for
18 DEM generation e.g. in the Khumbu region in the Nepal Himalaya by Bolch et al.
19 (2011) and Pieczonka et al. (2011).
- 20 • ALOS-PRISM (Advanced Land Observing Satellite - Panchromatic Remote-Sensing
21 Instrument for Stereo Mapping) was an optical sensor mounted on a Japanese satellite
22 system which operated from January 2006 to April 2011 (Bignone and Umakawa,
23 2008; Tadono and Shimada, 2009; Lamsal et al., 2011; Holzer et al., 2015). We
24 purchased a radiometrically calibrated along-track triplet mode scene from December
25 2010.
- 26 • SPOT6/7 (Système pour l’Observation de la Terre) along-track tri-stereo images
27 were acquired upon request in April 2014, May 2015 and October 2015. SPOT6 and 7
28 are the newest satellites of the SPOT series which have been frequently used for
29 geodetic glacier mass balance studies (e.g. Berthier et al., 2007, 2014; Pieczonka et al.,
30 2013). We acquired stereoscopic images in panchromatic mode corrected for

1 radiometric and sensor distortions. Two of the three SPOT6/7 scenes used in this study
2 were acquired in April/May which means that limited amounts of winter snow is still
3 present on the images. However, the imagery has a high spation resolution (1.5 m) and
4 high radiometric depth of 12bit which leads to good correlation results also over
5 snowy parts.

- 6 • Overlapping pairs of high-resolution images acquired by the WorldView-2 and 3
7 satellites in February 2014 provide the basis of 8m DEMs downloaded from
8 <http://www.pgc.umn.edu/elevation> (Noh and Howat, 2015).

9 **3.2 DEMs and elevation changes**

10 **3.2.1 DEM generation**

11 The Hexagon DEM used here was generated for the study by Pellicciotti et al. (2015). We
12 therefore refer to this study for further technical details regarding the Hexagon DEM. The
13 SPOT6/7, Cartosat-1 and ALOS PRISM DEMs were generated for this study using the
14 OrthoEngine module of PCI Geomatica 2015. We used the same parameters for DEM
15 generation as proposed by Berthier et al. (2014) except setting the parameter ‘DEM detail’ to
16 ‘very high’ instead of ‘low’, which provided better results for the rugged debris-covered
17 glacier surfaces. The basis for the georectification were six differential global positioning
18 system (dGPS) points collected on Lirung Glacier on 23 October 2014 (Brun et al., 2016).
19 Because glacier motion and ablation have to be accounted for when using on-glacier dGPS
20 points, we first generated a DEM from an across-track Pléiades stereo image pair from 1 and
21 9 November 2014 using the available dGPS points as ground-control points (GCPs). Glacier
22 melt between 23 October and the acquisition dates of the Pléiades scenes is negligible due to
23 the low temperatures during this period. The horizontal shift due to glacier motion during this
24 period is less than the grid size of the Pléiades image (0.5 m) and is therefore also negligible.
25 Subsequently, we determined 17 off-glacier GCPs on the basis of the Pléiades scene which
26 were then used to derive a DEM from the SPOT6 April 2014 tri-stereo scene. The Pléiades
27 DEM itself is not used in the following to calculate glacier elevation changes since it covers
28 only a small part of the catchment and since only low stereo matching scores were achieved at
29 elevations higher than 4300 m a.s.l. due to snowfall onset between 1 and 9 November 2014.
30 To guarantee high quality GCPs, only pixels with correlation scores higher than 0.7 were
31 considered for GCPs. Since the Pléiades scene covers only about one fourth of the upper

1 Langtang catchment, an additional 60 GCPs were determined on the basis of the April 2014
2 SPOT6 scene for the DEM extraction from the Cartosat-1, ALOS Prism and SPOT7 scenes.
3 In addition to the GCPs, approximately 100 tie points for each scene were used to match
4 stereo pairs before DEM extraction.

5 The WorldView DEMs are 8 m posting DEMs produced using the Surface Extraction with
6 TIN-based Search-space Minimization (SETSM) by Noh and Howat (2015). The WorldView
7 DEMs rely on the satellite positioning model to locate the surface in space. The scenes from
8 February 2015 which provide the basis of the two WorldView DEMs used in this study were
9 acquired only 20 days apart (Table 2) and are adjacent to each other. The Worldview-2 DEM
10 covers the western part of the study catchment and the WorldView-3 DEM the eastern part.
11 Those DEMs were merged for this study and in the following are referred to as one single
12 DEM representative of February 2015.

13 In addition to the DEMs discussed above, the 2000 SRTM (Shuttle Radar Topography
14 Mission) 1 Arc-Second Global DEM (30 m spatial resolution) was used to calculate slopes
15 and accumulation area ratios (AARs) of glaciers (Table 1) and to define 50 m altitude bands.
16 However, the SRTM DEM was not used for DEM differencing because of the uncertainty
17 regarding the penetration depth of the radar signal into snow and ice (Gardelle et al., 2013;
18 Kääb et al., 2015; Pellicciotti et al., 2015). Only DEMs extracted from optical stereo imagery
19 are therefore employed to calculate elevation changes in this study.

20 **3.2.2 Co-registration and DEM differencing**

21 Co-registration of DEM-pairs is applied in order to minimize the errors associated with shifts.
22 Systematic errors in the elevation change maps due to tectonic uplift which could be relevant
23 after the April 2015 Nepal earthquake are also corrected with the co-registration. For this
24 purpose we exclude from each DEM the non-stable terrain such as glaciers and in general all
25 off-glacier area at elevations higher than 5400 m a.s.l. (which is the estimated equilibrium line
26 altitude (ELA) in the Upper Langtang catchment (Ragettli et al., 2015)). The correlation score
27 maps, indicating which pixels have been matched successfully during the DEM extraction
28 process, are used to exclude all DEM grid cells with a correlation score below 0.5. Then,
29 horizontal shifts are determined by minimizing the aspect-dependent bias of elevation
30 differences (Nuth and Kääb, 2011) between each DEM pair. Because of the slope dependency
31 of the method all terrain below a slope of 10° is excluded. The 'older' DEM is then resampled

1 (bilinear interpolation) according to the determined horizontal shift. In a second step the
2 vertical DEM shifts and possible tilts are corrected using second order trend surfaces fitted to
3 all gently inclined ($\leq 15^\circ$) stable terrain (Bolch et al., 2008; Pieczonka et al., 2011; Pieczonka
4 and Bolch, 2015).

5 We resample all DEMs bilinearly to the grid size of the coarsest DEM (30 m) to reduce the
6 effect of different resolutions. Elevation differences are calculated by subtracting the older
7 from the younger DEM (such that glacier thickening values are positive) and are converted to
8 elevation change rates by dividing by the number of ablation seasons between the acquisition
9 dates. Seasonal effects on elevation change rates are neglected when discussing time intervals
10 between DEMs of 4 years or longer, since elevation changes during the winter half-year are
11 usually minor. On average, less than 20% of annual precipitation occur during post-monsoon
12 and winter (Immerzeel et al., 2014b), and less than 3% of annual glacier ice-melt (Ragettli et
13 al., 2015). Area-average glacier elevation change rates are calculated using always the
14 maximum glacier extent between two acquisition dates.

15 **3.2.3 Processing of elevation change maps**

16 Processing of the elevation change ($\Delta h/\Delta t$) maps involves two main steps: i) removal of pixel
17 values identified as outliers and ii) filling of gaps.

18 Outlier removal

19 The stereo matching score maps provided by PCI Geomatica are used to identify elevation
20 data that can be considered for elevation change calculations. If the correlation score of a
21 given DEM pixel is below 0.5, this indicates a poor matching score (Pieczonka et al. 2011)
22 and therefore the corresponding $\Delta h/\Delta t$ values are treated as ‘no data’. Very unrealistic
23 elevation change data (exceeding ± 150 m) are also excluded from the analysis.

24 We use the standard deviation (σ) of observed elevation changes to identify $\Delta h/\Delta t$ outliers.
25 Outliers are defined separately for debris-covered glacier areas and debris-free glacier areas.
26 For the latter we additionally distinguish between glacier area below and above the ELA
27 (estimated at 5400 m a.s.l., see above). σ -levels are thus calculated for each of the three area
28 types in every $\Delta h/\Delta t$ map. Below the ELA (both debris-free and debris-covered area), pixels
29 are defined as outliers if $\Delta h/\Delta t$ values differ from the average by $> 3\sigma$ (e.g. Gardelle et al.,
30 2013). This means that only very few data are classified as outliers, since three standard
31 deviations account for 99.7% of the sample (assuming the distribution is normal). The

1 conservative outlier definitions are justified by the shallow slopes and high contrast, which
2 also explains why stereo matching scores are generally higher below the ELA (Figure 2c).
3 Above the ELA, steep terrain or featureless snow surfaces lead to low DEM accuracy and
4 therefore the outlier criteria should be more restrictive (e.g. Pieczonka et al., 2013; Pieczonka
5 and Bolch, 2015). On debris-free glacier area above the ELA, pixels are therefore defined as
6 outliers if $\Delta h/\Delta t$ values differ from the average by $>1\sigma$ (which applies to approximately 32%
7 of the values if the distribution is normal). A stricter criterion for the accumulation area is also
8 justified by the fact that it can be assumed that elevation changes in the accumulation areas
9 over periods of several years are on average small (Schwitter and Raymond, 1993; Huss et al.,
10 2010). Because we use different σ thresholds above and below the ELA we test the sensitivity
11 of calculated glacier volume changes to a ± 100 m ELA uncertainty. Furthermore, we test the
12 sensitivity to different outlier definitions by comparing our results to the results obtained with
13 a 2σ -level applied to all area types.

14 Gap filling

15 On the glacier areas below the ELA, with only very few data gaps, missing data are replaced
16 using inverse distance weighting (IDW). In the accumulation areas, on the other hand, data
17 gaps can extend over a wide elevation range if the terrain is steep or if the gaps are very large.
18 Because of the elevation dependency of $\Delta h/\Delta t$ values (e.g. Huss et al., 2010) only values from
19 the same altitudinal range should be used to fill data gaps. We thus replace missing data in the
20 accumulation areas by median $\Delta h/\Delta t$ values per 50-m elevation band considering all available
21 data for a given glacier (also from $\Delta h/\Delta t$ maps representative of different periods). For this,
22 we first calculate the mean elevation change rates per 50-m elevation band of each glacier and
23 every $\Delta h/\Delta t$ map and then determine the median of the ensemble. $\Delta h/\Delta t$ maps that are
24 rejected from the ensemble (see Section 3.2.5 below) and in general all values representative
25 of short periods ($\Delta t < 4$ years) are not considered to calculate the ensemble-median values.

26 **3.2.4 Uncertainty**

27 Elevation change uncertainty estimates are based on the standard error $E_{\Delta h}$ calculated per
28 elevation band (Gardelle et al., 2013). The standard error quantifies the effect of random
29 errors on uncertainty according to the standard principles of error propagation:

$$30 \quad E_{\Delta h} = \frac{\sigma_{\Delta h, \text{noglac}}}{\sqrt{N_{\text{eff}}}} \quad (1)$$

$$N_{eff} = \frac{N_{tot} \times PS}{2d} \quad (2)$$

$\sigma_{\Delta h, noglac}$ is the standard deviation of the mean elevation change of non-glacierized terrain per elevation band, N_{eff} is the effective and N_{tot} the total number of observations. PS is the pixel size (30 m) and d is the distance of spatial autocorrelation. d is equal to the range of the spherical semivariogram obtained by least squares fit to the experimental, isotropic variogram of all off-glacier elevation differences (Wang and Kääb, 2015; Magnússon et al., 2016). The distance of spatial autocorrelation of the elevation change maps varies between 260 m and 730 m with an average of 495 m.

To quantify the elevation change uncertainty of glacier area spanning several elevation bands, weighted averages of $E_{\Delta h}$ are calculated. $E_{\Delta h}$ of each individual elevation band is weighted by the glacier hypsometry. Elevation change uncertainties therefore vary for each individual glacier because of the different glacier area-elevation distributions. $E_{\Delta h}$ tends to increase with altitude (Figure 3, Figure 4) due to steeper slopes, snow and deep shadows, which are factors that decrease the accuracy of DEMs derived from stereo data (e.g. Nuimura et al., 2011). Uncertainty estimates for each individual glacier therefore account for the spatially non-uniform distribution of uncertainty. Elevation change uncertainties of glaciers with a high accumulation area such as Kimoshung and Lirung Glaciers (Table 1) are 50%-100% higher than those of other glaciers, in accordance with lower DEM matching scores (Figure 2). The low uncertainty associated to debris-covered areas agrees with the 30%-100% lower off-glacier errors on shallow slopes ($s < 18^\circ$, 95th percentile of debris-covered glacier slopes) than on steeper slopes ($s < 45^\circ$, 95th percentile of glacier slopes; Figure S1).

The standard error can be interpreted as the 68% confidence interval of the sample mean if the distribution is normal. Since we are conservatively assuming no error compensation across elevation bands the approximate confidence level in our uncertainty estimates per glacier is higher than 68%.

This study aims at obtaining an ensemble of results about elevation change rates from the set of seven DEMs available for the period 2006-2015 and we thus calculate an ensemble uncertainty. The uncertainty in a sample mean is different from the uncertainty in individual observations about recent volume change rates. To identify the range of ensemble values (hereafter ‘ensemble uncertainty’) we use the standard deviation of the ensemble values

1 multiplied by 1.96. By multiplication with 1.96 we obtain 95% confidence levels, assuming
2 normal distribution.

3 For overall mass budget uncertainties we assume an ice density of 850 kg/m^3 to convert the
4 volume change into mass balance (Sapiano et al., 1998; Huss, 2013) and consider the
5 elevation change rate uncertainties and an ice density uncertainty of 60 kg/m^3 .

6 **3.2.5 Ensemble selection**

7 The possible two-fold combinations of all available DEMs are classified in two groups: maps
8 that involve the Hexagon 1974 DEM and maps that represent only 21st century elevation
9 changes (2006-2015). From the first group we only use the 1974-2006 $\Delta h/\Delta t$ map, to strictly
10 separate our two main study periods 1974-2006 and 2006-2015. The redundancy of
11 information between 2006 and 2015 allows extracting 12 maps of $\Delta h/\Delta t$ with a time interval
12 larger than 4 years. Since $\Delta h/\Delta t$ uncertainties increase with shorter time intervals between
13 DEMs (Figure 5, Table 3) $\Delta h/\Delta t$ maps with $\Delta t < 4$ years are not considered for the 2006-2015
14 ensemble. After careful evaluation of the DEMs in terms of $\Delta h/\Delta t$ uncertainties and the 2015
15 Nepal earthquake impact we discard from the 2006-2015 ensemble also all $\Delta h/\Delta t$ maps
16 involving the ALOS PRISM 2010 and the SPOT7 May 2015 scenes. The 2006-2015
17 ensemble consists therefore of six maps of $\Delta h/\Delta t$ (Figure S2).

18 The ALOS PRISM DEM is discarded because uncertainties associated to $\Delta h/\Delta t$ maps
19 involving this DEM are 30-100% higher than if other DEMs are involved (Table 3). The
20 ALOS-PRISM sensor has a radiometric resolution of 8-bit, which means that in comparison
21 to a 12-bit image (SPOT6/7, Table 2), $2^4=16$ times less information is provided per
22 panchromatic image pixel. The image contrast is therefore lower, which decreases the
23 accuracy of this DEM.

24 The May 2015 DEM is not considered for the 2006-2015 ensemble because of massive
25 deposits of avalanched snow and ice as a consequence of the April 2015 Nepal earthquake,
26 which strongly limits the representativeness of this DEM for the 2006-2015 period. The
27 October 2015 SPOT7 DEM is still considered for the 2006-2015 ensemble because until six
28 months after the earthquake most avalanche material disappeared from the glacier (section
29 4.6).

1 Due to the incomplete representation of Langtang Glacier on the SPOT6 Apr 2014 scene (the
2 scene does not cover the area north of 28°19'N, Figure S2a and d), $\Delta h/\Delta t$ maps involving this
3 DEM are excluded when discussing ensemble results for Langtang Glacier.

4 **3.3 Delineation of glaciers, debris-covered areas, and supraglacial cliffs/lakes**

5 The glacier outlines were manually delineated. We used the orthorectified satellite images
6 with the least snow cover (the Cartosat-1 2006 and 2009 scenes) to delineate the accumulation
7 areas, and assumed no changes in the accumulation area over time. The tongues of the seven
8 studied glaciers and debris extents were re-delineated for every year for which satellite images
9 are available (1974, 2006, 2009, 2010, 2014 and 2015), using the corresponding orthorectified
10 satellite images. A first operator delineated the outlines and a second operator provided
11 feedback in order to improve delineation accuracy. To quantify the uncertainty in derived
12 glacier area changes we consider a 0.5 pixel size delineation uncertainty (Paul et al., 2013).

13 The four largest glaciers in the valley were already delineated manually by Pellicciotti et al.
14 (2015) for the years 1974 and 2000. However, we decided not to use those outlines because of
15 the considerably higher resolution of the images that are available for this study and for
16 consistency in the procedure applied for different outlines. We also re-delineated the
17 catchment boundaries using the SRTM 30 m DEM and flow accumulation to accurately
18 identify the ice divides between neighboring catchments. As a result, the calculated glacier
19 areas (Table 1) changed considerably with respect to Pellicciotti et al. (2015). The 1974
20 glacier area of Langshisha Glacier changed by -40.4% (Figure S3), mostly due to clipping
21 with the catchment mask which reduced the extent of the accumulation areas. The 1974 areas
22 of Langtang, Shalbachum and Lirung changed by -8.7%, -9.5% and +8.0%, respectively.

23 To identify glacier area associated to small glaciers in the catchment that are not discussed in
24 this study we used the glacier outlines provided by the GAMDAM glacier inventory
25 (Nuimura et al., 2015). Those areas were masked out from off-glacier terrain for the co-
26 registration of the DEMs and stable terrain accuracy assessments.

27 Six quality checked maps of supraglacial cliffs and lakes are used to characterize debris-
28 covered glacier surfaces (Steiner et al., 2016). The cliff and lake inventories were generated
29 based on the available satellite imagery for the period 2006-2015 (Oct 2006, Nov 2009, Dec
30 2010, Apr 2014, May 2015 and Oct 2015). As for the glacier outlines, cliff and lake outlines
31 have been delineated by two independent operators. To further improve the accuracy of the

1 inventories, a third operator used slope and elevation change maps to identify potential cliff
2 and lake locations. The first two operators then used these indications to review the
3 inventories. All outlines have been obtained by manual delineation on the basis of the
4 orthorectified satellite images.

5 We calculated the fraction of pixels including lakes and cliffs per 50 m elevation band of each
6 debris-covered tongue (excluding tributary branches, Figure 6). In the following, we only
7 discuss median 2006-2015 cliff and lake area fractions to minimize seasonal effects. Large
8 avalanche cones, such as those present on Lirung and Langtang Glacier after the April 2015
9 earthquake, are masked out from the inventories before calculating median values.

10 **3.4 Surface velocities**

11 To assist with the interpretation of volumetric changes, we use glacier velocities determined
12 with the COSI-Corr cross-correlation feature-tracking algorithm (Leprince et al., 2007) and
13 the available satellite imagery. The orthorectified Cartosat-1 Nov 2009 and ALOS-PRISM
14 Dec 2010 images were used for this purpose. Other image pairs were not considered due to
15 longer periods between acquisitions (leading to image decorrelation) or the presence of snow
16 patches at lower elevations (SPOT6 April 2014, SPOT7 May 2015). The selected
17 orthorectified images (5 m resolution) were adjusted according to the shifts determined by co-
18 registration (Section 3.2.2). Since the window size must be large enough to avoid correlating
19 only noise but small enough not to degrade the output resolution (Dehecq et al., 2015), we
20 tested several configurations. The best results for the COSI-Corr correlation analysis were
21 achieved using multiscale window sizes of 128 down to 32 pixels, as also proposed by
22 Scherler et al. (2008). To post-process the velocity data we removed pixels with x- or y-
23 velocity values greater than 40 m/a, since these were identified as errors by manually
24 measuring the surface displacement on the basis of the orthorectified images and prominent
25 features. We then ran a median filter on the data to remove areas which show a local reversal
26 in x or y directions. Missing values were then filled with the mean of the adjacent 8 values.
27 Finally, the velocity map was resampled to 30 m resolution with a bicubic algorithm.

28 **3.5 Assessment of the April 2015 earthquake impact**

29 We quantify the impact of the avalanche events after the April 2015 earthquake on volume
30 changes of debris-covered tongues. For this purpose we use the April 2014 - May 2015 Δh

1 map to quantify the accumulated volumes less than two weeks after the earthquake, and the
2 April 2014 - Oct 2015 Δh map to quantify the remaining volumes after one ablation season.
3 To identify glacier area where avalanche material accumulated we consider all glacier grid
4 cells with significant positive elevation changes ($\Delta h > 5$ m). Approximately 7.9% (1.9 km²) of
5 all debris-covered areas were affected by avalanches according to this definition. To account
6 for pre-earthquake volume losses we first subtract from the DEM differences the elevation
7 changes between April 2014 and April 2015 determined on the basis of the Oct 2006 - Feb
8 2015 mean thinning rates. Note that we do not use the Feb 2015 - May 2015 and the Feb 2015
9 - Oct 2015 Δh maps to quantify avalanche debris volumes because the calculated uncertainties
10 associated to these maps are up to 300% higher than the uncertainties associated to
11 differential DEMs involving the Apr 2014 scene (Table S1).

12

13 **4 Results**

14 **4.1 Mean glacier surface elevation changes**

15 The 2006-2015 ensemble consistently indicates an increase in mean glacier thinning rates in
16 comparison to the period 1974-2006 (Figure 7h). For 2006-2015 we calculate an ensemble-
17 mean thinning rate of -0.45 ± 0.18 m a⁻¹, while for the period 1974-2006 we identify a
18 thinning rate of -0.24 ± 0.08 m a⁻¹ (Table 4). This corresponds to an increase in determined
19 mean thinning rates by 0.21 m a⁻¹ or 87.5%. The error bounds associated to the two periods
20 are overlapping at the extremes. However, error bounds are not overlapping at 80%
21 confidence levels (assuming normal distribution). Given the probability of less than 10% for
22 1974-2006 and 2006-2015 thinning rates for being above or below this confidence interval,
23 the estimated confidence level of accelerated thinning rates is higher than 99%.

24 From the seven studied glaciers in the valley, the thinning rates of Langtang, Langshisha and
25 Yala Glaciers have accelerated at 99% confidence levels (Figure 7, Table 4). At Shalbachum
26 Glacier the error bounds are overlapping but the estimated probability of thinning acceleration
27 is more than 90%. At Lirung and Kimoshung Glaciers the mean elevation change rates have
28 likely remained approximately constant (Table 4). Mean thinning rates of these glaciers
29 increased by less than 0.10 m a⁻¹ between 1974-2006 and 2006-2015. Also at Ghanna Glacier
30 the 1974-2006 value and the 2006-2015 ensemble mean differ by only 0.05 m a⁻¹ (Table 4).
31 However, the scatter in the 2006-2015 values is such that no trend can be identified. Ghanna

1 Glacier is the only glacier where the ensemble of values available for the period 2006-2015
2 did not narrow down the uncertainty associated to individual periods (Figure 7).

3 The most negative elevation change for 1974-2006 was observed at Shalbachum (-0.43 ± 0.08
4 m a^{-1} , Table 4) and Ghanna Glacier ($-0.51 \pm 0.05 \text{ m a}^{-1}$). The least negative values were
5 calculated for Langshisha ($-0.12 \pm 0.09 \text{ m a}^{-1}$) and Kimoshung Glaciers ($0.06 \pm 0.13 \text{ m a}^{-1}$).
6 Comparing the period 1974-2006 and the 2006-2015 ensemble mean values, the strongest
7 thinning acceleration took place at Yala Glacier (from $-0.33 \pm 0.06 \text{ m a}^{-1}$ to $-0.89 \pm 0.23 \text{ m a}^{-1}$,
8 Table 4). Yala Glacier was also the glacier with the highest 2006-2015 ensemble mean
9 thinning rate.

10 Elevation change rates are also calculated separately for the five debris-covered tongues
11 (Figure 8, Table 4). An increase in volume loss rates is evident on the Langtang, Langshisha,
12 Shalbachum and Lirung tongues. Thinning rates increased between 15% (Langtang tongue)
13 and 68% (Langshisha and Shalbachum tongues). Changes in thinning rates are not significant
14 for Ghanna tongue, but five out of six members of the 2006-2015 ensemble decreased as
15 compared to the previous period.

16 Of all debris-covered areas, the downwasting rates on Lirung tongue are the highest. This
17 applies to both the period 1974-2006 ($-1.03 \pm 0.05 \text{ m a}^{-1}$, Table 4) and to the 2006-2015
18 ensemble mean ($-1.67 \pm 0.59 \text{ m a}^{-1}$, Table 4). The 2006-2015 ensemble uncertainty is very
19 large on Lirung tongue ($\pm 0.59 \text{ m a}^{-1}$), which we believe is due to systematic errors in the
20 2009-2014 differential DEM that represents an outlier in the ensemble (Figure 8). However,
21 neither on Lirung nor on Langtang tongue (the two glaciers most affected by post-earthquake
22 avalanches, see section 4.6) post-earthquake elevation changes (2006-Oct 2015 or 2009-Oct
23 2015) represent outliers with respect to other 2006-2015 multi-annual periods. The lowest
24 volume loss rates are identified for Ghanna tongue (Figure 8, Table 4). Here, the 2006-2015
25 ensemble mean value ($-0.50 \pm 0.20 \text{ m a}^{-1}$) is about 30% of the thinning rate at Lirung tongue.

26 **4.2 Sensitivity to outlier correction and ELA definitions**

27 Mean elevation change values are most sensitive to outlier definitions for Langshisha Glacier
28 1974-2006 (Table 5). If a 2σ -level is used to define outliers for all area types (instead of a 3σ -
29 level above and a 1σ -level below the ELA, Section 3.2.3), $\Delta h/\Delta t_{1974-2006}$ for Langshisha
30 Glacier changes by -0.09 m a^{-1} from $-0.12 \pm 0.09 \text{ m a}^{-1}$ to $-0.21 \pm 0.09 \text{ m a}^{-1}$. If we compare
31 the results obtained with an estimated ELA at 5300 m a.s.l. to the results obtained with an

1 ELA at 5500 m a.s.l., mean elevation changes of individual glaciers differ by up to -0.23 m a^{-1}
2 (Shalbachum Glacier 1974-2006). However, only for two glaciers the sensitivity values
3 exceed the uncertainty values estimated from off-glacier elevation change errors (at
4 Shalbachum and Yala Glacier 1974-2006, Table 5). In both cases the differences can be
5 explained by unrealistic patterns (strongly negative elevation changes above 5400 m a.s.l.),
6 that are not identified as outliers with a 3σ threshold applied to areas below 5500 m a.s.l. Our
7 analysis thus shows that elevation change estimates are in most cases not significantly
8 influenced by outlier definitions and ELA estimates.

9 **4.3 Altitudinal distribution of elevation changes**

10 The altitudinal distribution of mean elevation changes clearly show that the thinning patterns
11 of all debris-covered tongues have changed over time (Figure 9, Figure 10). Areas with clear
12 increases in thinning rates can be identified for Langtang Glacier 5000-5150 m a.s.l.
13 (25%-100% thinning rate increase), for Langshisha Glacier 4650-5100 m a.s.l. (25%-260%),
14 for Shalbachum Glacier 4500-4800 m a.s.l. (25%-180%) and for Lirung Glacier 4300-4350
15 m a.s.l. (80%-170%). Thinning rates have remained mostly constant in the lower third of the
16 elevation ranges of the tongues (Langtang, Shalbachum and Lirung Glaciers). At Ghanna
17 Glacier, thinning rates have recently declined near the glacier terminus at 4800-4850 m a.s.l.
18 (60-90% thinning rate decrease, Figure 9e). This pattern contrasts with all other temporal
19 patterns for debris-covered glacier areas.

20 On Langshisha Glacier thinning patterns in 1974-2006 and 2006-2015 are different near the
21 terminus (Figure 9b). Here, the glacier tongue became very narrow in the last decade and
22 ultimately a small part below 4500 m a.s.l. disconnected from the main tongue (Figure 1)
23 between 2010 and 2014. The fragmentation of the tongue lead to mean thinning rates close to
24 zero at elevation bands where a substantial part of the glacier area disappeared.

25 Overall, the thinning profiles of 2006-2015 ensemble members show very similar
26 characteristics (Figure 9, Figure 10). The profiles diverge for the uppermost elevation bands
27 of the tongues and in the accumulation areas. This agrees with the larger error that is
28 attributed to higher elevations (Figure 3). Above 5500 m a.s.l. it is impossible to separate
29 uncertainty from actual differences in thinning rates.

30 On Yala Glacier there has been a three-fold increase in thinning rates below 5400 m a.s.l.,
31 comparing 1974-2006 to the 2006-2015 ensemble results (Figure 10d). Maximal thinning

1 takes place at the terminus and then decreases nearly linearly with altitude until it reaches
2 values close to zero. This is in clear contrast to the much less uniform patterns on debris-
3 covered glaciers (Figure 10a-c). On debris-covered glaciers, the elevation corresponding to
4 the maximum thinning rates is different from glacier to glacier. On Shalbachum and Lirung
5 Glaciers the maximum is reached somewhere close to the upper end of the tongue (4650-4750
6 m a.s.l. and 4300-4400 m a.s.l., respectively, Figure 9c and d), on Langtang and Ghanna
7 Glaciers more in the middle part (4950 – 5150 m a.s.l. and 4900-5000 m a.s.l., respectively,
8 Figure 9a and e) and on Langshisha Glacier closer to the terminus (4450-4700 m a.s.l., Figure
9 9b). On the large debris-covered glaciers, areas of maximum thinning seem to have shifted
10 and extended to higher elevations only at Langtang Glacier, where during the period 1974-
11 2006 maximum thinning occurred between 4850 and 4950 m a.s.l. (Figure 9a). On
12 Shalbachum Glacier maximum thinning during the period 1974-2006 occurred slightly higher
13 up at 4750 – 4800 m a.s.l. (Figure 9c).

14 Note that the altitudinal $\Delta h/\Delta t$ profiles (Figure 9, Figure 10) always refer to the same position
15 in space, since 50 m elevation bands were delimited only once on the basis of the SRTM
16 1 Arc-Second Global DEM. To account for the up-valley movement of on-glacier elevation
17 bands over time due to surface lowering, profiles would have to be slightly shifted relative to
18 each other. However, given the maximum thinning rates of 1-1.5 ma^{-1} in 1974-2006, the
19 maximum relative adjustment of values in Figure 9 and Figure 10 would never exceed one
20 50 m elevation band. Accounting for the shifting of elevation bands over time would therefore
21 not lead to different conclusions regarding changes in spatial $\Delta h/\Delta t$ patterns.

22 **4.4 Glacier area changes**

23 Debris-free Yala Glacier experienced the strongest increase in relative annual area loss of all
24 studied glaciers (1974-2006: $-0.43 \pm 0.05\% \text{ a}^{-1}$, 2006-2015: $-1.77 \pm 0.16\% \text{ a}^{-1}$, Table 6).
25 During the same two time intervals Kimoshung Glacier shrank only at rates of $0.08 \pm$
26 $0.01\% \text{ a}^{-1}$ and $0.05 \pm 0.02\% \text{ a}^{-1}$, respectively. This represents significantly lower retreat rates
27 for the second period than at Yala Glacier. The differences in area change rates are consistent
28 with the identified differences in mean glacier surface elevation changes, where the two
29 glaciers also represent opposite extremes (Section 4.1).

30 In comparison to the current retreat rates of Yala Glacier, all debris-covered glaciers are
31 shrinking at a much slower pace, with retreat rates between $-0.04 \pm 0.04\% \text{ a}^{-1}$ and $-0.40 \pm$

1 0.12% a⁻¹ (Table 6). Also debris-covered glaciers for which we observe high annual volume
2 losses have nearly stationary fronts (e.g. Shalbachum Glacier: 2006-2015 thinning rate $-0.53 \pm$
3 0.19 m a^{-1} , 2006-2015 area loss $-0.04 \pm 0.04 \text{ \% a}^{-1}$). Ghanna Glacier in contrast shows a
4 slightly more significant retreat ($-0.40 \pm 0.12\% \text{ a}^{-1}$, Table 6), although the mean thinning rates
5 are the least negative of all debris-covered areas (Figure 8).

6 **4.5 Surface velocities and supraglacial cliff/lake areas**

7 Approximately 10% of all grid cells for the three largest debris-covered tongues (Langtang,
8 Langshisha, Shalbachum) contain supraglacial cliff features ('Cliff Area' in Table 7). At
9 Lirung and Ghanna tongues this value decreases to 8% and 3%, respectively. For Ghanna
10 tongue practically no supraglacial lakes could be identified, while at the other debris-covered
11 tongues 'Lake Area' is between 2.3% and 3.3%.

12 The mean surface velocities of the tongues range between 1.6 m a^{-1} (Ghanna tongue) and
13 7 m a^{-1} (Langshisha tongue). The mean and the standard deviation of off-glacier surface
14 velocities are 1.3 m a^{-1} and 1.9 m a^{-1} , respectively. At Ghanna and Lirung tongue, which both
15 have a mean surface velocity below 3 m a^{-1} , it is therefore practically impossible to
16 discriminate moving ice from quasi-stagnant ice. Following Scherler et al. (2011b), all glacier
17 grid cells with a surface velocity of less than 2.5 m a^{-1} are therefore termed 'stagnant' for
18 simplicity. According to this definition, the tongue area classified as 'stagnant' (Table 7)
19 ranges from 20% (Langshisha tongue) to 85% (Ghanna tongue).

20 In our sample of five debris-covered glaciers, cliffs and lakes seem to appear more frequently
21 on glaciers which are dynamically active. We identify a highly significant negative correlation
22 (Pearson's linear correlation coefficient $r=-0.99$) between cliff area fraction per tongue and the
23 percentage of stagnant tongue area. 'Lake Area' and '% stagnant area' are also negatively
24 correlated ($r=-0.87$). At the scale of individual tongues, a correlation between surface
25 velocities and cliff appearance is evident at Shalbachum Glacier (Figure 11c). Here we
26 identify a correlation of 0.85 (respectively 0.68) between the altitudinal velocity profile and
27 cliff (respectively lake) areas per 50 m elevation band. Also on the two other large debris-
28 covered tongues in the valley, on Langtang and Langshisha tongues, cliff appearance clearly
29 decreases towards the termini where the glaciers are quasi-stagnant. However, here the
30 surface velocities and cliff appearance are not linearly correlated since the highest cliff area

1 densities are identified 200-300 m below the altitude ranges corresponding to maximum
2 surface velocity.

3 To investigate a possible link between accelerated thinning and the presence of supraglacial
4 lakes and cliffs we compare 'Cliff Area' and 'Lake Area' (as provided in Table 7) to changes
5 in mean thinning rates per tongue ($\Delta \Delta h/\Delta t$, difference between '1974-2006' and 'ensemble
6 mean 2006-2015' as provided in Table 4). Overall, the correlation coefficient between
7 fractional cliff area per tongue and $\Delta \Delta h/\Delta t$ is -0.62 (and -0.50 between lake area and
8 $\Delta \Delta h/\Delta t$). The likely reduced thinning rates on Ghanna tongue (Figure 8e) indeed correspond
9 to low cliff and lake area fractions (3.2% and 0.4%, respectively). On Lirung, Shalbachum
10 and Langshisha tongues thinning accelerated by 0.47-0.64 m a^{-1} , whereas fractional cliff and
11 lake areas are similar (cliff area: 8.0-10.5%, lake area: 2.3-2.6%). Also Langtang tongue is
12 characterized by relatively high cliff and lake area fractions (10% and 3.3%, respectively,
13 Table 7) but the identified changes in thinning rates are only minor. The acceleration of mean
14 thinning rates at Langtang tongue is significant at the 95% confidence level (Figure 8a), but
15 the difference in mean thinning rates 1974-2006 and 2006-2015 is only -0.12 m a^{-1} (Table 4).

16 At locations where thinning rates did not increase significantly we mostly identify low cliff
17 area fractions below 10% (e.g. on Langtang tongue below 4750 m a.s.l. and above 5150 m
18 a.s.l., at Shalbachum below 5500 m a.s.l. and at Ghanna tongue). Conversely, cliff area
19 fractions are generally higher than 10% where the 2006-2015 ensemble consistently indicates
20 thinning acceleration (Figure 11). Exception to this observation are the high cliff area
21 fractions at Langtang Glacier 4750-4900 m a.s.l., where thinning rates did not change
22 significantly (Figure 11a), and low cliff area fractions at Shalbachum Glacier 4750-4800 m
23 a.s.l., where thinning rates increased (Figure 11c). Lirung tongue also shows a different
24 behavior, except at the lowest elevation band. However, maximum thinning acceleration at
25 4300 m a.s.l. corresponds to a relatively high lake area fraction of 6% (Figure 11d).

26 Altitude bands with no significant increases in thinning rates on Langtang Glacier consistently
27 coincides with relatively low surface velocities below 5 m a^{-1} . At Langhisha and Shalbachum
28 tongues this is also the case (Figure 11). Across all debris-covered glacier tongues, 77% of all
29 elevation bands where thinning accelerated ($\Delta \Delta h/\Delta t < -0.2 \text{ m a}^{-1}$) are not stagnating
30 (velocities above 2.5 m a^{-1}), and in 72% of all elevation bands where thinning rates remained
31 constant or declined ($\Delta \Delta h/\Delta t \geq -0.2 \text{ m a}^{-1}$) we observe velocities below 2.5 m a^{-1} .

1 **4.6 Impacts of the April 2015 earthquake**

2 We calculate a total volume of post-earthquake avalanche debris in May 2015 of $2.5 \cdot 10^7 \text{ m}^3$,
3 which is equivalent to a cube length of 292 m. 40% of the avalanche material remained until 6
4 Oct 2015 (Table 8). The two glaciers which were most affected by avalanches were Langtang
5 Glacier (receiving 58% of the total volume) and Lirung Glacier (29%). The avalanche cone at
6 Lirung Glacier piled up to a height of nearly 60 m, while the avalanche material at Langtang
7 Glacier was more spread (Figure 12). Consequently, more material remained until 6 Oct 2015
8 at Lirung Glacier (57%), while at Langtang Glacier 31% remained (Table 8). Field visits at
9 the end of October 2015 to Lirung and Langtang Glaciers revealed that a smooth debris layer
10 melted out of the avalanche material and covered the surface uniformly with a thickness of a
11 few centimeters (P. Buri and P. Egli, personal communication).

12 Volume loss from glacier area where avalanche material accumulated between May and
13 October 2015 was 30 times higher than during an average ablation season (May 2015 – Oct
14 2015: $-1.5 \cdot 10^7 \text{ m}^3$, Oct 2006 – Apr 2014: $-5 \cdot 10^5 \text{ m}^3 \text{ a}^{-1}$). After this rapid initial downwasting
15 the avalanche deposits diminished to a volume of 10^7 m^3 , equivalent to an average positive
16 surface elevation change over all debris-covered glacier area of $0.52 \pm 0.19 \text{ m}$ (Table 8, or
17 $0.06 - 0.09 \text{ m a}^{-1}$ if divided over six to nine years). The avalanche impact on the Oct 2015
18 DEM is thus within the uncertainty range associated to multi-annual $\Delta h/\Delta t$ values (± 0.12
19 m a^{-1} , Table 3) and justifies why the October 2015 DEM is considered for the 2006-2015
20 ensemble.

21 The avalanche traces are still visible six months after the earthquake at Lirung Glacier (4350-
22 4400 m a.s.l.), at Langtang Glacier (4500-4900 m a.s.l.), at Langshisha Glacier (4800 m a.s.l.)
23 and at Shalbachum Glacier (4750 m a.s.l.) (Figure 9). Except for Lirung Glacier at 4350
24 m a.s.l. the 2006-Oct 2015 and 2009-Oct 2015 thinning profiles are within the error bounds
25 associated to other multi-annual periods.

26

27 **5 Discussion**

28 **5.1 Elevation changes of debris-covered glaciers**

29 Elevation changes in the debris-covered area are primarily independent of elevation (Figure
30 9), as previously identified in the Langtang catchment (Pellicciotti et al., 2015) and elsewhere

1 in high-mountain Asia (e.g. Bolch et al., 2011; Dobhal et al., 2013; Pieczonka et al., 2013;
2 Pieczonka and Bolch, 2015; Ye et al., 2015). Such patterns have usually been explained by
3 downglacier increase of debris thickness and by ablation associated with supraglacial lakes
4 and exposed ice cliffs. Our analysis shows that, with few exceptions, the highest thinning
5 rates and the strongest increase in thinning rates can be associated to areas with a high
6 concentration of ice cliffs and supraglacial ponds (Figure 11, Figure S4). While previous
7 studies have pointed out that debris-covered areas with a large presence of supraglacial cliffs
8 and lakes make a disproportionately large contribution to ablation (Reid and Brock, 2014;
9 Buri et al., 2016; Miles et al., 2016a; Thompson et al., 2016), this is the first study which
10 documents the relation between accelerations in volume loss rates and the large presence of
11 supraglacial cliffs and lakes.

12 Supraglacial cliffs seem to appear more frequently on slowly moving ice ($5\text{-}10\text{ m a}^{-1}$, Figure
13 11) and not where the glacier is stagnant (Sakai et al., 2002; Bolch et al., 2008; Thompson et
14 al., 2016). This can be explained by compressive stresses associated with flow deceleration
15 that may initiate fracturing (Benn et al., 2009). Such stresses are usually not large enough to
16 initiate open surface crevasses, but in combination with elevated water pressure due to local
17 water inputs lead to hydrologically driven fracture propagation (hydrofracturing) and
18 englacial conduit formation (Benn et al., 2009). The collapse of large englacial voids
19 destabilizes the debris layers and leads to the formation of new ice cliffs. Accordingly,
20 accelerated thinning of debris-covered area in the Upper Langtang catchment does not take
21 place on stagnating parts of the tongues, but where the transition between the active and the
22 stagnant ice can be expected (Figure 11).

23 The appearance of supraglacial lakes, on the other hand, is strongly related to the surface
24 gradient (Sakai and Fujita, 2010; Miles et al., 2016b). Large supraglacial lakes can only form
25 where the slope is less than 2° (Reynolds, 2000) and where local water input is high. These
26 conditions are not met on debris-covered glacier sections in the Upper Langtang catchment,
27 since local surface slope is consistently above 5° (Pellicciotti et al., 2015). It is interesting to
28 note that the highest lake area fractions (Lake Area $> 6\%$) are found on avalanche deposition
29 zones at Langtang Glacier (4750-4800 m a.s.l., Figure 7a and Figure 11a) and at Lirung
30 Glacier (4300 m a.s.l., Figure 7d and Figure 11d). This is likely related to high local surface
31 water inputs from melting of avalanche snow and ice. On Langtang Glacier frequent
32 avalanche inputs may explain why thinning did not accelerate at the altitude range between

1 4750 m a.s.l. and 4900 m a.s.l., in spite of the presence of exposed ice (Cliff Area > 13%,
2 Figure 11a).

3 Several studies show that lakes and cliffs are important but cannot explain the mass loss alone
4 (e.g. Sakai et al., 2002; Juen et al., 2014). The high thinning magnitudes on the upper sections
5 of Shalbachum tongue (4750-4800 m a.s.l.) likely cannot be attributed to lakes and cliffs
6 (cliff/lake area fractions are below 5%, Figure S4c), and thin layers of deposited debris in the
7 upper sections of the glacier tongue could explain such patterns.

8 Reduced ice fluxes also contribute to thinning accelerations. To assess how much this factor
9 contributes to the observed accelerations in thinning it would be necessary to quantify
10 changes in ice flux over time (e.g. Nuimura et al., 2011; Berthier and Vincent, 2012; Nuth et
11 al., 2012). However, information about the evolution of surface velocities over long time
12 periods would be required, which our dataset cannot provide. Over the timescales considered
13 in this study, on the other hand, high warming rates have been identified in this part of the
14 Himalaya (Shrestha et al., 1999; Lau et al., 2010). The rise in air temperatures directly
15 impacts glacier melt rates, and can explain rapid acceleration of thinning where ice is not
16 insulated from warming by thick debris.

17 Banerjee and Shankar (2013) numerically investigated the response of extensively debris-
18 covered glaciers to rising air-temperatures and describe the dynamical response as follows:
19 during an initial period the fronts remain almost stationary and in the ablation region a slow-
20 flowing quasi-stagnant tongue develops. During this period, which may last more than 100
21 years, glaciers lose volume by thinning. After this initial period glaciers start to retreat with a
22 higher rate, while annual volume loss decreases because of thickening debris layers. Since
23 thinning rates near the fronts of the large debris-covered glaciers in the valley (Langtang,
24 Langshisha and Shalbachum Glaciers) have not yet started to significantly decrease (Figure
25 11a-c) and the glacier tongues are still dynamically active (Figure 13) it can be assumed that
26 the quasi-stationary length period will persist for these glaciers in the near future. The model
27 of Banerjee and Shankar (2013) does not account for supraglacial cliffs and lakes, which
28 likely contribute to thinning acceleration (Figure 11). However, we have shown that they
29 primarily appear on parts of the glacier tongues which are still dynamically active (Table 7).
30 Our results suggest that they become less abundant with decreasing flow. The presence of
31 cliffs and lakes therefore does not interfere with the dynamical response of debris-covered
32 glaciers as described by Banerjee and Shankar (2013).

1 Near the snout of Ghanna Glacier a deceleration in thinning rates by -80% can be clearly
2 identified (Figure 9e, Figure 11e, 4800-4850 m a.s.l.). Previous studies have provided
3 evidence that ablation rates of debris-covered ice may decrease over time as a consequence of
4 thickening debris cover, in spite of rising air-temperatures (Banerjee and Shankar, 2013;
5 Rowan et al., 2015).

6 **5.2 Elevation changes of debris-free glaciers**

7 2006-2015 downwasting rates on Yala Glacier are 0.5-1.2 m a⁻¹ higher than on Kimoshung
8 Glacier (Table 4). However, the two glaciers have a very different hypsometry (Figure S5).
9 Kimoshung Glacier has a very steep tongue that reaches to similarly low elevations as the
10 debris-covered glacier tongues (Table 1). The glacier is nearly in equilibrium with the climate
11 (Table 4), which explains the low thinning rates at low elevations (Figure S5). Currently the
12 estimated AAR of Yala Glacier is 40% (Table 1), which is a common value in the HKH
13 region (Kääb et al., 2012). The estimated AAR of 86% at Kimoshung Glacier, on the other
14 hand, corresponds to an exceptionally high value for the HKH (Khan et al., 2015). The
15 differences in volume loss rates point to the role of glacier hypsometry for the response of
16 debris-free glaciers to climatic changes (e.g. Jiskoot et al., 2009). Almost balanced mass
17 budgets in recent years (Table 4) and only minor area changes (Table 6) are associated to
18 Kimoshung Glacier. Thinning did not increase significantly with respect to the period 1974-
19 2006 (Figure 7g). Due to the steep tongue of this glacier the AAR is not sensitive to changes
20 in the ELA (Table 5). Only a small fraction of area is therefore additionally exposed to
21 temperatures above freezing level in case of atmospheric warming, which causes the glacier
22 to be less sensitive to observed warming trends in the region (Shrestha et al., 1999; Lau et al.,
23 2010). One possible explanation for the balanced conditions of Kimoshung Glacier could
24 therefore be that precipitation in recent decades remained approximately stable, which agrees
25 with the findings of studies on precipitation trends in this part of the Himalaya (Shrestha et
26 al., 2000; Immerzeel, 2008; Singh et al., 2008). However, further analysis is required for
27 justification. The mass balance of Yala Glacier, on the other hand, is sensitive to fluctuations
28 in temperature. A hypothetical rise of the ELA by 100 m at this glacier causes 30% of its area
29 to turn from accumulation into ablation area (Table 5). Accordingly, thinning of Yala Glacier
30 is accelerating rapidly (Figure 9f). Due to the common AAR of Yala Glacier and the extreme
31 topography of Kimoshung Glacier it can be assumed that other debris-free glaciers in the

1 region are also thinning and that balanced conditions such as observed on Kimoshung Glacier
2 are exceptional.

3 **5.3 Differences between debris-free and debris-covered glaciers**

4 The response of debris-covered and debris-free glaciers to warming is substantially different,
5 as described in the two sections above and exemplified by the altitudinal elevation change
6 profiles in Figure 10. Our observations do not support the findings of previous studies about
7 similar present-day lowering rates of debris-covered and debris-free glacier areas at the same
8 elevation (Kääb et al., 2012; Nuimura et al., 2012; Gardelle et al., 2013). Also for debris-
9 covered elevation bands where up to 18% of the area is covered by supraglacial cliffs and
10 lakes (e.g. at Langtang tongue 5050 m a.s.l. or at Langshisha tongue 4750 m a.s.l.) thinning
11 rates do not exceed 1.8 m a^{-1} , while for Yala Glacier the lowering rates are already above this
12 value at 5250 m a.s.l. and further increase downglacier (Figure 10). Within the same
13 altitudinal range (5200-5300 m a.s.l.) thinning rates of debris-covered glaciers do not exceed
14 35%-75% of the thinning rates of Yala Glacier.

15 Regarding the mean surface elevation changes (Table 4), our observations reveal a
16 heterogeneous response to climate of both the debris-free and the debris-covered glaciers. As
17 discussed in the two sections above, there are examples for both types of glaciers where
18 thinning has increased significantly or where thinning remained approximately constant. A
19 significant difference in thinning trends between debris-free and debris-covered glaciers in
20 our sample cannot be identified. In our sample, the best predictor for thinning accelerations
21 seems to be the altitude distributions of glaciers. Glaciers with a high AAR (Kimoshung) or
22 which reach the highest elevations (Lirung) have the most balanced mass budgets and show
23 no significant changes in volume loss over time (Figure 7, Table 4). Glaciers which are most
24 sensitive to ELA changes (more than $\pm 10\%$ AAR change in response to $\pm 100 \text{ m}$ ELA
25 uncertainty, Table 5) such as Yala, Langtang and Langshisha Glaciers reveal the most
26 significant thinning accelerations (Figure 7, Table 4). However, debris-free Yala Glacier is
27 currently downwasting at 60%-100% higher rates than the large debris-covered glaciers in the
28 valley. Considering the common characteristics of Yala Glacier and given that this glacier has
29 been denominated as a benchmark glacier for the Nepal Himalayas (Fujita and Nuimura,
30 2011) it seems important that future geodetic or field based studies extend our analysis to
31 larger glacier samples.

1 **5.4 Post-earthquake avalanche impacts**

2 Accumulation by debris-laden avalanches is one of the most important processes for debris-
3 covered glacier formation (Scherler et al., 2011a). The tongue of Lirung Glacier would likely
4 not exist without accumulation through avalanches (Ragetli et al., 2015). It is detached from
5 the accumulation area (Figure 1) and reaches 200-700 m lower elevations than all other
6 debris-covered glaciers (Table 1). Our volume calculations of the post-earthquake avalanche
7 impact allow quantifying the avalanche impact on mass balance and comparing it to mass loss
8 during an average year. Given the avalanche deposits remaining on Lirung tongue by 6 Oct
9 2015 (divided by the area of the tongue: 3.87 ± 0.23 m, Table 8) and the average $\Delta h/\Delta t$ rates
10 between Oct 2006 and Feb 2015 of -1.64 ± 0.10 m a⁻¹ (Figure 8d), the avalanche after the
11 earthquake compensated by 240% the volume loss of one average year. At the scale of all
12 debris-covered area in the valley this value amounts to 50% (0.52 ± 0.19 m avalanche
13 deposits and -1.02 ± 0.08 m a⁻¹ average thinning). According to Scally and Gardner (1989)
14 avalanche deposit density increases until the end of the ablation season to about 80% of ice
15 density. The mass deposits therefore compensate mass loss during a normal year by about
16 180% at Lirung tongue (40% at the catchment scale). Still, our analysis has revealed that the
17 impacts are not significant in comparison to the 2006-2015 ensemble uncertainty (Section 4.6,
18 Figure 8d and f).

19 **5.5 Comparison to other studies**

20 The four largest debris-covered glaciers in the valley (Langtang, Langshisha, Shalbachum,
21 Lirung) have been the focus of a recent geodetic mass balance study by Pellicciotti et al.
22 (2015), who reconstructed elevation and mass changes using the 1974 Hexagon DEM which
23 is also used in this study (spatial resolution 30 m) and the 2000 SRTM3 DEM (90 m). They
24 found that all four glaciers lost mass over the study period but with different rates (on average
25 -0.32 ± 0.18 m w.e. a⁻¹). We find an overall glacier mass balance for the period 1974-2006 of
26 the four glaciers which is slightly less negative (-0.22 ± 0.08 m w.e. a⁻¹). However, the results
27 match within the uncertainties. The lower uncertainty estimates by our study are justified by
28 the high resolution and quality of the 2006 Cartosat-1 DEM (Table 3). Differences in the mass
29 balance of Langtang, Lirung and Shalbachum Glacier are within uncertainty bounds and can
30 be attributed to differences in used glacier masks, study period, outlier correction approaches,
31 density assumptions and uncertainties regarding the penetration depth of the SRTM radar
32 signal (Kääb et al., 2015). However, for Langshisha Glacier we calculate a mass balance

1 which is substantially less negative than in Pellicciotti et al. (2015). While we identify almost
2 balanced conditions for the period 1974-2006 (-0.10 ± 0.08 m w.e. a^{-1} , Table 4), the mass
3 balance indicated by Pellicciotti et al. (2015) is very negative (-0.79 ± 0.18 m w.e. a^{-1}). The
4 discrepancy can be explained by the overestimated extent of the accumulation areas by
5 Pellicciotti et al. (2015) (Figure S3) in combination with unrealistic lowering rates of up to
6 -2 m a^{-1} at about 6000 m a.s.l. (Figure 4d in Pellicciotti et al., 2015). The more realistic
7 elevation change values obtained by the present study for the accumulation areas (ranging
8 between -0.4 and $+0.4$ m a^{-1} , Figure 10b) point to the need of restrictive outlier definitions and
9 the advantage of having information from multiple datasets available for gap filling.

10 Yala Glacier has been frequently visited for field measurements in the last 25 years.
11 Sugiyama et al. (2013) calculated mean thinning rates of Yala Glacier for the periods 1982-
12 1996 (-0.69 ± 0.25 m a^{-1}) and 1996-2009 (-0.75 ± 0.24 m a^{-1}) on the basis of ground
13 photogrammetry and GPS surveys. The values suggest a more moderate acceleration of
14 volume loss rates than in our study (-0.33 ± 0.06 m a^{-1} 1974-2006 to -0.89 ± 0.23 m a^{-1} 2006-
15 2015, Table 4). However, similarly to our study Sugiyama et al. (2013) identified a rapid
16 acceleration of thinning rates at the lowest elevations. At higher elevations the uncertainty of
17 photogrammetric surveys increases because of low contrast due to homogeneous snow layers.

18 The acceleration in mass loss in recent periods identified by this study agrees with other
19 studies from the Nepalese Himalaya which assess multi-temporal elevation changes (Bolch et
20 al., 2011; Nuimura et al., 2012). Bolch et al. (2011) identify an increase in mass loss rates by
21 0.47 m w.e. a^{-1} comparing the two periods 1970-2007 (-0.32 ± 0.08 m w.e. a^{-1}) and 2002-2007
22 (-0.79 ± 0.52 m w.e. a^{-1}). Nuimura et al. (2012) calculate increasing mass losses in the same
23 study region between 1992-2008 (-0.26 ± 0.24 m w.e. a^{-1}) and 2000-2008 (-0.45 ± 0.60 m
24 w.e. a^{-1}). However, the identified acceleration in glacier thinning is not significant given the
25 largely overlapping error bounds. Moreover, the mass loss estimates of Gardelle et al. (2013)
26 for the Khumbu region and the period 2001-2011 (average of -0.41 ± 0.21 m w.e. a^{-1}) are in
27 the same order as calculated by Bolch et al. (2011) for 1970-2007. The ensemble approach of
28 this study can therefore substantially strengthen previous conclusions that mass loss of
29 glaciers in the Central Himalaya is accelerating. The volume changes calculated over several
30 multi-year periods between 2006 and 2015 consistently indicate that glacier thinning has
31 indeed accelerated (Figure 7h).

32

1 **6 Conclusions**

2 This study presents glacier volume changes of seven glaciers (five partially debris-covered,
3 two debris-free) in the upper Langtang catchment in Nepal, using a digital elevation model
4 (DEM) from 1974 stereo Hexagon satellite data and seven DEMs derived from 2006-2015
5 stereo or tri-stereo satellite imagery. We carefully selected elevation change maps which are
6 least affected by uncertainty to obtain multiple independent DEM differences for the period
7 2006-2015.

8 Our results point to increasing thinning rates, from $-0.24 \pm 0.08 \text{ m a}^{-1}$ in 1974-2006 to $-0.45 \pm$
9 0.18 m a^{-1} in 2006-2015, where the estimated confidence level of accelerated thinning rates is
10 higher than 99%. This study therefore supports the findings of previous studies (Bolch et al.,
11 2011; Nuimura et al., 2012) that glacier wastage in the Central Himalaya is accelerating.
12 However, whereas a majority of glaciers in the study region are thinning rapidly, glaciers with
13 a high accumulation area have almost balanced mass budgets and experience no or only
14 insignificant accelerations in thinning.

15 Our observations also reveal that thinning has mostly accelerated in the upper reaches of the
16 tongues (up to +150%, comparing the periods 1974-2006 and 2006-2015), while the nearly
17 stagnant areas near the terminus show constant or decreasing thinning rates (up to -80%). The
18 highest thinning rates and the strongest increase in thinning rates can be associated to areas
19 with a high concentration of ice cliffs and supraglacial ponds. Constant or decelerating
20 thinning rates can be associated to areas with relatively homogeneous debris layers near the
21 termini of glaciers. We conclude that the response of extensively debris-covered glaciers to
22 global warming is largely determined by feedback processes associated to different surface
23 characteristics.

24 The behavior of glaciers in the study area is highly heterogeneous, and the presence or
25 absence of debris is not a good predictor for mass balance trends. However, the spatial
26 thinning patterns on debris-covered glaciers are fundamentally different than those on debris-
27 free glaciers. Debris-free glaciers in our sample present thinning rates that are linearly
28 dependent on elevation, while debris-covered glaciers have highly non-linear altitudinal
29 elevation change profiles. Our observations do not provide evidence for the existence of a so-
30 called debris-cover anomaly, where the insulating effect of thick supraglacial debris is
31 compensated by enhanced melt from exposed ice cliffs or due to high energy absorption at
32 supraglacial ponds. Within the same altitudinal range, lowering rates on debris-free Yala

1 Glacier are 35%-300% higher than on debris-covered glacier area. On debris-free Kimoshung
2 Glacier the thinning rates are similar to those of debris-covered area, but this result must be
3 explained by compressive flows that compensate for surface lowering by ablation due to its
4 exceptionally balanced conditions.

5 Geodetic mass balance studies such as this have been increasingly revealing heterogeneous
6 patterns of changes and a complex response of debris-covered glaciers that call for an
7 enhanced understanding of processes over debris-covered glaciers. Their ablation, mass
8 balance and response to climate is modulated by debris supply, transport, glacier flow, lakes
9 and cliffs developments and a complex subglacial hydrology and hydraulics that all need to be
10 understood in the future to be able to predict future changes of these glaciers over multiple
11 time scales.

12 **Acknowledgements**

13 This study is funded by the Swiss National Science Foundation (SNF) project UNCOMUN
14 (Understanding Contrasts in High Mountain Hydrology in Asia). T. Bolch acknowledges
15 funding through German Research Foundation (DFG, code BO 3199/2-1) and European
16 Space Agency (Glaciers_cci project, code 400010177810IAM). We thank Evan Miles for
17 helping with the glacier delineations and the post-processing of surface velocity data. Jakob
18 Steiner and Pascal Buri manually delineated cliffs and lakes for the inventories used in this
19 study, and they are gratefully acknowledged. We thank Etienne Berthier for the Pléiades
20 image data and Fanny Brun for her help with the identification of GCPs. DigitalGlobe
21 imagery was used to produce the WorldView-1 and 2 digital elevation models. We thank
22 Etienne Berthier (Scientific Editor), Eyjolfur Magnússon and one anonymous reviewer for
23 extensive and thorough comments that considerably helped to improve the paper.

24

1 **References**

- 2 Banerjee, A. and Shankar, R.: On the response of Himalayan glaciers to climate change, *J.*
3 *Glaciol.*, 59(215), 480–490, doi:10.3189/2013JoG12J130, 2013.
- 4 Benn, D., Gulley, J., Luckman, A., Adamek, A. and Glowacki, P. S.: Englacial drainage
5 systems formed by hydrologically driven crevasse propagation, *J. Glaciol.*, 55(191), 513–523,
6 doi:10.3189/002214309788816669, 2009.
- 7 Berthier, E. and Vincent, C.: Relative contribution of surface mass-balance and ice-flux
8 changes to the accelerated thinning of Mer de Glace, French Alps, over 1979–2008, *J.*
9 *Glaciol.*, 58(209), 501–512, doi:10.3189/2012JoG11J083, 2012.
- 10 Berthier, E., Arnaud, Y., Kumar, R., Ahmad, S., Wagnon, P. and Chevallier, P.: Remote
11 sensing estimates of glacier mass balances in the Himachal Pradesh (Western Himalaya,
12 India), *Remote Sens. Environ.*, 108(3), 327–338, doi:10.1016/j.rse.2006.11.017, 2007.
- 13 Berthier, E., Vincent, C., Magnússon, E., Gunnlaugsson, Á. Þ., Pitte, P., Le Meur, E.,
14 Masiokas, M., Ruiz, L., Pálsson, F., Belart, J. M. C. and Wagnon, P.: Glacier topography and
15 elevation changes derived from Pléiades sub-meter stereo images, *Cryosph.*, 8(6), 2275–2291,
16 doi:10.5194/tc-8-2275-2014, 2014.
- 17 Bignone, F. and Umakawa, H.: Assessment of ALOS PRISM digital elevation model
18 extraction over Japan, *Int. Arch. Photogramm. Remote Sens. Spat. Inf. Sci.*, 37, 1135–1138,
19 2008.
- 20 Bolch, T., Buchroithner, M. and Pieczonka, T.: Planimetric and volumetric glacier changes in
21 the Khumbu Himal, Nepal, since 1962 using Corona, Landsat TM and ASTER data, *J.*
22 *Glaciol.*, 54(187), 592–600, doi:10.3189/002214308786570782, 2008.
- 23 Bolch, T., Pieczonka, T. and Benn, D. I.: Multi-decadal mass loss of glaciers in the Everest
24 area (Nepal Himalaya) derived from stereo imagery, *Cryosph.*, 5(2), 349–358, doi:10.5194/tc-
25 5-349-2011, 2011.
- 26 Bolch, T., Kulkarni, A., Käab, A., Huggel, C., Paul, F., Cogley, J. G., Frey, H., Kargel, J. S.,
27 Fujita, K., Scheel, M., Bajracharya, S. and Stoffel, M.: The state and fate of Himalayan
28 glaciers., *Science*, 336(6079), 310–314, doi:10.1126/science.1215828, 2012.
- 29 Brun, F., Buri, P., Miles, E. S., Wagnon, P., Steiner, J., Berthier, E., Ragettli, S., Immerzeel,
30 W. W. and Pellicciotti, F.: Quantifying volume loss from ice cliffs on debris-covered glaciers
31 using high resolution terrestrial and aerial photogrammetry, *J. Glaciol.*, in press, 1–12,
32 doi:10.1017/jog.2016.54, 2016.
- 33 Buri, P., Pellicciotti, F., Steiner, J. F., Miles, E. S. and Immerzeel, W. W.: A grid-based model
34 of backwasting of supraglacial ice cliffs on debris-covered glaciers, *Ann. Glaciol.*, 57(71),
35 199–211, doi:10.3189/2016AoG71A059, 2016.

- 1 Burnett, M.: Hexagon (KH-9) Mapping Program and Evolution, National Reconnaissance
2 Office, Chantilly, Virginia., 2012.
- 3 Collier, E. and Immerzeel, W.: High-resolution modeling of atmospheric dynamics in the
4 Nepalese Himalaya, *J. Geophys. Res. Atmos.*, 120(19), 9882–9896,
5 doi:10.1002/2015JD023266.Received, 2015.
- 6 Dehecq, A., Gourmelen, N. and Trouve, E.: Deriving large-scale glacier velocities from a
7 complete satellite archive: Application to the Pamir–Karakoram–Himalaya, *Remote Sens.*
8 *Environ.*, 162, 55–66, doi:10.1016/j.rse.2015.01.031, 2015.
- 9 Dobhal, D. P., Mehta, M. and Srivastava, D.: Influence of debris cover on terminus retreat
10 and mass changes of Chorabari Glacier, Garhwal region, central Himalaya, India, *J. Glaciol.*,
11 59(217), 961–971, doi:10.3189/2013JoG12J180, 2013.
- 12 Fujita, K.: Effect of precipitation seasonality on climatic sensitivity of glacier mass balance,
13 *Earth Planet. Sci. Lett.*, 276(1-2), 14–19, doi:10.1016/j.epsl.2008.08.028, 2008.
- 14 Fujita, K. and Nuimura, T.: Spatially heterogeneous wastage of Himalayan glaciers., *Proc.*
15 *Natl. Acad. Sci. U. S. A.*, 108(34), 14011–14014, doi:10.1073/pnas.1106242108, 2011.
- 16 Gardelle, J., Berthier, E., Arnaud, Y. and Kääb, a.: Region-wide glacier mass balances over
17 the Pamir-Karakoram-Himalaya during 1999–2011, *Cryosph.*, 7(4), 1263–1286,
18 doi:10.5194/tc-7-1263-2013, 2013.
- 19 Holzer, N., Vijay, S., Yao, T., Xu, B., Buchroithner, M. and Bolch, T.: Four decades of
20 glacier variations at Muztagh Ata (eastern Pamir): a multi-sensor study including Hexagon
21 KH-9 and Pléiades data, *Cryosph.*, 9(6), 2071–2088, doi:10.5194/tc-9-2071-2015, 2015.
- 22 Huss, M.: Density assumptions for converting geodetic glacier volume change to mass
23 change, *Cryosph.*, 7(3), 877–887, doi:10.5194/tc-7-877-2013, 2013.
- 24 Huss, M., Jouvett, G., Farinotti, D. and Bauder, A.: Future high-mountain hydrology: a new
25 parameterization of glacier retreat, *Hydrol. Earth Syst. Sci.*, 14(5), 815–829,
26 doi:10.5194/hess-14-815-2010, 2010.
- 27 Immerzeel, W.: Historical trends and future predictions of climate variability in the
28 Brahmaputra basin, *Int. J. Climatol.*, 28(2), 243–254, doi:10.1002/joc.1528, 2008.
- 29 Immerzeel, W. W., Kraaijenbrink, P. D. a., Shea, J. M., Shrestha, A. B., Pellicciotti, F.,
30 Bierkens, M. F. P. and de Jong, S. M.: High-resolution monitoring of Himalayan glacier
31 dynamics using unmanned aerial vehicles, *Remote Sens. Environ.*, 150, 93–103,
32 doi:10.1016/j.rse.2014.04.025, 2014a.
- 33 Immerzeel, W. W., Petersen, L., Ragettli, S. and Pellicciotti, F.: The importance of observed
34 gradients of air temperature and precipitation for modeling runoff from a glacierized
35 watershed in the Nepalese Himalayas, *Water Resour. Res.*, 50(3), 2212–2226,
36 doi:10.1002/2013WR014506, 2014b.

- 1 Jiskoot, H., Curran, C. J., Tessler, D. L. and Shenton, L. R.: Changes in Clemenceau Icefield
2 and Chaba Group glaciers, Canada, related to hypsometry, tributary detachment, length–slope
3 and area–aspect relations, *Ann. Glaciol.*, 50(53), 133–143,
4 doi:10.3189/172756410790595796, 2009.
- 5 Juen, M., Mayer, C., Lambrecht, a., Han, H. and Liu, S.: Impact of varying debris cover
6 thickness on ablation: a case study for Koxkar Glacier in the Tien Shan, *Cryosph.*, 8(2), 377–
7 386, doi:10.5194/tc-8-377-2014, 2014.
- 8 Kääh, A., Berthier, E., Nuth, C., Gardelle, J. and Arnaud, Y.: Contrasting patterns of early
9 twenty-first-century glacier mass change in the Himalayas, *Nature*, 488(7412), 495–498,
10 doi:10.1038/nature11324, 2012.
- 11 Kääh, A., Treichler, D., Nuth, C. and Berthier, E.: Brief Communication: Contending
12 estimates of 2003–2008 glacier mass balance over the Pamir–Karakoram–Himalaya,
13 *Cryosph.*, 9(2), 557–564, doi:10.5194/tc-9-557-2015, 2015.
- 14 Kargel, J., Leonard, G., Shugar, D. et al.: Geomorphic and geologic controls of geohazards
15 induced by Nepal’s 2015 Gorkha earthquake, *Science*, 351(6269), 1–18,
16 doi:10.1126/science.aac8353, 2016.
- 17 Khan, A., Naz, B. S. and Bowling, L. C.: Separating snow, clean and debris covered ice in the
18 Upper Indus Basin, Hindukush-Karakoram-Himalayas, using Landsat images between 1998
19 and 2002, *J. Hydrol.*, 521, 46–64, doi:10.1016/j.jhydrol.2014.11.048, 2015.
- 20 Lamsal, D., Sawagaki, T. and Watanabe, T.: Digital terrain modelling using Corona and
21 ALOS PRISM data to investigate the distal part of Imja Glacier, Khumbu Himal, Nepal, *J.*
22 *Mt. Sci.*, 8(3), 390–402, doi:10.1007/s11629-011-2064-0, 2011.
- 23 Lau, W. K. M., Kim, M.-K., Kim, K.-M. and Lee, W.-S.: Enhanced surface warming and
24 accelerated snow melt in the Himalayas and Tibetan Plateau induced by absorbing aerosols,
25 *Environ. Res. Lett.*, 5(2), 025204, doi:10.1088/1748-9326/5/2/025204, 2010.
- 26 Leprince, S., Barbot, S., Ayoub, F. and Avouac, J.-P.: Automatic and precise
27 orthorectification, coregistration, and subpixel correlation of satellite images, application to
28 ground deformation measurements, *IEEE Trans. Geosci. Remote Sens.*, 45(6), 1529–1558,
29 2007.
- 30 Magnússon, E., Muñoz-Cobo Belart, J., Pálsson, F., Ágústsson, H. and Crochet, P.: Geodetic
31 mass balance record with rigorous uncertainty estimates deduced from aerial photographs and
32 lidar data – Case study from Drangajökull ice cap, NW Iceland, *Cryosph.*, 10(1), 159–177,
33 doi:10.5194/tc-10-159-2016, 2016.
- 34 Mattson, L. E., Gardner, J. S. and Young, G. J.: Ablation on Debris Covered Glaciers: an
35 Example from the Rakhiot Glacier, Punjab, Himalaya, *IAHS Publ.*, 218, 289–296, 1993.
- 36 Maurer, J. and Rupper, S.: Tapping into the Hexagon spy imagery database: A new automated
37 pipeline for geomorphic change detection, *ISPRS J. Photogramm. Remote Sens.*, 108, 113–
38 127, doi:10.1016/j.isprsjprs.2015.06.008, 2015.

- 1 Mayer, C., Lambrecht, A., Belo, M., Smiraglia, C. and Diolaiuti, G.: Glaciological
2 characteristics of the ablation zone of Baltoro glacier, Karakoram, Pakistan, *Ann. Glaciol.*,
3 43(1), 123–131, doi:10.3189/172756406781812087, 2006.
- 4 Mihalcea, C., Mayer, C., Diolaiuti, G., Lambrecht, A., Smiraglia, C. and Tartari, G.: Ice
5 ablation and meteorological conditions on the debris-covered area of Baltoro glacier,
6 Karakoram, Pakistan, *Ann. Glaciol.*, 43(1), 292–300, doi:10.3189/172756406781812104,
7 2006.
- 8 Mihalcea, C., Mayer, C., Diolaiuti, G., Agata, C. D., Smiraglia, C., Lambrecht, A.,
9 Vuillermoz, E. and Tartari, G.: Spatial distribution of debris thickness and melting from
10 remote-sensing and meteorological data, at debris-covered Baltoro glacier, Karakoram,
11 Pakistan, *Ann. Glaciol.*, 48(1), 49–57, doi:10.3189/172756408784700680, 2008.
- 12 Miles, E. S., Pellicciotti, F., Willis, I. C., Steiner, J. F., Buri, P. and Arnold, N. S.: Refined
13 energy-balance modelling of a supraglacial pond, Langtang Khola, Nepal, *Ann. Glaciol.*,
14 57(71), 29–40, doi:10.3189/2016AoG71A421, 2016a.
- 15 Miles, E. S., Willis, I. C., Arnold, N. S. and Pellicciotti, F.: Spatial, seasonal, and interannual
16 variability of supraglacial ponds in the Langtang Valley of Nepal, 1999 to 2013, *J. Glaciol.*,
17 under revision, 2016b.
- 18 Noh, M.-J. and Howat, I. M.: Automated stereo-photogrammetric DEM generation at high
19 latitudes: Surface Extraction with TIN-based Search-space Minimization (SETSM) validation
20 and demonstration over glaciated regions, *GIScience Remote Sens.*, 52(2), 198–217,
21 doi:10.1080/15481603.2015.1008621, 2015.
- 22 Nuimura, T., Fujita, K., Fukui, K., Asahi, K., Aryal, R. and Ageta, Y.: Temporal Changes in
23 Elevation of the Debris-Covered Ablation Area of Khumbu Glacier in the Nepal Himalaya
24 since 1978, *Arctic, Antarct. Alp. Res.*, 43(2), 246–255, doi:10.1657/1938-4246-43.2.246,
25 2011.
- 26 Nuimura, T., Fujita, K., Yamaguchi, S. and Sharma, R. R.: Elevation changes of glaciers
27 revealed by multitemporal digital elevation models calibrated by GPS survey in the Khumbu
28 region, Nepal Himalaya, 1992–2008, *J. Glaciol.*, 58(210), 648–656,
29 doi:10.3189/2012JoG11J061, 2012.
- 30 Nuimura, T., Sakai, A., Taniguchi, K., Nagai, H., Lamsal, D., Tsutaki, S., Kozawa, A.,
31 Hoshina, Y., Takenaka, S., Omiya, S., Tsunematsu, K., Tshering, P. and Fujita, K.: The
32 GAMDAM glacier inventory: a quality-controlled inventory of Asian glaciers, *Cryosph.*, 9(3),
33 849–864, doi:10.5194/tc-9-849-2015, 2015.
- 34 Nuth, C. and Kääb, A.: Co-registration and bias corrections of satellite elevation data sets for
35 quantifying glacier thickness change, *Cryosph.*, 5(1), 271–290, doi:10.5194/tc-5-271-2011,
36 2011.
- 37 Nuth, C., Schuler, T. V., Kohler, J., Altena, B. and Hagen, J. O.: Estimating the long-term
38 calving flux of Kronebreen, Svalbard, from geodetic elevation changes and mass-balance
39 modelling, *J. Glaciol.*, 58(207), 119–133, doi:10.3189/2012JoG11J036, 2012.

- 1 Östrem, G.: Ice melting under a thin layer of moraine, and the existence of ice cores in
2 moraine ridges, *Geogr. Ann.*, 41(4), 228–230, 1959.
- 3 Paul, F., Barrand, N. E., Baumann, S., Berthier, E., Bolch, T., Casey, K., Frey, H., Joshi, S.
4 P., Konovalov, V., Bris, R. Le, Mölg, N., Nosenko, G., Nuth, C., Pope, a., Racoviteanu, a.,
5 Rastner, P., Raup, B., Scharrer, K., Steffen, S. and Winsvold, S.: On the accuracy of glacier
6 outlines derived from remote-sensing data, *Ann. Glaciol.*, 54(63), 171–182,
7 doi:10.3189/2013AoG63A296, 2013.
- 8 Pellicciotti, F., Stephan, C., Miles, E., Immerzeel, W. W. and Bolch, T.: Mass-balance
9 changes of the debris-covered glaciers in the Langtang Himal, Nepal, 1974–99, *J. Glaciol.*,
10 61(225), doi:10.3189/2015JoG13J237, 2015.
- 11 Pieczonka, T. and Bolch, T.: Region-wide glacier mass budgets and area changes for the
12 Central Tien Shan between ~1975 and 1999 using Hexagon KH-9 imagery, *Glob. Planet.*
13 *Change*, 128, 1–13, doi:10.1016/j.gloplacha.2014.11.014, 2015.
- 14 Pieczonka, T., Bolch, T. and Buchroithner, M.: Generation and evaluation of multitemporal
15 digital terrain models of the Mt. Everest area from different optical sensors, *ISPRS J.*
16 *Photogramm. Remote Sens.*, 66(6), 927–940, doi:10.1016/j.isprsjprs.2011.07.003, 2011.
- 17 Pieczonka, T., Bolch, T., Junfeng, W. and Shiyin, L.: Heterogeneous mass loss of glaciers in
18 the Aksu-Tarim Catchment (Central Tien Shan) revealed by 1976 KH-9 Hexagon and 2009
19 SPOT-5 stereo imagery, *Remote Sens. Environ.*, 130, 233–244,
20 doi:10.1016/j.rse.2012.11.020, 2013.
- 21 Pratap, B., Dobhal, D. P., Mehta, M. and Bhambri, R.: Influence of debris cover and altitude
22 on glacier surface melting: a case study on Dokriani Glacier, central Himalaya, India, *Ann.*
23 *Glaciol.*, 56(70), 9–16, doi:10.3189/2015AoG70A971, 2015.
- 24 Racoviteanu, A. E., Arnaud, Y., Williams, M. W. and Manley, W. F.: Spatial patterns in
25 glacier characteristics and area changes from 1962 to 2006 in the Kanchenjunga–Sikkim area,
26 eastern Himalaya, *Cryosph.*, 9(2), 505–523, doi:10.5194/tc-9-505-2015, 2015.
- 27 Ragettli, S., Pellicciotti, F., Immerzeel, W. W., Miles, E. S., Petersen, L., Heynen, M., Shea,
28 J. M., Stumm, D., Joshi, S. and Shrestha, A.: Unraveling the hydrology of a Himalayan
29 catchment through integration of high resolution in situ data and remote sensing with an
30 advanced simulation model, *Adv. Water Resour.*, 78, 94–111,
31 doi:10.1016/j.advwatres.2015.01.013, 2015.
- 32 Reid, T. D. and Brock, B. W.: Assessing ice-cliff backwasting and its contribution to total
33 ablation of debris-covered Miage glacier, Mont Blanc massif, Italy, *J. Glaciol.*, 60(219), 3–13,
34 doi:10.3189/2014JoG13J045, 2014.
- 35 Reynolds, J.: On the formation of supraglacial lakes on debris-covered glaciers, *IAHS Publ.*,
36 (264), 153–161, 2000.
- 37 Rowan, A. V., Egholm, D. L., Quincey, D. J. and Glasser, N. F.: Modelling the feedbacks
38 between mass balance, ice flow and debris transport to predict the response to climate change

- 1 of debris-covered glaciers in the Himalaya, *Earth Planet. Sci. Lett.*, 430, 427–438,
2 doi:10.1016/j.epsl.2015.09.004, 2015.
- 3 Sakai, A. and Fujita, K.: Formation conditions of supraglacial lakes on debris-covered
4 glaciers in the Himalaya, *J. Glaciol.*, 56(195), 177–181, doi:10.3189/002214310791190785,
5 2010.
- 6 Sakai, A., Nakawo, M. and Fujita, K.: Melt rate of ice cliffs on the Lirung Glacier, Nepal
7 Himalayas, 1996, *Bull. Glacier Res.*, 16, 57–66, 1998.
- 8 Sakai, A., Takeuchi, N., Fujita, K. and Nakawo, M.: Role of supraglacial ponds in the
9 ablation process of a debris-covered glacier in the Nepal Himalayas, *Debris-Covered Glaciers*,
10 IAHS Publ., 265, 119–130, 2000.
- 11 Sakai, A., Nakawo, M. and Fujita, K.: Distribution characteristics and energy balance of ice
12 cliffs on debris-covered glaciers, Nepal Himalaya, *Arctic, Antarct. Alp. Res.*, 34(1), 12–19,
13 doi:10.2307/1552503, 2002.
- 14 Sapiano, J., Harrison, W. and Echelmeyer, K.: Elevation, volume and terminus changes of
15 nine glaciers in North America, *J. Glaciol.*, 44(146), 119–135, doi:10.3198/1998JoG44-146-
16 119-135, 1998.
- 17 Scally, F. De and Gardner, J.: Evaluation of avalanche mass determination approaches: an
18 example from the Himalaya, Pakistan, *J. Glaciol.*, 35(120), 248–252, 1989.
- 19 Scherler, D., Leprince, S. and Strecker, M.: Glacier-surface velocities in alpine terrain from
20 optical satellite imagery—Accuracy improvement and quality assessment, *Remote Sens.*
21 *Environ.*, 112(10), 3806–3819, doi:10.1016/j.rse.2008.05.018, 2008.
- 22 Scherler, D., Bookhagen, B. and Strecker, M. R.: Hillslope-glacier coupling: The interplay of
23 topography and glacial dynamics in High Asia, *J. Geophys. Res.*, 116(F02019), 1–21,
24 doi:10.1029/2010JF001751, 2011a.
- 25 Scherler, D., Bookhagen, B. and Strecker, M. R.: Spatially variable response of Himalayan
26 glaciers to climate change affected by debris cover, *Nat. Geosci.*, 4(3), 156–159,
27 doi:10.1038/ngeo1068, 2011b.
- 28 Schwitter, M. and Raymond, C.: Changes in the longitudinal profiles of glaciers during
29 advance and retreat, *J. Glaciol.*, 39(133), 582–590, 1993.
- 30 Shrestha, A., Wake, C., Mayewski, P. and Dibb, J.: Maximum temperature trends in the
31 Himalaya and its vicinity: An analysis based on temperature records from Nepal for the
32 period 1971–94, *J. Clim.*, 12, 2775–2786, doi:10.1175/1520-
33 0442(1999)012<2775:MTTITH>2.0.CO;2, 1999.
- 34 Shrestha, A., Wake, C., Dibb, J. and Mayewski, P.: Precipitation fluctuations in the Nepal
35 Himalaya and its vicinity and relationship with some large scale climatological parameters,
36 *Int. J. Climatol.*, 20(3), 317–327, doi:10.1002/(SICI)1097-0088(20000315)20:3<317::AID-
37 JOC476>3.0.CO;2-G, 2000.

- 1 Singh, P., Kumar, V., Thomas, T. and Arora, M.: Changes in rainfall and relative humidity in
2 river basins in northwest and central India, *Hydrol. Process.*, 22(16), 2982–2992,
3 doi:10.1002/hyp.6871, 2008.
- 4 Steiner, J. F., Pellicciotti, F., Buri, P., Miles, E. S., Immerzeel, W. W. and Reid, T. D.:
5 Modelling ice-cliff backwasting on a debris-covered glacier in the Nepalese Himalaya, *J.*
6 *Glaciol.*, 61(229), 889–907, doi:10.3189/2015JoG14J194, 2015.
- 7 Steiner, J. F., Buri, P., Miles, E. S., Ragetti, S. and Pellicciotti, F.: Life and death of ice cliffs
8 and lakes on debris covered glaciers - insights from a new dataset, *Geophys. Res. Abstr.*,
9 18(EGU2016-13922), 2016.
- 10 Sugiyama, S., Fukui, K., Fujita, K., Tone, K. and Yamaguchi, S.: Changes in ice thickness
11 and flow velocity of Yala Glacier, Langtang Himal, Nepal, from 1982 to 2009, *Ann. Glaciol.*,
12 54(64), 157–162, doi:10.3189/2013AoG64A111, 2013.
- 13 Surazakov, A. and Aizen, V.: Positional Accuracy Evaluation of Declassified Hexagon KH-9
14 Mapping Camera Imagery, *Photogramm. Eng. Remote Sens.*, 76(5), 603–608,
15 doi:10.14358/PERS.76.5.603, 2010.
- 16 Tadono, T. and Shimada, M.: Calibration of PRISM and AVNIR-2 onboard ALOS “Daichi,”
17 *IEEE Trans. Geosci. Remote Sens. Sens.*, 47(12), 4042–4050,
18 doi:10.1109/TGRS.2009.2025270, 2009.
- 19 Thompson, S., Benn, D. I., Mertes, J. and Luckman, A.: Stagnation and mass loss on a
20 Himalayan debris-covered glacier: processes, patterns and rates, *J. Glaciol.*, 1–19,
21 doi:10.1017/jog.2016.37, 2016.
- 22 Tiwari, P., Pande, H., Punia, M. and Dadhwal, V. K.: Cartosat-I: Evaluating mapping
23 capabilities, *Int. J. Geoinformatics*, 4(1), 51–56 [online] Available from:
24 <http://creativecommons.gssc.osaka-cu.ac.jp/IJG/article/view/609> (Accessed 26 January 2016),
25 2008.
- 26 Wang, D. and Käab, A.: Modeling Glacier Elevation Change from DEM Time Series, *Remote*
27 *Sens.*, 7(8), 10117–10142, doi:10.3390/rs70810117, 2015.
- 28 Yao, T., Thompson, L., Yang, W., Yu, W., Gao, Y., Guo, X., Yang, X., Duan, K., Zhao, H.,
29 Xu, B., Pu, J., Lu, A., Xiang, Y., Kattel, D. B. and Joswiak, D.: Different glacier status with
30 atmospheric circulations in Tibetan Plateau and surroundings, *Nat. Clim. Chang.*, 2(9), 663–
31 667, doi:10.1038/nclimate1580, 2012.
- 32 Ye, Q., Bolch, T., Naruse, R., Wang, Y., Zong, J., Wang, Z., Zhao, R., Yang, D. and Kang,
33 S.: Glacier mass changes in Rongbuk catchment on Mt. Qomolangma from 1974 to 2006
34 based on topographic maps and ALOS PRISM data, *J. Hydrol.*, 530, 273–280,
35 doi:10.1016/j.jhydrol.2015.09.014, 2015.

36

1 Figures and Tables

2 Table 1. Characteristics of the studied glaciers in the upper Langtang catchment. The
 3 measures are based on the SRTM 1 Arc-Second Global DEM and glacier outlines of 2006.

Name	Area km ²	Debris cover km ²	Mean slope %	Mean slope glacier tongue* %	AAR**	Elevation range m a.s.l.
1 Langtang	46.5	15.5	17.1	7.2	52%	4479-6615
2 Langshisha	16.3	4.5	17.7	7.5	55%	4415-6771
3 Shalbachum	10.2	2.6	16.9	9.1	52%	4231-6458
4 Lirung	6.5	1.1	34.0	9.9	49%	4044-7120
5 Ghanna	1.4	0.7	20.9	15.5	15%	4721-5881
6 Kimoshung	4.4	-	24.4	32.1	86%	4385-6648
7 Yala	1.9	-	22.7	20.3	40%	5122-5676

*Here we consider the debris-covered area for glaciers with debris-covered tongues and all glacier area below 5400 m a.s.l. for debris-free glaciers.

**Assuming an equilibrium line altitude of 5400 m a.s.l. (Sugiyama et al., 2013; Ragettli et al., 2015)

4 Table 2. Remote-sensing data used

Sensor	Date of acquisition	Stereo mode (b/h-ratio)	Spatial/radiometric Resolution	Role
Hexagon KH-9	23 Nov 1974	Stereo (0.4)	6-9m/8-bits	DEM differencing, glacier outlines
Cartosat-1	15 Oct 2006	Stereo (0.62)	2.5m/10-bits	DEM differencing, glacier outlines
Cartosat-1	9 Nov 2009	Stereo (0.62)	2.5m/10-bits	DEM differencing, velocities, glacier outlines
ALOS PRISM	3 Dec 2010	Tri-stereo (0.5)	2.5m/8-bits	DEM differencing, velocities, glacier outlines
SPOT6	21 Apr 2014	Tri-stereo (0.5)	1.5m/12-bits	DEM differencing, glacier outlines
WorldView-2	2 Feb 2015	Stereo (0.5)	0.46m/11-bits	DEM differencing
WorldView-3	22 Feb 2015	Stereo (0.5)	0.31m/11-bits	DEM differencing
SPOT7	7 Mai 2015	Tri-stereo (0.64)	1.5m/12-bits	DEM differencing, glacier outlines
SPOT7	6 Oct 2015	Tri-stereo (0.68)	1.5m/12-bits	DEM differencing
Pléiades	1 and 9 Nov 2014	Across track stereo (0.4)	0.5m/12-bits	Basis for georectification

5

1 Table 3. Mean* uncertainties associated to different sets of elevation change ($\Delta h/\Delta t$) maps.

	No. of maps in category	All glacier area	Debris-covered glacier area
All $\Delta h/\Delta t$ maps, $\Delta t < 4$ a	9	1.18 m a ⁻¹	0.47 m a ⁻¹
All $\Delta h/\Delta t$ maps, $4 \text{ a} \leq \Delta t < 10$ a	12	0.29 m a ⁻¹	0.12 m a ⁻¹
All $\Delta h/\Delta t$ maps involving Hexagon 1974 DEM	7	0.07 m a ⁻¹	0.03 m a ⁻¹
DEM involved, $4 \text{ a} \leq \Delta t < 10$ a			
Cartosat-1 Oct 2006	5	0.22 m a ⁻¹	0.09 m a ⁻¹
Cartosat-1 Nov 2009	4	0.29 m a ⁻¹	0.11 m a ⁻¹
ALOS-PRISM Dec 2010	4	0.43 m a ⁻¹	0.19 m a ⁻¹
SPOT6 April 2014	2	0.24 m a ⁻¹	0.09 m a ⁻¹
WorldView Feb 2015	3	0.32 m a ⁻¹	0.14 m a ⁻¹
SPOT7 May 2015	3	0.31 m a ⁻¹	0.12 m a ⁻¹
SPOT7 October 2015	3	0.24 m a ⁻¹	0.10 m a ⁻¹

* Uncertainties associated to individual maps are shown in Table S1

2 Table 4. Glacier volume and mass changes 1974-2006, ensemble mean 2006-2015*.

	Average elevation differences (m a ⁻¹)		Average mass balance (m w.e. a ⁻¹)	
	Nov1974-Oct2006	Ensemble mean* 2006-2015	Nov1974-Oct2006	Ensemble mean* 2006-2015
Glaciers				
Langtang	-0.28 ± 0.08	-0.55 ± 0.13	-0.24 ± 0.08	-0.47 ± 0.13
Langshisha	-0.12 ± 0.09	-0.45 ± 0.19	-0.10 ± 0.08	-0.38 ± 0.18
Shalbachum	-0.43 ± 0.08	-0.53 ± 0.19	-0.36 ± 0.09	-0.45 ± 0.18
Lirung	-0.17 ± 0.13	-0.22 ± 0.16	-0.14 ± 0.11	-0.19 ± 0.14
Ghanna	-0.51 ± 0.05	-0.46 ± 0.43	-0.43 ± 0.07	-0.39 ± 0.36
Kimoshung	0.07 ± 0.13	-0.02 ± 0.17	0.05 ± 0.10	-0.02 ± 0.13
Yala	-0.33 ± 0.06	-0.89 ± 0.23	-0.28 ± 0.07	-0.76 ± 0.24
Average	-0.24 ± 0.08	-0.45 ± 0.18	-0.21 ± 0.08	-0.38 ± 0.17
Debris-covered areas				
Langtang	-0.79 ± 0.03	-0.91 ± 0.05	-0.67 ± 0.07	-0.78 ± 0.10
Langshisha	-0.69 ± 0.03	-1.16 ± 0.23	-0.58 ± 0.07	-0.98 ± 0.25
Shalbachum	-0.78 ± 0.04	-1.30 ± 0.20	-0.66 ± 0.08	-1.10 ± 0.23
Lirung	-1.03 ± 0.05	-1.67 ± 0.59	-0.87 ± 0.10	-1.42 ± 0.56
Ghanna	-0.58 ± 0.03	-0.50 ± 0.20	-0.49 ± 0.06	-0.43 ± 0.19
Average	-0.78 ± 0.03	-1.02 ± 0.18	-0.66 ± 0.07	-0.87 ± 0.20

*Average of 6 overlapping periods between Oct 2006 and Oct 2015 (Figure S4)

3

4

1 Table 5. Sensitivity to outlier correction and Equilibrium Line Altitude (ELA) definitions. $\Delta_{2\sigma}$
 2 is the difference in results if a 2σ -level is used to define outliers at all area types, instead of a
 3 3σ -level above and a 1σ -level below the ELA. The estimated ELA is 5400 m a.s.l. (Sugiyama
 4 et al., 2013; Ragetti et al., 2015). $\Delta_{\text{ELA} \pm 100\text{m}}$ represents the differences in results obtained with
 5 an ELA at 5500 m a.s.l. in comparison to results obtained with an ELA at 5300 m a.s.l.

Name	AAR		Elevation differences (m a^{-1})			
			Nov1974-Oct2006		Ensemble mean 2006-2015	
			ELA -100 m	ELA +100 m	$\Delta_{\text{ELA} \pm 100\text{m}}$	$\Delta_{2\sigma}$
1 Langtang	61%	43%	0.02	-0.01	-0.01	0.01
2 Langshisha	60%	45%	0.06	-0.09	0.02	-0.02
3 Shalbachum	60%	37%	-0.23	0.08	0.04	-0.02
4 Lirung	52%	46%	-0.01	0.01	0.05	-0.01
5 Ghanna	20%	12%	-0.01	0.02	0.00	0.04
6 Kimoshung	88%	80%	0.07	0.06	0.05	-0.16
7 Yala	70%	13%	-0.13	-0.04	-0.11	0.02
All Glacier Area	61%	44%	0.00	-0.01	0.02	-0.01

Sensitivity values that exceed uncertainty ranges as indicated in Table 4 are printed in bold letters.

6

7 Table 6. Glacier area changes over the periods 1974-2006 and 2006-2015.

ID	Glacier name	1974-2006		2006-2015	
		km^2	$\% \text{ a}^{-1}$	km^2	$\% \text{ a}^{-1}$
1	Langtang	-2.65 ± 0.03	-0.17 ± 0.01	-0.45 ± 0.07	-0.11 ± 0.02
2	Langshisha	-0.48 ± 0.09	-0.09 ± 0.02	-0.13 ± 0.05	-0.09 ± 0.04
3	Shalbachum	-0.28 ± 0.06	-0.08 ± 0.02	-0.03 ± 0.04	-0.04 ± 0.04
4	Lirung	-0.45 ± 0.08	-0.20 ± 0.03	-0.05 ± 0.05	-0.08 ± 0.08
5	Ghanna	-0.16 ± 0.03	-0.33 ± 0.05	-0.05 ± 0.01	-0.40 ± 0.12
6	Kimoshung	-0.11 ± 0.01	-0.08 ± 0.01	-0.02 ± 0.01	-0.05 ± 0.02
7	Yala	-0.31 ± 0.03	-0.43 ± 0.05	-0.31 ± 0.03	-1.77 ± 0.16

8

9

1 Table 7. Characteristics of the debris-covered tongues (debris-covered glacier area excluding
 2 tributary branches).

	Cliff Area	Lake Area	Mean velocity	% stagnant
Langtang	10.0%	3.3%	5.9 m a ⁻¹	31%
Langshisha	10.5%	2.3%	7.0 m a ⁻¹	20%
Shalbachum	10.3%	2.6%	5.4 m a ⁻¹	29%
Lirung	8.0%	2.3%	2.8 m a ⁻¹	48%
Ghanna	3.2%	0.4%	1.6 m a ⁻¹	85%


Cliff and lake area corresponds to the percentage of 30-m pixels containing cliffs/lakes (median of 6 available cliff and lake maps from the period 2006-2015). Mean velocity is calculated on the basis of 2009-2010 surface velocities (Figure 13). To discriminate moving ice from quasi-stagnant ice we use a threshold of 2.5 m a⁻¹ (cf. Scherler et al., 2011b), which also corresponds to the approximate uncertainty of remote-sensing derived surface velocity.

3 Table 8. Elevation changes of debris-covered glacier tongues due to avalanches triggered by
 4 the Nepal earthquake on 25 April 2015. The first three data columns provide the volume
 5 changes of avalanche affected area ($\Delta h > 5$ m in May 2015) divided by the total debris-cover
 6 area (Table 1).

	21 Apr 2014- 25 Apr 2015* (m)	25 Apr 2015- 7 May 2015 (m)	25 Apr 2015- 6 Oct 2015 (m)	6 Oct 2015, volume remaining (%)
Langtang**	-0.10 ±0.05	1.33 ±0.42	0.42 ±0.20	31.3%
Langshisha	-0.04 ±0.05	0.32 ±0.37	0.10 ±0.19	31.6%
Shalbachum	-0.11 ±0.05	0.74 ±0.35	0.31 ±0.20	42.5%
Lirung	-0.87 ±0.06	6.79 ±0.38	3.87 ±0.23	57.0%
Average	-0.13 ±0.05	1.31 ±0.35	0.52 ±0.19	39.5%

*Estimation based on average annual melt Oct 2006 – Apr 2014


**Only lower part (south of 28°19'N), upper part not on April 2014 scene



1


2 Figure 1. Map of the upper Langtang catchment. The numbers on the map correspond to the
 3 glaciers listed in Table 1. Monsoon snow-cover frequency is based on Landsat 1999 to 2013
 4 land cover classifications (Miles et al., 2016b). 1974 glacier area (dotted lines) is shown for
 5 the seven studied glaciers only.

6




1
 2 Figure 2. a) Uncertainty estimates of average elevation change rates ($\Delta h/\Delta t$) per individual
 3 glacier and per debris-covered tongue. The central mark is the median of the ensemble ($\Delta h/\Delta t$
 4 maps that are rejected according to Section 3.2.5 are excluded). b) Ensemble of stereo
 5 matching scores per individual glacier and debris-covered tongue and c) per glacier area
 6 above and below the estimated ELA. The central marks correspond to the median of all
 7 DEMs (except Hexagon 1974 and WorldView 2015 DEMs for which matching scores are not
 8 available). The edges of each box are the 25th and 75th percentiles. The whiskers extend to
 9 the most extreme data points.


10




1
 2 Figure 3. Off-glacier elevation change error ($\Delta h/\Delta t$) per 50-m elevation band. The black line
 3 represents the median error in the ensemble of $\Delta h/\Delta t$ maps (excluding $\Delta h/\Delta t$ maps that are
 4 rejected according to Section 3.2.5). Error bars represent 95% confidence intervals. The color
 5 bars represent hypsometries of glacier area, off-glacier area and debris-covered glacier areas,
 6 respectively.



7
 8 Figure 4. Elevation change rates ($\Delta h/\Delta t$) derived from a) Hexagon Nov 1974 and Cartosat-1
 9 Oct 2006 DEMs and (b) Cartosat-1 Oct 2006 and WorldView Feb 2015 DEMs.




1
 2 Figure 5. Uncertainties in elevation change rates ($\Delta h/\Delta t$) in function of the time interval
 3 between DEMs (Δt). Median results of all available 28 $\Delta h/\Delta t$ maps. Error bars extend to the
 4 most extreme data points.




5
 6 Figure 6. Supraglacial cliffs and lakes as identified from the Oct 2006 Cartosat-1 satellite
 7 image: a) Langtang and Ghanna Glaciers, b) Shalbachum Glacier, c) Lirung Glacier, d)
 8 Langshisha Glacier. Cliff area shows the median fraction (%) of 30-m pixels per 50m
 9 elevation band that contain cliffs, considering all 6 available cliff maps from 2006-2015.
 10

1



2

3 Figure 7. Mean elevation change rates ($\Delta h/\Delta t$) per period and glacier. For better readability,
4 only the maximum width of error bounds corresponding to individual periods 2006-2015 are
5 shown.



6

7 Figure 8. Mean elevation change rates ($\Delta h/\Delta t$) per period and debris-covered glacier area. For
8 better readability, only the maximum width of error bounds corresponding to individual
9 periods 2006-2015 are shown.






Figure 9. Altitudinal distribution of mean annual elevation change ($\Delta h/\Delta t$) over 50 m elevation bands of debris-covered tongues (debris-covered area of each glacier excluding tributary branches). Uncertainty bounds correspond to uncertainty in function of elevation derived for each $\Delta h/\Delta t$ map individually (Figure 3).




1
 2 Figure 10. Altitudinal distribution of mean annual elevation change ($\Delta h/\Delta t$) and altitudinal
 3 distribution of glacier area (%) over 50 m elevation bands of selected glaciers. Uncertainty
 4 bounds correspond to uncertainty in function of elevation derived for each $\Delta h/\Delta t$ map
 5 individually (Figure 3). Ensemble median values shown here are used to replace missing data
 6 in the accumulation areas of glaciers after outlier exclusion (Section 3.2.3). Note that the
 7 x-axis ranges are different for each sub-figure.

8




1
2 Figure 11: Altitudinal distribution of cliff and lake area fractions, glacier velocity and changes
3 in thinning rates. Cliff and lake area is shown as % of 30-m pixels containing cliffs/lakes per
4 50 m elevation band, whereas the values represent the median of 6 available cliff and lake
5 maps from the period 2006-2015. Glacier velocities (m/a) represent the median per 50 m
6 elevation band of data shown in Figure 11 and error bars represent the standard deviation in
7 pixel values per elevation band. Changes in thinning rates ($\Delta(\Delta h/\Delta t)$ [m/a]) are calculated
8 comparing 1974-2006 and the 2006-2015 ensemble-mean. Negative $\Delta(\Delta h/\Delta t)$ values
9 represent thinning accelerations. Error bars represent the maximum variations in $\Delta(\Delta h/\Delta t)$
10 considering all individual periods within the 2006-2015 ensemble.



1
 2
 3
 4
 5

Figure 12. Avalanche affected sections of Lirung and Langtang glacier, pre- and after the earthquake on 25 April 2015, and corresponding surface elevation changes (Δh). Imagery ©Airbus DS 2014/2015.



1

2 Figure 13: Surface velocities 2009-2010 cropped to catchment boundaries. Values have units
 3 of meters per year and are derived by cross-correlation feature tracking. Off-glacier velocities
 4 are shown in transparent color.

5