

# GPS-derived estimates of surface mass balance and ocean-induced basal melt for Pine Island Glacier ice shelf, Antarctica

David E. Shean<sup>1,2</sup>, Knut Christianson<sup>2</sup>, Kristine M. Larson<sup>3</sup>, Stefan R.M. Ligtenberg<sup>4</sup>, Ian R. Joughin<sup>1</sup>, Ben E. Smith<sup>1</sup>, C. Max Stevens<sup>2</sup>, [M. Bushuk<sup>5</sup>](#), [D.M. Holland<sup>6,7</sup>](#)

<sup>1</sup>Applied Physics Laboratory Polar Science Center, University of Washington, Seattle, WA, USA

<sup>2</sup>Department of Earth and Space Sciences, University of Washington, Seattle, WA, USA

<sup>3</sup>Department of Aerospace Engineering Sciences, University of Colorado, Boulder, CO, USA

<sup>4</sup>Institute for Marine and Atmospheric research Utrecht, Utrecht University, Netherlands

<sup>5</sup>[Geophysical Fluid Dynamics Laboratory, Princeton University, Princeton, NJ, USA](#)

<sup>6</sup>[Courant Institute of Mathematical Sciences, New York University, New York, NY, USA](#)

<sup>7</sup>[Center for Global Sea-Level Change, New York University, Abu Dhabi, UAE](#)

Correspondence to: David Shean (dshean@uw.edu)

## Abstract

In the last two decades, Pine Island Glacier (PIG) experienced marked speedup, thinning, and grounding-line retreat, likely due to marine ice-sheet instability, and ice-shelf basal melt. To better understand these processes, we combined 2008–2010 and 2012–2014 GPS records with dynamic firm model output to constrain local surface and basal mass balance for PIG. We used GPS interferometric reflectometry to precisely measure absolute surface elevation ( $z_{surf}$ ) and Lagrangian surface elevation change ( $Dz_{surf}/Dt$ ). Observed surface elevation relative to a firm layer tracer for the initial surface ( $z_{surf} - z_{surf}^0$ ) is consistent with model estimates of surface mass balance (SMB, primarily snow accumulation) and downward velocity due to firm compaction. A relatively abrupt  $\sim 0.2$ – $0.3$  m surface elevation decrease, likely due to surface melt and increased compaction rates, is observed during a period of warm atmospheric temperatures from December 2012 to January 2013. Observed  $Dz_{surf}/Dt$  trends ( $-1$  to  $-4$  m/yr) for the PIG shelf sites are all highly linear. Corresponding basal melt rate estimates range from  $\sim 10$  to  $40$  m/yr, in good agreement with those derived from ice-bottom acoustic ranging, phase-sensitive ice-penetrating radar, and high-resolution stereo DEM records. The GPS and DEM records document higher melt rates within and near features associated with longitudinal extension (i.e., transverse surface depressions, rifts). Basal melt rates for the 2012–2014 period show limited temporal variability, despite large changes in ocean temperature recorded by moorings in Pine Island Bay. Our results demonstrate the value of long-term GPS records and interferometric reflectometry for ice-shelf mass balance studies, with implications for the sensitivity of ice-ocean interaction at PIG.

## 1 Introduction

The widespread availability of precise Global Positioning System (GPS) measurements has revolutionized the study of ice dynamics and glacier mass balance (e.g., Gao and Liu, 2001). Continuously operating dual-frequency GPS receivers provide high-frequency (1 Hz or less), highly accurate ( $< 1$ – $3$  cm) measurements of position, which can be used to derive surface velocity and elevation change. For applications involving ice dynamics, these measurements

Deleted: *In-situ*

Deleted: records

Deleted: E.

Deleted: ice-shelf basal melt and

Deleted: .

Deleted: analyzed

Deleted: *in-situ*

Deleted: for PIG

Deleted: , firm compaction, and basal melt.

Deleted: computed time series of horizontal velocity, strain rate, antenna height,

Deleted: .

Deleted: Dh

Deleted: The antenna height time series show a

Deleted: increase of  $\sim 0.7$ – $1.0$  m/yr, which

Deleted: for

Deleted: ) of  $\sim 0.7$ – $0.9$  mwe/yr

Deleted:  $\sim 0.7$ – $0.8$  m/yr

Deleted: An

Deleted: Dh

Deleted: all

Deleted: is

Deleted: , with trends of  $-1$  to  $-4$  m/yr and residuals of  $< 0.4$  m

Deleted: significant changes in ocean heat content, suggesting that sub-shelf melt rates may be less sensitive to ocean heat content than previously reported, at least for these locations and time periods.

Deleted:

offer important constraints for the mass continuity equation, which equates surface elevation change with ice flux divergence, surface mass balance, and basal mass balance. Here, we explore a methodology to constrain each of these components directly from GPS observables.

**Deleted:** , and ice flux divergence.

Surface mass balance (SMB) processes include precipitation, sublimation, wind redistribution of surface snow, and melt water runoff. Regional climate models forced by reanalysis output now provide daily estimates of Antarctic SMB on a relatively coarse grid (~5.5 to 27 km). *In-situ* SMB measurements are, however, still essential for model calibration and validation. Traditionally, SMB is measured using stake networks, automated weather stations, near-surface radar surveys, and firn/ice cores, all of which require substantial field operations in remote locations. These measurements also tend to bias model calibration towards accessible locations, and recent studies indicate that these biases can significantly affect mass balance results, often resulting in overestimates of cumulative balance due to poor sampling in dynamic areas (Andreassen et al., 2016).

**Deleted:** (e.g., RACMO, MAR)

**Deleted:** to monthly

**Deleted:** can be

**Deleted:** locations that are either

**Deleted:** or of high scientific interest for other reasons. Reanalysis of glacier mass balance records

**Deleted:** measurement

**Deleted:** , as

**Deleted:** are less well sampled

Antarctic firn/ice core records indicate that SMB variability over most of Antarctica during the last 800 years was statistically insignificant, but that accumulation increased more than 10% for high-accumulation coastal regions (e.g., the Amundsen Sea Embayment) since the 1960s (Frezzotti et al., 2013). Historically, these areas have been poorly sampled with traditional methods, providing limited data available for validation of modeled SMB.

**Deleted:** These

**Deleted:** are precisely those that that are most

**Deleted:** for SMB reanalysis calibration and validation using

**Deleted:** (i.e., firn/ice cores).

Accurate knowledge of firn compaction and its spatiotemporal variability is essential for interpreting observed surface elevation change in remote sensing data (e.g., satellite altimetry), and for partitioning this change into components related to ice dynamics and SMB (e.g., Shepherd et al., 2012; Wouters et al., 2015). Depth-dependent compaction rates can be estimated from a number of different methods, including vertical strain measurements (Arthern et al., 2010; Hamilton and Whillans, 1998), borehole optical stratigraphy (Hawley and Waddington, 2011), repeat phase-sensitive radio-echo sounding (pRES) measurements (e.g., Jenkins et al., 2006) and ice-penetrating radar observations of internal layers over time (e.g., Medley et al., 2014, 2015). In the absence of these measurements, dynamic firn models forced by modeled SMB can provide estimates of compaction rates throughout the firn column, which can be integrated to obtain estimates for the contribution of firn compaction to surface elevation change over time (e.g., Ligtenberg et al., 2011).

**Deleted:** *in-situ*

**Deleted:** modelled

Basal mass balance (BMB) for ice shelves (i.e., bottom melting, accretion) is driven by complex ice-ocean interaction. State-of-the-art ice-shelf cavity ocean circulation models offer some insight into sub-shelf ice-ocean interaction, but these models lack validation, as *in-situ* hydrographic observations are limited, especially within the sub-shelf cavity and the ice-ocean boundary layer. Some direct measurements are available from autonomous submersibles (e.g., Dutrieux et al., 2014) and instrumentation deployed through ice-shelf boreholes (e.g., Stanton et al., 2013), but available data are limited to short time periods and small spatial extents. Precise measurements of surface elevation change from remote sensing observations (e.g., laser altimetry, digital elevation models (DEMs)) can also be used to infer BMB (e.g., Dutrieux et al., 2013; Moholdt et al., 2014; Pritchard et al., 2012; Shean, 2016), but temporal resolution is limited, as time intervals between repeat observations are typically several months to years.

**Deleted:** beneath

**Deleted:** packages

**Deleted:** (e.g., Dutrieux et al., 2013; Pritchard et al., 2012; Shean, 2016), but

**Deleted:** *in-situ*

**Deleted:** ice shelf

**Deleted:** firn compaction,

**Deleted:** ,

**Deleted:** and antenna height

**Deleted:** and

**Deleted:** , and then use firn model results to estimate the time-variable downward velocity of the GPS poles.

Here, we use temporally dense GPS records from the Pine Island Glacier Ice Shelf to constrain local SMB, flux divergence and BMB. We use changes in observed GPS antenna elevation and reflectometry-derived surface elevation to validate SMB/firn model output. Flux divergence is estimated from observed strain rates between GPS stations.

These estimates are then used to isolate elevation change due to local BMB. This approach yields temporally dense records of basal melt rates at spatially sparse GPS locations, which are combined with high-resolution DEMs from the same time period to provide spatial context. These complementary results for the PIG ice shelf provide new information about the time-variable magnitude and spatial distribution of basal melting, offering indirect observations of ice-ocean interaction and BMB sensitivity to ocean heat content variability, with implications for other rapidly evolving "warm-cavity" Antarctic ice shelves.

**Deleted:** constraints

**Deleted:** estimates

**Deleted:** for the observed elevation change.

### 1.1 PIG background

Pine Island Glacier is one of the largest and most dynamic ice streams in West Antarctica. Since the 1970s, PIG has experienced ~30 km of grounding line retreat along its centerline (Rignot et al., 2014) (~8 km average retreat across full width of fast-flowing trunk (Joughin et al., 2016)), a ~75% increase in surface velocity (Mouginot et al., 2014), and >100 m of thinning (Bindschadler, 2002; Pritchard et al., 2009), with accelerated retreat beginning in the 1990s. These changes have been attributed to some combination of geometric instability (i.e., marine ice sheet instability) and external forcing (i.e., increased ocean heat content and/or changes in sub-shelf circulation) (Jacobs et al., 2011; Joughin et al., 2010).

**Deleted:** due to increased ocean heat content, circulation, and basal melt (Jacobs et al., 2011). Present-day surface velocities are ~4 km/yr, with annual discharge of ~130 Gt/yr

Present-day surface velocities are ~4 km/yr, with annual discharge of ~130 Gt (Medley et al., 2014; Mouginot et al., 2014) and net mass loss estimates of 40 to 50 Gt/yr (Medley et al., 2014; Rignot, 2008). This mass loss is responsible for ~0.11 mm/yr global sea level rise (SLR), or approximately 40–45% of the total ~0.26 mm/yr Antarctic SLR contribution (Church et al., 2013; Rietbroek et al., 2016; Shepherd et al., 2012).

**Deleted:** (Church et al., 2013; Rietbroek et al., 2016).

Figure 1 shows the fast-flowing portion of the PIG ice stream, which terminates in a large ice shelf ("main shelf") that is ~25 km wide, ~100 km long, and ~1–1.5 km thick across the grounding line. Basal melting accounts for ~70–80% of mass loss from the ice shelf, with estimated 2003-2008 melt rates of ~95–101 Gt/yr (Depoorter et al., 2013; Rignot et al., 2013) and 2008-2015 melt rates of ~80-90 Gt/yr (Shean, 2016).

**Deleted:** (Shean, 2016)

The main shelf has complex surface topography, including km-scale surface ridges and troughs that correspond to basal keels and channels, respectively (Bindschadler et al., 2011; Vaughan et al., 2012). A series of longitudinal (along-flow) ridges and troughs are present along the shelf centerline, with transverse (across-flow) ridges and troughs along the lateral margins (Figure 1). Local basal melt rates vary considerably across these features (Dutrieux et al., 2013; Shean, 2016).

**Deleted:** (Dutrieux et al., 2013; Shean, 2016)

Hydrographic observations seaward of the PIG calving front in Pine Island Bay suggest that basal melting declined by ~50% between 2010 and 2012 (Dutrieux et al., 2014). Long-term 2009–2015 mooring records seaward of the southern calving front (Figure 1) show a significant decrease in ocean temperature (~1–1.5°C) over ~450–770 m depths from late 2011 to early 2012, and then again from mid-2012 to early 2013 (Christianson et al., 2016; Webber et al., 2017). These observations show that the ocean heat content at the PIG ice-shelf front varies considerably over monthly to interannual timescales.

**Deleted:** (Christianson et al., 2016).

## 1.2 PIG GPS sites

Several long-term GPS stations were installed on the PIG shelf as part of a larger investigation of ice-sheet, ice-shelf, and ocean dynamics (Bindschadler et al., 2011; Stanton et al., 2013). During the early part of this effort, two GPS stations continuously collected data from January 2008 to January 2010: one on the southern PIG ice shelf (PIG2) and another on the fast-flowing, grounded ice upstream of the grounding line (PIG1) (Figure 1). In addition, a ~2x2 km array of five stations (SOW1–4, BOAR, Figure 2) was installed ~50 km downstream of the grounding line, near the center of the main shelf from January 2012 to late December 2013.

The stations used dual-frequency Trimble NetRS GPS receivers (2008–2010 sites) and NetR9 receivers (2012–2014 sites), with Trimble Zephyr Geodetic 2 antennas mounted on 12-foot (3.66 m) poles with insulating pole-base stoppers. The poles were driven into the snow by hand, with initial pole bases set ~0.5–1.0 m beneath the surface (M. Truffer, personal communication, 2016).

High-resolution optical imagery and DEM data (see Section 2.5) over the 2012–2014 sites show that SOW1, BOAR, and SOW3 were installed with along-flow orientation in a longitudinal surface trough (Figure 2) that overlies a longitudinal basal channel. An ice-penetrating radar profile with transverse orientation was collected upstream of the GPS array, providing ice thickness estimates of ~450–460 m near the apex of a longitudinal channel and ~540 m over adjacent keels (Stanton et al., 2013). Figure 2 shows estimated ice thickness for longitudinal and transverse profiles across the GPS array.

A borehole was drilled through the ice shelf approximately 1.34 km upstream of SOW1 (K. Riverman, personal communication, 2016), and an instrument package with an upward-facing ice-bottom altimeter (acoustic ranger) was deployed beneath the shelf from January to February 2012. Measurements from this bottom altimeter and complementary pRES experiments provided basal melt rate estimates of ~14–25 m/yr within the longitudinal channel (Christianson et al., 2016; Stanton et al., 2013).

The 2012–2014 GPS array was located near several transverse surface depressions (Figure 2), which are likely associated with transverse basal channels and/or rifts. Local surface slopes were ~0.6–0.9° within the largest of these depressions, immediately downstream of SOW3 and SOW4. A notable linear surface depression located approximately 1 km upstream of SOW1 (black arrow in Figure 2) opened as a rift in ~2014 (R1 in Jeong et al. (2016)), and was subsequently the site of a large iceberg calving event that occurred around July 2015. The fortuitous placement of the 2012–2014 GPS array near these features complicates interpretation of GPS records, but also provides new constraints on the spatiotemporal (co)evolution of strain rates and rift formation for the PIG shelf.

## 2 Data and Methods

### 2.1 GPS antenna position

As described in Christianson et al. (2016), GPS data were processed using differential-carrier-phase positioning relative to bedrock GPS sites (Backer Island [BACK, -74.26°N, -102.28°E] for 2012–2014 records; Howard Nunatak [HOWN, -77.31°N, -8.65°E] for 2008–2010 records) with epoch-by-epoch zenith tropospheric delay estimation. Daily-averaged positions of these base stations were calculated using GAMIT and stabilized relative to a fixed circum-

Deleted: involving

Deleted: and

Deleted: 45

Deleted: on

Deleted: [

Deleted: ]

Deleted: section

Deleted: 3.6

Deleted: oriented approximately

Deleted: a flowline within

Deleted: Transverse

Deleted: profiles across this channel provide

Deleted: hydrostatic

Deleted: ,

Deleted: )

Deleted: longitudinal

Deleted: /velocity processing

Antarctic reference frame using a Kalman filter (GLOBK, (Herring et al., 2015)). Antenna positions relative to the WGS84 ellipsoid were calculated every 30 seconds. We analyzed a subset of these positions sampled at 10-minute intervals, and removed any positions with uncertainty >8 cm. The BOAR record was curtailed on April 29, 2013 (1.31 year duration), when an abrupt ~2.0 m elevation decrease and corresponding horizontal offset occurred, suggesting that the pole fell over.

We estimate initial antenna position accuracy of ~1 cm. Positions were converted to a local Cartesian horizontal coordinate system with final antenna elevation values ( $z_{ant}$ ) as orthometric height above the EGM2008 geoid (Pavlis et al., 2012). Absolute geoid errors are poorly constrained for coastal Antarctica, but relative geoid error along the ~8 km GPS paths should be <-1.2 cm. A constant offset of 3.71 m (3.66 m pole length + 0.053 m phase center to bottom of antenna) was removed from antenna elevation ( $z_{ant}$ ) to estimate corresponding pole-base elevation.

We estimated vertical tidal displacement for all GPS positions on the PIG ice shelf using the CATS2008A model (Padman et al., 2002). We used mean sea level pressure values from the 0.75°-grid-cell ERA-Interim reanalysis products (Dee et al., 2011) to correct for vertical displacement due to the inverse barometer effect (IBE, e.g., Padman et al., 2003). To do this, we removed the 2002–2016 median (985.21 hPa) from 6-hour sea level pressure and scaled the residuals by ~1 cm/hPa. Figure 3 shows that tidal amplitudes for the GPS sites range from approximately -0.9 to +1.3 m and IBE amplitudes range from -0.3 to +0.3 m. These signals were removed from the GPS antenna elevation ( $z_{ant}$ ), and residual high-frequency noise was removed with a low-pass filter (1.5-day cutoff), yielding smoothed time series for further analysis (Figure 3). We conservatively estimate final  $z_{ant}$  absolute accuracy of ~0.1 m for the 10-minute GPS positions.

## 2.2 Antenna-surface distance

The GPS interferometric reflectometry method provides a precise measurement of antenna phase-center height above the reflecting surface (Larson, 2016). The reflecting surface for PIG is the interface between the atmosphere and the snow/firn surface, and we define the antenna height above this interface as the “antenna-surface distance” ( $h_{ant-surf}$ ). Figure 4 shows a schematic of this GPS site geometry.

Assuming that the GPS pole base remains fixed within its original firn layer (see Section 1.1 for further discussion), observed decreases in the antenna-surface distance ( $h_{ant-surf}$ ) can be attributed to surface accumulation (e.g. snowfall, deposition of snow by wind). Conversely, an increase in antenna-surface distance can be attributed to surface ablation (e.g. melt, sublimation, removal of snow by wind) and compaction of snow/firn above the pole base.

We computed mean daily antenna-surface distance for all sites using L1 C/A code multipath surface reflections and the GPS interferometric reflectometry methodology outlined in Larson et al. (2015). This method takes advantage of the fact that the interference between the direct and reflected GPS signals produces characteristic frequencies in signal-to-noise ratio data recorded by the GPS receiver; these frequencies are directly related to the distance between the GPS antenna phase center and the reflecting surface. Geodetic antennas are designed to suppress multipath, so these interference patterns are best resolved at low GPS satellite elevation angles. Reflector height solutions were calculated for elevation angles of 5–25°, which sample the surface within a radial extent of ~5–50 m. Local surface slopes at

**Deleted:** Geodetic GPS

**Deleted:** ; we

**Deleted:** . All position data were converted to a local Cartesian horizontal coordinate system with orthometric heights above the EGM2008 geoid (Pavlis et al., 2012). Positions

**Deleted:** were removed.

**Deleted:** years

**Deleted:**

**Deleted:** Horizontal velocities for each GPS station were computed from daily mean positions. Relative distances between stations were used to calculate strain, with linear fits to estimate strain rates. ... [1]

**Deleted:** (e.g., Padman et al., 2003).

**Deleted:** .

**Deleted:** phase center elevations,

**Moved (insertion) [1]**

**Deleted:** The constant 3.66 m pole length and 5.32 cm antenna phase center offset was removed from filtered absolute antenna elevation ( $h_a$ ) to estimate corresponding absolute pole-base elevation ( $h_{pb}$ ) relative to the EGM2008 geoid. Figure 4 shows a schematic of this geometry with additional observables as described below ... [2]

**Deleted:** height above the

**Deleted:** ( $z_a$ ) from

**Deleted:** using

**Deleted:** (Figure 4).

**Deleted:** .

**Continuous antenna height**

each site are negligible, eliminating the need for an azimuthal correction (e.g., Larson and Nievinski, 2013). ~~Daily antenna-surface distance ( $h_{ant-surf}$ ) accuracy is estimated to be ~1 cm (Larson et al., 2015).~~

### 2.3 GPS-derived surface elevation

The antenna-surface distance ( $h_{ant-surf}$ ) was subtracted from antenna elevation ( $z_{ant}$ ) to obtain daily records of surface elevation  $z_{surf}$  (i.e., elevation of air-snow/firn interface above EGM2008 geoid), with resulting relative accuracy of ~1-2 cm (absolute accuracy subject to the same ~0.1 m  $z_{ant}$  uncertainty due to tidal, IBE, and geoid corrections). The  $z_{surf}$  surface elevation values are directly comparable with satellite/airborne laser altimetry data and stereo DEM products. We use variable name  $z_{surf}$  rather than the more traditional glaciological variable name  $h$  to limit potential confusion between different “height” and “elevation” variables.

Continuous  $z_{surf}$  time series were generated for all seven PIG GPS sites. The SOW3 record was curtailed on August 22, 2013, when antenna-surface distance decreased below the minimum threshold of ~0.5 m (Nievinski, 2013).

### 2.4 GPS velocity and strain rate

Horizontal velocities for each GPS station were computed from daily mean antenna positions. We calculated principal strain rates for 8 different triangular sections within the array (each defined by unique combination of 3 sites), using the methods outlined by Savage et al. (2001). We tested multiple time intervals for these strain rate calculations, from 2 to 120 days, and use 42 days as a compromise between temporal resolution and uncertainty (assuming uncorrelated daily position error of ~1 cm). We used observed horizontal strain rates to estimate elevation change related to local flux divergence.

Some component of observed GPS surface elevation change may also be related to deformation due to local gradients in the driving stress and surface-parallel flow due to advection over basal topography. The vertical component of surface-parallel flow ( $V_0$  in Larson et al. (2015)) can be estimated using observed horizontal GPS paths and surface gradients from an independent DEM. Advection over bed topography is irrelevant for a freely-floating ice shelf, and we attempt to estimate an upper bound for  $V_0$  due to local deformation by considering local surface gradients and observed relative horizontal displacements within the 2012–2014 GPS array.

### 2.5 High-resolution DEMs

In addition to the GPS elevation data, we generated WorldView/GeoEye stereo DEMs (Shean et al., 2016) with 32-m posting over the PIG shelf (Shean, 2016) to provide spatial context for the GPS time series. A total of 7 WorldView DEMs intersected the GPS positions between 2012–2014. We sampled DEM surface elevation at corresponding GPS positions and compared with GPS-derived surface elevation where possible.

High-resolution Lagrangian  $Dz_{surf}/Dt$  maps (see methodology in Shean, 2016; note we use  $D/Dt$  to indicate a Lagrangian differential operator) were computed for the 2012–2014 GPS sites by forward-propagating 32-m DEM pixels from two initial DEM products (February 2, 2012 and October 23, 2012) using interpolated, time-variable surface velocity maps from Joughin et al. (2010) and Christianson et al. (2016). Lagrangian  $Dz_{surf}/Dt$  maps were

**Deleted:** Continuous antenna height  
**Deleted:** , with resulting surface elevation accuracy of ~1-2 cm. These surface elevation values are directly comparable with satellite/airborne laser altimetry data and stereo DEM products. The multipath antenna height above surface ( $z_a$ ) was also used to estimate pole-base depth below the surface ( $z_{pb}$ ) at each site.  
**Deleted:** Continuous antenna height

**Deleted:** heights  
**Deleted:** ~0.5 m  
**Deleted:** SMB  
**Deleted:** Temperature data

Moved (insertion) [2]

Moved (insertion) [3]

Moved (insertion) [4]

generated for all valid combinations of these initial DEMs and ~15 subsequent DEMs (~0.5–2.5 years later). Composite products were generated, with median  $Dz_{surf}/Dt$  values assigned to initial DEM pixel locations.

## 2.6 Surface mass balance

We analyzed estimates of 1979–2015 monthly and 2010–2013 daily SMB for three 27-km grid cells over the PIG shelf from the Regional Atmospheric Climate Model (RACMO) v2.3 (Ettema et al., 2009; Lenaerts et al., 2012; Van Meijgaard et al., 2008; Van Wessem et al., 2014). The average 1979–2015 SMB ( $\bar{a}$ ) is 0.91 m w.e./yr for the grid cell closest to the 2012–2014 GPS array (-75.07°N, -100.80°E, Figure 1). The values for adjacent grid cells are 0.74 m w.e./yr near the grounding line of the main shelf (-75.15°N, -99.88°E) and 0.84 m w.e./yr over the south shelf (-75.30°N, -101.14°E), providing some information on large-scale spatial variability. These values are consistent with SMB estimates of ~0.5–1.0 m w.e./yr derived from CRESESIS Snow Radar data collected upstream of the PIG grounding line (Medley et al., 2014, 2015) and SMB estimates of 0.99 and 1.06 m w.e./yr for stake measurements near 2006–2008 GPS sites over the upstream PIG trunk (Scott et al., 2009). We conservatively estimate SMB uncertainty of 0.2 m w.e./yr.

## 2.7 AWS temperature data

To provide context for surface elevation change due to surface melt events, we analyzed continuous 2011–2015 temperature data (3-hour interval) from the Evans Knoll (-74.85°N, -100.40°E, Figure 1) automated weather station (AWS) (Lazzara et al., 2012), located at an elevation of ~178 m (height above EGM2008 geoid) on a bedrock outcrop approximately 40 km north of the 2012–2014 GPS array (Figure 1). We also analyzed local New York University (NYU) AWS temperature data available near PIG2 from January 9, 2008 to November 7, 2009, and near BOAR from January 19, 2013 to May 26, 2015. Unfortunately, no AWS data were collected on the PIG shelf during 2012. An analysis of overlapping time periods for the Evans Knoll and 2013–2015 NYU AWS temperature records shows a median offset of +1.24°C (Evans Knoll warmer than NYU, with normalized median absolute deviation [NMAD] of 2.76°C), which is consistent with a dry adiabatic lapse rate and local site conditions. This offset was removed from the Evans Knoll temperature data to provide a continuous temperature estimate for the GPS sites over the full 2012–2014 period.

To provide historical context, we extracted 2-m air temperature from 1979–2015 for grid cells over the PIG shelf in 0.75°-resolution ERA-Interim reanalysis products (Dee et al., 2011) with 6-hour interval. The median offset between the ERA-Interim temperature data and the 2013–2015 NYU AWS temperature data was +0.10°C (ERA-Interim warmer than NYU) with NMAD 2.78°C. This median offset was removed from the ERA-Interim temperatures. We did not attempt to correct any seasonal bias in ERA-Interim products (e.g., Jones et al., 2016).

## 2.8 Firn model

We use a dynamic firn model to simulate elevation change related to SMB and firn processes, which can be used to isolate the component of observed elevation change related to ice dynamics and basal mass balance. Model SMB output from RACMO2.3 (Section 2.6) was used to force the semi-empirical 1-D IMAU-FDM dynamic firn model

Moved (insertion) [5]

Deleted: regional atmospheric climate model RACMO

Deleted: mwe/

Deleted: mwe/

Deleted: mwe/

Deleted: mwe/

Deleted: mwe/

Deleted: We also analysed

Deleted: with

Deleted: The AWS temperature values were scaled to the surface elevation of the GPS array (~66 m above EGM2008 geoid) assuming a dry lapse rate of 9.8° C/km

Deleted: The ERA-Interim temperature data display a bias of -2.81°C (median of offsets) compared to the scaled 2011–2015 AWS data, which is consistent with previous evaluations (Jones et al., 2016). This bias was removed from the ERA-Interim temperature record, with residual offsets displaying root-mean-square error (RMSE) of 3.26 °C and normalized median absolute deviation (NMAD) of 2.77 °C.

Deleted: densification

Deleted: Model SMB output from RACMO2.3

(Ligtenberg et al., 2011) with 3-hour ~~time step, and~~ IMAU-FDM output ~~was generated~~ at 2-day intervals. Velocities ( $v_{ice}$ ) across the firm-ice transition (defined as the layer with 917 kg/m<sup>3</sup> density) were assumed to be in equilibrium with average 1979–2015 SMB ( $\bar{a}$ ). Vertical velocity components for surface accumulation, surface sublimation, surface snow drift erosion/deposition, surface melt, dry firm compaction ( $v_{fc}$ ), and a vertical ~~buoyancy correction (over floating ice shelf grid cells) were computed for the 2008–2010 and 2012–2014 periods (see Ligtenberg et al., 2011 for model details). These components were combined to provide time series of simulated surface elevation ( $\widehat{z}_{surf}$ ) and a tracer for pole-base elevation at each GPS station. We conservatively estimate IMAU-FDM surface elevation uncertainty of ~10%, which corresponds to ~0.05 m for  $\widehat{z}_{surf}$  and ~0.1 m/yr for  $v_{fc}$ .~~

**Deleted:** timesteps. The

**Deleted:** is available

**Deleted:** mass difference buoyancy correction were computed for the 2008–2010 and 2012–2014 periods. These estimates were combined to provide time series of simulated surface ( $\hat{h}$ ) and pole-base elevations ( $\hat{h}_{pb}$ ) for each GPS station

### 3 Derivation of basal mass balance

**Deleted:** <#>Basal melt rate .

We integrate the above observations and model output to estimate basal mass balance for the PIG GPS sites. Mass conservation for a column with ice-equivalent thickness  $H_{ice}$  (after removing a thickness correction  $d$  that accounts for total air content in the firm column) relates Eulerian thickness change ( $\frac{dH_{ice}}{dt}$ ) with dynamic thinning/thickening due to flux divergence ( $\nabla \cdot H_{ice} \mathbf{u}$ , positive for extension), surface mass balance  $\dot{a}$  (meters ice equivalent), and basal mass balance  $\dot{b}$  (meters ice equivalent, defined as positive for melt):

**Deleted:** H

**Deleted:**  $dH$

**Deleted:** local

**Deleted:**  $Hu$

$$\frac{\partial H_{ice}}{\partial t} = -\nabla \cdot (H_{ice} \mathbf{u}) + \dot{a} - \dot{b} \quad (1)$$

**Deleted:**  $\frac{\partial H}{\partial t}$

**Deleted:** (H)

The material derivative definition relates Eulerian (fixed reference frame) and Lagrangian (reference frame moving with the ice column) thickness change:

**Deleted:** This approach assumes that the total firm air content ( $d \approx 12$  m for the PIG shelf (Shean, 2016)) remains constant for the period  $dt$ , which precludes the need to account for thickness change due to firm compaction, while allowing for variable ice-equivalent SMB input  $\dot{a}$ . This assumption is supported by the limited variability (~1-3% or +/-0.3 m) in modelled IMAU-FDM total firm air content for the three PIG shelf grid cells during the relevant ~2-year study periods.

$$\frac{DH_{ice}}{Dt} = \frac{\partial H_{ice}}{\partial t} + \mathbf{u} \cdot (\nabla H_{ice}) \quad (2)$$

Rearranging Equation 2 and substituting into Equation 1, we obtain the mass conservation equation for Lagrangian thickness change:

**Deleted:**  $\frac{DH}{Dt} = \frac{\partial H}{\partial t} + \mathbf{u} \cdot (\nabla H)$

**Deleted:**  $\frac{DH}{Dt}$

**Deleted:**  $H(\nabla \cdot \mathbf{u})$

$$\frac{DH_{ice}}{Dt} = -H_{ice}(\nabla \cdot \mathbf{u}) + \dot{a} - \dot{b} \quad (3)$$

For a floating ice shelf in hydrostatic equilibrium, we can estimate ice-equivalent thickness from ice-equivalent surface elevation ( $z_{surf} - d$ ), where  $z_{surf}$  is measured surface elevation and  $d$  is total firm air content;

**Deleted:** freeboard

**Deleted:**  $h_f = (h -$

**Deleted:**  $h$

**Deleted:**

**Deleted:** . We can then estimate total ice-equivalent thickness H

**Deleted:**  $H \approx h_f \left( \frac{\rho_w}{\rho_w - \rho_i} \right)$

**Deleted:** Assuming

**Deleted:** ), we can

**Deleted:** Lagrangian

**Deleted:**  $\left( \frac{\partial h_f}{\partial t} + h_f(\nabla \cdot \mathbf{u}) \right)$

**Deleted:** Equation 5 depends on ice-equivalent surface elevation change.

$$\dot{b} = - \left( \frac{Dz_{surf}}{Dt} + (z_{surf} - d)(\nabla \cdot \mathbf{u}) \right) \left( \frac{\rho_w}{\rho_w - \rho_i} \right) + \dot{a} \quad (5)$$

We assume that the total firm-air content ( $d \approx 12$  for the PIG shelf [see appendix in Shean (2016)], with uncertainty of ~2 m) remains constant for the period  $dt$ , and drop the constant  $d$  from the material derivative term. This simplification is supported by the limited temporal variability (+/-0.3 m or ~1-3%) in modeled IMAU-FDM total firm air content for the three PIG shelf grid cells during the relevant ~2-year study periods.

We now consider elevation change for the GPS pole base within the firm column of this simplified ice shelf. The effects of processes that drive short-term surface-elevation change (e.g., accumulation, melting) are largely absent below the upper few meters of the firm column. Thus, the elevation change of the GPS pole base (equivalent to  $Dz_{ant}/Dt$  for constant pole length) is much more sensitive to the velocity across the firm-ice transition ( $v_{ice} = \bar{a}/\rho_i$ , see Section 2.8), and compaction within the underlying firm. The downward velocity of the pole base due to firm compaction ( $v_{fc}$ ) varies as a function of pole-base depth within the firm column. Values for  $v_{fc}$  can be estimated by integrating firm model compaction rates from the firm-ice transition to the pole-base tracer at each timestep. If SMB ( $\dot{a}$ ) for the time period  $dt$  is approximately equal to the long-term average SMB ( $\bar{a}$ ), then:

$$\frac{Dz_{ant}}{Dt} \approx \frac{Dz_{surf}}{Dt} + v_{fc} \quad (6)$$

which can be substituted into Equation 5 to estimate basal melt rate from observed antenna elevation change:

$$\dot{b} = - \left( \frac{Dz_{ant}}{Dt} - v_{fc} + (z_{surf} - d)(\nabla \cdot \mathbf{u}) \right) \left( \frac{\rho_w}{\rho_w - \rho_i} \right) + \dot{a} \quad (7)$$

The pole-base depth below the surface is negligible compared to total ice thickness ( $z_{surf} - z_{polebase} \ll H_{ice}$ ), and we neglect the small difference in associated firm air content. If modeled downward velocity due to firm compaction is correct, then basal melt rate estimates from Equation 5 and Equation 7 should be similar. We also note that estimated basal melt rates are ~9 times more sensitive to surface elevation change ( $Dz_{surf}/Dt$ ) than SMB ( $\dot{a}$ ) for a floating ice shelf.

## 4 Results

### 4.1 Horizontal velocity

Figure 5A shows the horizontal surface velocities of the PIG1 and PIG2 stations. On the floating ice at PIG2, velocity increased from ~355 m/yr to ~380 m/yr between 2008 and 2010 as the GPS moved downstream. Velocities for grounded ice at PIG1 increased at a relatively steady rate from ~420 m/yr to ~460 m/yr as the station moved toward the fast-flowing PIG trunk (Figure 1).

Figure 5B shows the 2012–2014 velocities for the GPS array, which varied from ~3830–4040 m/yr (Christianson et al., 2016). Velocities at SOW1, BOAR, and SOW3 were similar, while SOW4 (closer to shelf centerline) consistently moved ~20 m/yr faster than these three sites, and SOW2 consistently moved ~15 m/yr slower. Thus, there appears to be ~30–40 m/yr dextral (right-handed) shear across the ~2.4 km distance between the SOW4 and SOW2 sites. This transverse velocity gradient is also apparent in velocity mosaics (e.g., Christianson et al., 2016).

The velocity of all five stations varied by ~2–4% from 2012–2014, as described in detail by Christianson et al. (2016). In general, the stations displayed similar relative velocity evolution, with several abrupt >0.1–0.2 m/day velocity changes during the ~2-year period (Figure 5).

- Deleted: .
- Deleted: short-term
- Deleted: surface
- Deleted: rate
- Deleted:  $Dh_{pb}$
- Deleted:  $z_{pb}$
- Deleted:  $h_{pb}$
- Deleted: ( $\dot{a}$ )
- Deleted:  $\frac{Dh_{pb}}{Dt}$
- Deleted:  $\frac{Dh_f}{Dt}$
- Deleted: pole-base
- Deleted:  $\left( \frac{Dh_{pb}}{Dt} - v_{fc} + h_f(\nabla \cdot \mathbf{u}) \right)$
- Deleted: We neglect the slight reduction in total ice-equivalent thickness at the pole base ( $h_{pb}$ ) vs. surface ( $h$ ), as pole-base depth is negligible compared to total ice thickness ( $z_{pb} \ll H$ ). For the local flux divergence term, we use local freeboard thickness (as in Equation 7) or  $H$  from radar measurements (Stanton et al., 2013). Uncertainty is estimated as ~0.15 m/yr for downward firm-compaction velocity (Ligtenberg et al., 2011), ~0.1 m/yr for elevation change associated with local flux divergence, and ~5 kg/m<sup>3</sup> for the density of ice and ocean water. ... [3]
- Deleted:  $Dh/Dt$  (
- Deleted: )
- Deleted:  $Dh_{pb}/Dt$  (
- Deleted: )
- Deleted: inferred
- Deleted: pole-base
- Deleted:  $Dh_{pb}$
- Deleted: and local  $v_{fc}$
- Deleted: <#>Velocity .
- Deleted: surface
- Deleted: migrated
- Deleted: , with increased speedup beginning in late 2008.
- Deleted: GPS
- Deleted: at SOW1, SOW2, BOAR, SOW3, and SOW4
- Deleted: subtle
- Deleted: 10–20 cm

## 4.2 Strain rate

Figure 6 shows strain rate magnitude and direction for 8 different strain triangles within the 2012–2014 GPS array. Mean principal strain rates were  $+0.0018 \text{ yr}^{-1}$  (extension approximately in the along-flow direction) and  $-0.0001 \text{ yr}^{-1}$  (compression approximately in the across-flow/transverse direction). The array displayed a clockwise rotation rate of  $\sim 1^\circ/\text{yr}$ .

Spatial variations of strain rates within the array are small (Figure 6). Strain triangles including SOW1 experienced higher strain rates, while triangles including SOW3 experienced lower strain rates, despite its location within the large transverse depression (Figure 2). Strain rate temporal variability is also limited, but there do appear to be significant changes correlated with shelf-wide velocity changes. In general, increased (decreased) extensional strain rates were observed following an increase (decrease) in absolute GPS array velocity.

Local surface slopes near SOW1, SOW2, and BOAR are negligible ( $<0.2^\circ$ ), so we assume no surface-parallel vertical motion for these stations (i.e.,  $V_0 = 0$ ). If all of the observed  $\sim 3.4 \text{ m/yr}$  relative displacement between SOW1 and SOW3 was attributed to flow down  $\sim 0.6^\circ$  local surface slopes at SOW3, then the associated  $V_0$  magnitude would only be  $\sim 0.03 \text{ m/yr}$ , which is negligible compared to the observed  $\sim 5.2 \text{ m/yr } Dz_{\text{surf}}/Dt$ .

For estimated ice-equivalent thickness of  $\sim 430\text{--}500 \text{ m}$ , the observed strain rates correspond to shelf thinning rates ( $DH_{\text{ice}}/Dt$ ) of  $\sim 0.5\text{--}0.9 \text{ m/yr}$ , with expected surface elevation change ( $Dz_{\text{surf}}/Dt$ ) of only  $<0.07\text{--}0.13 \text{ m/yr}$ . Based on these estimates, we assume a value of  $-0.1 \pm 0.03 \text{ m/yr}$  for the divergence term in Equations 5 and 7 when calculating basal melt rates.

## 4.3 Antenna-surface distance

Initial antenna-surface distances ( $h_{\text{ant-surf}}$ ) were  $\sim 2.5$  to  $3.1 \text{ m}$ , indicating that initial pole-base depths were  $\sim 0.6$  to  $1.2 \text{ m}$  below the initial surface (Figure 7A, Figure 8). Antenna-surface distance decreased over time at all sites.

At both PIG1 and PIG2, there were periods of relatively rapid antenna-surface distance decrease (e.g., from May to August 2008), followed by a steady increase (e.g., August 2008 to February 2009). These changes are consistent with periods of snow accumulation followed by several months of ongoing firm compaction with limited snowfall. The 2012–2014 records show similar periods of abrupt antenna-surface distance decrease and steady increase, with more limited duration.

All GPS array records show an abrupt antenna-surface distance increase ( $\sim 0.2\text{--}0.3 \text{ m}$ ) between December 2012 and January 2013, which is consistent with surface melting and enhanced firm compaction rates in the upper few meters of the firm column.

## 4.4 GPS antenna and surface elevation change

Trends in observed antenna elevation change ( $Dz_{\text{ant}}/Dt$ ) are negative and highly linear ( $R^2 0.98\text{--}1.00$ ) for all PIG shelf sites, with rates of  $-1.6$  to  $-2.1 \text{ m/yr}$  at SOW1, SOW2, and BOAR, and higher rates of  $-5.2 \text{ m/yr}$  and  $-3.8 \text{ m/yr}$  at SOW3 and SOW4, respectively (Figure 7B, Table 1). Observed  $Dz_{\text{ant}}/Dt$  over grounded ice at PIG1 is  $-7.6 \text{ m/yr}$ , with apparent concave-downward curvature. This is consistent with  $V_0$  expected for surface-parallel flow (see Section 2.4) and

**Deleted:** For this analysis, we assume that SOW1, BOAR,  
**Deleted:** SOW3 GPS stations  
**Deleted:** oriented  
**Deleted:** a flowline,  
**Deleted:** that the observed displacements represent longitudinal strain. Figure 6A shows that the observed cumulative displacement between SOW1 and SOW3 (initial distance 2073.0 m) was  $\sim 6.7 \text{ m}$  ( $\sim 3.4 \text{ m/yr}$ ), with extensional strain rates of  $\sim$   
**Deleted:** 0017  
**Deleted:** Observed strain rates between SOW1 and BOAR ( $\sim 2.1 \text{ m/yr}$  over 1045.2 m,  $0.0020 \text{ yr}^{-1}$ ) were greater than those between BOAR and SOW3 ( $\sim 1.5 \text{ m/yr}$  over 1029.1 m,  $0.0014 \text{ yr}^{-1}$ ). Transverse  
**Deleted:** were relatively low, with compression ( $-0.0004 \text{ yr}^{-1}$ ) between BOAR and SOW2, and extension ( $0.0007 \text{ yr}^{-1}$ ) between BOAR and SOW4. Subtle changes in  
**Deleted:** between SOW1 and SOW3 were observed from 2012–2014 (Figure 6C).  
**Deleted:** occurred between SOW1 and SOW3  
**Deleted:** These longitudinal strain rates correspond to shelf thinning rates of  $\sim 0.5\text{--}0.9 \text{ m/yr}$  assuming no lateral spreading, with limited expected surface  $Dh/Dt$  ( $<0.07\text{--}0.13 \text{ m/yr}$ ) for measured ice thickness of  $\sim 450\text{--}460 \text{ m}$  (Stanton et al., 2013) or for thickness estimated from local freeboard ( $\sim 430\text{--}500 \text{ m}$ ).  
**Deleted:**  $\langle \# \rangle$  GPS antenna and surface elevation change ... [4]  
**Deleted:**  $V_0 = 0$   
**Deleted:** .  
**Deleted:**  $5.2 \text{ m/yr } Dh/Dt$ . Considering that much of the observed relative displacement appears to be related to longitudinal extension (Section 3.2), we neglect any potential  $Dh/Dt$  contribution related to  $V_0$   
**Deleted:**  $\langle \# \rangle$  GPS antenna and surface elevation change ... [5]

**Deleted:** ( $h_a$ )  
**Deleted:** relative to the initial absolute  
**Deleted:** ( $h_{a0}$ ) at each station. These results show  
**Deleted:** .  
**Deleted:**  $Dh/Dt$  (Table 1),  
**Deleted:** . The PIG1 rates  
**Deleted:** are  
**Deleted:** from local  
**Deleted:** topography

dynamic thinning over the PIG trunk associated with velocity increases in 2006–2008 GPS observations (Scott et al., 2009) and satellite records (Joughin et al., 2010; Mouginot et al., 2014).

The 2008–2010 surface elevation change ( $Dz_{surf}/Dt$ ) at PIG2 is negligible (-0.13 m/yr). By contrast, surface elevations decreased significantly at all 2012–2014 GPS array sites, with rates of -0.9 to -1.3 m/yr for SOW1, SOW2 and BOAR, and rates of -4.1 m/yr and -3.0 m/yr at SOW3 and SOW4, respectively.

Residuals about these linear fits (Figure 9A+B) are small for PIG shelf sites (root mean squared error (RMSE) of 0.095 m for  $Dz_{ant}/Dt$ , and RMSE of 0.143 m for  $Dz_{surf}/Dt$ ), with some seasonal to annual variability. We also note relatively abrupt (~days-weeks) elevation changes that occurred across all stations in the 2012–2014 array (e.g., -0.3 to +0.3 m anomaly during June 2012).

#### 4.5 Surface mass balance

We consider surface elevation relative to a firm layer tracer for the initial surface elevation ( $z_{surf} - z_{surf(0)}$ ) to estimate cumulative elevation change due to SMB after GPS installation. Observed rates were  $-0.9-1.1$  m/yr for 2008–2010 sites and  $-0.8-0.9$  m/yr for 2012–2014 sites (with SOW3 at  $-1.1$  m/yr) (Figure 9C).

The 1979–2015 RACMO average SMB over the central PIG shelf is  $\sim 0.9$  m w.e./yr. Monthly SMB climatology shows low accumulation rates of  $\sim 0.01-0.04$  m w.e./month over the PIG shelf during the austral summer (November to February), and high accumulation rates of  $\sim 0.08-0.1$  m w.e./month during austral winter (March to October) (Figure 9D). Daily SMB products show periods of days to weeks with increased accumulation (e.g., March 2013) that can be correlated with abrupt decreases in antenna-surface distance.

The  $\sim 3-4$  week period between December 24, 2012 and January 17, 2013 was relatively warm, with scaled AWS temperatures of  $\sim 1-5^\circ\text{C}$  for most days (Figure 9E). We note that these are 2-m air temperatures, and that once surface melting commenced, actual surface temperatures would be lower, but still above freezing. The daily RACMO SMB data also show two accumulation events during the last week of December 2012 (Figure 9D), which involved rain on snow (M. Truffer, personal communication, 2016). Surface elevations decreased by  $\sim 0.2-0.3$  m across the entire GPS array during this warm/rainy period (Figure 9B), which is consistent with surface melting, refreezing, and/or enhanced firm compaction rates. No corresponding short-term changes were recorded by the antenna elevations during the  $\sim 3-4$  week period (Figure 9A), suggesting that the processes responsible for the observed surface changes did not affect the firm layers near the pole base ( $\sim 1.5$  m depth). We note that there are many warm periods between 1979-2015 with greater magnitude and duration than the December 2012 to January 2013 period in the ERA-Interim 2-m air temperatures over the PIG shelf.

#### 4.6 Firm model

Simulated downward pole-base velocities due to firm compaction ( $v_c$ ) were  $\sim 0.70-0.75$  m/yr (Figure 7, Table 1) from 2008–2010 and 2012–2014, despite variable initial pole-base depth. A slight decrease in the compaction rate occurred over time, but the curves appear approximately linear (Figure 7B). These firm compaction rates are consistent with observed  $Dz_{surf}/Dt$  during extended periods with little or no surface accumulation.

**Deleted:** Figure 9C shows corresponding surface elevation ( $h$ ) change relative to initial surface elevation ( $h_0$ ) at each station.

**Deleted:** D+E

**Deleted:**  $(Dh_s/Dt)$

**Deleted:**  $Dh/Dt$

**Deleted:** sites on the PIG shelf

**Deleted:** observed at all sites.

**Deleted:** apparent

**Deleted:**  $\langle \# \rangle$  Antenna height evolution ... [6]

**Deleted:** ( $z_s$  in Figure 4, Section 2.3) can be attributed to surface accumulation (e.g., snowfall, deposition of snow by wind). Conversely, an increase in antenna height

**Moved up [1]:** can be attributed to surface ablation (e.g., melt, sublimation, removal of snow by wind) and compaction of snow/firm above the pole base.

**Deleted:** For convenience, we define the reflector height ( $z_r$ ) to track cumulative accumulation and ablation

**Deleted:** ( $h_0$ ) ... [7]

**Deleted:** of  $\sim$

**Deleted:** 0

**Deleted:** 7

**Deleted:** 8

**Deleted:** 0

**Deleted:** .)

**Moved (insertion) [6]**

**Deleted:** At both PIG1 and PIG2, there are periods of relatively rapid reflector height increase (e.g., from May–August 2008), followed by a steady decrease (e.g., August 2008–February 2009). This is consistent with periods of increased snow accumulation followed by several months of ongoing firm compaction with limited snowfall. The 2012–2014 records show similar periods of abrupt increase and steady decrease, with more limited duration. ... [8]

**Moved up [2]:** ... [9]

**Deleted:** (Shean, 2016)

**Moved up [3]:** to provide spatial context for the GPS time series. A total of 7 WorldView DEMs intersected the GPS positions between 2012–2014. We sampled DEM surface elevation at corresponding GPS positions and compared with GPS-derived surface elevation

Figure 7C shows that the IMAU-FDM simulated surface elevation ( $\bar{z}_{surf}$ ) ranges from -0.1 to +0.4 m from 2008–2010 and -0.2 to +0.2 m from 2012–2014. The observed  $D\bar{z}_{surf}/Dt$  trend is +0.17 m/yr from 2008–2010, with no significant trend from 2012–2014. The magnitude and timing of the simulated surface elevation variability is consistent with the detrended observed surface elevation change (Figure 9B). The observed  $Dz_{surf}/Dt$  trends (-1 to -4 m/yr), however, cannot be explained by simulated elevation change due to SMB and firm processes (Figure 7C).

#### 4.7 Basal melt rates

We computed basal melt rates from surface  $Dz_{surf}/Dt$  and antenna  $Dz_{ant}/Dt$  elevation change using Equations 5 and 7, respectively. The resulting melt rate estimates range from ~2–4 m/yr at PIG2 to ~39–43 m/yr at SOW3, with good agreement between the two approaches (Table 1).

The 2012–2014 melt rate estimates show significant spatial variability. The three upstream stations (SOW1, SOW2 and BOAR) experienced lower melt rates of ~9–13 m/yr, while the downstream stations near the transverse depression (SOW3 and SOW4) experienced higher rates of ~29–43 m/yr for the same time period.

#### 4.8 High-resolution DEMs

Figure 8 shows sampled DEM elevation compared with GPS surface elevation at each site, with statistics provided in Table 2. In general, we observe good agreement between the two datasets, with RMSE of 0.72 m and NMAD of 0.57 m for the full sample (n=25). The DEMs display a slight bias (+0.43 m) relative to the GPS surface elevation.

We observe good agreement between GPS-derived (Table 1) and DEM-derived (Table 2)  $Dz_{surf}/Dt$  trends. The shorter DEM  $Dz_{surf}/Dt$  intervals (e.g., ~1 year for SOW1 and BOAR) display larger errors than longer DEM intervals (~2 years for SOW2 and SOW3).

Figure 10 shows the composite DEM-derived  $Dz_{surf}/Dt$  maps, which provide spatial context for the GPS-derived  $Dz_{surf}/Dt$  records. Little or no elevation change was observed over longitudinal ridges, while areas within and near transverse depressions experienced enhanced thinning. This thinning was concentrated on the upstream side of the transverse depressions. The  $Dz_{surf}/Dt$  products relative to the October 23, 2012 DEM (Figure 10D) also show the spatial pattern of thinning associated with the rift that opened upstream of SOW1 in ~2014 (Jeong et al., 2016).

### 5 Assumptions

The methods presented in Section 2 relied on several simplifying assumptions. We now offer further discussion of these assumptions and their potential influence on our results.

#### 5.1 Hydrostatic equilibrium

In the absence of direct ice thickness measurements (e.g., radar profiles near PIG GPS sites), we assume hydrostatic equilibrium and use surface elevation to estimate freeboard ice thickness - a standard practice for ice shelf studies. While this assumption can lead to increased uncertainty within a few ice thicknesses of the grounding line (Brunet et

<b>Moved (insertion) [7]</b>
<b>Moved (insertion) [8]</b>
<b>Deleted:</b> (as described in Section 2.3) where possible. Table 2 shows statistics for the sampled DEM elevation compared with GPS surface elevation at each site (Figure 8).
<b>Deleted:</b> We note that the January 14, 2012 WorldView DEM is anomalously high, which biases DEM Dh/Dt trends with limited sample count (Table 2).
<b>Deleted:</b> High-resolution Lagrangian Dh/Dt elevation-change maps (see methodology in Shean, 2016) were computed for the 2012–2014 GPS sites by forward propagating 32-m DEM pixels from two initial DEM products (February 2, 2012 and October 23, 2012) using interpolated, time-variable surface velocity maps from Joughin et al.
<b>Moved up [4]:</b> (2010) and Christianson et al. (2016).
<b>Deleted:</b> Lagrangian Dh/Dt maps were generated for all valid DEM combinations. Products with time interval between 0.5–2.5 years were aggregated, and median Dh/Dt values were assigned to initial DEM pixel locations.
<b>Deleted:</b> DEM Dh/Dt trends
<b>Deleted:</b> and GPS surface Dh/Dt trends (Table 1), with values of -1 to -2 m/yr near SOW1, SOW2, and BOAR, and -4 to -5 m/yr near SOW3 and SOW4.
<b>Deleted:</b> Dh
<b>Deleted:</b> Lagrangian
<b>Deleted:</b> Dh
<b>Deleted:</b> Dh
<b>Deleted:</b> estimates
<b>Deleted:</b> (Figure 10).
<b>Deleted:</b> Dh
<b>Moved up [5]:</b> <#>Surface mass balance
<b>Deleted:</b> <#> ... [10]
<b>Moved up [6]:</b> <#> Daily SMB products show periods of days to weeks with increased accumulation (e.g., March 2013) that can be correlated with abrupt
<b>Deleted:</b> <#>increases in surface reflector height. By contrast, the reflector height records show a steady decrease due to ongoing near-surface firm compaction during extended periods with little/no accumulation. ... [11]
<b>Moved up [7]:</b> <#> ranges from -0.1 to +0.4 m from 2008–2010 and -0.2 to +0.2 m from 2012–2014.
<b>Deleted:</b> <#>The observed $D\bar{h}/Dt$ trend is +0.17 m/yr from 2008–2010, with no significant trend from 2012–2014. The magnitude and timing of the modelled surface elevation variability is consistent with the detrended observed surface elevation change (Figure 9D). The observed $Dh/Dt$ trends (-1 to -4 m/yr) (Figure 9C), however, cannot be explained by modelled elevation change due to only SMB and firm. ... [12]
<b>Moved up [8]:</b> <#>The 2012–2014 melt rate estimates show significant spatial variability.
<b>Deleted:</b> <#>The three upstream stations (SOW1, SOW2 and BOAR) show lower melt rates of ~9–14 m/yr, while the downstream stations near the transverse depression (SOW ... [13]
<b>Deleted:</b> results
<b>Deleted:</b> the
<b>Deleted:</b> 3

al., 2010; Griggs and Bamber, 2011), it is reasonable for the mid-shelf location of the GPS array, which in ~2012, had been approaching hydrostatic equilibrium for over 10–12 years since crossing the grounding line.

Previous studies using airborne ice-penetrating radar data have noted that most of the PIG shelf is generally near hydrostatic equilibrium (Bindschadler et al., 2011; Dutrieux et al., 2013; Vaughan et al., 2012). Dense radar grids, however, reveal narrow shelf-bottom channels, crevasses, and other features with horizontal length scales of ~10s–100s of meters that are not apparent in ice shelf surface topography (Langley et al., 2014; Vaughan et al., 2012). The thinner ice above these narrow features is partially supported by lateral bridging stresses, so that the corresponding surface elevation will appear higher than the expected freeboard surface elevation, providing erroneously large ice thickness estimates using Equation 4 (Drews, 2015; Shabtaie and Bentley, 1982; Vaughan et al., 2012).

Experiments with a high-resolution ice-flow model show that wider basal channels tend to be near equilibrium, while increased bridging stresses support ice over narrow basal channels (Drews, 2015). The PIG GPS array is ~2 km across, which is >4-5x the local ice thickness (~350-500 m). The ~1-2 km length scale of nearby longitudinal channels/keels is >2-3x the local ice thickness, with typical surface elevation difference between trough floors and adjacent ridge crests of <10 m. For the observed ice thickness, magnitude, and length scale of surface variations, and the relatively long timescales involved, we argue that the hydrostatic assumption is reasonable, and any vertical elevation change due to evolving bridging stresses should be negligible compared to the magnitude of observed  $Dz_{surf}/Dt$  and our conservative error estimates.

## 5.2 SMB spatial variability

We used modeled SMB from a single RACMO2.3 grid cell to drive the IMAU-FDM dynamic firm model, and applied the result to all GPS stations. We expect SMB to vary spatially (e.g., Medley et al., 2015) due to local environmental conditions (e.g. PIG2 elevation is >400 m higher than SOW1-4 stations on the shelf) and local surface topography (e.g., km-scale ridges/troughs), which will affect near-surface winds and snow redistribution.

The larger  $z_{surf} - z_{surf}'$  values (a proxy for surface accumulation) at SOW3 (Figure 9C) indicate that greater local accumulation occurred at this site within the transverse depression (Figure 2), potentially due to preferential deposition of wind-blown snow. However, we also note that the accumulation histories of SOW4, which sits near a surface ridge crest, and the three sites located on the floor of a broad, flat surface trough (SOW1, SOW2, BOAR) appear similar (Figure 9C).

The IMAU-FDM values do not account for horizontal advection of the firm column through spatially-variable RACMO fields (accumulation, surface temperature, etc.) over time. The GPS sites over the PIG shelf are moving ~4 km/yr (Figures 1 and 5), which is nearly double the observed PIG shelf velocities from the mid-1970s (Mouginot et al., 2014). Thus, the local firm columns beneath the GPS sites likely experienced variable SMB input over their ~50-100 km horizontal path during the corresponding 1979-2015 time period. This suggests that the true firm column thickness and compaction rates may differ from the IMAU-FDM estimates. For this reason, we use a constant firm air content estimate ( $d \approx 12 \pm 2$  m) derived from available airborne ice-penetrating radar two-way travel time and altimetry surface elevation measurements (see appendix A of Shean, 2016).

Deleted: modelled

Deleted: these parameters

Deleted: In addition, the

Deleted: 1-D

Deleted: northwestward at

Deleted: (see appendix A of Shean, 2016)

### 5.3 Pole settling/tilting

We now consider the possibility that some of the observed  $Dz_{surf}/Dt$  could be related to settling or tilting of the GPS poles over time. We assume that the poles froze in place shortly after installation, and the contact area (~1200 cm<sup>2</sup> for a ~1-meter-long cylinder with ~3.8 cm diameter) with surrounding firn should be sufficient to counter the downward gravitational force. Thus, we expect that antenna elevation change ( $Dz_{ant}/Dt$ ) represents rates at the base of the pole, rather than rates within an overlying firn layer.

A related consideration involves heating of the exposed pole during summer, which might lead to decoupling from the surrounding snow/firn and allow for additional penetration of the pole base within the firn. The pole base stoppers have prevented this penetration. In addition, we do not see any indication of such settling from December 2012 to January 2013, when surface elevations decreased by ~0.2–0.3 m, but pole base elevations showed little change (Figure 9A+B). The lack of pole-base elevation change also suggests that surface meltwater did not percolate more than ~1–2 m below the surface.

Finally, we assume that the poles were installed with vertical orientation and did not tilt over time. For an initially vertical pole with length of 3.71 m (including antenna phase-center offset), a 10° tilt would introduce a -0.06 m vertical antenna elevation error (-0.03 m/yr for a 2-year period), while a 20° tilt would introduce a -0.22 m vertical error (-0.11 m/yr). Thus, we expect vertical error associated with any tilting to be negligible compared to the large observed  $Dz_{ant}/Dt$  (-1.12 to -7.60 m/yr). These 10° and 20° tilts could, however, introduce horizontal errors of up to 0.64 and 1.27 m, respectively, which would affect intra-network displacement estimates and strain rate estimates. While it is possible that some minor tilting could have occurred (especially during initial months), this was not noted during servicing/removal, and the reflectometry results do not indicate any systematic change in directional antenna-surface offset.

### 5.4 Strain rate length scales

We estimated ~0.1 m/yr surface elevation change due to local flux divergence, assuming that the observed relative horizontal displacements are evenly distributed across the strain triangles, which have ~1–2 km edges between GPS stations. This assumption is supported by the ~1-km spatial extent of thinning signals within/near transverse depressions in the  $Dz_{surf}/Dt$  maps (Figure 10). Even if this strain is concentrated over a shorter distance (e.g., ~200 m), this contribution only increases to ~0.5 m/yr, which is still small compared to observed  $Dz_{surf}/Dt$  signals of ~3–4 m/yr. The relatively large spatial variability in  $Dz_{surf}/Dt$  values (~1 to ~4 m/yr) and lack of spatial variability in strain rates supports the assumption that the observed  $Dz_{surf}/Dt$  is primarily caused by basal melt.

## 6 Discussion

### 6.1 SMB and firn compaction

The evolution of GPS-derived surface elevation relative to a tracer for the initial surface ( $z_{surf} - z_{surf0}$ ) is consistent with SMB estimates ( $\dot{a}$ ) combined with estimates of downward velocity due to firn compaction ( $v_{fc}$ ), providing qualitative validation for the RACMO SMB and IMAU-FDM results. If we assume a near-surface bulk snow/firn

**Deleted:** We also expect local SMB and firn column thickness variability within the array. The greater  $z_{rh}$  surface reflector heights (a proxy for surface accumulation) at SOW3 indicate that enhanced local accumulation occurred within the transverse depression (Figure 2), likely due to wind-blown snow. However, we note that all of the remaining 2012–2014 stations display similar  $z_{rh}$  reflector height evolution, as do the two 2008–2010 stations (Figure 9A). Future RACMO and IMAU-FDM products with improved resolution should provide additional constraints on spatial variability. [... [14]

**Deleted:** snow/

**Deleted:** the  $Dh/Dt$  recorded by the GPS

**Deleted:** pole

**Deleted:** thus,

**Deleted:** However

**Deleted:** 20–30 cm and

**Deleted:** .

**Deleted:** below

**Deleted:** depth into

**Deleted:** snow/firn

**Deleted:** As discussed in Section 3.2, the expected surface  $Dh/Dt$  from

**Deleted:** is <0.1 m/yr

**Deleted:** ~1.5 m/yr

**Deleted:** displacement is

**Deleted:** ~1

**Deleted:** distance

**Deleted:** (BOAR and SOW3).

**Deleted:** <

**Deleted:** significantly less than the

**Deleted:**  $Dh/Dt$  signals. Interestingly, despite similar extensional strain rate estimates for SOW1-BOAR (0.0020 yr<sup>-1</sup>) and BOAR-SOW3 (0.0014 yr<sup>-1</sup>), we note a large difference in local  $Dh$

**Deleted:** at these stations (Figures 9 and 10). This

**Deleted:** local flux divergence for these sites is a minor component of the observed  $Dh/Dt$ , and consequently, that

**Deleted:**  $Dh$

**Deleted:** Long-term

**Deleted:** observed reflector height increase ( $z_{rh}$ )

**Deleted:** ~0.7–1.0 m/yr at all

**Deleted:** sites (Figure 9A)

**Deleted:** observed long-term

**Deleted:**  $\bar{a}$ ) and downward near-surface velocity

**Deleted:** .

**Deleted:** mean

density of  $\sim 500 \text{ kg/m}^3$ , the  $\sim 0.7\text{--}0.9 \text{ m w.e./yr}$  RACMO SMB estimates correspond to a surface elevation increase of  $\sim 1.4\text{--}1.8 \text{ m/yr}$ . Combining these rates with a  $\sim 0.7\text{--}0.75 \text{ m/yr}$  surface elevation decrease due to firm compaction provides estimates similar to observed  $z_{surf} - z_{surf0}$  trends of  $\sim 0.8\text{--}1.1 \text{ m/yr}$ . Based on these results, we suggest that it may be possible to extract detailed SMB records for other sites using only observed GPS antenna-surface distance and simple assumptions about firm densification (e.g., Herron and Langway, 1980). The problem is further simplified for grounded ice with negligible basal mass balance rates.

- Deleted: 0.5
- Deleted: mwe/
- Deleted: Removing the expected
- Deleted: associated with
- Deleted: arrive at expected relative reflector height increases of  $\sim 0.7\text{--}1.0 \text{ m/yr}$ , which matches observed  $Dz_{ref}/Dt$  values (Figure 9A+B).

The fact that we observe similar trends for  $Dz_{surf}/Dt$  and  $(Dz_{ant}/Dt - v_{\hat{e}})$  (Table 1) also supports the assumption that 2008–2010 and 2012–2014 SMB ( $\hat{a}$ ) is consistent with long-term 1979–2015 SMB ( $\bar{a}$ ) and associated firm-compaction rates.

- Deleted: surface  $Dh$
- Deleted: and pole-base  $Dh_{p}/Dt$

The limited variability in surface elevation at PIG2 (Figure 7C) suggests that the observed 2008–2010 SMB over the South PIG shelf was approximately in balance with ongoing firm compaction and basal melt during this period. We observed large surface elevation trends for the 2012–2014 GPS sites with no significant  $Dz_{surf}/Dt$  trend, suggesting that SMB and firm compaction during this period were consistent with long-term values, and that the large observed  $Dz_{surf}/Dt$  must be attributed to other processes, specifically basal melting.

- Deleted: Figure 9
- Deleted: . The 2012–2014 sites, however, show significant
- Deleted: decrease that greatly exceeds simulated IMAU-FDM surface elevation ( $\hat{h}$ ) variability ( $\pm 0.2 \text{ m}$ ) due to SMB and firm compaction. No
- Deleted:  $\hat{h}/$
- Deleted: is expected during this period

6.2 Residual elevation variability

The detrended surface (Figure 9A) and antenna (Figure 9B) elevation residuals appear to be uncorrelated. This suggests that seasonal surface processes (e.g., accumulation influencing near-surface compaction rates) are not responsible for driving antenna elevation variability. We considered several possible sources for the observed sub-annual elevation variability, including ocean (e.g., currents, sea surface height), atmospheric (e.g. pressure, temperature), and dynamic processes (e.g, resistive stress from sea ice and/or mélange in shear margins). Unfortunately, we were unable to definitively determine the cause(s) for these variations in the  $\sim 2$  year GPS records. Some of the short-term (days-weeks) variability (e.g. June 2012) observed across all five 2012–2014 stations (Figure 9B) could be related to insufficient or incorrect IBE correction. The magnitude and timing of these systematic anomalies, however, suggests that they are likely related to grounding/ungrounding events (e.g., Joughin et al., 2016).

- Deleted: are
- Deleted: Dh
- Deleted: values
- Deleted: melt
- Deleted: Dh/Dt
- Deleted: D
- Deleted: pole-base
- Deleted: E)
- Deleted: the observed pole-base
- Deleted: in the surface and pole-base records
- Deleted: D+E
- Deleted: (Joughin et al., 2016)

6.3 Strain rate history, rifting, and grounding evolution

The lateral shear across the GPS array is consistent with increased longitudinal extension closer to the PIG centerline, potentially due to locally enhanced ductile deformation (i.e., “necking” (Bassis and Ma, 2015)) across transverse depressions, and/or expansion of basal/surface crevasses and rifts. The SOW3 station, which lies within a large transverse depression (Figure 2), displays a slight acceleration in antenna elevation change (Figure 9A), potentially due to increased local extension within the depression.

- Deleted: related to
- Deleted: likely
- Deleted: negative
- Deleted: E

An upstream regrounding event would slow ice upstream of the GPS array, initially resulting in increased extensional strain rates across the transverse rifts/depressions, followed by a velocity decrease at the GPS array. Conversely, an upstream ungrounding event would initially lead to decreased extensional strain rates across the transverse rifts/depressions, followed by an increase in observed GPS velocities. We suggest that an upstream regrounding event (Joughin et al., 2016) in  $\sim$ June 2012 could be responsible for increased strain rates across the GPS array (Figure 6).

- Deleted: The fact that the rift immediately upstream of SOW1 ultimately became the site of the 2015 calving event (Jeong et al., 2016) suggests that the velocities observed at the GPS array are not necessarily representative of the velocities near the PIG grounding line.
- Deleted: .
- Deleted: in  $\sim$ June 2012

Similarly, an ungrounding event in ~April 2013 followed by a grounding event in ~November 2013 could explain the decrease and subsequent increase in strain rates.

There is an abrupt  $\sim 0.1\text{--}0.2\text{ m}$  antenna elevation ( $z_{ant}$ ) decrease at both SOW3 and SOW4 in late 2013, near the end of the records (Figure 9A). No surface elevation ( $z_{surf}$ ) information is available at SOW3 due to missing antenna surface distance data for this period, but a corresponding surface elevation decrease is observed at SOW4 (Figure 9B). These elevation decreases do not appear to be related to site servicing. Rather, these observations are consistent with relatively abrupt local extension within the transverse depression affecting SOW3 and SOW4, but not the upstream GPS sites. The timing of this event corresponds with observed lengthening of the large rift (R1) upstream of SOW1 (Jeong et al., 2016), supporting the hypothesis that relatively rapid, localized extension occurred across the transverse depressions and rifts during this period.

#### 6.4 Comparison with *in-situ* basal melt rate observations

The GPS-derived basal melt rate estimates ( $\sim 9\text{--}13\text{ m/yr}$  for SOW1, SOW2 and BOAR sites) appear consistent with those from bottom altimeter ( $\sim 14.7\text{ m/yr}$  from January–February 2012) and pRES ( $\sim 15\text{--}25\text{ m/yr}$ ) measurements of Stanton et al. (2013). These measurements provide some validation for the GPS results, as they are not influenced by surface mass balance and firn processes. A direct comparison may be imprudent, however, as the Stanton et al. (2013) borehole was  $\sim 1.34\text{ km}$  upstream of SOW1 (near the R1 rift), which likely affected local melt rates, and we observe considerable  $\sim\text{km-scale}$  spatial variability in melt rates across GPS array. Furthermore, the bottom altimeter sampled a  $\sim 5\text{ cm}$  diameter spot with unknown upstream/downstream orientation, approximately 30–40 cm from the edge of the 20 cm borehole. Aside from local melt variability expected due to turbulent flow near the altimeter pole or borehole edge, the altimeter provided a relatively small spatial sample compared to the GPS results, which are sensitive to changes in a column of ice with much larger footprint (100s of meters).

#### 6.5 Basal melt rate spatial variability

The GPS records at SOW1, SOW2, and BOAR show similar  $Dz_{surf}/Dt$  rates and residuals, which is consistent with their apparent orientation on the same “block” between transverse rifts/depressions (Figure 2), and supports the hypothesis that they were exposed to similar sub-shelf circulation. The DEM  $Dz_{surf}/Dt$  maps show enhanced surface elevation change rates, and thus higher basal melt rates, on the upstream side of transverse depressions (Figure 10), which is consistent with increased  $Dz_{surf}/Dt$  observed at the SOW3 and SOW4 sites.

This relationship is potentially related to enhanced buoyant flow and/or turbulence over increased basal slopes (e.g., Jenkins, 2011) beneath transverse surface depressions. We also suggest that these transverse basal channels may offer conduits for meltwater flow between adjacent longitudinal channels, potentially leading to increased circulation velocity and higher melt rates within the transverse depressions.

#### 6.6 Basal melt rate sensitivity to ocean temperature variability

The 2012–2014 GPS data reveal subtle ( $\sim 2\text{--}4\%$ ) changes in velocity over time that appear to display a lagged correlation with observed variations in ocean temperature records from moorings in Pine Island Bay (Christianson et

Deleted: 10–20 cm pole-base

Deleted: E

Deleted: height

Deleted: D

Deleted: at

Deleted: 14

Deleted: These measurements provide some validation for the GPS results, but a direct comparison may be imprudent. [... [15]

Deleted: measurements were taken at a site approximately

Deleted: K. Riverman, personal communication, 2016), within the same longitudinal channel, but much closer to the site of

Deleted: (Figure 2A). Given the complex and evolving

Deleted: surface topography, we also expect local variation in basal topography and associated

Deleted: during the 2012–2014 period.

Deleted: spatial extent (10s to

Deleted:

Deleted: Dh

Deleted: . They are also located over the same set of longitudinal basal channels, and should be exposed to similar sub-shelf circulation.

Deleted: The DEM Dh/Dt maps show enhanced surface elevation change rates, and thus higher basal melt rates, on the north side (left side in Figure 10) of longitudinal basal channels and on the upstream side of transverse depressions. The latter is consistent with increased Dh/Dt observed at the SOW3 and SOW4 sites, located on the upstream side of a transverse depression. .

Deleted: and/or turbulence as water within the upstream meltwater channel first enters the “cavern” of the transverse depression.

Deleted: the

Deleted: depressions

Deleted: serve as

Deleted: connecting

Deleted: the

Deleted: Implications for

Deleted: changes

Deleted: correlated

Deleted: (Figure 1)

al., 2016), (see Figure 1 for location). Our analysis supports the alternative Christianson et al. (2016) hypothesis that these velocity variations are primarily related to upstream grounding evolution (Joughin et al., 2016), and extension across a series of transverse depressions.

Rates of antenna and surface elevation change ( $Dz_{ant}/Dt$  and  $Dz_{surf}/Dt$ ) were essentially constant in time, with no significant variation in inferred basal melt rates for this 2-year time period. If sub-shelf melt rates beneath the GPS array covaried with ocean heat content beyond the shelf front in Pine Island Bay, and had decreased by ~50% as suggested by Dutrieux et al. (2014), a significant change in both  $Dz_{ant}/Dt$  and  $Dz_{surf}/Dt$  would be expected during this period. The lack of any significant deviation suggests that melt rates at these sites were not noticeably affected by observed ocean temperature variability. This finding suggests that either: 1) these sites are not representative of melt rates for the inner shelf (e.g., those near the grounding line), 2) the oceanographic measurements near the PIG ice front are not representative of water circulating beneath these ice-shelf sites, and/or 3) local melt rates are less sensitive to the observed oceanographic changes than previously assumed.

### 6.7 Future work

High-resolution velocity maps derived from sub-meter imagery could potentially constrain local velocity divergence and length scales for observed strain between GPS receivers. In addition, seismic data from stations deployed near the GPS array and regional sites could help constrain the timing and location of rift propagation and grounding/ungrounding events.

High-resolution (<1-km grid) SMB output and improved dynamic firn model output would likely offer an improved understanding of local variability across the GPS array. It may also be possible to further constrain firn-compaction rates, and thus long-term SMB, using relative layer thicknesses observed in CReSIS snow radar measurements (e.g., Medley et al., 2015) or *in-situ* pRES observations (e.g., Jenkins et al., 2006). However, airborne radar data over the PIG shelf suffer from clutter due to km-scale surface/basal topography and crevasses, while the available intermittent pRES records (Stanton et al., 2013) likely lack the sensitivity to detect small changes in firn layer thickness during the ~3-week observation period.

These limitations highlight the current value of long-term GPS records to constrain surface evolution where observations are sparse and model results are poorly constrained. Expanding the scope of our study to include the full archive of geodetic GPS data for the Antarctic and Greenland ice sheets would offer a valuable new dataset for cal/val of models and remote sensing data.

We offer the following recommendations to improve GPS interferometric reflectometry results for future GPS deployments: 1) set GPS elevation mask to 0°, 2) track all possible signals (L2C, L5, Galileo, and GLONASS), 3) ensure that antenna-surface distance will remain >0.5 m between servicing visits, and 4) document and photograph GPS sites, noting antenna-surface distance and any pole tilt during install and servicing.

### 7 Summary and conclusions

We analyzed GPS records for the PIG shelf from 2008–2010 and 2012–2014. We produced daily time series of antenna-surface distance ( $h_{ant-surf}$ ) and antenna elevation ( $z_{ant}$ , relative to EGM2008 geoid), which were combined to

- Deleted: .
- Deleted: ,
- Deleted: (Figure 6).
- Deleted: Both the GPS pole-base  $Dh_{pb}$
- Deleted: surface Dh
- Deleted: fits are highly linear
- Deleted: were directly related to
- Deleted: observed Dh
- Deleted: Dh/Dt
- Deleted: are
- Deleted: significantly
- Deleted: -
- Deleted: greater
- Deleted: (Figure 1)

- Deleted: It may

- Deleted: ice-sheet SMB
- Deleted: model validation.

- Deleted: 2008–2010 and 2012–2014
- Deleted: PIG. These data provide validation for surface elevation and Dh/Dt derived from

accurately measure surface elevation ( $z_{surf}$ ) at each site. The surface elevation data can be directly compared with remote-sensing measurements, providing independent validation for high-resolution WorldView stereo DEM records (RMSE of  $\sim 0.72$  m, NMAD of  $\sim 0.57$  m).

The GPS-derived surface elevation data provide new information about local SMB that can be compared with coarse-resolution model output and AWS data. Surface elevation relative to a firm layer tracer for the initial surface ( $z_{surf} - z_{surf0}$ ) increased at rates of  $\sim 0.8$ – $1.1$  m/yr for all GPS sites, which is consistent with modeled SMB of  $\sim 0.7$ – $0.9$  m w.e./yr and modeled downward firm compaction velocities ( $v_c$ ) of  $\sim 0.70$ – $0.75$  m/yr. An abrupt  $\sim 0.2$ – $0.3$  m surface elevation decrease, likely due to surface melt and/or enhanced firm compaction, is observed across all GPS sites during a period of warmer atmospheric temperatures from December 2012 to January 2013.

Trends in observed surface elevation change ( $Dz_{surf}/Dt$ ) were highly linear for all GPS sites on the PIG shelf. Similar rates with reduced residual variability were obtained after removing simulated pole-base firm compaction velocity ( $v_c$ ) from antenna elevation change ( $Dz_{ant}/Dt$ ), which provides further validation for firm model results. Extensional strain rates were  $\sim 0.001$ – $0.002$  yr<sup>-1</sup> for the 2012–2014 GPS array, which corresponds to only  $\sim 0.1$  m/yr surface elevation change due to local flux divergence.

An alternative form of the mass conservation equation was used to estimate BMB from observed Lagrangian surface elevation change, strain rates, and SMB. Basal melt rates were  $\sim 10$  to  $\sim 40$  m/yr near the center of the fast-flowing PIG shelf, and  $\sim 2$ – $4$  m/yr for the southern shelf. These melt rates are similar to those derived from complementary *in-situ* instrument records (Stanton et al., 2013) and high-resolution stereo DEMs (Shean, 2016).

Both GPS and DEM records show higher basal melt rates within and near transverse surface depressions and rifts associated with longitudinal extension. Basal melt rates for the 2012–2014 period show limited temporal variability, despite significant changes in ocean heat content at the ice front and likely in the ice-shelf cavity. Residual elevation change variability is likely related to upstream grounding/ungrounding events and the local evolution of transverse depressions/rifts. Our results demonstrate the value of long-term GPS records and interferometric reflectometry for constraining ice shelf mass balance estimates.

## 25 Acknowledgements

D. Shean was supported by a NASA NESSF fellowship (NNX12AN36H). K. Christianson was supported by NASA grants NNX16AM01G and NNX12AB69G and NSF grant 0732869. K. Larson was supported by NSF AGS-1449554. S.R.M. Ligtenberg was supported by an NWO ALW Veni grant (865.15.023). An NSF OPP grant to CReSIS (ANT-0424589) provided support for I. Joughin and additional support for D. Shean, D. Holland and M. Bushuk acknowledge support from NSF grant PLR- 0732869 and NYU Abu Dhabi grant G1204. H. Conway provided useful feedback on an earlier version of this manuscript and discussions with P. Dutrieux helped guide interpretation. M. van Wessem, P. Kuipers Munneke, and M. van den Broeke provided RACMO SMB products. The AWS data are available from the University of Wisconsin-Madison Automatic Weather Station Program (NSF ANT-1245663). We acknowledge the substantial effort required to obtain the GPS data used in this study, involving multiple PIG field campaigns led by R. Bindshadler and M. Truffer, with significant contributions from many others. We acknowledge GPS data collection and archiving provided by the UNAVCO Facility with support from NSF and NASA under NSF

Deleted: , with sampled DEM

Deleted: and

Deleted: .

Deleted: antenna height records document a relative surface increase of  $\sim 0.7$ – $1.0$

Deleted: modelled

Deleted: mwe/

Deleted: 7

Deleted: Observed longitudinal strain rates for the 2012–2014 GPS array are  $\sim 0.001$ – $0.002$  yr<sup>-1</sup>, with negligible associated

Deleted: . . . [16]

Deleted: is

Deleted: the PIG shelf sites, with trends of  $-1$  to  $-4$  m/yr and residual RMSE of  $\sim 0.1$  m for the 2012–2014 sites.

Deleted: estimates

Deleted: are

Deleted: downward GPS

Deleted: firm compaction from GPS antenna elevation records. Estimated basal

Deleted: are

Deleted: (Shean, 2016)

Deleted: variable strain rates across

Deleted: S.

Deleted: C.

Deleted: L.

Deleted: L.

Deleted: J.

Deleted: S.

Deleted: , D. Holland,

Cooperative Agreement No. EAR-0735156. Resources supporting the DEM generation were provided by the NASA High-End Computing (HEC) Program through the NASA Advanced Supercomputing (NAS) Division at Ames Research Center. [We thank L. Padman and an anonymous reviewer for their comments/suggestions, which significantly improved this manuscript.](#)

## 5 References

- Andreassen, L. M., Elvehøy, H., Kjølmoen, B. and Engeset, R. V.: Reanalysis of long-term series of glaciological and geodetic mass balance for 10 Norwegian glaciers, *The Cryosphere*, 10(2), 535–552, doi:10.5194/tc-10-535-2016, 2016.
- Arthern, R. J., Vaughan, D. G., Rankin, A. M., Mulvaney, R. and Thomas, E. R.: *In situ* measurements of Antarctic snow compaction compared with predictions of models, *J. Geophys. Res.*, 115(F3), doi:10.1029/2009JF001306, 2010.
- Bassis, J. N. and Ma, Y.: Evolution of basal crevasses links ice shelf stability to ocean forcing, *Earth Planet. Sci. Lett.*, 409, 203–211, doi:10.1016/j.epsl.2014.11.003, 2015.
- Bindschadler, R., Vaughan, D. G. and Vornberger, P.: Variability of basal melt beneath the Pine Island Glacier ice shelf, West Antarctica, *J. Glaciol.*, 57(204), 581–595, 2011.
- Bindschadler, R. A.: History of lower Pine Island Glacier, West Antarctica, from Landsat imagery, *J. Glaciol.*, 48(163), 536–544, 2002.
- [Brunt, K. M., Fricker, H. A., Padman, L., Scambos, T. A. and O’Neel, S.: Mapping the grounding zone of the Ross Ice Shelf, Antarctica, using ICESat laser altimetry, \*Ann. Glaciol.\*, 51\(55\), 71–79, 2010.](#)
- Christianson, K., Bushuk, M., Dutrieux, P., Parizek, B. R., Joughin, I. R., Alley, R. B., Shean, D. E., Abrahamsen, E. P., Anandakrishnan, S., Heywood, K. J., Kim, T.-W., Lee, S. H., Nicholls, K., Stanton, T., Truffer, M., Webber, B. G. M., Jenkins, A., Jacobs, S., Bindschadler, R. and Holland, D. M.: Sensitivity of Pine Island Glacier to observed ocean forcing: PIG response to ocean forcing, *Geophys. Res. Lett.*, 43(20), 10,817–10,825, doi:10.1002/2016GL070500, 2016.
- Church, J. A., Cazenave, A., Gregory, J. M., Jevrejeva, S., Levermann, A., Merrifield, M. A., Milne, G. A., Nerem, R. S., Nunn, P. D., Payne, A. J., Pfeffer, W. T., Stammer, D. and Unnikrishnan, A. S.: Sea Level Change. [online] Available from: [http://www.ipcc.ch/pdf/assessment-report/ar5/wg1/WG1AR5\\_Chapter13\\_FINAL.pdf](http://www.ipcc.ch/pdf/assessment-report/ar5/wg1/WG1AR5_Chapter13_FINAL.pdf) (Accessed 7 January 2015), 2013.
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, I., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J.-N. and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Q. J. R. Meteorol. Soc.*, 137(656), 553–597, doi:10.1002/qj.828, 2011.

Deleted: -

- Depoorter, M. A., Bamber, J. L., Griggs, J. A., Lenaerts, J. T. M., Ligtenberg, S. R. M., van den Broeke, M. R. and Moholdt, G.: Calving fluxes and basal melt rates of Antarctic ice shelves, *Nature*, doi:10.1038/nature12567, 2013.
- [Drews, R.: Evolution of ice-shelf channels in Antarctic ice shelves, \*The Cryosphere\*, 9\(3\), 1169–1181, doi:10.5194/tc-9-1169-2015, 2015.](#)
- 5 Dutriex, P., Vaughan, D. G., Corr, H. F. J., Jenkins, A., Holland, P. R., Joughin, I. and Fleming, A. H.: Pine Island glacier ice shelf melt distributed at kilometre scales, *The Cryosphere*, 7(5), 1543–1555, doi:10.5194/tc-7-1543-2013, 2013.
- Dutriex, P., De Rydt, J., Jenkins, A., Holland, P. R., Ha, H. K., Lee, S. H., Steig, E. J., Ding, Q., Abrahamsen, E. P. and Schroder, M.: Strong Sensitivity of Pine Island Ice-Shelf Melting to Climatic Variability, *Science*, 343(6167), 174–178, doi:10.1126/science.1244341, 2014.
- 10 Ettema, J., van den Broeke, M. R., van Meijgaard, E., van de Berg, W. J., Bamber, J. L., Box, J. E. and Bales, R. C.: Higher surface mass balance of the Greenland ice sheet revealed by high-resolution climate modeling, *Geophys. Res. Lett.*, 36(12), doi:10.1029/2009GL038110, 2009.
- Frezzotti, M., Searchilli, C., Becagli, S., Proposito, M. and Urbini, S.: A synthesis of the Antarctic surface mass balance during the last 800 yr, *The Cryosphere*, 7(1), 303–319, doi:10.5194/tc-7-303-2013, 2013.
- 15 Gao, J. and Liu, Y.: Applications of remote sensing, GIS and GPS in glaciology: a review, *Prog. Phys. Geogr.*, 25(4), 520–540, 2001.
- [Griggs, J. A. and Bamber, J. L.: Antarctic ice-shelf thickness from satellite radar altimetry, \*J. Glaciol.\*, 57\(203\), 485, 2011.](#)
- 20 Hamilton, G. S. and Whillans, I. M.: First point measurements of ice-sheet thickness change in Antarctica, *Ann. Glaciol.*, 27(1), 125–129, 1998.
- Hawley, R. L. and Waddington, E. D.: Instruments and Methods *In situ* measurements of firn compaction profiles using borehole optical stratigraphy, *J. Glaciol.*, 57(202), 289–294, 2011.
- Herring, T. A., King, R. W. and McClusky, S. C.: Introduction to GAMIT/GLOBK, [online] Available from: [http://chandler.mit.edu/~simon/gtgk/Intro\\_GG.pdf](http://chandler.mit.edu/~simon/gtgk/Intro_GG.pdf) (Accessed 23 November 2016), 2015.
- 25 [Herron, M. M. and Langway, C. C., Jr.: Firn densification: an empirical model, \*J. Glaciol.\*, 25, 373–385, 1980.](#)
- Jacobs, S. S., Jenkins, A., Giulivi, C. F. and Dutriex, P.: Stronger ocean circulation and increased melting under Pine Island Glacier ice shelf, *Nat. Geosci.*, 4(8), 519–523, doi:10.1038/ngeo1188, 2011.
- Jenkins, A.: Convection-Driven Melting near the Grounding Lines of Ice Shelves and Tidewater Glaciers, *J. Phys. Oceanogr.*, 41(12), 2279–2294, doi:10.1175/JPO-D-11-03.1, 2011.
- 30 Jenkins, A., Corr, H. F., Nicholls, K. W., Stewart, C. L. and Doake, C. S.: Interactions between ice and ocean observed with phase-sensitive radar near an Antarctic ice-shelf grounding line, *J. Glaciol.*, 52(178), 325–346, 2006.
- Jeong, S., Howat, I. M. and Bassis, J. N.: Accelerated ice shelf rifting and retreat at Pine Island Glacier, West Antarctica: Central PIG rifting and calving, *Geophys. Res. Lett.*, doi:10.1002/2016GL071360, 2016.
- 35 Jones, R. W., Renfrew, I. A., Orr, A., Webber, B. G. M., Holland, D. M. and Lazzara, M. A.: Evaluation of four global reanalysis products using *in situ* observations in the Amundsen Sea Embayment, Antarctica: Amundsen Sea Reanalyses Evaluation, *J. Geophys. Res. Atmospheres*, 121(11), 6240–6257, doi:10.1002/2015JD024680, 2016.

Deleted: -

Deleted: -

- Joughin, I., Smith, B. E. and Holland, D. M.: Sensitivity of 21st century sea level to ocean-induced thinning of Pine Island Glacier, Antarctica, *Geophys. Res. Lett.*, 37(20), n/a-n/a, doi:10.1029/2010GL044819, 2010.
- Joughin, I., Shean, D. E., Smith, B. E. and Dutrieux, P.: Grounding line variability and subglacial lake drainage on Pine Island Glacier, Antarctica, *Geophys. Res. Lett.*, 43(17), 9093–9102, doi:10.1002/2016GL070259, 2016.
- 5 [Langley, K., von Deschanden, A., Kohler, J., Sinisalo, A., Matsuoka, K., Hattermann, T., Humbert, A., Nøst, O. A. and Isaksson, E.: Complex network of channels beneath an Antarctic ice shelf, \*Geophys. Res. Lett.\*, n/a-n/a, doi:10.1002/2013GL058947, 2014.](#)
- [Larson, K. M.: GPS interferometric reflectometry: applications to surface soil moisture, snow depth, and vegetation water content in the western United States: GPS interferometric reflectometry, \*Wiley Interdiscip. Rev. Water\*, 3\(6\), 775–787, doi:10.1002/wat2.1167, 2016.](#)
- 10 Larson, K. M. and Nievinski, F. G.: GPS snow sensing: results from the EarthScope Plate Boundary Observatory, *GPS Solut.*, 17(1), 41–52, doi:10.1007/s10291-012-0259-7, 2013.
- Larson, K. M., Wahr, J. and Kuipers Munneke, P.: Constraints on snow accumulation and firn density in Greenland using GPS receivers, *J. Glaciol.*, 61(225), 101–114, doi:10.3189/2015JoG14J130, 2015.
- 15 Lazzara, M. A., Weidner, G. A., Keller, L. M., Thom, J. E. and Cassano, J. J.: Antarctic Automatic Weather Station Program: 30 Years of Polar Observation, *Bull. Am. Meteorol. Soc.*, 93(10), 1519–1537, doi:10.1175/BAMS-D-11-00015.1, 2012.
- Lenaerts, J. T. M., van den Broeke, M. R., van de Berg, W. J., van Meijgaard, E. and Kuipers Munneke, P.: A new, high-resolution surface mass balance map of Antarctica (1979–2010) based on regional atmospheric climate modeling, *Geophys. Res. Lett.*, 39(4), n/a-n/a, doi:10.1029/2011GL050713, 2012.
- 20 Ligtenberg, S. R. M., Helsen, M. M. and van den Broeke, M. R.: An improved semi-empirical model for the densification of Antarctic firn, *The Cryosphere*, 5(4), 809–819, doi:10.5194/tc-5-809-2011, 2011.
- Medley, B., Joughin, I., Smith, B. E., Das, S. B., Steig, E. J., Conway, H., Gogineni, S., Lewis, C., Criscitiello, A. S., McConnell, J. R., van den Broeke, M. R., Lenaerts, J. T. M., Bromwich, D. H., Nicolas, J. P. and Leuschen, C.:
- 25 Constraining the recent mass balance of Pine Island and Thwaites glaciers, West Antarctica, with airborne observations of snow accumulation, *The Cryosphere*, 8(4), 1375–1392, doi:10.5194/tc-8-1375-2014, 2014.
- Medley, B., Ligtenberg, S. R. M., Joughin, I., Van Den Broeke, M. R., Gogineni, S. and Nowicki, S.: Antarctic firn compaction rates from repeat-track airborne radar data: I. Methods, *Ann. Glaciol.*, 56(70), 155–166, doi:10.3189/2015AoG70A203, 2015.
- 30 [Moholdt, G., Padman, L. and Fricker, H. A.: Basal mass budget of Ross and Filchner-Ronne ice shelves, Antarctica, derived from Lagrangian analysis of ICESat altimetry: Ice shelf basal melting from altimetry, \*J. Geophys. Res. Earth Surf.\*, n/a-n/a, doi:10.1002/2014JF003171, 2014.](#)
- Mouginot, J., Rignot, E. and Scheuchl, B.: Sustained increase in ice discharge from the Amundsen Sea Embayment, West Antarctica, from 1973 to 2013, *Geophys. Res. Lett.*, 41(5), 1576–1584, doi:10.1002/2013GL059069, 2014.
- 35 Nievinski, F. G.: Forward and inverse modeling of GPS multipath for snow monitoring, University of Colorado. [online] Available from: [http://xenon.colorado.edu/portal/publications/Nievinski\\_2013\\_PhD.pdf](http://xenon.colorado.edu/portal/publications/Nievinski_2013_PhD.pdf) (Accessed 9 December 2016), 2013.

- Padman, L., Fricker, H. A., Coleman, R., Howard, S. and Erofeeva, L.: A new tide model for the Antarctic ice shelves and seas, *Ann. Glaciol.*, 34(1), 247–254, 2002.
- Padman, L., King, M., Goring, D., Corr, H. and Coleman, R.: Ice-shelf elevation changes due to atmospheric pressure variations, *J. Glaciol.*, 49(167), 521–526, doi:10.3189/172756503781830386, 2003.
- 5 Pavlis, N. K., Holmes, S. A., Kenyon, S. C. and Factor, J. K.: The development and evaluation of the Earth Gravitational Model 2008 (EGM2008), *J. Geophys. Res.*, 117(B4), doi:10.1029/2011JB008916, 2012.
- Pritchard, H. D., Arthern, R. J., Vaughan, D. G. and Edwards, L. A.: Extensive dynamic thinning on the margins of the Greenland and Antarctic ice sheets, *Nature*, 461(7266), 971–975, doi:10.1038/nature08471, 2009.
- Pritchard, H. D., Ligtenberg, S. R. M., Fricker, H. A., Vaughan, D. G., van den Broeke, M. R. and Padman, L.: Antarctic ice-sheet loss driven by basal melting of ice shelves, *Nature*, 484(7395), 502–505, doi:10.1038/nature10968, 2012.
- 10 Rietbroek, R., Brunnabend, S.-E., Kusche, J., Schröter, J. and Dahle, C.: Revisiting the contemporary sea-level budget on global and regional scales, *Proc. Natl. Acad. Sci.*, 201519132, 2016.
- Rignot, E.: Changes in West Antarctic ice stream dynamics observed with ALOS PALSAR data, *Geophys Res Lett.*, 35(12), L12505, doi:10.1029/2008GL033365, 2008.
- 15 Rignot, E., Jacobs, S., Mouginot, J. and Scheuchl, B.: Ice-Shelf Melting Around Antarctica, *Science*, 341(6143), 266–270, doi:10.1126/science.1235798, 2013.
- Rignot, E., Mouginot, J., Morlighem, M., Seroussi, H. and Scheuchl, B.: Widespread, rapid grounding line retreat of Pine Island, Thwaites, Smith, and Kohler glaciers, West Antarctica, from 1992 to 2011, *Geophys. Res. Lett.*, 20 41(10), 3502–3509, doi:10.1002/2014GL060140, 2014.
- [Savage, J. C., Gan, W. and Svarc, J. L.: Strain accumulation and rotation in the Eastern California Shear Zone, \*J. Geophys. Res. Solid Earth\*, 106\(B10\), 21995–22007, doi:10.1029/2000JB000127, 2001.](#)
- Scott, J., Gudmundsson, H., Smith, A., Bingham, R., Pritchard, H. and Vaughan, D.: Increased rate of acceleration on Pine Island Glacier strongly coupled to changes in gravitational driving stress, *The Cryosphere*, 3, 125–131, 2009.
- 25 [Shabtaie, S. and Bentley, C. R.: Tabular icebergs: implications from geophysical studies of ice shelves, \*J. Glaciol.\*, 28\(100\), 1982.](#)
- Shean, D.: Quantifying ice-shelf basal melt and ice-stream dynamics using high-resolution DEM and GPS time series, [Ph.D.](#) Thesis, University of Washington, Seattle, WA, 14 July. [online] Available from: <https://digital.lib.washington.edu:443/researchworks/handle/1773/36365> (Accessed 22 November 2016), 2016.
- 30 Shean, D. E., Alexandrov, O., Moratto, Z. M., Smith, B. E., Joughin, I. R., Porter, C. and Morin, P.: An automated, open-source pipeline for mass production of digital elevation models (DEMs) from very-high-resolution commercial stereo satellite imagery, *ISPRS J. Photogramm. Remote Sens.*, 116, 101–117, doi:10.1016/j.isprsjprs.2016.03.012, 2016.
- Shepherd, A., Ivins, E. R., A, G., Barletta, V. R., Bentley, M. J., Bettadpur, S., Briggs, K. H., Bromwich, D. H., Forsberg, R., Galin, N., Horwath, M., Jacobs, S., Joughin, I., King, M. A., Lenaerts, J. T. M., Li, J., Ligtenberg, S. R. M., Luckman, A., Luthcke, S. B., McMillan, M., Meister, R., Milne, G., Mouginot, J., Muir, A., Nicolas, J. P., Paden, J., Payne, A. J., Pritchard, H., Rignot, E., Rott, H., Sorensen, L. S., Scambos, T. A., Scheuchl, B.,
- 35

- Schrama, E. J. O., Smith, B., Sundal, A. V., van Angelen, J. H., van de Berg, W. J., van den Broeke, M. R., Vaughan, D. G., Velicogna, I., Wahr, J., Whitehouse, P. L., Wingham, D. J., Yi, D., Young, D. and Zwally, H. J.: A Reconciled Estimate of Ice-Sheet Mass Balance, *Science*, 338(6111), 1183–1189, doi:10.1126/science.1228102, 2012.
- 5 Stanton, T. P., Shaw, W. J., Truffer, M., Corr, H. F. J., Peters, L. E., Riverman, K. L., Bindschadler, R., Holland, D. M. and Anandakrishnan, S.: Channelized Ice Melting in the Ocean Boundary Layer Beneath Pine Island Glacier, Antarctica, *Science*, 341(6151), 1236–1239, doi:10.1126/science.1239373, 2013.
- Van Meijgaard, E., Van Uft, L. H., Van de Berg, W. J., Bosveld, F. C., Van den Hurk, B., Lenderink, G. and Siebesma, A. P.: The KNMI regional atmospheric climate model RACMO version 2.1, Koninklijk Nederlands  
10 Meteorologisch Instituut. [online] Available from:  
[http://www.weeralarm.nl/publications/fulltexts/tr302\\_racmo2v1.pdf](http://www.weeralarm.nl/publications/fulltexts/tr302_racmo2v1.pdf) (Accessed 31 May 2013), 2008.
- Van Wessem, J. M., Reijmer, C. H., Morlighem, M., Mougnot, J., Rignot, E., Medley, B., Joughin, I., Wouters, B., Depoorter, M. A., Bamber, J. L., Lenaerts, J. T. M., De Van Berg, W. J., Van Den Broeke, M. R. and Van  
15 Meijgaard, E.: Improved representation of East Antarctic surface mass balance in a regional atmospheric climate model, *J. Glaciol.*, 60(222), 761–770, doi:10.3189/2014JG14J051, 2014.
- Vaughan, D. G., Corr, H. F. J., Bindschadler, R. A., Dutrieux, P., Gudmundsson, G. H., Jenkins, A., Newman, T., Vornberger, P. and Wingham, D. J.: Subglacial melt channels and fracture in the floating part of Pine Island Glacier, Antarctica, *J. Geophys. Res.*, 117(F3), doi:10.1029/2012JF002360, 2012.
- [Webber, B. G. M., Heywood, K. J., Stevens, D. P., Dutrieux, P., Abrahamsen, E. P., Jenkins, A., Jacobs, S. S., Ha, H. K., Lee, S. H. and Kim, T. W.: Mechanisms driving variability in the ocean forcing of Pine Island Glacier, \*Nat. Commun.\*, 8, 14507, doi:10.1038/ncomms14507, 2017.](#)
- 20 Wouters, B., Martin-Espanol, A., Helm, V., Flament, T., van Wessem, J. M., Ligtenberg, S. R. M., van den Broeke, M. R. and Bamber, J. L.: Dynamic thinning of glaciers on the Southern Antarctic Peninsula, *Science*, 348(6237), 899–903, doi:10.1126/science.aaa5727, 2015.

25

Tables

Site	Time period	Days	$D(z_{surf} - z_{surf0})/Dt$ (m/yr)	$Dz_{ant}/Dt$ (m/yr)	$v_{fc}$ (m/yr)	$Dz_{surf}/Dt$ (m/yr)	$Dz_{ant}/Dt - v_{fc}$ (m/yr)	$b$ (m/yr)	$b$ (m/yr)
								from $Dz_{surf}/Dt$	from $Dz_{ant}/Dt$
PIG1	2008-1-13, 2009-9-4	601	0.93	-7.60*	-0.75	-6.76*	-6.85*	--	--
PIG2	2008-1-10, 2010-1-27	747	1.12	-1.12	-0.74	-0.13	-0.38	2.0 +/- 0.9	4.4 +/- 1.4
SOW1	2012-2-10, 2013-12-23	714	0.77	-1.81	-0.75	-1.13	-1.06	11.5 +/- 1.1	10.8 +/- 1.5
SOW2	2012-2-10, 2013-12-23	714	0.85	-2.08	-0.74	-1.33	-1.34	13.3 +/- 1.2	13.4 +/- 1.5
BOAR	2012-2-10, 2013-4-29	476	0.78	-1.58	-0.77	-0.91	-0.81	9.4 +/- 1.1	8.5 +/- 1.4
SOW4	2012-2-10, 2013-12-24	714	0.86	-3.76	-0.73	-3.00	-3.03	29.1 +/- 1.7	29.3 +/- 1.9
SOW3	2012-2-9, 2013-12-24	716	1.10	-5.23	-0.72	-4.10	-4.50	39.4 +/- 2.1	43.1 +/- 2.4

Table 1: GPS station data. Fields include surface elevation change relative to tracer for initial surface  $D(z_{surf} - z_{surf0})/Dt$ , antenna elevation change  $Dz_{ant}/Dt$  (equal to pole-base elevation change), modeled downward pole-base vertical velocity due to firn compaction  $v_{fc}$ , surface elevation change  $Dz_{surf}/Dt$ , and antenna  $Dz_{ant}/Dt$  after removing  $v_{fc}$  and corresponding ice-equivalent basal melt rates  $b$  computed using equations 5 and 7, respectively. \*Note: PIG1 values over grounded ice do not include correction to remove expected  $Dz_{surf}/Dt$  due to advection along local surface slopes ( $V_0$ ).

Site	DEM n	DEM dt (days)	DEM $Dz_{surf}/Dt$ (m/yr)	GPS-DEM RMSE (m)	GPS-DEM mean (m)	GPS-DEM std (m)
SOW1	5	302*	-2.30	0.69	-0.26	0.64
SOW2	5	619	-2.03	0.76	-0.46	0.60
BOAR	4	302*	-1.69	0.86	-0.55	0.66
SOW4	5	368*	-3.35	0.75	-0.61	0.44
SOW3	6	619	-4.32	0.54	-0.30	0.45

Table 2: Statistics for WorldView DEM accuracy from comparisons with measured GPS surface elevation data. Asterisks identify records with shorter time interval and increased uncertainty.

- Deleted:  $b_{ho}$  (m) ... [17]
- Deleted:  $z_{pb0}$  (m) ... [18]
- Deleted:  $Dz_{th}$  ... [18]
- Deleted:  $Dh_0/Dt, Dh_{pb}$
- Deleted:  $Dh_{pb}$
- Deleted:  $Dh$
- Deleted: Surf.  $b$
- Deleted: Pole-base  $b$
- Deleted: 484.28
- Deleted Cells ... [19]
- Deleted: 71
- Deleted: 0.83
- Deleted Cells ... [20]
- Deleted: 76.09
- Deleted Cells ... [21]
- Deleted: 0.78
- Deleted Cells ... [22]
- Deleted: 02
- Deleted: 1+/-1.
- Deleted: 5...+/-1.7 ... [23]
- Deleted: 67.78
- Deleted Cells ... [24]
- Deleted: 61
- Deleted: 0.67
- Deleted Cells ... [25]
- Deleted: 6...+/-1.2 ... [26]
- Deleted: 9...+/-1.8 ... [27]
- Deleted: 64.72
- Deleted: 89
- Deleted: 0.75
- Deleted: 4...+/-1.3 ... [28]
- Deleted: 5...+/-1.9 ... [29]
- Deleted: 66.44
- Deleted: 82
- Deleted: 0.68
- Deleted: 5
- Deleted: 6...+/-1.8 ... [30]
- Deleted: 69.66
- Deleted Cells ... [31]
- Deleted: 1.08
- Deleted Cells ... [32]
- Deleted: 76
- Deleted: 2+/-2.
- Deleted: 4+/-2.5
- Deleted: 63.16
- Deleted: 1.16
- Deleted Cells ... [33]
- Deleted: 1.00
- Deleted Cells ... [34]
- Deleted: 5...+/-2.7 ... [35]
- Deleted: +/-3.2
- Deleted: and  $Dh/Dt$  results.... Fields include surface e ... [36]
- Deleted: for both the surface  $Dh/Dt$  and corrected pole ... [37]
- Deleted:  $Dh$

Figures




Figure 1: Context for Pine Island Glacier ice shelf with 2006–2016 median surface velocity (Christianson et al., 2016; Joughin et al., 2010) over shaded relief map from October–December 2012 DEM mosaic. Black lines show ~2-year paths between initial (green) and final (red) GPS station locations. Yellow dot shows Evans Knoll AWS and blue squares show RACMO grid cell centers used during analysis. Purple triangles beyond shelf front show locations of ocean mooring temperature data from Christianson et al. (2016). White line shows approximate 2011 grounding line (Shean, 2016). Black rectangle shows location of Figure 2A.






Figure 2: A) WorldView DEM from November 11, 2012 with 2012–2014 GPS array positions overlaid. Note GPS positions relative to transverse depressions and location of R1 rift associated with 2015 calving event (black arrow). Ice flow direction indicated by white arrow. White lines show locations of profiles. B) Smoothed surface elevation (0.5 km window, approx.  $\sim 1H_{ice}$ ) and estimated freeboard thickness for longitudinal profile X-X' and C) transverse profile Y-Y'. Profile intersection is near BOAR (red point). Vertical exaggeration is 22x.



Deleted: -  
Deleted: Evan's  
Deleted: (Shean, 2016)



Deleted: is from top right corner to bottom left corner.  
Deleted: (~  
Deleted: kernel,  
Deleted: H




Figure 3: A) Original GPS antenna elevation (light gray), after tide correction (mid-gray), and after tide+IBE correction (black) for SOW4. Red line shows smoothed time series and yellow dashed line is linear fit (-3.76 m/yr). Sampled DEM elevations (cyan) show surface elevation, which is offset from GPS antenna elevation by antenna surface distance (see Figures 4 and 8). B) High-frequency (<1.5 days) component of GPS record and CATS2008A tide model prediction, showing excellent agreement. C) Estimated Inverse Barometer Effect (IBE) magnitude from scaled sea-level pressure.

Deleted: absolute

Deleted: height





Figure 4: Schematic of GPS station geometry. Surface elevation ( $z_{surf}$ , dark blue line) is computed by removing antenna surface distance ( $h_{ant-surf}$ , dotted black line) from antenna elevation ( $z_{ant}$ , black line). Pole-base elevation (red line) is computed from pole length and antenna phase-center offset. At time  $t_1$  (right panel), ongoing firn compaction resulted in decreased antenna and pole-base elevation, while new snow accumulation offset surface lowering. The layer within the firn column corresponding to the initial surface ( $z_{surf0}$ ) is represented by dotted blue line.



- Deleted:
- Deleted: Absolute surface
- Deleted: h
- Deleted: multi-path
- Deleted: height above
- Deleted: ( $z_a$ )
- Deleted: absolute
- Deleted:  $h_a$ , black line). With known pole length, we can calculate pole-base depth below the surface ( $z_{pb}$ ) and absolute pole
- Deleted:  $h_{pb}$ ,
- Deleted: ).
- Deleted: resulted in increased
- Deleted: elevation. Reflector height ( $z_a$ ) is measured relative
- Deleted:  $h_a$  (
- Deleted: ).







Figure 5: Station velocities derived from daily mean positions for A) 2008–2010 GPS sites, and B) 2012–2014 GPS sites. Note offset between SOW2 and SOW4, indicative of lateral shear across the ~2 km wide array, with greater extension near the center of the PIG shelf.



**Figure 6:** Horizontal strain rates over 42-day periods for 8 different triangular sections within the GPS array. **A)** First principal strain rate, positive for extension. Error bars calculated for uncorrelated GPS position error of 1 cm. **B)** Second principal strain rate. **C)** Annualized velocity magnitude. Diagrams in right column show color-coded triangular sections, with  $\sim 2$ -year mean of principal strain rates plotted at centroids.






Figure 7: **A)** Observed antenna-surface distance ( $h_{ant-surf}$ ) for each station. Legend lists original distance. **B)** Observed antenna elevation ( $z_{ant}$ ) relative to initial absolute antenna elevation values listed in legend. Dashed black lines show modeled IMAU-FDM downward velocity due to firn compaction ( $v_{fc}$ ) at the pole base. **C)** Calculated surface elevation ( $z_{surf}$ ) and simulated IMAU-FDM surface elevation from SMB/firn ( $z_{surf}^{FDM}$ , thin black lines), both relative to initial absolute surface elevation values listed in legend.





Figure 8: Time series of GPS antenna elevation (black), surface elevation (thick blue), tracer for initial surface (dotted blue), and pole-base elevation (red), all relative to initial absolute surface elevation. See schematic in Figure 4. Green points show sampled WorldView DEM surface elevation. Note surface elevation decrease at all sites but PIG2.

Deleted: Filtered (solid line) and original (transparent) GPS

Deleted: elevation ( $h_a$ )

Deleted: (first item in legend). Legend includes linear  $Dh_a/Dt$  fit to filtered antenna elevations.

Deleted: estimated



Deleted:

Deleted: pole-base anchor elevation (red),

Deleted: and


Deleted: the

Deleted: GPS


Deleted: for schematic of different values.

Deleted: , with observed DEM RMSE of 0.72 and NMAD of 0.57 m. Surface

Deleted: decreased



**Figure 9:** A) Detrended GPS antenna elevation ( $z_{ant}$ ; see Figure 7B for original records) after removing  $v_c$  with arbitrary y-axis offset. Legend lists linear trend, which can be compared with  $Dz_{surf}/Dt$  trend in panel B (see Equation 6). B) Detrended surface elevation ( $z_{surf}$ ; see Figure 7C) and detrended IMAU-FDM simulated surface elevation ( $\hat{z}_{surf}$ ), with arbitrary y-axis offset. Legend lists linear trend. Note limited residual magnitude and dampened seasonal signal of  $z_{ant}$  compared to  $\hat{z}_{surf}$ . Unlike  $\hat{z}_{surf}$ , no significant change is observed in  $z_{ant}$  from Dec. 2012 to Jan. 2013. C) Surface elevation relative to tracer for initial surface ( $z_{surf} - z_{surf0}$ ). As annotated, positive slopes are indicative of new snow accumulation, shallow negative slopes indicate ongoing compaction, and steep negative slopes likely indicate surface melt. Note  $-0.2$ – $0.3$  m surface decrease from December 2012 to January 2013. Legend values show linear fit at each site. D) Daily and monthly RACMO2.3 SMB. Note correlation of accumulation events and increases in panel C. E) Scaled 2-m temperature data from Evans Knoll AWS (black) and ERA-Interim (gray), with above-zero AWS temperatures plotted in red. Note extended warm period from mid-Dec. 2012 to mid-Jan. 2013, which corresponds to  $-0.2$ – $0.3$  m surface elevation decrease in Panels B and C.



Deleted:

Deleted: Plots of observed and modelled values for GPS sites (see Figure 4 for reference). A) Reflector height ( $z_{ant}$ ) relative to initial surface ( $h_0$ ). Positive values indicate surface height increase relative to GPS antenna. Legend indicates initial antenna height above surface. Note  $\sim 20$ – $30$  cm surface decrease from December 2012 to January 2013. B...) Detrended reflector height ( $z_{ant}$ ), with linear fit listed in legend. C) Surface elevation  $h$ , with initial surface height (m above EGM2008 geoid) in legend. Black lines show IMAU-FDM modelled surface elevation ( $\hat{h}$ ) due to SMB and firn compaction. Note that due to basal melting, observed  $dh/dt$  for 2012–2014 sites is significantly greater than modelled  $d\hat{h}/dt$ . D) Detrended surface elevation  $h$  and detrended IMAU-FDM  $\hat{h}$ , with arbitrary y-axis offset. Legend lists linear fit (corresponding values listed in Table 1). E) Detrended GPS pole-base ... [39]

Deleted:  $h_{pole}$ ... see Figure 7B for original records) after removing  $v_c$  (see ... with arbitrary y-axis offset. Legend lists linear trend, which can be compared with  $Dz_{surf}/Dt$  trend in panel B (see Equation 6). B) Detrended surface elevation ( $z_{surf}$  see Figure 7), ... [40]

Deleted: fit, which can be directly compared with  $Dh/Dt$  fits in D (see Table 1). Note reduced... trend. Note limited residual magnitude and dampened... seasonal signal of  $z_{ant}$  compared to  $D...$ . Unlike surface records... no significant change is observed in  $z_{ant}$  from Dec. 2012 to Jan. 2013. F) Daily ... [41]




Figure 10: WorldView DEMs and composite Lagrangian  $D_{z_{surf}}/Dt$  products generated using A-B) initial DEM from February 2, 2012, and C-D) initial DEM from October 23, 2012. Note enhanced thinning observed within transverse depressions and rift upstream of GPS array. The  $D_{z_{surf}}/Dt$  maps are used to calculate basal melt rates (scaling factor of  $\sim 9$ , e.g.,  $D_{z_{surf}}/Dt$  of  $\sim 1$  m/yr corresponds to a basal melt rate estimate of  $\sim 9$  m/yr).

Deleted: median

Deleted: Dh

Deleted: Dh

Deleted: .0

Deleted: Dh/Dt  $\approx$

Deleted: melt rate

5

Horizontal velocities for each GPS station were computed from daily mean positions. Relative distances between stations were used to calculate strain, with linear fits to estimate strain rates.

### GPS corrections

The constant 3.66 m pole length and 5.32 cm antenna phase center offset was removed from filtered absolute antenna elevation ( $h_a$ ) to estimate corresponding absolute pole-base elevation ( $h_{pb}$ ) relative to the EGM2008 geoid. Figure 4 shows a schematic of this geometry with additional observables as described below.

### GPS antenna height and surface elevation

We neglect the slight reduction in total ice-equivalent thickness at the pole base ( $h_{pb}$ ) vs. surface ( $h$ ), as pole-base depth is negligible compared to total ice thickness ( $z_{pb} \ll H$ ). For the local flux divergence term, we use local freeboard thickness (as in Equation 7) or  $H$  from radar measurements (Stanton et al., 2013). Uncertainty is estimated as  $\sim 0.15$  m/yr for downward firn-compaction velocity (Ligtenberg et al., 2011),  $\sim 0.1$  m/yr for elevation change associated with local flux divergence, and  $\sim 5$  kg/m<sup>3</sup> for the density of ice and ocean water.

If modelled pole-base velocities are

These longitudinal strain rates correspond to shelf thinning rates of  $\sim 0.5$ – $0.9$  m/yr assuming no lateral spreading, with limited expected surface  $Dh/Dt$  ( $< 0.07$ – $0.13$  m/yr) for measured ice thickness of  $\sim 450$ – $460$  m (Stanton et al., 2013) or for thickness estimated from local freeboard ( $\sim 430$ – $500$  m).

### Downslope flow

In addition to elevation change associated with strain rates and corresponding local flux divergence, some component of observed  $Dh/Dt$  may be related to deformation due to local gradients in the driving stress. Over grounded ice, this component of  $Dh/Dt$  is typically dominated by advection over basal topography, and the vertical component of the corresponding motion ( $V_0$ ) can be estimated using observed horizontal GPS displacement and local surface gradients from an independent DEM (e.g., Larson et al., 2015).

While this approach is irrelevant for a freely-floating ice shelf in hydrostatic equilibrium, we attempt to estimate an upper bound for the vertical component of  $Dh/Dt$  due to diffusion of local topography. To do this, we again consider observed relative horizontal displacements within the 2012–2014 GPS array.

### GPS antenna and surface elevation change

Figure 7 shows

### Antenna height evolution

Assuming that the pole base remains fixed within its original firn layer (effectively behaving as an isochron tracer in the firn column), any observed decrease in antenna height above the

1.1

( $h_0$ ).

Initial antenna heights ( $z_a$ ) were ~2.5 to 3.1 m above the surface, indicating that initial pole-base depths ( $z_{pb}$ ) were ~0.6 to 1.2 m below the surface (Figure 8). Antenna height above the surface at all sites decreased over time (Figure 8), with observed  $Dz_{rh}/Dt$

At both PIG1 and PIG2, there are periods of relatively rapid reflector height increase (e.g., from May–August 2008), followed by a steady decrease (e.g., August 2008–February 2009). This is consistent with periods of increased snow accumulation followed by several months of ongoing firn compaction with limited snowfall. The 2012–2014 records show similar periods of abrupt increase and steady decrease, with more limited duration.

All records show an abrupt reflector height decrease (~0.2–0.3 m) between December 2012 and January 2013, which is consistent with surface melting and/or enhanced firn compaction rates in the upper few meters of the firn column (see Section 3.7).

### High-resolution DEMs

In addition to the GPS elevation data, we generated WorldView/GeoEye stereo DEMs (Shean et al., 2016) with 32-m posting over the PIG shelf

The 1979–2015 RACMO average SMB over the central PIG shelf is ~0.9 mwe/yr. Monthly SMB climatology shows low accumulation rates of ~0.01–0.04 mwe/month over the PIG shelf during the austral summer (November to February), and high accumulation rates of ~0.08–0.1 mwe/month during austral winter (March to October) (Figure 9F).

2

increases in surface reflector height. By contrast, the reflector height records show a steady decrease due to ongoing near-surface firn compaction during extended periods with little/no accumulation.

The ~3-4 week period between December 24, 2012 and January 17, 2013 was relatively warm, with scaled AWS atmospheric temperatures of ~1–5°C for most days (Figure 9F). Surface elevations decreased by ~0.2–0.3 m across the entire GPS array during this period (Figure 9A), which is consistent with significant surface melting. The daily RACMO SMB data also show two accumulation events during the last week of December 2012 (Figure 9E), which likely involve rain on snow. Some component of the observed surface elevation decrease could be associated with enhanced near-surface melting and firn compaction during these events. No corresponding short-term changes were recorded by the pole-base GPS elevations during the ~3-4 week period, suggesting that the processes responsible for the observed surface changes were limited to the upper ~1.5 m.

An analysis of 1979-2015 ERA-Interim 2-m air temperatures over the PIG shelf shows many previous warm periods with greater magnitude and duration than the 2012/2013 period. However, the 2012/2013 warm period stands out in the past decade, with the most recent comparable periods during 2005/2006 and 2006/2007.

### Firn model results

Estimated downward pole-base velocities due to firn compaction ( $v_{fc}$ ) were all ~0.70–0.75 m/yr (Figure 7, Table 1) from 2008–2010 and 2012–2014, despite the range of initial pole depths. A slight decrease in the compaction rate occurred over time, but the curves appear linear at these depths (Figure 7). These values are consistent with observed  $Dz_{rh}/Dt$  and  $Dh/Dt$  during periods with little or no surface accumulation

**Figure 9C shows that the IMAU-FDM simulated surface elevation ( $\tilde{h}$ )**

3

**Page 12: [12] Deleted** **David Shean** **7/1/17 5:59:00 PM**

The observed  $D\tilde{h}/Dt$  trend is +0.17 m/yr from 2008-2010, with no significant trend from 2012-2014. The magnitude and timing of the modelled surface elevation variability is consistent with the detrended observed surface elevation change (Figure 9D). The observed  $Dh/Dt$  trends (-1 to -4 m/yr) (Figure 9C), however, cannot be explained by modelled elevation change due to only SMB and firn.

### Basal melt rates

We computed basal melt rates from surface  $Dh/Dt$  and pole-base  $Dh_{pb}/Dt$  using Equations 5 and 7, respectively. The resulting values range from ~2-4 m/yr at PIG2 to ~40-43 m/yr at SOW3, with good agreement between the rates obtained by surface and pole-base elevation change (Table 1).

**Page 12: [13] Deleted** **David Shean** **7/1/17 5:59:00 PM**

The three upstream stations (SOW1, SOW2 and BOAR) show lower melt rates of ~9–14 m/yr, while the downstream stations near the transverse depression (SOW3 and SOW4) have higher rates of ~29–43 m/yr for the same time period.

**Page 14: [14] Deleted** **David Shean** **7/1/17 5:59:00 PM**

We also expect local SMB and firn column thickness variability within the array. The greater  $z_{rh}$  surface reflector heights (a proxy for surface accumulation) at SOW3 indicate that enhanced local accumulation occurred within the transverse depression (Figure 2), likely due to wind-blown snow. However, we note that all of the remaining 2012–

2014 stations display similar  $z_{rh}$  reflector height evolution, as do the two 2008–2010 stations (Figure 9A). Future RACMO and IMAU-FDM products with improved resolution should provide additional constraints on spatial variability.

### Pole-base settling

**Some of the reflector height increase and/or observed negative Dh/Dt could be related to settling of the poles within the firn.**

3.1

**Page 16: [15] Deleted** **David Shean** **7/1/17 5:59:00 PM**

These measurements provide some validation for the GPS results, but a direct comparison may be imprudent.

The Stanton et al. (2013)

**Page 18: [16] Deleted** **David Shean** **7/1/17 5:59:00 PM**

Observed surface Dh

**Page 24: [17] Deleted Cells** **David Shean** **7/1/17 5:59:00 PM**

Deleted Cells

**Page 24: [18] Deleted Cells** **David Shean** **7/1/17 5:59:00 PM**

Deleted Cells

**Page 24: [19] Deleted Cells** **David Shean** **7/1/17 5:59:00 PM**

Deleted Cells

**Page 24: [20] Deleted Cells** **David Shean** **7/1/17 5:59:00 PM**

Deleted Cells

**Page 24: [21] Deleted Cells** **David Shean** **7/1/17 5:59:00 PM**

Deleted Cells

**Page 24: [22] Deleted Cells** **David Shean** **7/1/17 5:59:00 PM**

Deleted Cells

**Page 24: [23] Deleted** **David Shean** **7/1/17 5:59:00 PM**

5

**Page 24: [23] Deleted** **David Shean** **7/1/17 5:59:00 PM**

5

**Page 24: [24] Deleted Cells** **David Shean** **7/1/17 5:59:00 PM**

Deleted Cells

**Page 24: [25] Deleted Cells** **David Shean** **7/1/17 5:59:00 PM**

Deleted Cells

**Page 24: [26] Deleted** **David Shean** **7/1/17 5:59:00 PM**

6

<b>Page 24: [26] Deleted</b>	<b>David Shean</b>	<b>7/1/17 5:59:00 PM</b>
------------------------------	--------------------	--------------------------

6

<b>Page 24: [27] Deleted</b>	<b>David Shean</b>	<b>7/1/17 5:59:00 PM</b>
------------------------------	--------------------	--------------------------

9

<b>Page 24: [27] Deleted</b>	<b>David Shean</b>	<b>7/1/17 5:59:00 PM</b>
------------------------------	--------------------	--------------------------

9

<b>Page 24: [28] Deleted</b>	<b>David Shean</b>	<b>7/1/17 5:59:00 PM</b>
------------------------------	--------------------	--------------------------

4

<b>Page 24: [28] Deleted</b>	<b>David Shean</b>	<b>7/1/17 5:59:00 PM</b>
------------------------------	--------------------	--------------------------

4

<b>Page 24: [29] Deleted</b>	<b>David Shean</b>	<b>7/1/17 5:59:00 PM</b>
------------------------------	--------------------	--------------------------

5

<b>Page 24: [29] Deleted</b>	<b>David Shean</b>	<b>7/1/17 5:59:00 PM</b>
------------------------------	--------------------	--------------------------

5

<b>Page 24: [30] Deleted</b>	<b>David Shean</b>	<b>7/1/17 5:59:00 PM</b>
------------------------------	--------------------	--------------------------

6

<b>Page 24: [30] Deleted</b>	<b>David Shean</b>	<b>7/1/17 5:59:00 PM</b>
------------------------------	--------------------	--------------------------

6

<b>Page 24: [31] Deleted Cells</b>	<b>David Shean</b>	<b>7/1/17 5:59:00 PM</b>
------------------------------------	--------------------	--------------------------

Deleted Cells

<b>Page 24: [32] Deleted Cells</b>	<b>David Shean</b>	<b>7/1/17 5:59:00 PM</b>
------------------------------------	--------------------	--------------------------

Deleted Cells

<b>Page 24: [33] Deleted Cells</b>	<b>David Shean</b>	<b>7/1/17 5:59:00 PM</b>
------------------------------------	--------------------	--------------------------

Deleted Cells

<b>Page 24: [34] Deleted Cells</b>	<b>David Shean</b>	<b>7/1/17 5:59:00 PM</b>
------------------------------------	--------------------	--------------------------

Deleted Cells

<b>Page 24: [35] Deleted</b>	<b>David Shean</b>	<b>7/1/17 5:59:00 PM</b>
------------------------------	--------------------	--------------------------

5

<b>Page 24: [35] Deleted</b>	<b>David Shean</b>	<b>7/1/17 5:59:00 PM</b>
------------------------------	--------------------	--------------------------

5

<b>Page 24: [36] Deleted</b>	<b>David Shean</b>	<b>7/1/17 5:59:00 PM</b>
------------------------------	--------------------	--------------------------

and Dh/Dt results.

<b>Page 24: [36] Deleted</b>	<b>David Shean</b>	<b>7/1/17 5:59:00 PM</b>
------------------------------	--------------------	--------------------------

and Dh/Dt results.

<b>Page 24: [36] Deleted</b>	<b>David Shean</b>	<b>7/1/17 5:59:00 PM</b>
------------------------------	--------------------	--------------------------

and Dh/Dt results.

Page 24: [36] Deleted David Shean 7/1/17 5:59:00 PM

and Dh/Dt results.

Page 24: [36] Deleted David Shean 7/1/17 5:59:00 PM

and Dh/Dt results.

Page 24: [36] Deleted David Shean 7/1/17 5:59:00 PM

and Dh/Dt results.

Page 24: [36] Deleted David Shean 7/1/17 5:59:00 PM

and Dh/Dt results.

Page 24: [37] Deleted David Shean 7/1/17 5:59:00 PM

for both the surface Dh/Dt and corrected pole-base  $Dh_{pb}/Dt$  (

Page 24: [37] Deleted David Shean 7/1/17 5:59:00 PM

for both the surface Dh/Dt and corrected pole-base  $Dh_{pb}/Dt$  (

Page 24: [37] Deleted David Shean 7/1/17 5:59:00 PM

for both the surface Dh/Dt and corrected pole-base  $Dh_{pb}/Dt$  (

Page 24: [37] Deleted David Shean 7/1/17 5:59:00 PM

for both the surface Dh/Dt and corrected pole-base  $Dh_{pb}/Dt$  (

Page 29: [38] Deleted David Shean 7/1/17 5:59:00 PM

: A) Observed cumulative displacement between SOW1 and other 2012–2014 stations. Legend lists initial distances. B) Observed cumulative strain, with best fit strain rate listed in legend. C) Smoothed residuals from linear fit, highlighting subtle variations and inflections in strain rates. D) Daily GPS velocity. Note timing of abrupt absolute velocity changes observed at all sites and inflection points in observed strain across the array.

