

1 **Grounding and Calving Cycle of Mertz Ice Tongue**

2 **Revealed by Shallow Mertz Bank**

3 Xianwei Wang^{1,2}, David M. Holland^{1,3}, Xiao Cheng^{2,5} and Peng Gong^{4,5}

4 1. Center for Global Sea Level Change, New York University Abu Dhabi. Abu Dhabi, United Arab Emirates.

5 2. State Key Laboratory of Remote Sensing Science, and College of Global Change and Earth System Science,
6 Beijing Normal University. Beijing 100875, China.

7 3. Courant Institute of Mathematical Sciences, New York University. New York 10012, United States of America.

8 4. Ministry of Education Key Laboratory for Earth System Modeling, and Center for Earth System Science,
9 Tsinghua University, Beijing, China 100084.

10 5. Joint Centre for Global Change Studies, Beijing, China.

11
12 *Correspondence to: Xianwei Wang: wangxianwei0304@163.com or Xiao Cheng: xcheng@bnu.edu.cn*

13 **Abstract**

14 A recent study, using remote sensing, provided evidence that a seafloor shoal influenced
15 the 2010 calving event of the Mertz Ice Tongue (MIT), by partially grounding the MIT several
16 years earlier. In this paper, we start by proposing a method to calculate Firn Air Content (FAC)
17 around Mertz from seafloor-touching icebergs. Our calculations indicate the FAC around Mertz
18 region as 4.87 ± 1.31 m. We then design an indirect method of using freeboard and sea surface
19 height data extracted from ICESat/GLAS, FAC, and relatively accurate seafloor topography to
20 detect grounding sections of the MIT between 2002 and 2008 and analyze the process of
21 grounding prior to the calving event. By synthesizing remote sensing data, we point out that the
22 grounding position was localized northeast of the Mertz ice front close to the Mertz Bank. The
23 grounding outlines of the tongue caused by the Mertz Bank are extracted as well. From 2002 to
24 2008, the grounding area increased and the grounding became more pronounced. Additionally,
25 the ice tongue could not effectively climb over the Mertz Bank in following the upstream ice
26 flow direction and that is why MIT rotated clockwise after late 2002. Furthermore, we
27 demonstrate that the area-increasing trend of the MIT changed little after calving ($\sim 36 \text{ km}^2/\text{a}$),
28 thus allowing us to use remote sensing to estimate the elapsed time until the MIT can reground
29 on and be bent by the shoal. This period is approximately 70 years. Our observations suggest that
30 the calving of the MIT is a cyclical process controlled by the presence of the shallow Mertz Bank
31 location and the flow rate of the tongue. This calving cycle also explains the cyclic variations in
32 sea-surface conditions around the Mertz detected by earlier studies.

33 **Keywords:** Mertz Ice Tongue, firn air content, grounding, Mertz Bank, calving cycle.

34 **1. Introduction**

35 Surface-warming induced calving or disintegration of floating ice has occurred in
36 Antarctica, such as the Larsen B ice shelf (Scambos et al., 2000, 2003; Domack et al., 2005;
37 Shepherd et al., 2003). While surface or sub-surface melting has largely been recognized to
38 contribute to floating ice loss in Antarctica (Depoorter et al., 2013), calving caused by interaction
39 with the seafloor has not been widely considered. The Mertz Ice Tongue (MIT) was reported to
40 have calved in 2010, subsequent to being rammed by a large iceberg, B-9B (Legresy et al. 2010).
41 After the calving, the areal coverage of Mertz polynya, sea ice production and dense, shelf water
42 formation in the region changed (Kusahara et al. 2011; Tamura et al. 2012). However, the
43 iceberg collision may have only been an apparent cause of the calving as other factors had not
44 been fully considered such as seafloor interactions (Massom et al., 2015; Wang. 2014). By
45 comparing inverted ice thickness to surrounding bathymetry, and combining remote sensing
46 analysis, Massom et al., (2015) considered that the seabed contact may have held the glacier
47 tongue in place to delay calving by ~8 years. The interaction of the MIT with seafloor, the exact
48 grounding location of the MIT before calving and the extent of grounding are still not well-
49 known.

50 The MIT (66°S-68°S, 144°E-150°E, Fig. 1), located in King George V Land, East
51 Antarctica, extended over 140 km from its grounding line to the tongue front and is
52 approximately 30 km wide at the front (Legresy et al., 2004). Much field exploration has been
53 conducted around Mertz and the increasing availability over the last decade of remote sensing,
54 hydrographic surveying, and bathymetric data allows to investigate the mechanism of the ice
55 tongue instability and calving. From satellite altimetry, a modest elevation change rate of 0.03
56 m/a (Pritchard et al., 2012) and a freeboard change rate of -0.06 m/a (Wang et al., 2014) were

57 found, which implied that the combined effects of surface accumulation and basal melt were not
58 dramatic for this ice tongue. Investigations of tidal effects, surface velocity, rift propagation, and
59 ice front propagation (Berthier et al., 2003; Frezzotti et al., 1998; Legresy et al., 2004;
60 Lescarmonier et al., 2012; Massom et al., 2010, 2015) have been conducted with an objective of
61 detecting underlying factors affecting the stability of the MIT. Grounding has been suggested as
62 a potential mechanism to affect the stability of the MIT by delaying calving (Massom et al.
63 2015). However, without highly accurate bathymetric data, it is impossible to carry out such a
64 study. Fortunately, In 2010, a new and high resolution bathymetry model, with a resolution of
65 100 m was released for the Terra Adelie and George V continental margin (Beaman et al., 2011),
66 and it has later been used to generate Bedmap-2 (Fretwell et al., 2013). This accurate data set
67 (Fig 3) provides an opportunity for better exploring seafloor shoals and their impacts on the
68 instability of the MIT. In this study, we focus on grounding events of the MIT from 2002 to 2008.
69 A method for grounding detection is proposed and grounding of the MIT before the calving is
70 investigated. A calving cycle of the MIT caused by grounding on seafloor shoal, Mertz Bank is
71 discussed as well.

72 **2. Data**

73 The primary data used to investigate grounding of the MIT in this study are elevation
74 data from Geoscience Laser Altimeter System (GLAS) onboard the Ice, Cloud and land
75 Elevation Satellite (ICESat) and the seafloor bathymetry data mentioned above. In this section,
76 the ICESat/GLAS and bathymetry data, as well as some preprocessing are introduced.

77 **2.1 ICESat/GLAS**

78 ICESat is the first spaceborne laser altimetry satellite orbiting the Earth, launched by the
79 National Aeronautics and Space Administration (NASA) in 2003 (Zwally et al. 2002) with

80 GLAS as the primary payload onboard. ICESat/GLAS was operated in an orbit of ~600 km and
81 had a geographical coverage from 86° S to 86° N. ICESat/GLAS usually observed in nadir
82 viewing geometry and employed laser pulses of both 532 nm and 1064 nm to measure the
83 distance from the sensor to ground (Zwally et al. 2002). On the ground, ICESat/GLAS's
84 footprint covered an area of approximately 70 m in diameter, with adjacent footprints spaced by
85 ~170 m. The horizontal location accuracy of the footprint was approximately 6 m (Abshire et al.
86 2005). The accuracy and precision of ICESat/GLAS altimetry data were 14 cm and 2 cm
87 respectively (Shuman et al. 2006). ICESat/GLAS usually made two or three campaigns a year
88 from 2003 to the end of 2009, each campaign lasted for approximately one month. 15 different
89 types of data were produced for various scientific applications, named as GLA01, GLA02, ...
90 GLA15. In this study, GLA12 data (elevation data for polar ice sheet) covering Mertz from
91 release 33 between 2003 and 2009 is used (Fig. 2).

92 **2.2 Seafloor Topography**

93 Detailed bathymetry maps are fundamental spatial data for marine science studies
94 (Beaman et al., 2003, 2011) and crucially needed in the data-sparse Antarctic coastal region
95 (Massom et al. 2015). Regionally, around Mertz, a large archive of ship track single-beam and
96 multi-beam bathymetry data from 2000 to 2008 were used to generate a high resolution Digital
97 Elevation Model (DEM) for which the spatial coverage can be found in Figs. 3(b) and 3(c). The
98 DEM product was reported to have a vertical accuracy of approximately 11.5 m (500 m depth)
99 and a horizontal accuracy of 70 m (500 m depth) in the poorest situation (Beaman et al. 2011).
100 As can be seen from Figs. 3(b) and 3(c), there is no bathymetry data under the MIT, which may
101 result in large uncertainty for seafloor interpolation. The oldest bathymetry data collected along
102 the margin of the MIT was from 2000 (Beaman et al. 2011). Additionally, around the Mertz ice

103 front, for both the east and west flanks, bathymetry data does exist. Since the ice front has a
104 width of ~34 km (Wang et al. 2014), the accuracy of seafloor DEM under the MIT varies
105 depending on distance to margin. Inside the 2000 boundary of the MIT, the closer to the dash-
106 dotted polygon (Figs. 6 and 7), the better accuracy the seafloor DEM. Outside of that boundary,
107 the quality of the seafloor DEM data is much better because of high density of single-beam or
108 multi-beam bathymetric measurements.

109 Around Antarctica, the seafloor topography data from Bedmap-2 was produced by
110 Fretwell et al. (2013) which adopted the DEM from Beaman et al. (2011). In this study,
111 Bedmap-2 seafloor topography data covering Mertz is employed to detect the contact between
112 seafloor and the MIT. Because of inconsistent elevation systems for ICESat/GLAS and the
113 seafloor topography data, the Earth Gravitational Model 2008 (EGM08) geoid (Pavlis et al. 2012)
114 with respect to World Geodetic System 1984 (WGS-84) ellipsoid is taken as reference. Since the
115 seafloor topography from Bedmap-2 is referenced to the so-called g104c geoid, an elevation
116 transformation is required and can be implemented through:

$$117 \quad E_{sf} = E_{seafloor} + gl04c_{to_wgs84} - EGM2008 \quad (1)$$

118 where E_{sf} and $E_{seafloor}$ is the seafloor topography under the EGM08 and g104c geoid
119 respectively, $gl04c_{to_wgs84}$ is the value needed to convert height relative to the g104c geoid to
120 that under the WGS-84, and $EGM2008$ is the geoid undulation with respect to the WGS-84.

121 **3. Methods**

122 **3.1 Grounding Detection Methods**

123 ICESat/GLAS data has been widely used to determine ice freeboard, or ice thickness,
124 since its launch in 2003 (Kwok et al., 2007; Wang et al., 2011, 2014; Yi et al., 2011; Zwally et
125 al., 2002, 2008). The methods we designed for grounding detection of the MIT using the

126 ICESat/GLAS data are introduced here. First, assuming a floating MIT, based on freeboard data
127 extracted in different observation dates, ice draft of the MIT is inverted. Next, ice bottom
128 elevation is calculated based on the inverted ice draft and the lowest sea-surface height. Finally,
129 the ice bottom is compared with seafloor bathymetry to detect ice grounding. The underlying
130 logic for grounding detection is that if the inverted ice bottom is lower than seafloor, we can
131 draw a conclusion that the ice tongue is grounding rather than floating.

132 The method for extracting a freeboard map using ICESat/GLAS from multiple campaigns
133 over the MIT was described in Wang et al. (2014). Without providing details, here we only
134 introduce it schematically. Four steps are included in freeboard map production for each of the
135 datasets from November 14, 2002, March 8, 2004, December 27, 2006 and January 31, 2008.

136 The first step involves data preprocessing, saturation correction, data quality control, and
137 tidal correction removal. The magnitude of the ICESat/GLAS waveform can become saturated
138 because of different gain setting, or high reflection from natural surfaces. Thus, saturated
139 waveforms with *i_satElevCorr* (i.e. an attribute from GLA12 data record) greater than or equal
140 to 0.50 m are ignored and only those measurements with *i_satElevCorr* less than 0.50 m are
141 corrected following the procedures in Wang et al. (2012, 2013). Additionally, measurements
142 with *i_reflctUC* greater than or equal to one are ignored. Furthermore, the tidal correction from
143 TPX07.1 tide model in GLA12 data record is removed to obtain estimates for the instantaneous
144 sea surface height. Finally, elevation data from ICESat/GLAS related to the WGS-84 ellipsoid
145 and EGM 08 geoid from 2003 to 2009 is available for subsequent use.

146 The second step is to derive sea-surface height according to each track and to calculate
147 freeboard for each campaign. Because of tidal variations near the MIT, surface elevations of the
148 MIT can vary as well. To derive sea-surface height from ICESat/GLAS and provide a reference

149 for freeboard calculation for different campaigns, the ICESat/GLAS data over the MIT within a
150 buffer region (with 10 km as buffer radius of MIT boundary in 2007) are selected and sea-
151 surface height is determined as the lowest elevation measurement along each track (Wang et al.
152 2014). Freeboard is then calculated by subtracting the corresponding sea-surface height from
153 elevation measurements of the MIT according to different tracks from the same campaign. Thus
154 freeboard data for different campaigns from 2003 to 2009 is obtained.

155 The third step is to relocate footprints using estimated ice velocity. ICESat observed the
156 MIT almost repeatedly along different tracks in different campaigns (Fig. 2). However,
157 observations from only one campaign cannot provide good coverage of the MIT. All
158 observations from 2003 to 2009 are combined together to produce a freeboard map of the MIT.
159 Fig. 2 shows the spatial coverage of ICESat/GLAS from 2003 to 2009 over Mertz, but the
160 geometric relation between tracks is not correct over the MIT because the tongue was fast
161 moving and observed in different years by ICESat. Regions observed in an earlier campaign
162 would move downstream later (Wang et al. 2014). For example, consider ICESat data from track
163 T31 from March 22, 2003 and T165 (Fig. 2) from November 1, 2003 respectively. Fig. 2 shows
164 that the distance between track T165 and T31 is ~ 7.5 km without accounting for ice advection
165 between observation dates. However because of the fast moving ice tongue, the distance of their
166 actual ground tracks on surface of the MIT should be longer because T165 was located upstream
167 and observed later. Thus footprints relocation using ice velocity is critical to obtain accurate
168 geometric relations among different tracks. The ice velocity data from Rignot et al. (2011)
169 generated from InSAR data from 2006 to 2010 is used to relocate the footprints of ICESat/GLAS.
170 The correct geospatial relations between observations from different campaigns can be achieved
171 on November 14, 2002, March 8, 2004, December 27, 2006, and January 31, 2008, through:

172
$$X = x + \sum_{i=1}^n v_{xi}\Delta t + v_{xm}t_m \quad (2)$$

173
$$Y = y + \sum_{i=1}^n v_{yi}\Delta t + v_{ym}t_m \quad (t_m = t_2 - t_1 - n\Delta t) \quad (3)$$

174 where x and y are the horizontal positions directly from the ICESat measurements, and X
 175 and Y are the horizontal positions after relocation respectively; v_x and v_y are the horizontal
 176 components of the ice velocities; t_1 and t_2 are the start and end times; Δt is the time interval and
 177 n indicates the largest integer time steps for time interval between t_1 and t_2 ; t_m is the residual
 178 time; In this work, Δt is set as 10 days; v_{xi} and v_{yi} is derived from ice velocity field according
 179 to different locations during relocation and may change in different time intervals.

180 Freeboard changes with time should be considered as well, but it is neglected because
 181 comparison of freeboard from crossing tracks showed a slightly decreasing trend of -0.06 m/a on
 182 average (Wang et al. 2014). The spatial distribution of freeboard data over the MIT for
 183 November 14, 2002, is shown in Fig. 5(a).

184 The forth step is to interpolate the freeboard map using the relocated freeboard data from
 185 the third step. Kriging interpolation in ArcGIS is selected in this study to produce freeboard
 186 maps of the MIT because it can provide an optimal interpolation estimate for a given coordinate
 187 location by considering the spatial relationships of a data set. With this method, freeboard maps
 188 of the MIT are produced for November 14, 2002, March 8, 2004, December 27, 2006, and
 189 January 31, 2008 respectively when the ice tongue outline can be delineated from Landsat
 190 images.

191 Ice draft is calculated with Eq. (4) assuming hydrostatic equilibrium and using the lowest
 192 sea-surface height E_{sea_level} as reference for the sea surface elevation.

193
$$\rho_w D = \rho_i (H_f + D - FAC) \quad (4)$$

194 where D is the ice draft, i.e. vertical distance from the sea surface to the bottom of the ice;
 195 H_f is the freeboard, i.e. the vertical distance from the sea surface to the top of the snow; ρ_w and
 196 ρ_i are the densities of ocean water and ice, respectively. In this study, the ice and sea water
 197 density are taken as 915 kg/m^3 and 1024 kg/m^3 , respectively (Wang et al., 2014); FAC is the firm
 198 air content which corresponds to the decrease in thickness (in meters) that occurs when the firm
 199 column is compressed to the density of glacier ice, as defined in Holland et al., (2011) and
 200 Ligtenberg et al. (2014).

201 The sea surface is taken as the lowest sea surface height (E_{sea_level}) and is derived from
 202 the minimum of all sea surface heights from the different ICESat/GLAS tracks between 2003
 203 and 2009 and amounts in our case to -3.35 m . For time varying sea-surface heights caused by
 204 tides, the minimum sea-surface height can allow ice with a given draft to ground to the seafloor.
 205 Then, the ice bottom elevation is calculated by considering the ice draft and the lowest sea-
 206 surface height. Elevation difference of the ice bottom and the seafloor is calculated. A negative
 207 value indicates that the ice bottom is lower than the seafloor, which suggests grounding.

208 The elevation of the underside (bottom) of the tongue E_{ice_bottom} is calculated from:

$$209 \quad E_{ice_bottom} = E_{sea_level} - D \quad (5)$$

210 Similarly, the elevation difference of ice tongue bottom and seafloor is defined as E_{dif} ,

211 which can be calculated by:

$$212 \quad E_{dif} = E_{ice_bottom} - E_{sf} \quad (6)$$

213 where E_{sf} is the seafloor elevation as defined in Eq. (1).

214 **3.2. Firm Air Content Estimation Method**

215 The Antarctic ice sheet is covered by a dry, thick firm layer which represents an
 216 intermediate stage between fresh snow and glacial ice, having varying density from Antarctic

217 inland to the coast (van den Broeke, 2008). The density and depth of the Antarctic firn layer has
218 been modeled (e.g., van den Broeke, 2008) using a combination of regional climate model output
219 and a steady-state firn compaction model. However, for ice thickness inversion, Firn Air Content
220 (FAC) is usually used to make the calculation convenient (Rignot and Jacobs, 2002). FAC is
221 defined as the decrease in thickness (in meters) that occurs when the firn column is compressed
222 to the density of glacier ice (Holland et al., 2011). Time-dependent FAC has also been modeled
223 by considering the physical process of the firn layer (e.g., Ligtenberg et al. 2014). For the MIT,
224 there are some in-situ measurements of snow thickness available from Massom et al. (2010) who
225 used a snow layer depth of 1 m to derive the thickness of surrounding multi-year, fast sea ice.
226 However on the surface of the MIT, no in-situ measurements of density or depth of firn layer are
227 available.

228 Because of different density and thickness of the firn layer on the top of an ice tongue, it
229 is challenging to simulate the density profile of the MIT without in-situ measurements as control
230 points. In this study, we use FAC extracted from adjacent seafloor-touching icebergs rather than
231 that from modeling to investigate the grounding of the MIT. The MIT may be composed of pure
232 ice, water, air, firn or snow that will influence the density of the ice tongue. However, if
233 assuming a pure ice density only to calculate ice mass, the thickness of MIT must be corrected
234 by the FAC. The FAC can be inferred from surrounding icebergs that are slightly grounded
235 under the assumption of hydrostatic equilibrium and known ice draft and freeboard. It is,
236 however, critical to target and use icebergs that fulfil the condition of slight grounding. From
237 Smith (2011), icebergs can be divided into three categories based on bathymetry and seasonal
238 pack ice distributions: grounded, constrained, and free-drifting icebergs. Without pack ice, an
239 iceberg can be free-drifting or grounded. Free-drifting icebergs can move several tens of

240 kilometers a day, such as iceberg A-52 (Smith et al. 2007). Grounded icebergs can be heavily or
241 lightly anchored. Heavily grounded icebergs have firm contact with the seafloor and can be kept
242 stationary for a long time, such as iceberg B-9B (Massom. 2003). However, slightly grounded
243 icebergs may have less contact with the seafloor and can possibly move slowly under the
244 influence of ocean tide, ocean currents, or winds, but much slower than free-drifting icebergs.
245 The relation of grounded iceberg to the drifting velocity is not well-known. However, slowly
246 drifting or nearly stationary icebergs in open water are good indicators for slight grounding and
247 therefore are used to infer FAC.

248 Because of the heavily grounded iceberg B-9B to the east of the MIT blocking the
249 drifting of pack ice or icebergs from the east, icebergs located between B-9B and the MIT are
250 most likely generated from the Mertz or Ninnis glaciers. Some icebergs may be slightly
251 grounded as can be detected from remote sensing. We calculate the FAC from these slightly
252 grounded icebergs and later apply it to grounding event detection of the MIT. Around the MIT,
253 the locations of three icebergs ('A', 'B' and 'C') were investigated using MODIS and Landsat
254 images in austral summer of 2006 and 2008 respectively and shown in Fig. 4. Fortunately,
255 ICESat/GLAS observed these icebergs on February 23, 2006 (54th day of 2006) and February 18,
256 2008 (49th day of 2008) which allows us to analyze the behavior of these icebergs three-
257 dimensionally. Fig. 4a shows that icebergs 'A', 'B' and 'C' were almost stagnant and only
258 slightly changed their positions and orientation over two months (from 28 to 85 day of 2006).
259 Thus we can consider these icebergs slightly grounded. For these slightly grounded icebergs,
260 hydrostatic equilibrium should still apply, so the ice draft inverted from freeboard measurement
261 assuming hydrostatic equilibrium should be equal to the water depth. Based on this analysis, we
262 can take water depth as the draft to calculate the FAC.

263 Because only icebergs ‘A’ and ‘C’ were observed by track T1289 of ICESat/GLAS in
264 2006, the FAC is inverted using freeboard and water depth from bathymetry from both
265 icebergs(Figs. 3b, 3c, 4, and Table 1). However, the icebergs were not stationary, which
266 indicates that only some parts were slightly grounded. Therefore, only the top two largest
267 freeboard measurements of icebergs ‘A’ and ‘C’ from T1289 in 2006 are used to calculate the
268 FAC with Eq. (7) with a least-squares method under hydrostatic equilibrium.

$$269 \quad FAC = H_{f_k} + D_k - \frac{\rho_w}{\rho_i} D_k + \varepsilon_k \quad (7)$$

270 where k refers to the icebergs ‘A’ or ‘C’, H_f is the top two largest freeboard measurement of
271 each iceberg, D is the ice draft which is the same as sea water depth and is taken from the
272 seafloor bathymetry directly, ε is the residual of FAC.

273 Table 1 shows the freeboard of iceberg ‘A’ and ‘C’ from 2006 and seafloor bathymetry
274 for FAC inversion and grounding detection of icebergs ‘A’ and ‘B’ in 2008 (detailed freeboard
275 values for these icebergs can be found from S-Fig. 1). With the freeboard from 2006 and seafloor
276 bathymetry (Table 1), the FAC is calculated as 4.87 ± 1.31 m. Icebergs ‘A’ and ‘B’ were observed
277 by the same track T1289 on February 18, 2008 and thus are taken to evaluate the grounding
278 detection by using the inverted FAC. From iceberg trajectories observed by remote sensing (Fig.
279 4b), we know, iceberg ‘A’ drifted away from its original position. Thus it was not grounded.
280 However, iceberg ‘B’ were kept rotating in this period without drifting away, indicating a slight
281 grounding. Such grounding status determined from remote sensing can also be detected with our
282 method since the elevation difference of the ice bottom and seafloor from Table 1 does clearly
283 indicate a slightly grounded iceberg ‘B’ and a floating iceberg ‘A’. Thus, our FAC estimation
284 works well around Mertz.

285 FAC varies across the Antarctica ice sheet, usually decreasing from the interior to the
286 coast. For Mertz we obtain a FAC of 4.87 ± 1.31 m. Other studies, using a time variable approach,
287 modelled FAC values between 5 and 10 m (Ligtenberg et al. 2014) and in the absence of in-situ
288 measurements our estimate seems consistent, but there are some shortcomings which should be
289 further explored.

290 First, for FAC calculation, icebergs just touching the seafloor should be used in which
291 case the FAC calculated assuming hydrostatic equilibrium is the same as its actual value.
292 However, it is difficult to ascertain whether an iceberg is just touching the seafloor from remote
293 sensing images. The near stationary or slowly rotating icebergs detected with remote sensing
294 may be grounded more than just touching the seafloor, which may result in a inverted FAC
295 theoretically greater than its actual value. Thus, using this FAC value to detect grounding can
296 potentially lead to smaller grounding results. However, once a grounded iceberg or ice tongue is
297 detected using this FAC, the result is more convincing.

298 Second, limited observations from ICESat/GLAS may not catch the same and the thickest
299 section of a slight grounding iceberg. Because ICESat/GLAS observed only several times a year
300 on repeat tracks and icebergs were rotating slowly, the elevation profile in 2006 and 2008 along
301 the same track T1289 may not refer to the same ground surface. S-Fig. 1 shows the freeboard of
302 icebergs 'A', 'B' and 'C' derived from ICESat/GLAS from 2006 and 2008 respectively. By
303 comparing the freeboard of iceberg 'A' in 2006 (S-Fig. 1a), and 2008 (S-Fig. 1c), we find the
304 larger freeboard and the longer freeboard profile in 2006. Comparatively, the smaller freeboard
305 in 2008 may be caused by basal melting or observing a different portion of iceberg 'A' by
306 ICESat. Since the larger freeboard measured in 2006 indicates a high possibility of capturing the
307 thickest portion, it is reasonable to use it to invert the FAC. Additionally, icebergs 'A' and 'C'

308 did show a similar maximum freeboard (Table 1), which is another important reason to select the
 309 measurements of 2006 for the inversion.

310 **4. Accuracy of Grounding Detection**

311 The accuracy of E_{dif} is critical to grounding detection of the MIT. From Eq. (1) to (6),
 312 we find different components of error sources, such as from sea surface height determination, ice
 313 draft, seafloor bathymetry, and elevation transformation. Meanwhile, the uncertainty of ice draft
 314 is primarily depending on that of freeboard and FAC . Furthermore, the uncertainty of freeboard
 315 is influenced by the footprint relocation and freeboard changing rates. Considering all that
 316 mentioned above, the error sources of elevation difference E_{dif} can be synthesized by:

$$317 \Delta E_{dif} = \Delta E_{sl} + a(\Delta H_f + \Delta E_{re} + \Delta E_{fb_c} + \Delta FAC + \Delta E_{krig}) + \Delta E_{sf} + \Delta E_{trans} \quad (8)$$

318 where $a = \frac{\rho_i}{\rho_w - \rho_i}$; Δ stands for error of each variable; ΔE_{dif} stands for the error of the final
 319 elevation difference of ice bottom and seafloor; ΔE_{sl} , ΔH_f , ΔE_{re} , ΔE_{fb_c} , ΔFAC , ΔE_{sf} , ΔE_{krig} ,
 320 and ΔE_{trans} stand for errors caused by the sea surface height extraction, freeboard extraction,
 321 freeboard relocation, freeboard changing rates, FAC calculation, seafloor bathymetry, kriging
 322 interpolation and elevation system transformation, respectively.

323 The influence of elevation system transformation on final elevation difference can be
 324 neglected. Based on the error propagation, the uncertainty of elevation difference E_{dif} can be
 325 described by:

$$326 \varepsilon E_{dif} = \sqrt{(\varepsilon E_{sl})^2 + a^2[(\varepsilon H_f)^2 + (\varepsilon E_{re})^2 + (\varepsilon E_{fb_c})^2 + (\varepsilon FAC)^2 + (\varepsilon E_{krig})^2] + (\varepsilon E_{sf})^2} \quad (9)$$

327 where ε indicates the uncertainty of each parameter.

328 **4.1 Uncertainty of Kriging Interpolation**

329 Fig. 5a shows the spatial distribution of freeboard data over the MIT used for grounding
330 detection from November 14, 2002. The spatial difference of the ICESat/GLAS data between
331 Fig. 2 and Fig. 5 is caused by the footprint relocation, after which the spatial geometry between
332 different tracks is reasonably correct. In the lower right of the Mertz ice front (Fig. 5a), the
333 crossing-track distance between T1289 and T165 is approximately 7 km. In these data gaps, the
334 freeboard data used for grounding detection is interpolated using kriging. Thus, knowing the
335 uncertainty of kriging interpolation is critical to the final grounding detection.

336 To investigate the uncertainty of kriging interpolation method, freeboard measurements
337 from ICESat/GLAS should be compared with the interpolated freeboard estimates. A testing
338 region with freeboard measurements is selected (dashed blue square in Fig. 5a, 7 km×7 km in
339 size). A freeboard map is first interpolated with the gray dots (Fig. 5a) using kriging. The
340 freeboard measurements (284 of green dots in Fig. 5a) are then compared with the interpolation
341 from the square. The spatial distribution and the histogram of freeboard difference derived by
342 subtracting the krigged freeboard from the freeboard derived from ICESat/GLAS are shown in
343 Fig. 5b.

344 The freeboard measurement varies from 31.6 m to 40.0 m with an average of 36.6 m.
345 However, the interpolated freeboard varies from 32.9 m to 39.6 m with an average of 35.9 m.
346 From the freeboard difference (Fig. 5b), we find that the interpolated freeboard shows similar
347 results compared with the freeboard derived from ICESat/GLAS. The interpolated freeboard has
348 an accuracy of -0.7 ± 1.8 m indicating that the interpolated freeboard using kriging can reflect the
349 actual freeboard well.

350 **4.2 Grounding Detection Robustness**

351 Since the sea surface height is extracted from the ICESat/GLAS data track by track, we
352 use ± 0.15 m (Zwally et al. 2002) as the uncertainty of elevation data (εE_{st}). Also from Wang et
353 al. (2014), we can find that the uncertainty of freeboard extraction (εH_f) was ± 0.50 m. From
354 Rignot et al. (2011), the error of the ice velocity ranged from 5 m/a to 17 m/a. Assuming that the
355 ice velocity varied by 17 m/a (an upper threshold), the relocation error horizontally could reach
356 ± 54 m when considering a three-year period. Wang et al. (2014) extracted the average slope of
357 the MIT along the ice flow direction as 0.00024. However, because of large crevasses on the
358 surface, we use 50 times of this value as a conservative estimate of the average slope. In this way,
359 we can estimate εE_{re} as ± 0.65 m when considering a three-year period. The annual rate of
360 freeboard changes from 2003 to 2009 was -0.06 m/a (Wang et al. 2014). Therefore, we consider
361 the freeboard stable over this period. However when combining data from different time periods,
362 εE_{fb_c} is estimated to be ± 0.18 m. From Beaman et al. (2011), considering the elevation
363 uncertainty at the worst situation when water depth reaches 500 m, εE_{g104c} is ± 1.5 m. Using Eq.
364 (9) and kriging interpolation, from the analysis from Section 4.1, 1.8 m is taken as the
365 uncertainty. Using all these errors above, we calculate the final uncertainty of the elevation
366 difference as ± 23 m.

367 From the calculations above, a less than -23 m E_{dif} indicates a robust grounding event.
368 However, if E_{dif} is greater than 23 m, grounding cannot be confirmed. E_{dif} between -23 m and
369 23 m corresponds to slight grounding or floating. We can also determine different contributions
370 of each separate factor to the overall accuracy. Seafloor bathymetry contributes the greatest part
371 and is the dominant factor affecting the accuracy of grounding detection.

372 **5. Grounding Detection Results**

373 The spatial distribution of the elevation difference E_{dif} and the outlines of the MIT from
374 2002 to 2008 are shown in Fig. 6. Since the moving trajectory of the Mertz ice front changed by
375 more than 40 degrees clockwise (Massom et al. 2015; Wang. 2014), a buffer region with radius
376 of 2 km (region between black and grey lines in Fig. 6) is introduced to investigate grounding
377 potential of the MIT. The freeboard in the buffer region is extrapolated using the kriging
378 interpolation method and the elevation difference is calculated. The elevation difference less than
379 46 m (twice the uncertainty of the elevation difference εE_{dif}) both inside and outside the outline
380 is extracted and the statistics are shown in Table 2. Since the uncertainty to determine a
381 grounding event is ± 23 m, if some grids of the MIT have elevation difference E_{dif} less than -23
382 m, we can conclude that this section of the tongue is strongly grounded. The smaller the E_{dif} , the
383 more robust the grounding.

384 As illustrated in Table 2 and Fig 6, the minimum E_{dif} inside the MIT in 2002 was 11.9 m
385 and the minimum E_{dif} inside the MIT was less than -23 m after 2002. The minimum E_{dif} in the
386 buffer region were all less than -23 m from 2002 to 2008. This suggests that the MIT had
387 grounded on the shallow Mertz Bank at least since November 14, 2002. This result coincides
388 with the findings from Massom et al. (2015) who considered that the northwestern extremity of
389 the MIT started to touch a seafloor shoal in late 2002 to early 2003. Also, it would have been
390 difficult for the MIT to approach the buffer region (indicated with yellow to red colors in Fig. 6)
391 as the surrounding Mertz Bank gets shallower and steeper, suggesting substantive grounding
392 potentials. Inside the MIT, the minimum E_{dif} was just 11.9 m on November 14, 2002, which
393 indicates slight grounding. However on March 8, 2004, December 27, 2006, and January 31,
394 2008, the minimum E_{dif} reached -46.0 m, -52.3 m and -34.8m respectively, which indicates that
395 strong grounding occurred in some regions. From 2002 to 2008, more regions under the MIT had

396 E_{dif} less than 46 m, the area of which increased from 8 km² to 17 km². Additionally, the mean
397 of those E_{dif} less than 46 m gradually decreased from 28.8 m to 12.3m, according to which we
398 can conclude that the ice front became more firmly grounded as time passed on. Since the
399 grounding area increased from 8 km² to 17 km² (Table 2) and the mean of E_{dif} decreased, we
400 conclude that during the period from 2002 to 2008, the grounding of the northwest tip of the MIT
401 became more widespread.

402 Based on the calculated elevation difference, the grounding outlines of the MIT are
403 delineated for November 14, 2002, March 8, 2004, December 27, 2006 and January 31, 2008
404 respectively (Fig. 7). For the grounded part of the outlines in different years, the starting and
405 ending location and the perimeter are also extracted (Table 3), from which we conclude that the
406 length of the grounding outline on the Mertz Bank was only limited to a few kilometers. We find
407 that the lower right (northwest) section of the MIT was always grounded and grounding did not
408 occur in other regions (Fig. 6). The shallowest seafloor that the Mertz ice front touched was ~ -
409 290 m in November 2002. In 2004, 2006, and 2008, the lower right (northwest) of the MIT even
410 approached the contour of -220 m.

411 **6. Discussion**

412 **6.1 Area Changing Rate and ~70-year Calving Cycle of MIT**

413 Using Landsat TM/ETM+ images from 1989 to 2013, outlines of the MIT are extracted
414 manually. Assuming a fixed grounding line position, the area of the MIT over this period is
415 calculated. Using these data, from 1989 to 2007, an increasing are-change trend of the MIT was
416 obtained (from 5453 km² to 6126 km²) in Fig. 8. However, the area of the MIT was almost
417 constant from 2007 to 2010, before calving. The largest area of the MIT was 6113 km² closest to
418 the calving event in 2010. After the calving, the area decreased to 3617 km² in November 2010.

419 The average area-change trend of the MIT from 1989 to 2007 was also obtained using a
420 least-squares method, corresponding to 35.3 km²/a. However, after the calving a slightly higher
421 area-change trend of 36.9 km²/a, was found (Fig. 8). On average, the area-change trend of the
422 MIT was approximately 36 km²/a.

423 The surface dynamics of the MIT such as ice flow direction changes and middle rift
424 changes caused by grounding was analyzed by Massom et al. (2015). In the history of the MIT,
425 one or two large calving events were suspected to have happened between 1912 and 1956
426 (Frezzotti et al., 1998). Based on the interactions between the MIT and Mertz Bank suggested by
427 our observations and description below, it is likely that only one large calving event occurred
428 between 1912 and 1956. When the MIT touched Mertz Bank, the bank started to affect its
429 stability by bending it clockwise to the east, as can be found from velocity changes from
430 Massom et al. (2015). With continuous advection of the ice and flux input from the upstream, a
431 large rift from the west flank of the tongue would ultimately have to occur and could potentially
432 calve the MIT. A sudden length shortening of the MIT can be caused by such ice tongue calving
433 as indeed had happened in February, 2010. We also consider that even without a sudden collision
434 of iceberg B-9B in 2010, the MIT would eventually have calved because of the effect of the
435 shallow Mertz Bank.

436 When considering 6127 km² as the maximum area of the MIT and assuming a constant
437 area-changing trend of 36.9 km²/a after 2010, the MIT will take approximately 68 years to calve
438 again. When assuming an area changing trend of 35.3 km²/a as before 2010, the MIT will take a
439 little longer, approximately 71 years to calve. Therefore, without considering an accidental
440 collision with other large icebergs, the MIT is predicted to calve again in ~70 years. Because of

441 continuous advection of the ice from upstream and the fixed location of the shallow Mertz Bank,
442 the calving is likely repeatable and a cycle therefore exists.

443 After the MIT calved in February, 2010, the Mertz polynya size, sea-ice production, sea-
444 ice coverage and high-salinity shelf water formation changed as well. A sea-ice production
445 decrease of approximately 14-20% was found by Tamura et al. (2012) using satellite data and the
446 high-salinity shelf water export was reported to reduce up to 23% using a state-of-the-art ice-
447 ocean model (Kusahara et al. 2010). Recently, Campagne et al. (2015) pointed out a ~70-year
448 cycle of surface ocean condition and high-salinity shelf water production around the Mertz
449 through analyzing some reconstructed sea ice and ocean data over the last 250 years. They also
450 mentioned that this cycle was closely related to the presence and activity of the Mertz polynya.
451 However, the reason for this cycle was not fully understood.

452 From these findings addressed above and the MIT calving cycle we find that the calving
453 cycle of the MIT leads to the ~70-year cycle of surface ocean condition and high-salinity shelf
454 water production around the Mertz. Variations in length of the MIT will prevent sea ice drifting
455 from the east side to a variable degree. A long MIT contributes to maintain a large polynya
456 because sea ice from the east side cannot drift to the west side. The sea ice produced on the west
457 side is blown seaward by the katabatic wind and thereby maintains a polynya and stable sea ice
458 production. The sudden shortening of the MIT after a calving event therefore reduces the size of
459 Mertz Polynya formed by Antarctic katabatic winds, resulting in a lower sea-ice production and
460 further lessens high-salinity shelf water production. Therefore, the cycle of ocean conditions
461 around the Mertz found by Campagne et al. (2015) is likely dominated by the calving of the MIT.
462 Additionally, the 70-year cycle of the MIT calving coincides well with the change of surface
463 ocean condition around the Mertz which makes the explanation much more compelling.

464 **6.2 Seafloor DEM**

465 High accuracy seafloor is critical to the final success of the grounding detection.
466 According to our best knowledge, Beaman et al. (2011) provided the most accurate seafloor
467 DEM over the Mertz, so the seafloor DEM inside the dash-dotted polygon (Fig. 7) was kept and
468 the grounding detection was conducted there (Fig. 6). Additionally, the ice tongue continued to
469 advance out into the ocean, where the bathymetry observation density is good. From the results
470 shown in Fig. 6 all grounding sections of the MIT boundary were located outside of the 2000
471 boundary. Thus the analysis of the grounding detection near the ice front in 2002, 2004, 2006,
472 and 2008 is convincing. Inside the 2000 boundary, most of the grounding detection results were
473 above 100 m, indicating a floating status of the corresponding ice. Only abnormal seafloor
474 features higher than this seafloor DEM by more than 100 m could result in wide grounding
475 inside. Actually, no matter whether the MIT inside the 2000 boundary was grounded or not,
476 gradual grounding on the shallow Mertz Bank of the MIT since late 2002 is well supported by
477 observations and which we take as evidence to infer the primary cause of the instability of the
478 MIT.

479 **6.3 Influence of Mertz Bank on MIT**

480 Fig. 7 shows the extension line of the west flank in November, 2002, from which we can
481 find that if the MIT advected along the former direction, the ice flow would be seriously
482 obstructed when approaching the Mertz Bank. The shallowest region of the Mertz Bank has an
483 elevation of approximately -140 m and the MIT would have to climb the 140 m obstacle to cross
484 it. The shallow Mertz Bank would have caused strong grounding during the climbing. This
485 special feature of the seafloor shoal facing the MIT can further explain why the ice velocity
486 differed along the east and west flanks of the MIT before calving and why the ice tongue was

487 deflected clockwise to the east, as suggested by Massom et al. (2015). However, because of
488 sparsely-distributed bathymetric data in the Mertz region used in Massom et al. (2015), this
489 effect could not be easily seen. Here, from our grounding detection results and surrounding high-
490 accuracy bathymetry data, this effect is more clearly observed.

491 **7. Conclusion**

492 In this study, a method of FAC calculation from seafloor-touching icebergs around the
493 Mertz region is presented as an important element in understanding the MIT grounding. The
494 FAC around the Mertz is 4.87 ± 1.31 m. This FAC is used to calculate ice draft based on the sea
495 surface height and freeboard extracted from ICESat/GLAS and is performing well. A method to
496 extract the grounding sections of the MIT is described based on comparison of the inverted ice
497 draft assuming hydrostatic equilibrium with the seafloor bathymetry. The final grounding results
498 explain the dynamic behavior of the MIT. Previous work by Massom et al. (2015) has also
499 provided some evidence for seafloor interaction, in showing that the MIT front had an
500 approximate 280 m draft with the nearby seafloor as shallow as 285 m, suggesting the possibility
501 of grounding. In our work, we have provided ample detailed bathymetry and ice draft
502 calculations. Specifically, the ice bottom elevation of the MIT is inverted using the
503 ICESat/GLAS data and compared with seafloor bathymetry from 2002, 2004, 2006, and 2008
504 respectively. From these calculations we show conclusively that the MIT was indeed grounded
505 along a specific portion of its northwest tip over a limited region. We also point out that even
506 without collision by iceberg B-9B in early 2010 the ice tongue would eventually have calved
507 because of the ice advection from the upstream and the glacier flow increasingly diverted by the
508 obstructing seafloor shoal of the Mertz Bank.

509 From remote sensing images we are able to quantify the trend of area increase of the MIT
510 before and after the 2010 calving. While the area-increase trend of the MIT after calving was
511 slightly greater than that before, we use the averaged trend to estimate a timescale required for
512 the MIT to re-advance to the area of the shoaling bathymetry from its retreated, calved position.
513 Our estimate is ~70-years, which is remarkably consistent with Campagne et al. (2015) who
514 found a similar period for variations in sea surface conditions using seafloor sediment data. Thus,
515 the shoaling on the Mertz Bank combined with the rate of advancing of the MIT determines the
516 70-year repeat cycle. Also the calving cycle of the MIT explains the observed cycle of the sea-
517 surface conditions change well, which indicates that the calving of the MIT is the dominant
518 factor for the sea-surface condition change. Understanding the mechanism underlying the
519 periodicity of the MIT calving is important as the presence or absence of the MIT has a profound
520 impact on the sea ice and hence of the bottom water formation in the local region.

521 **Acknowledgements**

522 This research was supported by the Center for Global Sea Level Change (CSLC) of NYU
523 Abu Dhabi (Grant: G1204), the Open Fund of State Key Laboratory of Remote Sensing Science
524 (Grant: OFSLRSS201414), the National Natural Science Foundation of China (Grant: 41176163),
525 the China Postdoctoral Science Foundation (Grant: 2012M520185, 2013T60077) and
526 Fundamental Research Fund for the Central University. We are grateful to the Chinese Arctic
527 and Antarctic Administration, the European Space Agency for free data supply under project
528 C1F.18243, the National Snow and Ice Data Center (NSIDC) for the availability of the
529 ICESat/GLAS data (<http://nsidc.org/data/order/icesat-glas-subsetter>) and MODIS image archive
530 over the Mertz glacier (http://nsidc.org/cgi-bin/modis_iceshelf_archive.pl), British Antarctica
531 Survey for providing Bedmap-2 seafloor topography data

532 (<https://secure.antarctica.ac.uk/data/bedmap2/>), the National Geospatial-Intelligence Agency for
533 publicly released EGM2008 GIS data ([http://earth-](http://earth-info.nga.mil/GandG/wgs84/gravitymod/egm2008/egm08_gis.html)
534 [info.nga.mil/GandG/wgs84/gravitymod/egm2008/egm08_gis.html](http://earth-info.nga.mil/GandG/wgs84/gravitymod/egm2008/egm08_gis.html)), and the USGS for Landsat
535 data (<http://glovis.usgs.gov/>). Fruitful discussions with M. Depoorter, P. Morin, T. Scambos and
536 R. Warner, and constructive suggestions from Editor Andreas Vieli and two anonymous
537 reviewers are acknowledged.

538 **References**

- 539 1. Abshire, J. B., Sun, X., Riris, H., Sirota, J. M., McGarry, J. F., Palm, S., ... & Liiva, P.
540 (2005). Geoscience laser altimeter system (GLAS) on the ICESat mission: on - orbit
541 measurement performance. *Geophysical Research Letters*, 32(21).
- 542 2. Beaman, R. J., & Harris, P. T. (2003). Seafloor morphology and acoustic facies of the
543 George V Land shelf. *Deep Sea Research Part II: Topical Studies in Oceanography*,
544 50(8), 1343-1355.
- 545 3. Beaman, R. J., O'Brien, P. E., Post, A. L., & De Santis, L. (2011). A new high-resolution
546 bathymetry model for the Terre Adélie and George V continental margin, East Antarctica.
547 *Antarctic Science*, 23(01), 95-103.
- 548 4. Berthier, E., Raup, B., & Scambos, T. (2003). New velocity map and mass-balance
549 estimate of Mertz Glacier, East Antarctica, derived from Landsat sequential imagery.
550 *Journal of Glaciology*, 49(167), 503-511.
- 551 5. Campagne, P., Crosta, X., Houssais, M. N., Swingedouw, D., Schmidt, S., Martin, A., ...
552 & Massé G. (2015). Glacial ice and atmospheric forcing on the Mertz Glacier Polynya
553 over the past 250 years. *Nature Communications*, 6.

- 554 6. Depoorter, M. A., Bamber, J. L., Griggs, J. A., Lenaerts, J. T. M., Ligtenberg, S. R. M.,
555 van den Broeke, M. R., & Moholdt, G. (2013). Calving fluxes and basal melt rates of
556 Antarctic ice shelves. *Nature*, 502(7469), 89-92.
- 557 7. Domack, E., Duran, D., Leventer, A., Ishman, S., Doane, S., McCallum, S., ... & Prentice,
558 M. (2005). Stability of the Larsen B ice shelf on the Antarctic Peninsula during the
559 Holocene epoch. *Nature*, 436(7051), 681-685.
- 560 8. Fretwell, P., Pritchard, H. D., Vaughan, D. G., Bamber, J. L., Barrand, N. E., Bell, R., ...
561 & Fujita, S. (2013). Bedmap2: improved ice bed, surface and thickness datasets for
562 Antarctica. *Cryosphere*, 7(1).
- 563 9. Frezzotti, M., Cimbelli, A., & Ferrigno, J. G. (1998). Ice-front change and iceberg
564 behaviour along Oates and George V Coasts, Antarctica, 1912-96. *Annals of Glaciology*,
565 27, 643-650.
- 566 10. Holland, P. R., Corr, H. F., Pritchard, H. D., Vaughan, D. G., Arthern, R. J., Jenkins, A.,
567 & Tedesco, M. (2011). The air content of Larsen ice shelf. *Geophysical Research Letters*,
568 38(10).
- 569 11. Kusahara, K., Hasumi, H. & Williams, G. D. (2011), Impact of the Mertz Glacier Tongue
570 calving on dense water formation and export. *Nature communications*, 2, 159.
- 571 12. Kwok, R. Cunningham, G. F., Zwally, H. J., & Yi, D. (2007). Ice, Cloud, and land
572 Elevation Satellite (ICESat) over Arctic sea ice: retrieval of freeboard. *Journal of*
573 *Geophysical Research*, 112, C12013, doi:10.1029/2006JC003978.
- 574 13. Legresy, B., Wendt, A., Tabacco, I. E., Remy, F., & Dietrich, R. (2004). Influence of
575 tides and tidal current on Mertz Glacier, Antarctica. *Journal of Glaciology*, 50(170), 427-
576 435.


- 577 14. Legresy, B., N. Young, L. Lescarmonier, R. Coleman, R. Massom, B. Giles, A. Fraser, R.
578 Warener, B. Galton-Fenzi, L. Testut, M. Houssais and G. Masse (2010), CRAC!!! in the
579 Mertz Glacier, Antarctica.
580 [http://www.antarctica.gov.au/__data/assets/pdf_file/0004/22549/ml_402353967939815_](http://www.antarctica.gov.au/__data/assets/pdf_file/0004/22549/ml_402353967939815_mertz_final_100226.pdf)
581 [mertz_final_100226.pdf](http://www.antarctica.gov.au/__data/assets/pdf_file/0004/22549/ml_402353967939815_mertz_final_100226.pdf)
- 582 15. Lescarmonier, L., Legrésy, B., Coleman, R., Perosanz, F., Mayet, C., & Testut, L. (2012).
583 Vibrations of Mertz glacier ice tongue, East Antarctica. *Journal of Glaciology*, 58(210),
584 665-676.
- 585 16. Ligtenberg, S., Kuipers Munneke, P., & Van Den Broeke, M. R. (2014). Present and
586 future variations in Antarctic firn air content. *The Cryosphere*, 8(5), 1711-1723.
- 587 17. Massom, R. A. (2003). Recent iceberg calving events in the Ninnis Glacier region, East
588 Antarctica. *Antarctic Science*, 15(02), 303-313.
- 589 18. Massom, R. A., Giles, A. B., Fricker, H. A., Warner, R. C., Legrésy, B., Hyland, G.,
590 Young, N., & Fraser, A. D. (2010). Examining the interaction between multi-year
591 landfast sea ice and the Mertz Glacier Tongue, East Antarctica: Another factor in ice
592 sheet stability? *Journal of Geophysical Research*, 115, C12027,
593 doi:10.1029/2009JC006083.
- 594 19. Massom, R. A., Giles, A. B., Warner, R. C., Fricker, H. A., Legrésy, B., Hyland, G., ... &
595 Young, N. (2015). External influences on the Mertz Glacier Tongue (East Antarctica) in
596 the decade leading up to its calving in 2010. *Journal of Geophysical Research: Earth*
597 *Surface*, 120(3), 490-506.

- 598 20. Pavlis, N. K., Holmes S. A., Kenyon, S. C., & Factor, J. K. (2012). The development and
599 evaluation of the Earth Gravitational Model 2008 (EGM2008), *Journal of Geophysical*
600 *Research*. 117, B04406, doi:10.1029/2011JB008916.
- 601 21. Pritchard, H. D., Ligtenberg, S. R. M., Fricker, H. A., Vaughan, D. G., Van den Broeke,
602 M. R., & Padman, L. (2012). Antarctic ice-sheet loss driven by basal melting of ice
603 shelves. *Nature*, 484(7395), 502-505.
- 604 22. Rignot, E., Mouginot, J. & Scheuchl, B. (2011), Ice flow of the Antarctic ice sheet.
605 *Science*, 333(6048), 1427-1430.
- 606 23. Rignot, E., & Jacobs, S. S. (2002). Rapid bottom melting widespread near Antarctic ice
607 sheet grounding lines. *Science*, 296(5575), 2020-2023.
- 608 24. Scambos, T. Hulbe, A., C. & Fahnestock, M. A. (2003). Climate-induced ice shelf
609 disintegration in the Antarctic Peninsula. *Antarctic Research Series*, 79, 79-92.
- 610 25. Scambos, T. Hulbe, A., C. Fahnestock, M. A. & Bohlander, J. (2000). The link between
611 climate warming and breakup of ice shelves in the Antarctic Peninsula. *Journal of*
612 *Glaciology*, 46(154), 516-530.
- 613 26. Shepherd, A., Wingham, D., Payne, T., & Skvarca, P. (2003). Larsen Ice Shelf has
614 progressively thinned. *Science*, 302(5646), 856-859.
- 615 27. Shuman, C. A., Zwally, H. J., Schutz, B. E., Brenner, A. C., DiMarzio, J. P., Suchdeo, V.
616 P., & Fricker, H. A. (2006). ICESat Antarctic elevation data: Preliminary precision and
617 accuracy assessment. *Geophysical Research Letters*, 33(7).
- 618 28. Smith, K. L., Robison, B. H., Helly, J. J., Kaufmann, R. S., Ruhl, H. A., Shaw, T. J., ... &
619 Vernet, M. (2007). Free-drifting icebergs: hot spots of chemical and biological
620 enrichment in the Weddell Sea. *Science*, 317(5837), 478-482.

- 621 29. Smith, K. L. (2011). Free-drifting icebergs in the Southern Ocean: an overview. *Deep Sea*
622 *Research Part II: Topical Studies in Oceanography*, 58(11), 1277-1284.
- 623 30. Tamura, T., Williams, G. D., Fraser, A. D. & Ohshima, K. I. (2012). Potential regime
624 shift in decreased sea ice production after the Mertz Glacier calving, *Nature*
625 *communications*, 3, 826.
- 626 31. Van den Broeke, M. (2008). Depth and density of the Antarctic firn layer. *Arctic,*
627 *Antarctic, and Alpine Research*, 40(2), 432-438.
- 628 32. Wang, X.W., Cheng, X., Gong, P., Huang, H. B., Li Z., & Li, X. W. (2011). Earth
629 Science Applications of ICESat/GLAS: a Review. *International Journal of Remote*
630 *Sensing*, 32, 23, 8837-8864, doi: 10.1080/01431161.2010.547533
- 631 33. Wang, X.W., Cheng, X., Gong, P., Shum, C. K., Holland, D.M., & Li, X.W. (2014).
632 Freeboard and mass extraction of the disintegrated Mertz Ice Tongue with remote sensing
633 and altimetry data. *Remote Sensing of Environment*, 144, 1-10.
- 634 34. Wang, X.W. (2014). Mertz ice tongue evolutions from satellite observed data,
635 Postdoctoral Research Report, College of Global Change and Earth System Science,
636 Beijing Normal University, China. doi: 10.13140/2.1.1006.1603
- 637 35. Wang, X., Cheng, X., Li, Z., Huang, H., Niu, Z., Li, X., & Gong, P. (2012). Lake water
638 footprint identification from time-series ICESat/GLAS data. *IEEE Geoscience and*
639 *Remote Sensing Letters*, 9(3), 333-337.
- 640 36. Wang, X., Gong, P., Zhao, Y., Xu, Y., Cheng, X., Niu, Z., ... & Li, X. (2013). Water-
641 level changes in China's large lakes determined from ICESat/GLAS data. *Remote Sensing*
642 *of Environment*, 132, 131-144.


- 643 37. Yi, D., Zwally, H.J., & Robbins, J. (2011). ICESat observations of seasonal and
644 interannual variations of sea-ice freeboard and estimated thickness in the Weddell Sea,
645 Antarctica (2003-2009). *Annals of Glaciology*, 52(57), 43-51.
- 646 38. Zwally, H. J., Schutz, B., Abdalati, W., Abshire, J., Bentley, C., Brenner, A., Buftona, J.,
647 Deziouf, J., Hancocka, D., Hardinga, D., Herringg, T., Minsterh, B., Quinng, K., Palmi,
648 S., Spinhirnea, J., & Thomasj, R. (2002). ICESat's laser measurements of polar ice,
649 atmosphere, ocean, and land. *Journal of Geodynamics*, 34, 405-445.
- 650 39. Zwally, H. J., Yi, D., Kwok, R., & Zhao, Y. (2008). ICESat measurements of sea ice
651 freeboard and estimates of sea ice thickness in the Weddell Sea. *Journal of Geophysical*
652 *Research*, 113, C02S15, doi:10.1029/2007JC004284.
- 653

Figures



655


656 **Figure 1.** Mertz Ice Tongue (MIT), East Antarctica. Landfast sea ice is attached to the east flank
 657 of the MIT and the Mertz Polynya is to the west. The background image corresponds to band 4
 658 Landsat 7, captured on February 2, 2003. The green square found in the upper left inset indicates
 659 the location of the MIT in East Antarctica. A polar stereographic projection with -71°S as
 660 standard latitude is used.



662

663 **Figure 2.** Spatial distribution of the ICESat/GLAS data from 2003 to 2009 covering the Mertz
 664 region. Ground tracks of ICESat/GLAS are indicated with gray lines. Track 1289 (T1289) is
 665 highlighted in red as is used in Fig. 4. The background image corresponds to band 4 Landsat 7,
 666 captured on February 2, 2003. A polar stereographic projection with -71°S as standard latitude is
 667 used.

668




669

670

671

(a)




672

673

674

(b)



(c)

675


676

677

678 **Figure 3.** (a) Seafloor topography from bathymetry around Mertz and the outlines of the MIT
 679 from 2002 to 2008 marked with the colored polygons for different years. The shallow Mertz
 680 Bank is located in the lower right (northeast). The yellow dash-dotted line indicates the shape of
 681 the MIT from January 25, 2000, which is used to identify the bathymetry gap under the ice
 682 tongue. The dashed red inset box corresponds to the location of Figs. 6 and 7. (b) : multi-beam
 683 bathymetry dataset coverage over the Mertz region. The embedded figure in the upper left is the
 684 zoom in of the dashed red rectangle which shows the positions of icebergs ‘A’ and ‘B’ (polygon
 685 filled in red) on February 19, 2008 (Fig. 4b). (c): single-beam bathymetry dataset coverage over
 686 the Mertz region. The light blue polylines show the contours around the Mertz Bank and the
 687 black dots are bathymetric measurement profiles. Both (b) and (c) are redrawn from Beaman et

688 al. (2011) because the original spatial coverage of the single and multi-beam bathymetry data is
689 not available. However, for being able to use the Figures from Beaman et al. (2011), we geo-
690 registered it and put the contour around the Mertz Bank and the location of icebergs used in the
691 text over it which illustrates the density of the bathymetry measurements. Through comparing
692 the grounding lines from (b) and (c), we can conclude that the geo-registration is successful as
693 the grounding line we obtained from the National Snow and Ice Data Center (NSIDC) coincides
694 with that from Beaman et al. (2011) well in most parts. This Figure is under a projection of polar
695 stereographic projection with -71°S as standard latitude.


696



697

698 **Figure 4.** Freeboard extracted from Track T1289, ICESat/GLAS, the location of which can be
 699 found from Fig. 2 and Fig. 3(b). (a) and (b) show the freeboard extracted from the ICESat/GLAS
 700 date from February 23, 2006 (2006054) and February 18, 2008 (2008049) respectively. In each
 701 image, the positions of three icebergs (with name labeled as ‘A’, ‘B’ and ‘C’) closest to the
 702 ICESat/GLAS observation date are plotted with green, red and blue polygons respectively. The
 703 observation dates of remote sensing images are indicated with seven numbers (yyyyddd) in the
 704 legend. ‘yyyyddd’ stands for day ‘ddd’ in year ‘yyyy’. ‘MODIS02’ and ‘LE7’ indicate that the
 705 images used to extract outlines of the icebergs are from MODIS and Landsat 7 ETM+,
 706 respectively.

707





709

710



711 **Figure 5.** Evaluation of kriging interpolation method over the MIT using freeboard data derived
 712 from the ICESat/GLAS data. (a) shows profile location of freeboard derived from the
 713 ICESat/GLAS data after relocation over the MIT. The gray dots indicate the ICESat/GLAS data
 714 used for interpolation using kriging method. The blue dashed square indicates the 7 km x 7 km
 715 region used to investigate the accuracy of kriging interpolation method. Inside the square, the
 716 freeboard data marked with green dots are used to check the accuracy of the freeboard
 717 interpolated with kriging. (b) is the freeboard comparison result derived by subtracting the
 718 kriged freeboard from the freeboard derived from the ICESat/GLAS. The spatial distribution
 719 and the histogram of the freeboard difference are shown in the lower left and upper right
 720 respectively. The black polygon filled with light blue shows the boundary of the MIT on
 721 November 14, 2002.

722



(a)

(b)



(c)

(d)

723

724

725

726


727 **Figure 6.** Elevation difference of Mertz ice bottom and seafloor topography. (a), (b), (c) and (d)

728 correspond to the elevation difference from November 14, 2002 , March 8, 2004, December 27,

729 2006, and January 31, 2008, respectively assuming hydrostatic equilibrium under the minimum


730 sea surface height -3.35 m . The contours at an interval of 20 m in the lower right indicate the

731 seafloor topography of the Mertz Bank. The solid black line indicates the boundary of the MIT
732 and the thick gray line outlines a buffer region of the boundary with 2 km as buffer radius. The
733 dash-dotted line indicates the shape of the MIT on January 25, 2000, which is used to identify
734 the bathymetry gap under the MIT. In the legend, the negative values mean that the ice bottom is
735 lower than the seafloor, which of course is impossible. Therefore, the initial assumption of a
736 floating ice tongue was incorrect in those locations (yellow to red colors), and the ice was
737 grounded. Regions with more negative values indicate heavier grounding inside the MIT or
738 grounding potential in the buffer region. Please note that no bathymetric data was available
739 under most of the ice tongue and for locations of the bathymetric data, please refer to Figs 3b and
740 3c.



741

742 **Figure 7.** Digital Elevation Map (DEM) of seafloor around the Mertz and grounding section of
 743 the boundaries extracted from 2002 to 2008. The grounding sections of the MIT boundary from
 744 2002, 2004, 2006 and 2008 are marked with thick red, purple, green and blue polylines
 745 respectively and the MIT boundaries are indicated with polygons with the same legend as in Fig.
 746 3a. Additionally, the MIT boundary from 2000 indicated with dash-dotted yellow polygon is
 747 used to show the different quality of the seafloor DEM. Inside this polygon no bathymetry data
 748 was collected or used. The dashed red line indicates the ‘extension line’ of the west flank of the
 749 MIT on November 14, 2002, passing the shallowest region of the Mertz Bank (approximately -
 750 140 m).



751

752 **Figure 8.** Average trend of the area change of the MIT. The area of the MIT is extracted from
 753 the Landsat images from 1988 to 2013.

754

755

Tables

756 **Table 1.** Statistics of icebergs used to invert FAC with a least-square method and validation of
 757 grounding iceberg detection using this FAC. Icebergs ‘A’, ‘B’ and ‘C’ are the same as what are
 758 used in Fig. 4 and S-Fig 1. The measurements from icebergs ‘A’ and ‘C’ in February, 2006 are
 759 used to derive the FAC with a least-squares method. However, the measurements from Icebergs
 760 ‘A’ and ‘B’ in 2008 are used for validation.

Icebergs	date	Latitude	Longitude	Freeboard	Seafloor	Sea Surface Height	ε	E_{aif}
		($^{\circ}$)	($^{\circ}$)	(m)	(m)	(m)	(m)	(m)
A	Feb 23, 2006	-67.1737	146.6595	66.88	-528.48	-1.92	0.89	
		-67.1752	146.6604	66.34	-527.01	-1.92	1.30	
C	Feb 23, 2006	-67.1085	146.6247	66.37	-505.84	-1.92	-1.25	
		-67.1100	146.6255	66.28	-507.08	-1.92	-1.01	
A	Feb 18, 2008	-67.1194	146.6303	58.88	-522.52	-2.08		69.14
		-67.1209	146.6311	59.58	-524.16	-2.08		64.88
B	Feb 18, 2008	-67.0906	146.6151	67.22	-500.92	-2.08		-22.45
		-67.0921	146.6159	66.10	-500.47	-2.08		-13.55

761

762 **Table 2.** Statistics of grounding grids inside the MIT or grounding potentials outside of the MIT
763 ('I': inside the thick black line, Fig. 6; Number in brackets indicates how many grids are located
764 inside the 2000 Mertz boundary; 'O': between the black and gray lines, Fig. 6) from November
765 14, 2002, March 8, 2004, December 27, 2006 and January 31, 2008 respectively. Each grid
766 covers an area of 1 km². The Mean, Minimum and Standard deviation is calculated without
767 considering those fallen inside the 2000 Mertz boundary and only include those out of the 2000
768 Mertz boundary with an elevation difference less than 46 m.

769

Elevation difference (subtracting seafloor from ice bottom)	2002-11-14		2004-03-08		2006-12-27		2008-01-31	
	I	O	I	O	I	O	I	O
23-46 (m)	9(3)	10(0)	6(0)	3(0)	10(1)	1(0)	10(3)	5(0)
0-23 (m)	2(0)	6(0)	1(0)	1(0)	9(0)	2(0)	4(0)	2(0)
<0 (m)	0(0)	8(0)	2(0)	5(0)	7(0)	21(0)	6(0)	18(0)
Mean (m)	28.8	9.8	15.8	-1.1	10.9	-41.9	12.3	-31.0
Minimum (m)	11.9	-81.5	-46.0	-44.5	-52.3	-102.8	-34.8	-103.0
Standard deviation (m)	9.2	36.8	29.6	31.4	24.7	37.6	27.3	38.0
Number of grids	8	24	9	9	25	24	17	25

770

771 **Table 3.** Statistics of grounding outlines of the MIT as shown with thick polylines in Fig. 7 from
 772 November 14, 2002, March 8, 2004, December 27, 2006 and January 31, 2008 respectively

	2002-11-14	2004-03-08	2006-12-27	2008-01-31
Start location (°)	146.124 °E, 66.696 °S	146.155 °E, 66.681 °S	146.093 °E, 66.700 °S	146.088 °E, 66.699 °S
End location (°)	146.240 °E, 66.693 °S	146.256 °E, 66.683 °S	146.304 °E, 66.669 °S	146.292 °E, 66.668 °S
Perimeter (km)	7.0	6.4	24.7	20.9

773